

COMPACT COMPOSITION OPERATORS WITH NON-LINEAR SYMBOLS ON THE H^2 SPACE OF DIRICHLET SERIES

FRÉDÉRIC BAYART AND OLE FREDRIK BREVIG

ABSTRACT. We investigate the compactness of composition operators on the Hardy space of Dirichlet series induced by a map $\varphi(s) = c_0 s + \varphi_0(s)$, where φ_0 is a Dirichlet polynomial. Our results depend heavily on the characteristic c_0 of φ and, when $c_0 = 0$, on both the degree of φ_0 and its local behaviour near a boundary point. We also study the approximation numbers for some of these operators. Our methods involve geometric estimates of Carleson measures and tools from differential geometry.

1. INTRODUCTION

A theorem of Gordon and Hedenmalm [9] describes the bounded composition operators on the Hilbert space \mathcal{H}^2 of Dirichlet series,

$$f(s) = \sum_{n=1}^{\infty} a_n n^{-s},$$

with square summable coefficients endowed with the norm $\|f\|_{\mathcal{H}^2}^2 := \sum_{n=1}^{\infty} |a_n|^2$. We let \mathbb{C}_θ denote the half-plane of complex numbers $s = \sigma + it$ with $\sigma > \theta$. The Dirichlet series in \mathcal{H}^2 represent analytic functions in $\mathbb{C}_{1/2}$ and a mapping φ of $\mathbb{C}_{1/2}$ into itself defines a function $\mathcal{C}_\varphi(f) := f \circ \varphi$ on $\mathbb{C}_{1/2}$, if $f \in \mathcal{H}^2$. The operator $\mathcal{C}_\varphi : \mathcal{H}^2 \rightarrow \mathcal{H}^2$ is well-defined and bounded if and only if φ is a member of the following class:

Definition. The *Gordon–Hedenmalm class*, denoted \mathcal{G} , is the set of functions $\varphi : \mathbb{C}_{1/2} \rightarrow \mathbb{C}_{1/2}$ of the form

$$(1) \quad \varphi(s) = c_0 s + \sum_{n=1}^{\infty} c_n n^{-s} =: c_0 s + \varphi_0(s),$$

where c_0 is a non-negative integer called the *characteristic* of φ , the Dirichlet series φ_0 converges uniformly in \mathbb{C}_ε ($\varepsilon > 0$) and has the following mapping properties:

- (a) If $c_0 = 0$, then $\varphi_0(\mathbb{C}_0) \subset \mathbb{C}_{1/2}$.
- (b) If $c_0 \geq 1$, then either $\varphi_0 \equiv 0$ or $\varphi_0(\mathbb{C}_0) \subset \mathbb{C}_0$.

Since the paper of Gordon and Hedenmalm, several authors have studied the properties of composition operators acting on \mathcal{H}^2 or on similar spaces of Dirichlet series (see for instance

Date: October 23, 2015.

2010 Mathematics Subject Classification. Primary 47B33. Secondary 30B50, 30H10.

The second author is supported by Grant 227768 of the Research Council of Norway.

[1, 2, 7, 8, 11]). In the present work, we are interested in the study of the compactness of \mathcal{C}_φ when φ is a polynomial symbol, say

$$(2) \quad \varphi(s) = c_0 s + c_1 + \sum_{n=2}^N c_n n^{-s},$$

and we implicitly assume that $\varphi \in \mathcal{G}$. The symbol φ is said to have *unrestricted range* if

$$\inf_{s \in \mathbb{C}_0} \operatorname{Re}(\varphi(s)) = \begin{cases} 1/2 & \text{if } c_0 = 0, \\ 0 & \text{if } c_0 \geq 1. \end{cases}$$

Correspondingly, if $\varphi(\mathbb{C}_0)$ is strictly contained in any smaller half-plane, we say that \mathcal{C}_φ has *restricted range*. It is well-known that the composition operator \mathcal{C}_φ is compact when φ has restricted range [1, Thm. 21]. In what follows, we will assume that φ has unrestricted range.

Definition. A set of integers $\Lambda \subseteq \mathbb{N} - \{1\}$ is called \mathbb{Q} -*independent* if the set $\{\log n : n \in \Lambda\}$ is linearly independent over \mathbb{Q} .

Symbols of the form (2) have been extensively studied in the *linear case*,

$$(3) \quad \varphi(s) = c_0 s + c_1 + \sum_{j=1}^d c_{q_j} q_j^{-s},$$

where the set $\{q_j\}$ is \mathbb{Q} -independent and $c_{q_j} \neq 0$. When $c_0 \geq 1$, it is proven in [2] that the operator \mathcal{C}_φ is compact if and only if φ has restricted range. Our first result extends this to the case of an arbitrary polynomial:

Theorem 1. *Let φ be a Dirichlet polynomial of the form (2) with $c_0 \geq 1$. Then \mathcal{C}_φ is compact if and only if φ has restricted range.*

As is to be expected when investigating composition operators on \mathcal{H}^2 , the symbols with $c_0 = 0$ are more difficult to handle and require different techniques. In this case, it is proven independently in [2] and [8] that composition operators induced by linear symbols (3) with $c_0 = 0$ are compact if and only if φ has restricted range or $d \geq 2$.

The main effort of this paper is dedicated to extending this result to general polynomials. We rely crucially on a geometric description of such compact composition operators found in [11] (see Lemma 5 below). Our second result is:

Theorem 2. *Suppose that $\{q_j\}_{j=1}^d$ are \mathbb{Q} -independent and that*

$$\varphi(s) = \sum_{j=1}^d P_j(q_j^{-s})$$

is in \mathcal{G} , and that the polynomials P_j are non-constant. Then \mathcal{C}_φ is compact if and only if φ has restricted range or $d \geq 2$.

Theorem 2 is truly a non-linear extension of the results for linear symbols, however it fails to handle the relatively simple cases

$$(4) \quad \varphi_1(s) = \frac{9}{2} - 2^{-s} - 3^{-s} - 2 \cdot 6^{-s} \quad \text{and} \quad \varphi_2(s) = \frac{13}{2} - 4 \cdot 2^{-s} - 4 \cdot 3^{-s} + 2 \cdot 6^{-s},$$

where “mixed terms” are present. However, the compactness of the associated operators can be decided by our main result. Before this result can be stated, we need to introduce some additional definitions.

Definition. Let $\Lambda \subseteq \mathbb{N} - \{1\}$. We let the *complex dimension* of Λ , denoted $\mathcal{D}(\Lambda)$, be the infimum of $\text{card}(\Lambda_0)$ where $\Lambda_0 \subset \mathbb{N} - \{1\}$ is \mathbb{Q} -independent and multiplicatively generates Λ .

At this point, we should mention that the set Λ_0 attaining such an infimum is not necessarily unique. This is easily seen by considering $\Lambda = \{2^2 \cdot 3^2, 2^4 \cdot 3^2, 2^2 \cdot 3^4, 2^4 \cdot 3^4\}$, where Λ_0 can be chosen as any of the following sets:

$$\{2, 3\} \quad \{2^2, 3\} \quad \{2, 3^2\} \quad \{2^2, 3^2\} \quad \{2^2 \cdot 3, 3\} \quad \{2, 2 \cdot 3^2\}$$

Now, we will rewrite (2) as

$$(5) \quad \varphi(s) = c_1 + \sum_{n \in \Lambda} c_n n^{-s}$$

with $c_n \neq 0$ for every $n \in \Lambda$. We pick some $\Lambda_0 = \{q_1, q_2, \dots, q_d\}$ where $d = \mathcal{D}(\Lambda)$. Since Λ_0 generates Λ , any $n \in \Lambda$ can be written uniquely as a product of elements in Λ_0 ,

$$n = \prod_{j=1}^d q_j^{\alpha_j}.$$

This associates to n the d -dimensional multi-index $\alpha(n)$. Clearly, $\alpha(n)$ depends on the choice of Λ_0 as the example considered above illustrates.

Definition. The *degree of φ with respect to Λ_0* is defined by

$$\text{deg}(\varphi, \Lambda_0) = \sup \{|\alpha(n)| = \alpha_1 + \alpha_2 + \dots + \alpha_d : n \in \Lambda\}.$$

Among the different Λ_0 which generate Λ and with $\text{card}(\Lambda_0) = \mathcal{D}(\Lambda)$, we choose an optimal Λ_0 in the sense that it minimizes $\text{deg}(\varphi, \Lambda_0)$. The *degree* of φ is then equal to the value of $\text{deg}(\varphi, \Lambda_0)$ where Λ_0 is optimal in the previous sense.

It is clear that there can be more than one optimal Λ_0 , as the example considered above again demonstrates, where the three final possibilities all have $\text{deg}(\varphi, \Lambda_0) = 4$ if φ is given by (5).

Remark. For maps of the form (3) as considered before, the complex dimension is equal to d and the degree is equal to 1, which justifies our terminology “linear case”.

The study of the Hardy space of Dirichlet series \mathcal{H}^2 is intimately related to function theory on polydiscs. In our concerns, the main tool will be the so-called Bohr lift. Indeed, consider an optimal Λ_0 and use the substitution $q_j^{-s} \mapsto z_j$. To simplify the expressions in what follows, we will also subtract $1/2$. Hence we obtain a polynomial in d variables with the same degree as φ ,

$$(6) \quad \Phi(z) = \left(c_1 - \frac{1}{2}\right) + \sum_{n \in \Lambda} c_n z^{\alpha(n)}.$$

The polynomial Φ will be called an *optimal Bohr lift* of φ . Using Kronecker's theorem (see for instance [10, Ch. 13]), the \mathbb{Q} -independence of Λ_0 implies that Φ maps \mathbb{D}^d into \mathbb{C}_0 . The polynomial Φ induces a map, denoted by ϕ , on \mathbb{R}^d defined by

$$\phi(\theta_1, \theta_2, \dots, \theta_d) = \Phi(e^{i\theta_1}, e^{i\theta_2}, \dots, e^{i\theta_d}).$$

Remark. We will sometimes need to define the Bohr lift when the map $\varphi(s) = \sum_{n \geq 1} c_n n^{-s}$ is not a Dirichlet polynomial. It is then defined as

$$\Phi(z) = \left(c_1 - \frac{1}{2} \right) + \sum_{n \geq 2} c_n z^{\alpha(n)}$$

where we use the substitution $p_j^{-s} \mapsto z_j$. If we assume that $\varphi \in \mathcal{G}$, its Bohr lift Φ is now well-defined on $\mathbb{D}^\infty \cap c_0$, and Kronecker's theorem shows that this set is mapped by Φ into \mathbb{C}_0 .

Let us come back to a polynomial $\varphi \in \mathcal{G}$. If we assume that φ has unrestricted range, there exists at least one point $w \in \mathbb{T}^d$ so that $\operatorname{Re}\Phi(w) = 0$, by the compactness of \mathbb{T}^d . Let $w = (e^{i\vartheta_1}, e^{i\vartheta_2}, \dots, e^{i\vartheta_d})$. Then $\vartheta = (\vartheta_1, \vartheta_2, \dots, \vartheta_d)$ has to be a critical point of $\operatorname{Re}\phi$ since this last map admits a minimum at ϑ . Moreover, the mapping properties of φ implies that the Hessian matrix of $\operatorname{Re}\phi$ at ϑ should be non-negative.

Definition. We define the *boundary index* of Φ at w as the non-negative integer $J(\Phi, w)$ such that the signature of the Hessian matrix of $\operatorname{Re}\phi$ at ϑ is equal to $(J(\Phi, w), 0)$.

With these definitions at hand, we are able to state our main theorem which shows that, when there are mixed terms, the complex dimension does not give enough information and that we need a more careful study of φ .

Theorem 3. *Let $\varphi(s) = c_1 + \sum_{n \geq 2} c_n n^{-s}$ be a Dirichlet polynomial in \mathcal{G} with unrestricted range. Suppose that its complex dimension d is greater than or equal to 2, and let Φ be a minimal Bohr lift of φ . Assume that*

- *either the degree of φ is equal to 1 or 2,*
- *or the degree of φ is at least 3 and for any $w \in \mathbb{T}^d$, either $\operatorname{Re}\Phi(w) > 0$ or $\operatorname{Re}\Phi(w) = 0$ and $J(\Phi, w) \geq 2$.*

Then \mathcal{C}_φ is compact on \mathcal{H}^2 . Moreover, the result is optimal in the following sense:

- *If the complex dimension of φ is equal to 1, then \mathcal{C}_φ is never compact.*
- *There exist polynomials $\varphi \in \mathcal{G}$ of arbitrary complex dimension and of arbitrary degree greater than or equal to 3 such that \mathcal{C}_φ is not compact.*

At this point we should mention that Theorem 3 does not encompass Theorem 2, and we will return to this point later (see Section 7). However, Theorem 3 allows us to conclude that for the Dirichlet polynomials φ given by (4), which have complex dimension and degree equal to 2, the induced composition operators are compact.

We are also interested in the degree of compactness of our operators, which may be estimated using their approximation numbers.

Definition. Let H be a Hilbert space and let $T \in \mathcal{L}(H)$. The n th approximation number of T , denoted $a_n(T)$, is the distance of T to the operators of rank $< n$.

The study of the behaviour of $a_n(\mathcal{C}_\varphi)$ when $\varphi \in \mathcal{G}$ is a linear symbol (3) has been done in [11]. In particular, it is shown there that

$$\left(\frac{1}{n}\right)^{(d-1)/2} \ll a_n(\mathcal{C}_\varphi) \ll \left(\frac{\log n}{n}\right)^{(d-1)/2}$$

where d is the complex dimension of φ . We will extend this result to a general context. To keep this introduction sufficiently short, we refer to Section 8 for our statement, and give only one striking consequence of it: We may distinguish the Schatten classes of linear operators on \mathcal{H}^2 using composition operators induced by polynomial symbols. By definition, a compact linear operator T belongs to the Schatten class S_p , for $0 < p < \infty$, if

$$\|T\|_p^p := \text{Tr}(|T|^p) = \sum_{n=1}^{\infty} a_n(T)^p < \infty.$$

Corollary 4. *Let $0 < p < q$. There exists a Dirichlet polynomial $\varphi \in \mathcal{G}$ such that $\mathcal{C}_\varphi \in S_q \setminus S_p$.*

Let us end this introduction by mentioning that the the composition operators induced by the maps φ_1 and φ_2 have different degrees of compactness. Indeed, we will show that

$$\left(\frac{1}{n}\right)^{1/2} \ll a_n(\mathcal{C}_{\varphi_1}) \ll \left(\frac{\log n}{n}\right)^{1/2} \quad \text{and} \quad \left(\frac{1}{n}\right)^{1/3} \ll a_n(\mathcal{C}_{\varphi_2}) \ll \left(\frac{\log n}{n}\right)^{1/3}.$$

Organization. The remainder of this paper is divided into seven sections.

- Section 2 contains the proof of Theorem 1. The content of this section is independent from that of the following sections.
- In Section 3 we introduce some necessary tools and results needed for the proof of Theorem 2 and Theorem 3.
- Section 4 is devoted to the proof of Theorem 2.
- Section 5 contains the proof of Theorem 3.
- In Section 6 we prove Lemma 12, which is the most technical part of Theorem 3.
- In Section 7 we discuss the case $\deg(\varphi) \geq 3$ and $J(\Phi, w) = 0$, its connection to Theorem 2 and some related examples.
- Finally, in Section 8, we discuss the decay of the sequence of approximation numbers for some of our operators.

Notation. The notation $f(\varepsilon) \ll g(\varepsilon)$ will mean that $f(\varepsilon) \leq Cg(\varepsilon)$ for some constant C which does not depend on ε . We will sometimes write $f(\varepsilon) \ll_a g(\varepsilon)$ to emphasize that C depends on a . As usual, we let $\{p_j\}$ denote the sequence of prime numbers written in increasing order. We let \mathbf{m}_d denote the normalized Lebesgue measure on \mathbb{T}^d . This measure is invariant under rotations. If we do not have a priori knowledge of the complex dimension d , we will often call this measure \mathbf{m}_∞ . For a point $z = e^{i\theta}$ on the unit circle \mathbb{T} , we will always assume that $\theta \in (-\pi, \pi]$. Finally, $\mathbf{0}$ will denote the point $(0, \dots, 0) \in \mathbb{C}^d$, and $\mathbf{1}$ will similarly denote the point $(1, \dots, 1)$.

2. PROOF OF THEOREM 1

Let $\varphi(s) = c_0 s + c_1 + \sum_{n=2}^N c_n n^{-s} \in \mathcal{G}$ such that $c_0 \geq 1$. We already know that if φ has restricted range, then \mathcal{C}_φ is compact. Let us therefore assume that \mathcal{C}_φ is compact and also assume that φ has unrestricted range, to argue by contradiction.

By [2, Thm. 3], we know that

$$(7) \quad \frac{\operatorname{Re} \varphi(s)}{\operatorname{Re}(s)} \xrightarrow{\operatorname{Re}(s) \rightarrow 0} +\infty.$$

Now, since φ has unrestricted range there exists a sequence $\{s_k = \sigma_k + i t_k\}_{k \geq 1}$ in \mathbb{C}_0 such that $\operatorname{Re} \varphi(s_k) \rightarrow 0$. It is well-known that this forces that $\sigma_k \rightarrow 0$ (see [2]). Then

$$\operatorname{Re} \varphi(s_k) = c_0 \sigma_k + \operatorname{Re}(c_1) + \sum_{n=2}^N n^{-\sigma_k} (\operatorname{Re}(c_n) \cos(t_k \log(n)) + \operatorname{Im}(c_n) \sin(t_k \log(n))).$$

By successive extraction of subsequences, we may assume that there exist real numbers a_n and b_n so that for $2 \leq n \leq N$ we have, as $k \rightarrow \infty$,

$$\cos(t_k \log(n)) \rightarrow a_n \quad \text{and} \quad \sin(t_k \log(n)) \rightarrow b_n.$$

Hence, we may write

$$\operatorname{Re} \varphi(s_k) = c_0 \sigma_k + \operatorname{Re}(c_1) + \sum_{n=2}^N n^{-\sigma_k} (\operatorname{Re}(c_n) a_n + \operatorname{Im}(c_n) b_n) + \sum_{n=2}^N n^{-\sigma_k} F_n(t_k),$$

where each $F_n(t_k) \rightarrow 0$ as $k \rightarrow \infty$. Since $\operatorname{Re} s_k = \sigma_k$ also goes to 0, we may deduce that

$$\operatorname{Re}(c_1) + \sum_{n=2}^N (\operatorname{Re}(c_n) a_n + \operatorname{Im}(c_n) b_n) = 0,$$

so that we have

$$\operatorname{Re} \varphi(s) = c_0 \sigma + \sum_{n=2}^N (n^{-\sigma} - 1) (\operatorname{Re}(c_n) a_n + \operatorname{Im}(c_n) b_n) + \sum_{n=2}^N n^{-\sigma} F_n(t).$$

We will now choose another sequence $\{s'_k = \sigma'_k + i t_k\}_{k \geq 1}$ where $\operatorname{Re}(s'_k) \rightarrow 0$ in order to obtain a contradiction with (7). More precisely, let $\{\sigma'_k\}_{k \geq 1}$ be any sequence of positive real numbers tending to 0 such that, for any $n = 2, \dots, N$ and every $k \geq 1$, we have $n^{-\sigma'_k} |F_n(t_k)| \leq \sigma'_k$. Then we obtain

$$\operatorname{Re} \varphi(s'_k) = c_0 \sigma'_k + \sum_{n=2}^N (n^{-\sigma'_k} - 1) (\operatorname{Re}(c_n) a_n + \operatorname{Im}(c_n) b_n) + \sum_{n=2}^N n^{-\sigma'_k} F_n(t_k) = \mathcal{O}(\sigma'_k) = \mathcal{O}(\operatorname{Re}(s'_k)),$$

and this contradicts (7). The assumption that φ has unrestricted range must be wrong. \square

Remark. An inspection of the proof reveals that the statement of Theorem 1 remains true if we assume that $\varphi(s) = c_0 s + c_1 + \sum_{n=2}^{\infty} c_n n^{-s} \in \mathcal{G}$ with $c_0 \geq 1$, $\sum_{n=1}^{\infty} |c_n| < +\infty$ and that the complex dimension of φ is finite. The latter assumption is needed to use (7).

3. PRELIMINARIES

As explained in the introduction, our main tool for proving or disproving compactness is a result from [11]. We formulate it in a more general context than for polynomials since it will be used under this form in Section 8. Recall that a *Carleson square* in \mathbb{C}_0 is a closed square in $\overline{\mathbb{C}_0}$ with one of its sides lying on the vertical line $i\mathbb{R}$; the side length of Q is denoted by $\ell(Q)$. A non-negative Borel measure μ on $\overline{\mathbb{C}_0}$ is called a *vanishing Carleson measure* if

$$\limsup_{\ell(Q) \rightarrow 0} \frac{\mu(Q)}{\ell(Q)} = 0.$$

Lemma 5. *Suppose that $\varphi(s) = \sum_{n \geq 1} c_n n^{-s} \in \mathcal{G}$ and that $\varphi(\mathbb{C}_0)$ is bounded. The corresponding composition operator \mathcal{C}_φ is compact on \mathcal{H}^2 if and only if the measure*

$$\mu_\varphi(E) := \mathbf{m}_\infty(\{z \in \mathbb{T}^\infty : \Phi(z) \in E\}), \quad E \subseteq \mathbb{C}_0.$$

is vanishing Carleson in \mathbb{C}_0 , where Φ denotes a Bohr lift of φ .

Proof. This is Corollary 4.1 in [11]. □

Hence we consider squares

$$Q = Q(\tau, \varepsilon) = [0, \varepsilon] \times [\tau - \varepsilon/2, \tau + \varepsilon/2],$$

and want to investigate whether $\mu_\varphi(Q) = o(\varepsilon)$ uniformly in $\tau \in \mathbb{R}$. Our next lemma points out that this depends only on the local behaviour of Φ .

Lemma 6. *Let φ be a Dirichlet polynomial (2) with $c_0 = 0$ mapping \mathbb{C}_0 into $\mathbb{C}_{1/2}$ and let Φ be a minimal Bohr lift of φ . If for every $w \in \mathbb{T}^d$ with $\operatorname{Re} \Phi(w) = 0$ there exists a neighbourhood $\mathcal{U}_w \ni w$ in \mathbb{T}^d , constants $C_w > 0$ and $\kappa_w > 1$ such that for every $\tau \in \mathbb{R}$ and every $\varepsilon > 0$ we have*

$$(8) \quad \mathbf{m}_d(\{z \in \mathcal{U}_w : \Phi(z) \in Q(\tau, \varepsilon)\}) \leq C_w \varepsilon^{\kappa_w},$$

then \mathcal{C}_φ is compact.

Proof. Since φ is a Dirichlet polynomial, it has finite complex dimension d .

We first observe that (8) is always satisfied for those $w \in \mathbb{T}^d$ with $\operatorname{Re} \Phi(w) > 0$. Indeed, by continuity of Φ , we may always find a neighbourhood $\mathcal{U}_w \ni w$ and $\varepsilon_0 > 0$ such that, for all $\varepsilon \in (0, \varepsilon_0)$ and all $\tau \in \mathbb{R}$, $\{z \in \mathcal{U}_w : \Phi(z) \in Q(\tau, \varepsilon)\}$ is empty. We may then take $\kappa_w > 1$ be arbitrary and choose C_w with $C_w \varepsilon_0^{\kappa_w} \geq 1$.

We will then use a compactness argument and Lemma 5. Indeed, there exists a finite number of points w_1, \dots, w_N such that \mathbb{T}^d is covered by $\mathcal{U}_{w_1}, \dots, \mathcal{U}_{w_N}$. Now, we may take $C = C_{w_1} + \dots + C_{w_N}$ and $\kappa = \min(\kappa_{w_1}, \dots, \kappa_{w_N})$. Hence, for all $\tau \in \mathbb{R}$ and all $\varepsilon > 0$,

$$\mathbf{m}_d(\{z \in \mathbb{T}^d : \Phi(z) \in Q(\tau, \varepsilon)\}) \leq C \varepsilon^\kappa$$

which achieves the proof of the compactness of \mathcal{C}_φ on \mathcal{H}^2 . □

Hence, we will require more information about the Taylor coefficients of Φ at a boundary point. Assume that $\Phi(w) = 0$ where $w = \mathbf{1}$. In this case, we will rewrite

$$(9) \quad \Phi(z) = \sum_{n \in \Lambda} \widetilde{c}_n \prod_{j=1}^d (1 - z_j)^{\alpha_j} = \sum_{\alpha \in \mathbb{N}^d} c_\alpha (1 - z)^\alpha,$$

where we have adopted the convention $c_\alpha = \widetilde{c}_n$, which is not generally equal to c_n . We shall need a kind of Julia–Caratheodory theorem for Φ of the form (9).

Lemma 7. *Let $\Phi: \mathbb{D}^d \rightarrow \mathbb{C}_0$ be of the form (9) and let $|\alpha| = 1$. Then $c_\alpha \geq 0$. Moreover, there exists at least one multi-index α with $|\alpha| = 1$ and $c_\alpha > 0$, unless $\Phi \equiv 0$.*

Proof. We may assume that $\alpha = (1, 0, \dots, 0)$. Consider the one-variable polynomial

$$\psi(w) = \Phi(w, 1, \dots, 1).$$

Clearly, ψ maps \mathbb{D} to \mathbb{C}_0 , and $\psi(1) = 0$. We write

$$\psi(w) = a(1 - w) + b(1 - w)^2 + \mathcal{O}((1 - w)^3).$$

We set $w = e^{i\theta}$ and obtain

$$\psi(e^{i\theta}) = a \left(\frac{\theta^2}{2} - i\theta \right) - b\theta^2 + \mathcal{O}(\theta^3).$$

In particular,

$$\operatorname{Re}(\psi(e^{i\theta})) = \theta \operatorname{Im}(a) + \theta^2 \left(\frac{\operatorname{Re}(a)}{2} - \operatorname{Re}(b) \right) + \mathcal{O}(\theta^3).$$

Since this should be non-negative, clearly $\operatorname{Im}(a) = 0$. We now set $w = 1 - \delta$ for $0 < \delta < 1$ and consider $\psi(\delta) = a\delta + \mathcal{O}(\delta^2)$. Since the real part of this also should be non-negative as $\delta \rightarrow 0^+$ we must have $a \geq 0$. Hence $c_\alpha \geq 0$ when $|\alpha| = 1$.

Now, consider the mapping

$$\alpha \mapsto n(\alpha) = \prod_{j=1}^d p_j^{\alpha_j}.$$

It defines a total order on \mathbb{N}^d by setting $\alpha \leq \beta$ if and only if $n(\alpha) \leq n(\beta)$. Assume that $\Phi \not\equiv 0$ and that $c_\alpha = 0$ whenever $|\alpha| = 1$. Consider

$$\beta = \inf\{\alpha : c_\alpha \neq 0\},$$

which exists since $\Phi \not\equiv 0$. There is $\theta \in (-\pi, \pi]$ so that $c_\beta = |c_\beta| e^{i\theta}$. Fix $\theta_j \in (-\pi/2, \pi/2)$ and define

$$z_j = 1 - p_j^{-\sigma} e^{i\theta_j},$$

where $\sigma > 0$. For large enough σ , clearly $z = (z_1, \dots, z_d) \in \mathbb{D}^d$. Moreover,

$$\Phi(z_1, \dots, z_d) = |c_\beta| e^{i\theta} [n(\beta)]^{-\sigma} e^{i(\beta_1\theta_1 + \dots + \beta_d\theta_d)} + o([n(\beta)]^{-\sigma}),$$

as $\sigma \rightarrow \infty$. This implies that

$$\operatorname{Re}(\Phi(z_1, \dots, z_d)) = |c_\beta| [n(\beta)]^{-\sigma} \cos(\theta + \beta_1\theta_1 + \dots + \beta_d\theta_d) + o([n(\beta)]^{-\sigma}).$$

Since $|\beta| \geq 2$, we can always choose $\theta_j \in (-\pi/2, \pi/2)$ such that $\cos(\theta + \beta_1\theta_1 + \dots + \beta_d\theta_d) < 0$. This contradicts the mapping properties of Φ , and hence the assumption that $c_\alpha = 0$ whenever $|\alpha| = 1$ is wrong. \square

We will also need two lemmas from differential geometry. The first one is the parametrized Morse lemma (see for instance [5, Sec. 4.44]).

Lemma (Parametrized Morse Lemma). *Let $\mathcal{U} \subset \mathbb{R}^J \times \mathbb{R}^{d-J}$ be a neighbourhood of $\mathbf{0} \in \mathbb{R}^d$ and let $F : \mathcal{U} \rightarrow \mathbb{R}$, $(u, v) \mapsto F(u, v)$ be a smooth function. Assume that $F(\mathbf{0}) = 0$, that $\partial F / \partial u_i(\mathbf{0}) = 0$ for all $i = 1, \dots, J$ and that the matrix*

$$\left(\frac{\partial^2 F}{\partial u_i \partial u_j}(\mathbf{0}) \right)_{1 \leq i, j \leq J}$$

is positive definite. Then there exist a neighbourhood $\mathcal{V} \ni \mathbf{0}$ with $\mathcal{V} \subset \mathcal{U}$, a smooth diffeomorphism $\Gamma : \mathcal{V} \rightarrow \mathbb{R}^d$, $(u, v) \mapsto (\gamma(u, v), v)$ with $\Gamma(\mathbf{0}) = \mathbf{0}$ and a smooth map $h : \mathbb{R}^{d-J} \rightarrow \mathbb{R}$ such that, for any $(u, v) \in \mathcal{V}$,

$$F(u, v) = \sum_{j=1}^J \gamma_j(u, v)^2 + h(v).$$

The second lemma reads as follows.

Lemma 8. *Let $p \geq 1$ be an integer, and let $f : I \rightarrow \mathbb{R}$ be a smooth function where I is an open interval containing 0 and $f(x) \sim_0 x^p$. Then there exist $C > 0$ and an open interval $I' \ni 0$ inside I such that, for any $\tau \in \mathbb{R}$ and any $\delta > 0$, the set $\{x \in I' : |f(x) - \tau| < \delta\}$ has Lebesgue measure less than $C\delta^{1/p}$.*

Proof. Assume first that $f(x) = x^p$. If $|\tau| \leq 2\delta$, then the result is clear. Otherwise, if $\tau \geq 2\delta$, then x has to live in $[(\tau - \delta)^{1/p}, (\tau + \delta)^{1/p}]$ and the length of this interval may be easily estimated using the mean value theorem.

The general case reduces to this one. For small values of x , set $y = [f(x)]^{1/p}$ if p is odd or $y = [f(x)]^{1/p}$ for $x > 0$, $y = -[f(x)]^{1/p}$ for $x < 0$ if p is even. In both cases, y is differentiable at 0 and $dy/dx > 0$. Hence, $x = \gamma(y)$ where γ is a smooth diffeomorphism. Now, for some small open interval $I' \ni 0$, we have

$$\{x \in I' : |f(x) - \tau| < \delta\} = \{x \in I' : |(\gamma^{-1}(x))^p - \tau| < \delta\}.$$

Since γ is a diffeomorphism, the latter set has Lebesgue measure less than $C\delta^{1/p}$. \square

4. PROOF OF THEOREM 2

We intend to apply Lemma 6. Hence, let $w \in \mathbb{T}^d$ with $\operatorname{Re} \Phi(w) = 0$. By the rotational invariance of \mathbf{m}_d , we may always assume that $w = \mathbf{1}$. Moreover, since the conditions in Lemma 6 are invariant by vertical translations, we may also assume that $\Phi(w) = 0$. In this case we have

$$\Phi(z_1, z_2, \dots, z_d) = \sum_{j=1}^d \Phi_j(z_j) = \sum_{j=1}^d \sum_k a_k^{(j)} (1 - z_j)^k.$$

Since Φ is a minimal Bohr lift of φ , inspecting the proof of Lemma 7, we may conclude that in this case $a_1^{(j)} > 0$ for every $j = 1, 2, \dots, d$. This means we have

$$\operatorname{Re} \Phi(e^{i\theta_1}, e^{i\theta_2}, \dots, e^{i\theta_d}) = \sum_{j=1}^d b_j \theta_j^{k_j} + o(\theta_j^{k_j}),$$

where the coefficients $b_j \neq 0$ are real numbers and the exponents $k_j \geq 2$ are integers. The fact that this quantity is supposed to be non-negative implies that $b_j > 0$ and that k_j is even, by similar considerations as those in the proof of Lemma 7. Moreover

$$\operatorname{Im} \Phi(e^{i\theta_1}, e^{i\theta_2}, \dots, e^{i\theta_d}) = - \sum_{j=1}^d a_1^{(j)} \theta_j + o(\theta_j).$$

Proof of the first part of Theorem 2. Let $\tau \in \mathbb{R}$ and $\varepsilon > 0$ be arbitrary. The preceding discussion means there is some neighbourhood $\mathcal{U} \ni (1, 1, \dots, 1)$ in \mathbb{T}^d so that

$$\frac{1}{2} \sum_{j=1}^d b_j \theta_j^{k_j} \leq \operatorname{Re} \Phi(e^{i\theta_1}, e^{i\theta_2}, \dots, e^{i\theta_d}) \leq 2 \sum_{j=1}^d b_j \theta_j^{k_j},$$

when $e^{i\theta} \in \mathcal{U}$. Hence if $\Phi(e^{i\theta}) \in Q(\tau, \varepsilon)$ and $e^{i\theta} \in \mathcal{U}$, we conclude from the real part that $|\theta_j| \ll \varepsilon^{1/k_j}$, for $j = 1, 2, \dots, d$. Now, fixing θ_j for $j = 2, \dots, d$ we conclude from the imaginary part and Lemma 8 that θ_1 can live in an interval of size at most $C\varepsilon$. Hence we have

$$\mathbf{m}_d(\{z \in \mathcal{U}_w : \Phi(z) \in Q(\tau, \varepsilon)\}) \ll_w \varepsilon^{1+1/k_2+\dots+1/k_d}.$$

In fact, we may choose

$$\kappa_w = 1 + \sum_{j=1}^d \frac{1}{k_j} - \min_{1 \leq j \leq d} \frac{1}{k_j},$$

and conclude by Lemma 6, since $d \geq 2$. □

Proof of the second part of Theorem 2. In this case $d = 1$, and the polynomial $\Phi(z)$ is of only one variable. We again consider some neighbourhood $\mathcal{U} \ni 1$ in \mathbb{T} , so that when $e^{i\theta} \in \mathcal{U}$ we have

$$0 \leq \operatorname{Re} \Phi(e^{i\theta}) \leq 2b\theta^k \quad \text{and} \quad |\operatorname{Im} \Phi(e^{i\theta})| \leq 2a|\theta|,$$

where $a = a_1$, $b = b_1$ and $k \geq 2$ is even. Now, we choose $\tau = 0$ and observe that $\varphi(e^{i\theta})$ belongs to $Q(\tau, \varepsilon)$ provided $|\theta| \ll \varepsilon$. Hence

$$\mathbf{m}_1(\{z \in \mathbb{T} : \Phi(z) \in Q(\tau, \varepsilon)\}) \geq \mathbf{m}_1(\{z \in \mathcal{U}_w : \Phi(z) \in Q(\tau, \varepsilon)\}) \gg \varepsilon,$$

and \mathcal{C}_φ cannot be compact by Lemma 5. □

Remark. Inspecting the proof of Theorem 2, we see that we may replace the polynomials P_j , by corresponding power series

$$P_j(q_j^{-s}) = \sum_{k=0}^{\infty} c_k^{(j)} q_j^{-ks},$$

provided $\sum_{k=0}^{\infty} |c_k^{(j)}| < \infty$. However, we still require the complex dimension d to be finite.

5. PROOF OF THEOREM 3

We begin by observing that the penultimate point of Theorem 3 follows from the second part of Theorem 2. Regarding the final part of Theorem 3, it is contained in the following result:

Lemma 9. *There are polynomials $\varphi \in \mathcal{G}$ of any complex dimension and of any degree ≥ 3 for which the corresponding composition operator \mathcal{C}_φ is non-compact.*

Proof. Let $P(z) = P(z_1, z_2, \dots, z_d)$ be any polynomial in d variables and define

$$\Phi(z) = (1 - z_1) + \delta(1 - z_1)^2 P(z),$$

for some $\delta > 0$ to be decided later. We compute

$$\operatorname{Re} \Phi(e^{i\theta_1}, \dots, e^{i\theta_d}) = (1 - \cos \theta_1) \left(1 - 2\delta \left(\cos(\theta_1) \operatorname{Re} P(e^{i\theta}) - \sin(\theta_1) \operatorname{Im} P(e^{i\theta}) \right) \right).$$

Pick δ small enough so that we have

$$\frac{1 - \cos \theta_1}{2} \leq \operatorname{Re} \Phi(e^{i\theta_1}, \dots, e^{i\theta_d}) \leq 2(1 - \cos \theta_1).$$

The first inequality tells us that Φ is a minimal Bohr lift of

$$\varphi(s) = (1 - p_1^{-s}) + \delta(1 - p_1^{-s})^2 P(p_1^{-s}, \dots, p_d^{-s}),$$

with $\varphi \in \mathcal{G}$ having unrestricted range. Using the second inequality and a Taylor expansion of $\operatorname{Im} \Phi$, we also get that near $\mathbf{1}$,

$$\begin{aligned} \operatorname{Re} \Phi(e^{i\theta_1}, \dots, e^{i\theta_d}) &= \mathcal{O}(\theta_1^2), \\ \operatorname{Im} \Phi(e^{i\theta_1}, \dots, e^{i\theta_d}) &= \mathcal{O}(\theta_1). \end{aligned}$$

Similar considerations as in the proof of the second part of Theorem 2 allow us to conclude that \mathcal{C}_φ is not compact. \square

Remark. The key point of Lemma 9 is that even if Φ involves d variables, its local behaviour near $\mathbf{1}$ depends too heavily on z_1 to ensure compactness.

Having now concluded the negative parts of Theorem 3, we turn to the positive parts. Let us fix a polynomial $\varphi \in \mathcal{G}$ and let Φ denote a minimal Bohr lift of φ . We can simplify how to write Φ around a point $w \in \mathbb{T}^d$ such that $\operatorname{Re} \Phi(w) = 0$. Without loss of generality, we may again assume that $w = \mathbf{1}$ and that $\Phi(w) = 0$. Then we may write

$$\Phi(z) = \sum_{j=1}^d a_j (1 - z_j) + \sum_{j=1}^d b_j (1 - z_j)^2 + \sum_{1 \leq j < k \leq d} c_{j,k} (1 - z_j)(1 - z_k) + o\left(\sum_{1 \leq j \leq d} |1 - z_j|^2\right).$$

We let $z_j = e^{i\theta_j}$ and since $a_j \geq 0$ by Lemma 7 we get

$$\operatorname{Re}(\Phi(z)) = \sum_{j=1}^d \left(\frac{a_j}{2} - \operatorname{Re}(b_j) \right) \theta_j^2 - \sum_{1 \leq j < k \leq d} \operatorname{Re}(c_{j,k}) \theta_j \theta_k + o\left(\sum_{1 \leq j \leq d} \theta_j^2\right).$$

The quadratic form appearing above is brought to standard form by a linear change of variables,

$$\operatorname{Re}(\Phi(z)) = \sum_{j=1}^d (\ell_j(\theta))^2 + o\left(\sum_{1 \leq j \leq d} \theta_j^2\right).$$

Next, we write

$$\operatorname{Im}(\Phi(z)) = -\sum_{j=1}^d a_j \theta_j + o\left(\sum_{j=1}^d |\theta_j|\right) = -\ell_{d+1}(\theta) + o\left(\sum_{j=1}^d |\theta_j|\right),$$

and by Lemma 7 we know that $\ell_{d+1} \neq 0$, since at least one $a_j > 0$. The last step to finish the proof of Theorem 3 is the following result:

Lemma 10. *Let $\Phi : \mathbb{D}^d \rightarrow \mathbb{C}_0$ be an optimal Bohr lift of $\varphi \in \mathcal{G}$, where φ has unrestricted range and complex dimension $d \geq 2$. Suppose that $w \in \mathbb{T}^d$ is such that $\operatorname{Re} \Phi(w) = 0$. Then there exist a neighbourhood $\mathcal{U}_w \ni w$ in \mathbb{T}^d , $\kappa = \kappa_w > 1$ and $C = C_w > 0$ such that, for any $\tau \in \mathbb{R}$ and for every $\varepsilon > 0$,*

$$\mathbf{m}_d(\{z \in \mathcal{U}_w : \Phi(z) \in Q(\tau, \varepsilon)\}) \leq C\varepsilon^\kappa$$

in the following cases:

- $J(\Phi, w) \geq 1$ and ℓ_{d+1} is independent from (ℓ_1, \dots, ℓ_j) . We may choose $\kappa = 1 + J(\Phi, w)/2$.
- $J(\Phi, w) \geq 2$ and ℓ_{d+1} belongs to $\operatorname{span}(\ell_1, \dots, \ell_j)$. We may choose $\kappa = (1 + J(\Phi, w))/2$.
- $J(\Phi, w) = 1$, ℓ_{d+1} is a multiple of ℓ_1 and Φ has degree 2. We may choose $\kappa = 9/8$.
- $J(\Phi, w) = 0$ and Φ has degree 2. We may choose $\kappa = (d + 3)/4$.

Before we prove the different cases of this lemma, let us make some comments. Firstly, it is clear that Lemma 10 and Lemma 6 imply Theorem 3 when the degree of φ is at least 2. When the degree of φ is equal to 1, then

$$\Phi(z) = \sum_{j=1}^d a_j(1 - z_j)$$

so that each a_j is positive. This implies that $J(\Phi, w) = d$ so that we may again apply Lemma 10 and Lemma 6.

It is also important to notice that Φ cannot be an arbitrary polynomial mapping of \mathbb{D}^d into \mathbb{C}_0 . It is an optimal Bohr lift of some $\varphi \in \mathcal{G}$ with complex dimension d . In particular, we shall use that $\frac{\partial \Phi}{\partial z_j} \neq 0$ for every $1 \leq j \leq d$. Moreover, the polynomial $\Phi(z) = \lambda(1 - z_1 z_2)$ is not an optimal Bohr lift. Otherwise, it would arise from $\varphi(s) = \lambda(1 - q_1^{-s} q_2^{-s})$, but the optimal Bohr lift of φ is $\lambda(1 - z)$.

We are now ready for the proof of Lemma 10. By similar considerations as before, we may again assume that $w = \mathbf{1}$ and that $\Phi(w) = 0$. We write J for $J(\Phi, w)$.

The case $J = 0$. This implies that

$$\frac{a_j}{2} - \operatorname{Re}(b_j) = \operatorname{Re}(c_{j,k}) = 0,$$

for $j, k = 1, \dots, d$. We set $z_j = e^{i\theta_j}$ and compute

$$\begin{aligned} \operatorname{Re}(a_j(1 - z_j)) &= a_j(1 - \cos\theta_j) \\ \operatorname{Re}(b_j(1 - z_j)^2) &= -a_j \cos\theta_j(1 - \cos\theta_j) + 2\operatorname{Im}(b_j) \sin\theta_j(1 - \cos\theta_j) \\ \operatorname{Re}(c_{j,k}(1 - z_j)(1 - z_k)) &= \operatorname{Im}(c_{j,k})(\sin\theta_j(1 - \cos\theta_k) + \sin\theta_k(1 - \cos\theta_j)) \end{aligned}$$

This means that

$$\operatorname{Re}(\Phi(z)) = \sum_{j=1}^d \operatorname{Im}(b_j)\theta_j^3 + \sum_{1 \leq j < k \leq d} \frac{\operatorname{Im}(c_{j,k})}{2} (\theta_j\theta_k^2 + \theta_k\theta_j^2) + o\left(\sum_{j=1}^d |\theta_j|^3\right).$$

However, the non-negativity of $\operatorname{Re} \Phi$ then implies that $\operatorname{Im}(b_j) = \operatorname{Im}(c_{j,k}) = 0$. Hence we in total have $b_j = a_j/2$ and $c_{j,k} = 0$, which means

$$\Phi(z) = \sum_{j=1}^d \left(a_j(1 - z_j) + \frac{a_j}{2}(1 - z_j)^2 \right).$$

In fact, this means that $a_j > 0$ for every j , by the assumption that the complex dimension is d and Lemma 7. We may now use (the proof of) Theorem 2 to conclude that there exists a neighbourhood $\mathcal{U}_w \ni w$ such that

$$\mathbf{m}_d(\{z \in \mathcal{U}_w : \Phi(z) \in Q(\tau, \varepsilon)\}) \ll \varepsilon \times \varepsilon^{\frac{d-1}{4}} = \varepsilon^{\frac{d+3}{4}},$$

since we now have

$$\operatorname{Re} \Phi(e^{i\theta_1}, \dots, e^{i\theta_d}) = \frac{1}{4} \sum_{j=1}^d a_j \theta_j^4 + o(\theta_j^4),$$

and we are done with this case. □

The case $J \geq 1$ and independence. After a linear change of variables, we may write $\operatorname{Re} \phi$ and $\operatorname{Im} \phi$ as

$$\begin{aligned} \operatorname{Re} \phi(\theta_1, \dots, \theta_d) &= u_1^2 + \dots + u_J^2 + o\left(\sum_{j=1}^d u_j^2\right) \\ \operatorname{Im} \phi(\theta_1, \dots, \theta_d) &= u_d + o\left(\sum_{j=1}^d |u_j|\right) \end{aligned}$$

Since a linear change of variables does not change the value of the volume up to constants, we may assume that ϕ depends on (u, v) with $u = (u_1, u_2, \dots, u_J)$ and $v = (u_{J+1}, \dots, u_d)$. Applying the parametrized Morse lemma to $\operatorname{Re} \phi$, we may write

$$\operatorname{Re} \phi(u, v) = \gamma_1(u, v)^2 + \dots + \gamma_J(u, v)^2 + h(v).$$

We also apply the change of variables $(u, v) \mapsto \Gamma(u, v)$ to $\text{Im}\phi$ and since $\Gamma_d(u, v) = u_d$, we find

$$\text{Im}\phi(u, v) = u_d + g(\Gamma(u, v)),$$

where g is a smooth function defined on \mathcal{V} such that $\partial g / \partial u_d(\mathbf{0}) = 0$.

Now, we know that $\text{Re}\phi(u, v) \geq 0$ for every $(u, v) \in \mathbb{R}^d$. Since Γ is a diffeomorphism, v can take any value in some neighbourhood of zero in \mathbb{R}^{d-J} even if we require that

$$\gamma_1(u, v) = \gamma_2(u, v) = \dots = \gamma_J(u, v) = 0,$$

and hence $h(v) \geq 0$.

This implies that we may find some neighbourhood $\mathcal{W} \ni \mathbf{0}$ in \mathcal{V} such that, for every $\tau \in \mathbb{R}$ and every $\varepsilon > 0$,

$$(u, v) \in \mathcal{W} \quad \text{and} \quad \phi(u, v) \in Q(\tau, \varepsilon) \quad \implies \quad |\gamma_j(u, v)| \leq \varepsilon^{1/2}.$$

Now, for if we fix $\gamma_1(u, v), \dots, \gamma_{d-1}(u, v)$, it follows from Lemma 8 with $p = 1$ that $\gamma_d(u, v) = u_d$ has to belong to some interval of size $C\varepsilon$, provided that (u, v) is sufficiently close to $\mathbf{0}$. This means that there exists a neighbourhood $\mathcal{O} \subset \mathcal{W}$ of $\mathbf{0}$ such that

$$\{(u, v) \in \mathcal{O} : \phi(u, v) \in Q(\tau, \varepsilon)\} \subset \{(u, v) \in \mathcal{O} : \Gamma(u, v) \in R(\tau, \varepsilon)\},$$

where the volume of $R(\tau, \varepsilon)$ is less than $C\varepsilon^{1+\frac{J}{2}}$. Since Γ is a diffeomorphism, we are done. \square

The case $J \geq 2$ and dependence. With a similar linear change of variables as in the previous case, we may write

$$\begin{aligned} \text{Re}\phi(u_1, \dots, u_d) &= u_1^2 + \dots + u_J^2 + o\left(\sum_{j=1}^d u_j^2\right) \\ \text{Im}\phi(u_1, \dots, u_d) &= \sum_{j=1}^J \alpha_j u_j + o\left(\sum_{j=1}^d |u_j|\right) \end{aligned}$$

We use again the parametrized Morse lemma with $\text{Re}\phi$, and it is again easy to show that $\gamma_j(u, v) = u_j + o\left(\sum_{j=1}^d |u_j|\right)$ so that

$$\text{Im}\phi(u, v) = \sum_{j=1}^J \alpha_j \gamma_j(u, v) + g(\Gamma(u, v))$$

with $\partial g / \partial u_j(\mathbf{0}) = 0$ for $j = 1, \dots, d$.

We argue as in the previous case. For every $j = 2, \dots, J$, for any $\tau \in \mathbb{R}$ and every $\varepsilon > 0$,

$$(u, v) \in \mathcal{W} \subset \mathcal{V} \quad \text{and} \quad \phi(u, v) \in Q(\tau, \varepsilon) \quad \implies \quad |\gamma_j(u, v)| \leq \varepsilon^{1/2}.$$

Now, for a fixed value of $\gamma_2(u, v), \dots, \gamma_d(u, v)$, it is again clear that $\gamma_1(u, v)$ has to belong to some interval of size $C\varepsilon$, provided (u, v) is sufficiently close to $\mathbf{0}$. This means that there exists a neighbourhood $\mathcal{O} \ni \mathbf{0}$ in \mathcal{W} such that

$$\{(u, v) \in \mathcal{O} : \phi(u, v) \in Q(\tau, \varepsilon)\} \subset \{(u, v) \in \mathcal{O} : \Gamma(u, v) \in R(\tau, \varepsilon)\},$$

where the volume of $R(\tau, \varepsilon)$ is less than $C\varepsilon^{1+\frac{J-1}{2}}$. We conclude as in the previous step. \square

The case $J = 1$ and dependence, $d = 2$. This is the most difficult case. At first, we do not assume that $d = 2$ but we always assume that the degree of φ is equal to 2. We know that there is constant $\lambda \in \mathbb{R}^*$ so that $\ell_1(\theta) = \lambda \ell_{d+1}(\theta)$, which means

$$\sqrt{\frac{a_j}{2} - \operatorname{Re}(b_j)} = \lambda a_j, \quad 1 \leq j \leq d$$

and that $\lambda > 0$ by the computations in the beginning of this section. We normalize $\Phi(z)$ as $\lambda^{-2}\Phi(z)$, so that we may assume that $\lambda = 1$. Hence

$$\ell_1(\theta) = \sum_{j=1}^d a_j \theta_j,$$

and this immediately implies that

$$(10) \quad \operatorname{Re}(b_j) = \frac{a_j}{2} - a_j^2 \quad \text{and} \quad \operatorname{Re}(c_{j,k}) = -2a_j a_k, \quad 1 \leq j, k \leq d.$$

Suppose that $a_1 = 0$. Then $\operatorname{Re}(b_1) = 0$ and $\operatorname{Re}(c_{1,k}) = 0$ for $2 \leq k \leq d$. We compute

$$\operatorname{Re}(\Phi(e^{ix}, 1, \dots, 1)) = -2\operatorname{Im}(b_1) \sin x (1 - \cos x) \geq 0,$$

which means that $\operatorname{Im}(b_1) = 0$, so that $b_1 = 0$. Next we compute

$$\begin{aligned} \Phi(e^{ix}, e^{iy}, 1, \dots, 1) &= a_2(1 - \cos y) + \left(\frac{a_2}{2} - a_2^2\right)(1 - 2\cos y + \cos 2y) \\ &\quad - \operatorname{Im}(c_{1,2})(-\sin x - \sin y + \sin(x+y)) \\ &= (1 - \cos y)(a_2(1 - \cos y) + 2a_2^2 \cos y + \operatorname{Im}(c_{1,2}) \sin x) \\ &\quad + \operatorname{Im}(c_{1,2}) \sin y(1 - \cos x). \end{aligned}$$

Taking $y = \pm\delta$ for small enough δ , we obtain that $\operatorname{Im}(c_{1,2}) = 0$. There is nothing special about z_2 , and hence we conclude that $\operatorname{Im}(c_{1,k}) = 0$, for $2 \leq k \leq d$. In particular, $c_{1,k} = 0$ for the same values of k . But this is impossible, since the variable z_1 no longer appear in our polynomial. Hence the assumption that $a_1 = 0$ must be wrong.

Arguing in the same way, we have that $a_j > 0$ for $1 \leq j \leq d$. Moreover, after renaming the variables, we may suppose $a_1 \geq a_2 \geq \dots \geq a_d > 0$. Finally,

$$0 \leq \operatorname{Re}(\Phi(-1, 1, \dots, 1)) = 2a_1 + 4\left(\frac{a_1}{2} - a_1^2\right) \implies a_1 \leq 1,$$

so without loss of generality, we may assume that

$$1 \geq a_1 \geq a_2 \geq \dots \geq a_d > 0.$$

From now on, we assume that $d = 2$ and that $1 \geq a_1 \geq a_2 > 0$. We need the following lemma.

Lemma 11. *We have $a_2 \leq 1 - a_1$.*

Proof. We compute

$$\Phi(-1, -1) = -4a_1^2 - 4a_2^2 - 8a_1 a_2 + 4a_1 + 4a_2 = 4(a_1 + a_2)(1 - a_1 - a_2).$$

Since this has to be non-negative, we get the result. \square

Remark. Lemma 11 immediately implies that $a_1 \in (0, 1)$ and $a_2 \in (0, 1/2]$ by the assumptions that $0 < a_2 \leq a_1 \leq 1$.

Let us apply the change of variables $\theta_1 = a_2 u + a_2 v$, $\theta_2 = a_1 u - a_1 v$ to ϕ :

$$(11) \quad \operatorname{Re} \phi(u, v) = -4a_1^2 a_2^2 u^2 + o(u^2 + v^2)$$

$$(12) \quad \operatorname{Im} \phi(u, v) = 2a_1 a_2 u + o(|u| + |v|).$$

As before, we intend to apply the parametrized Morse lemma to $\operatorname{Re} \phi$. Setting $\Psi = \Gamma^{-1}$, we get that, around $\mathbf{0}$,

$$\operatorname{Re} \phi \circ \Psi(u, v) = u^2 + h(v)$$

$$\operatorname{Im} \phi \circ \Psi(u, v) = u + g(u, v)$$

with h and g smooth functions which have no terms of order 1 at $\mathbf{0}$.

Assume first that $h \neq 0$. Let $p \geq 2$ be such that $h(v) \sim_0 \alpha_p v^p$ with $\alpha_p \neq 0$. Because $\phi \circ \Psi$ maps \mathbb{R}^2 into $\overline{\mathbb{C}_0}$, we must have that $\alpha_p > 0$ and that p is even. Now, if $\phi \circ \Psi(u, v) \in Q(\tau, \varepsilon)$ with (u, v) sufficiently close to $\mathbf{0}$, then $0 \leq h(v) \leq \varepsilon$ which implies by Lemma 8 that v belongs to some set of measure less than $C\varepsilon^{1/p}$. Moreover, for a fixed value of v , a look at the imaginary part and Lemma 8 yield that u has to belong to some interval of size $C\varepsilon$ and thus we are done with $\kappa = 1 + 1/p$.

Thus, we are lead to study what happens if $h \equiv 0$. The situation is easier if the Taylor expansion of $g(u, v)$ admits some term in v^p . In that case, we may write

$$\operatorname{Im} \phi \circ \Psi(u, v) = u g_1(u, v) + v^p g_2(v)$$

with smooth functions g_1 and g_2 , such that $g_1(0, 0) = 1$ and $g_2(0) \neq 0$. If $\phi \circ \Psi(u, v)$ belongs to $Q(\tau, \varepsilon)$, we conclude from the real part that then $|u| \leq \varepsilon^{1/2}$, and from the imaginary part, we get that, near $\mathbf{0}$,

$$|v^p g_2(v) - \tau| \leq C\varepsilon^{1/2}.$$

By appealing again to Lemma 8, we conclude that v belongs to some set of Lebesgue measure less than $C\varepsilon^{1/2p}$. For a fixed value of v , we look once more at the imaginary part, and obtain that u must belongs to some interval of size $C\varepsilon$. Hence, we are done with $\kappa = 1 + 1/(2p)$.

Therefore, it remains to show that we will always fall into one of the previous cases and compute the value of p . We again recall that the polynomial

$$\Phi(z) = \lambda(1 - z_1 z_2),$$

is a contradiction to the fact that Φ is a minimal Bohr lift of $\varphi \in \mathcal{G}$. More precisely, we are reduced to proving the following lemma.

Lemma 12. *Let $0 < a_2 \leq a_1 \leq 1$ and $a_2 \leq 1 - a_1$. Suppose that $\Phi : \mathbb{D}^2 \rightarrow \mathbb{C}_0$ is the polynomial*

$$(13) \quad \Phi(z) = a_1(1 - z_1) + a_2(1 - z_2) + b_1(1 - z_1)^2 + b_2(1 - z_2)^2 + c(1 - z_1)(1 - z_2),$$

where

$$\operatorname{Re}(b_1) = \frac{a_1}{2} - a_1^2, \quad \operatorname{Re}(b_2) = \frac{a_2}{2} - a_2^2 \quad \text{and} \quad \operatorname{Re}(c) = -2a_1 a_2.$$

Set $\theta_1 = a_2 u + a_2 v$, $\theta_2 = a_1 u - a_1 v$ and

$$\phi(u, v) = \Phi(e^{i\theta_1}, e^{i\theta_2}).$$

Then there does not exist smooth maps $\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $h : \mathbb{R}^2 \rightarrow \mathbb{R}$ so that

$$(14) \quad \operatorname{Re} \phi(u, v) = \gamma(u, v)^2$$

$$(15) \quad \operatorname{Im} \phi(u, v) = \gamma(u, v)h(u, v)$$

except if $\Phi(z) = \frac{1}{2}(1 - z_1 z_2)$. More precisely, if $\Phi(z) \neq \frac{1}{2}(1 - z_1 z_2)$, for any smooth maps $\gamma : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $h : \mathbb{R} \rightarrow \mathbb{R}$, then

- either the Taylor series of $\operatorname{Re} \phi - \gamma^2$ at $\mathbf{0}$ has a non-zero term of order ≤ 5 ,
- or the Taylor series of $\operatorname{Im} \phi - \gamma \cdot h$ at $\mathbf{0}$ has a non-zero term of order ≤ 4 .

The proof of this lemma is rather delicate and will be postponed in Section 6 in order to keep a clearer exposition of the proof of Lemma 10. However, using Lemma 12 we are able to finish this case. Indeed, if $\Gamma(u, v) = (\gamma(u, v), v)$ is the map given by the parametrized Morse lemma and if f_1, f_2 and f_3 are smooth functions such that

$$\operatorname{Re} \phi(u, v) = \gamma(u, v)^2 + f_1(v) \quad \text{and} \quad \operatorname{Im} \phi(u, v) = \gamma(u, v)f_2(u, v) + f_3(v),$$

then Lemma 12 implies that either $f_1(v) \sim_0 \alpha_p v^p$ with $p \leq 5$ or $f_3(v) \sim_0 \beta_p v^p$ with $p \leq 4$. By the considerations above we may conclude that $\kappa = 9/8$ is possible. \square

The case $J = 1$ and dependence, $d \geq 3$. It remains to consider the case $J = 1$, $d \geq 3$ and ℓ_1 is a multiple of ℓ_{d+1} . We shall deduce this case from the case $d = 2$ using the following lemma.

Lemma 13. Let $\varepsilon > 0$, $\tau \in \mathbb{R}$ and $z_3, \dots, z_d \in \mathbb{T}^{d-2}$. Consider the set

$$A_{z_3, \dots, z_d}(\tau, \varepsilon) = \{(z_1, z_2) \in \mathbb{T}^2 : \Phi(z) \in Q(\tau, \varepsilon)\}.$$

Then, for every $w_3, \dots, w_d \in \mathbb{T}^{d-2}$, there exists a neighbourhood $\mathcal{W} \ni (w_3, \dots, w_d)$ in \mathbb{T}^{d-2} such that, for all $(z_3, \dots, z_d) \in \mathcal{W}$ we have

$$A_{z_3, \dots, z_d}(\tau, \varepsilon) \subset A_{w_3, \dots, w_d}(\tau, 2\varepsilon).$$

Proof. Assume that this is not the case. Then there exists a sequence $(z_1^{(k)}, \dots, z_d^{(k)})$ in \mathbb{T}^d such that $z_j^{(k)} \rightarrow w_j$ for $3 \leq j \leq d$ and

$$(z_1^{(k)}, z_2^{(k)}) \in A_{z_3^{(k)}, \dots, z_d^{(k)}}(\tau, \varepsilon) \setminus A_{w_3, \dots, w_d}(\tau, 2\varepsilon).$$

Extracting a subsequence if necessary, we may assume that $z_1^{(k)} \rightarrow w_1$ and $z_2^{(k)} \rightarrow w_2$ for some $(w_1, w_2) \in \mathbb{T}^2$. By the continuity of Φ , this implies that $\Phi(w) \in Q(\tau, \varepsilon) \setminus Q(\tau, 2\varepsilon)$, which is a contradiction. \square

We now set

$$\Psi_{1,2}(z_1, z_2) = \Phi(z_1, z_2, 1, \dots, 1).$$

Since $J(\Phi, w) = 1$, we already know that $a_j > 0$ for all $j = 1, \dots, d$ and hence the variables z_1 and z_2 both appear in the polynomial $\Psi_{1,2}$. Provided $\Psi_{1,2}(z) \neq \lambda_{1,2}(1 - z_1 z_2)$ for some $\lambda_{1,2} \in \mathbb{R}^*$, we know from the case $d = 2$ that there exists a neighbourhood $\mathcal{V} \ni (z_1, z_2)$ in \mathbb{T}^2 and $C > 0$ such that, for any $\tau \in \mathbb{R}$ and every $\varepsilon > 0$,

$$\mathbf{m}_2(\{(z_1, z_2) \in \mathcal{V} : \Phi(z_1, z_2, 1, \dots, 1) \in Q(\tau, 2\varepsilon)\}) \leq C\varepsilon^\kappa$$

with $\kappa = 9/8$. By Lemma 13, there exists a neighbourhood $\mathcal{W} \ni \mathbf{1}$ in \mathbb{T}^{d-2} such that, for any $(z_3, \dots, z_d) \in \mathcal{W}$,

$$\{(z_1, z_2) \in \mathcal{V} : \Phi(z_1, \dots, z_d) \in Q(\tau, \varepsilon)\} \subset \{(z_1, z_2) \in \mathcal{V} : \Phi(z_1, z_2, 1, \dots, 1) \in Q(\tau, 2\varepsilon)\}.$$

This yields

$$\mathbf{m}_d(\{z \in \mathcal{V} \times \mathcal{W} : \Phi(z) \in Q(\tau, \varepsilon)\}) \leq C\varepsilon^\kappa.$$

So the result is proved except if, for every $j < k$, there exists some $\lambda_{j,k} > 0$ such that

$$(16) \quad \Phi(1, \dots, 1, z_j, 1, \dots, 1, z_k, 1, \dots, 1) = \lambda_{j,k}(1 - z_j z_k) = \lambda_{j,k}((1 - z_j) + (1 - z_k) - (1 - z_j)(1 - z_k)).$$

Comparing this with the expansion of Φ near $\mathbf{1}$, we get

$$a_j = a_k = \lambda_{j,k}, \quad b_j = 0, \quad c_{j,k} = -\lambda_{j,k}.$$

Using (10), we may conclude that $a_j = 1/2$ and $c_{j,k} = -1/2$. In total this means that

$$\Phi(z_1, z_2, \dots, z_d) = \frac{1}{2} \sum_{j=1}^d (1 - z_j) - \frac{1}{2} \sum_{1 \leq j < k \leq d} (1 - z_j)(1 - z_k).$$

However,

$$\Phi(-1, -1, \dots, -1) = \frac{2d - 2d(d-1)}{2} = d(2-d) < 0,$$

since $d \geq 3$. Hence (16) is not possible for every $j < k$ and we are done. \square

6. PROOF OF LEMMA 12

We intend to prove this result by contradiction. We require several tedious computations, which can be done either by hand or by a computer algebra system. We have used Xcas, and our file is available for download [3]. In the proof below, we will skip certain computations such as computing Taylor coefficients, simplifying algebraical expressions and solving simple equations. The proof consists of three steps, and in each step we refer to the lines in [3] where the computations are performed.

The idea of the argument is rather easy. We assume that we may factorize $\operatorname{Re} \phi(u, v)$ and $\operatorname{Im} \phi(u, v)$ as (14) and (15) and we write

$$\begin{aligned} \gamma(u, v) &= -2a_1 a_2 u + \gamma_{2,0} u^2 + \gamma_{1,1} uv + \gamma_{0,2} v^2 + \gamma_{3,0} u^3 + \gamma_{2,1} u^2 v + \gamma_{1,2} u v^2 + \gamma_{0,3} v^3 \\ &\quad + \gamma_{4,0} u^4 + \gamma_{3,1} u^3 v + \gamma_{2,2} u^2 v^2 + \gamma_{1,3} u v^3 + \gamma_{0,4} v^4 + o(|u|^5 + |v|^5), \\ h(u, v) &= 1 + h_{1,0} u + h_{0,1} v + h_{2,0} u^2 + h_{1,1} uv + h_{0,2} v^2 + o(|u|^2 + |v|^2). \end{aligned}$$

We already know the first coefficients of γ and h by (11) and (12). We will then compare the Taylor expansions of $\operatorname{Re} \phi(u, v)$ and $\operatorname{Im} \phi(u, v)$ obtained using (13) or using (14) and (15). Looking at all coefficients of a given order, we will get first the value of the coefficients of the Taylor expansions of γ and h of a certain order and also an equation for $\operatorname{Im}(b_1)$, $\operatorname{Im}(b_2)$ and $\operatorname{Im}(c)$.

At one point, we will have more equations than variables. These equations will have to be compatible, and will force $\Phi(z_1, z_2) = (1 - z_1 z_2)/2$, which is equivalent to saying $a_1 = a_2 = 1/2$ and $\operatorname{Im}(b_1) = \operatorname{Im}(b_2) = \operatorname{Im}(c) = 0$. This will imply the desired result.

Step 1. The goal of the first step is to show that if we have $a_1 = a_2 = 1/2$, then we also have $\text{Im}(b_1) = \text{Im}(b_2) = \text{Im}(c) = 0$. In addition to this, we obtain some useful equations for the following steps. [3, Lin. 1–14]

We begin by looking at the coefficients of uv^2 in the real part of $\Phi(u, v)$. Using on the one hand (13) and on the other hand (14) we conclude that

$$\gamma_{0,2} = \frac{-6a_1^3\text{Im}(b_2) - 6a_2^3\text{Im}(b_1) + a_1a_2(a_1 + a_2)\text{Im}(c)}{8a_1a_2}.$$

We then obtain the first equation for $\text{Im}(b_1)$, $\text{Im}(b_2)$ and $\text{Im}(c)$ by looking at the coefficients of v^3 in the real part:

$$(17) \quad a_2^3\text{Im}(b_1) - a_1^3\text{Im}(b_2) + \frac{a_1a_2(a_2 - a_1)}{2}\text{Im}(c) = 0.$$

Since we know the value of $\gamma_{0,2}$, we can get a second equation for $\text{Im}(b_1)$, $\text{Im}(b_2)$ and $\text{Im}(c)$ by looking at the coefficients of v^2 in the imaginary part. Hence we get

$$(18) \quad \frac{4a_1a_2^2 - 3a_2^2}{4a_1}\text{Im}(b_1) + \frac{4a_1^2a_2 - 3a_1^2}{4a_2}\text{Im}(b_2) + \frac{-8a_1a_2 + a_1 + a_2}{8}\text{Im}(c) = 0.$$

By the assumptions on a_1 and a_2 , we know that $2(a_1 + a_2) < 3$ and hence we can solve (17) and (18) with respect to $\text{Im}(b_1)$ and $\text{Im}(b_2)$ to obtain

$$\text{Im}(b_1) = \frac{a_1(2a_2^2 + 2a_1a_2 + a_1 - 2a_2)}{2a_2^2(2a_1 + 2a_2 - 3)}\text{Im}(c) \quad \text{and} \quad \text{Im}(b_2) = \frac{a_2(2a_1^2 + 2a_1a_2 - 2a_1 + a_2)}{2a_1^2(2a_1 + 2a_2 - 3)}\text{Im}(c).$$

In particular, we may conclude that if $\text{Im}(c) = 0$, then we also have $\text{Im}(b_1) = \text{Im}(b_2) = 0$. If we substitute these values into the expression for $\gamma_{0,2}$, we obtain

$$\gamma_{0,2} = \frac{-(a_1 + a_2)^2}{2(2a_1 + 2a_2 - 3)}\text{Im}(c).$$

Now, looking at the coefficient of v^4 in the real part shows that

$$(\gamma_{0,2})^2 = -\frac{a_1a_2(a_1 + a_2)(a_1a_2^2 + a_1^2a_2 - a_1^2 - a_2^2 + a_1a_2)}{4},$$

and this yields our first expression for $\text{Im}(c)^2$,

$$(19) \quad \text{Im}(c)^2 = \frac{-a_1a_2(2a_1 + 2a_2 - 3)^2(a_1a_2^2 + a_1^2a_2 - a_1^2 - a_2^2 + a_1a_2)}{(a_1 + a_2)^3}.$$

From (19), it is evident that if $a_1 = a_2 = 1/2$, then $\text{Im}(c) = 0$. □

Step 2. In this step, we want to show that $a_1 = a_2$. Our first goal is to compute another equation to compare with (19). [3, Lin. 15–21]

We begin by successively looking at the coefficients of u^2v in the real part, uv in the imaginary part and uv^3 in the real part, to obtain

$$\begin{aligned}\gamma_{1,1} &= \frac{a_1 - a_2}{2} \operatorname{Im}(c), \\ h_{0,1} &= \frac{(a_1 - a_2)(4a_1 + 4a_2 - 3)}{4a_1a_2(2a_1 + 2a_2 - 3)} \operatorname{Im}(c), \\ \gamma_{0,3} &= \frac{-a_1 + a_2}{24a_1a_2(2a_1 + 2a_2 - 3)} \times \\ &\quad (20a_1^2a_2^4 + 40a_1^3a_2^3 + 20a_1^4a_2^2 - 12a_1a_2^4 - 54a_1^2a_2^3 - 54a_1^3a_2^2 \\ &\quad - 12a_1^4a_2 + 18a_1a_2^3 + 18a_1^2a_2^2 + 18a_1^3a_2 + 3(a_1 + a_2)^2 \operatorname{Im}(c)^2).\end{aligned}$$

Using these values, we will investigate the coefficient of v^3 in the imaginary part. This term depends indeed only on $\gamma_{1,1}$, $h_{0,1}$ and $\gamma_{0,3}$. Using the above expression, we obtain our second equation on $\operatorname{Im}(c)^2$:

$$(20) \quad \frac{-3(a_1 - a_2)(a_1 + a_2)^2(a_1 + a_2 - 1)}{4a_1a_2(2a_1 + 2a_2 - 3)^2} \operatorname{Im}(c)^2 = \frac{-(a_1 - a_2)(3a_1a_2^2 + 3a_1^2a_2 - a_1^2 - a_2^2 - a_1a_2)}{4}.$$

At this stage, we have to consider several cases. Assuming that $a_1 - a_2 \neq 0$ and $a_1 + a_2 - 1 \neq 0$, we may compute

$$(21) \quad \operatorname{Im}(c)^2 = \frac{-a_1a_2(2a_1 + 2a_2 - 3)^2(3a_1a_2^2 + 3a_1^2a_2 - a_1^2 - a_2^2 - a_1a_2)}{3(a_1 + a_2)^2(a_1 + a_2 - 1)}.$$

The only possibility is that the two values for $\operatorname{Im}(c)^2$ have to coincide. Equating (19) and (21) and simplifying, we obtain

$$\frac{a_1a_2(2a_1 + 2a_2 - 3)^2P(a_1, a_2)}{3(a_1 + a_2)^2(a_1 + a_2 - 1)} = 0,$$

where

$$P(a_1, a_2) = 2a_1^3 + 2a_2^3 + a_1a_2^2 + a_1^2a_2 - 3a_1^2 - 3a_2^2 + 3a_1a_2.$$

Since $2(a_1 + a_2) < 3$, the only possibility is that $P(a_1, a_2)$ vanishes somewhere in the domain

$$\Omega = \{(a_1, a_2) \in (0, 1)^2 : a_2 < a_1, a_2 < 1 - a_1\}.$$

We first look at what happens on the boundary, where we have

$$\begin{aligned}P(a_1, 0) &= 2a_1^3 - 3a_1^2 < 0 \quad \text{provided } a_1 \in (0, 1), \\ P(a_1, a_1) &= 3a_1^2(2a_1 - 1) < 0 \quad \text{provided } a_1 \in (0, 1/2), \\ P(a_1, 1 - a_1) &= -(2a_1 - 1) < 0 \quad \text{provided } a_1 \in (1/2, 1).\end{aligned}$$

Hence, P is negative on the boundary of Ω , except at $(1/2, 1/2)$. Hence, if P vanishes in Ω , then it admits a critical point there. Now, we consider the system

$$\begin{cases} 0 = \frac{\partial P}{\partial a_1}(a_1, a_2) = 6a_1^2 + a_2^2 + 2a_1a_2 - 6a_1 + 3a_2, \\ 0 = \frac{\partial P}{\partial a_2}(a_1, a_2) = a_1^2 + 6a_2^2 + 2a_1a_2 + 3a_1 - 6a_2. \end{cases}$$

The solutions of this system are easily found to be at the intersection of two distinct ellipses. We cannot have more than two points of intersection and we have two trivial solutions, $(0, 0)$ and $(1/3, 1/3)$. Hence, none of the critical points of P are inside Ω . Hence we get a contradiction, and we have finished this case.

Hence we must have $a_1 + a_2 = 1$ or $a_1 = a_2$. Let us now investigate the case $a_1 + a_2 = 1$. Looking at (20), this means that either $a_1 = a_2 = 1/2$ (and we are done) or

$$3a_1a_2^2 + 3a_1^2a_2 - a_1^2 - a_2^2 - a_1a_2 = 0.$$

Taking into account that $a_2 = 1 - a_1$, we get the equation $-4a_1^2 + 4a_1 - 1 = 0$ which admits the single solution $a_1 = 1/2$ and we get the same conclusion. Hence, the only remaining possibility is that $a_1 = a_2$. \square

Step 3. It remains to deal with the case $a_1 = a_2 = a \in (0, 1/2]$. We can no longer use (21) and need to find another equation for $\text{Im}(c)^2$. [3, Lin. 21–30]

We will be looking at the coefficient before v^4 in the imaginary part. By considering $\gamma \times h$, we see that this coefficient is equal to

$$\gamma_{0,4} + \gamma_{0,3}h_{0,1} + \gamma_{0,2}h_{0,2}.$$

Hence, it remains to compute $\gamma_{0,4}$ and $h_{0,2}$. First, we compute some auxiliary values. By looking at u^3 in the real part, u^2 in the imaginary part and u^2v^2 in the real part, respectively, we obtain

$$\begin{aligned} \gamma_{2,0} &= -\frac{a(2a-1)}{3a-4} \text{Im}(c), \\ h_{1,0} &= \frac{(2a-1)(4a-1)}{2a(4a-3)} \text{Im}(c), \\ \gamma_{1,2} &= \frac{a(2a-1)(48a^4 - 72a^3 + 27a^2 + 4\text{Im}(c)^2)}{4(4a-3)^2}. \end{aligned}$$

Knowing these values, we look at the coefficients of uv^4 in the real part and uv^2 and the imaginary part, respectively, to obtain

$$\begin{aligned} \gamma_{0,4} &= -a\text{Im}(c) \frac{32a^5 - 96a^4 + 90a^3 + 12a\text{Im}(c)^2 - 27a^2 - 6\text{Im}(c)^2}{6(4a-3)^3}, \\ h_{0,2} &= (-2a+1) \frac{128a^5 - 240a^4 + 144a^3 + 16a\text{Im}(c)^2 - 27a^2 - 8\text{Im}(c)^2}{8a(4a-3)^2}. \end{aligned}$$

Finally, we investigate the coefficient of v^4 in the imaginary part to obtain the equation

$$a(2a-1) \frac{16a^4 - 32a^3 + 15a^2 + 4\text{Im}(c)^2}{4(4a-3)^2} \text{Im}(c) = 0.$$

Now, if $\text{Im}(c) = 0$ and $a_1 = a_2 = a$, it follows at once from (19) that $a = 1/2$, since $a \in (0, 1/2]$. Conversely, we divide away $\text{Im}(c)$ and solve for $\text{Im}(c)^2$ to obtain the equation

$$\text{Im}(c)^2 = -a^2 \frac{(4a-3)(4a-5)}{4}.$$

Now, this has to be equal to (19), and we find

$$\frac{a(2a-1)(4a-3)^2}{8} = -\frac{a^2(4a-3)(4a-5)}{4}.$$

Here the only solutions are $a = 0$ and $a = 3/4$, neither of which belong to $(0, 1/2]$. Hence the assumption $\text{Im}(c) \neq 0$ must be wrong and we conclude $a_1 = a_2 = 1/2$. \square

7. REMARKS AND FURTHER EXAMPLES

If we look more closely at the map Φ defined in Lemma 9 (the negative part of Theorem 3), then we may observe that these counterexamples all satisfy

$$\begin{aligned} \text{Re}\Phi(e^{i\theta_1}, \dots, e^{i\theta_d}) &= \frac{\theta_1^2}{2} + o(\theta_1^2) \\ \text{Im}\Phi(e^{i\theta_1}, \dots, e^{i\theta_d}) &= -\theta_1 + o(|\theta_1|). \end{aligned}$$

Hence, $J(\Phi, \mathbf{1}) = 1$ and, using the terminology of the Section 5, we have dependence. Our next results shows that we may also have non-compactness if $J(\Phi, w) = 0$ for some $w \in \mathbb{T}^d$.

Theorem 14. *There are polynomials φ with unrestricted range, of any complex dimension $d \geq 2$ and of any complex degree ≥ 4 for which the corresponding composition operator \mathcal{C}_φ is non-compact and such that they admit a minimal Bohr lift Φ satisfying $J(\Phi, w) = 0$ for any $w \in \mathbb{T}^d$ with $\text{Re}\Phi(w) = 0$.*

Proof. Let $\delta > 0$ and define

$$\Phi(z_1, z_2) = 2(1 - z_1) + (1 - z_1)^2(1 - \delta(1 - z_2) - \delta(1 - z_1)(1 - z_2)).$$

Let $\varphi(s) = \Phi(p_1^{-s}, p_2^s)$. Clearly Φ is a minimal Bohr lift of φ . Then a computation shows that

$$\text{Re}\Phi(z_1, z_2) = 2(1 - \cos x) \left((1 - \cos x)(1 + 2\delta(\sin x \sin y + (1 - \cos y) \cos x)) + \delta(1 - \cos y) \right).$$

Clearly, for small enough $\delta > 0$ this quantity is non-negative. Hence $\varphi \in \mathcal{G}$ and $J(\Phi, (1, 1)) = 0$ because of the relation between a_1 and b_1 . Considerations similar to those of Lemma 9 show that \mathcal{C}_φ cannot be compact. To produce a counterexample with degree 3 and a bigger complex dimension d , we may simply replace $(1 - z_2)$ by

$$\frac{1}{d-1} \cdot \sum_{j=2}^d (1 - z_j)$$

in the definition of Φ . The production of examples with degree ≥ 5 is easier. We may just set

$$\Phi(z) = (1 - z_1) + \frac{1}{2}(1 - z_1)^2 + \delta(1 - z_1)^4 P(z),$$

where $P(z) = P(z_1, z_2, \dots, z_d)$ is any polynomial. The proof follows now that of Lemma 9. \square

The reason the first counterexample in Theorem 14 works is that we have a cancellation of the term $(1 - \cos x) \sin x \sin y$. It seems difficult to obtain the same cancellation if we restrict ourself to degree 3 and require $J = 0$.

Question. Is it possible to construct a counterexample of degree 3 with $J = 0$?

An answer to the question would in a certain sense improve the optimality of Lemma 10, but it would not yield the complete answer to which Dirichlet polynomials in \mathcal{G} induce compact composition operators. Indeed, the natural next point of investigation would be this: What happens when the “quartic form” is degenerate?

In this case, terms of degree 5 also have to disappear. This follows by the mapping properties and the argument is identical to the one used to show that degree 3 terms disappear in the case $J = 0$ given above. Hence we are reduced to studying a “sextic form”.

Our counterexamples can be modified to work in this case, but they now have degree 6 and 7. Degree ≤ 3 will also easily reduce to the case of Theorem 2 in the same manner as $J = 0$ did for degree ≤ 2 . However, the cases with degree 4 and 5 would need further investigation. Even if we could solve this case, we would need to investigate the case when the “sextic form” is degenerate and this leads to the “octic form” and so on.

Remark. The previous counterexample shows that we cannot deduce Theorem 2 from Theorem 3. Indeed, it is easy to construct symbols $\varphi \in \mathcal{G}$ which may be written $\varphi(s) = \sum_{j=1}^d P_j(p_j^{-s})$ and such that $J(\Phi, \mathbf{1}) = 0$ for Φ a minimal Bohr lift of φ . Indeed, we may consider

$$P_j(z) = (1 - z) + \frac{1}{2}(1 - z)^2 + \delta(1 - z)^4 Q_j(z)$$

where Q_j is an arbitrary polynomial and $\delta > 0$ is sufficiently small. Then \mathcal{C}_φ is compact by Theorem 2 if $d \geq 2$ but this cannot be deduced from Lemma 10.

This construction can be generalized to show that Theorem 2 can handle a variety of different interesting cases not covered by Theorem 3. In fact, given any d positive integers k_j , we may find a polynomial $\Phi(z) = \sum_{j=1}^d \Phi_j(z_j)$ which is a minimal Bohr lift of some $\varphi \in \mathcal{G}$, with $\operatorname{Re} \Phi(z_1, \dots, z_d) = 0$ if and only if $z = \mathbf{1}$ and here we have the expansion

$$\begin{aligned} \operatorname{Re} \Phi(e^{i\theta_1}, \dots, e^{i\theta_d}) &= \sum_{j=1}^d \theta_j^{2k_j} + o\left(\sum_{j=1}^d \theta_j^{2k_j}\right), \\ \operatorname{Im} \Phi(e^{i\theta_1}, \dots, e^{i\theta_d}) &= \sum_{j=1}^d a_1^{(j)} \theta_j + o\left(\sum_{j=1}^d |\theta_j|\right). \end{aligned}$$

As remarked upon in the proof of Theorem 2, we must have $a_1^{(j)} > 0$. The construction of such a polynomial is immediate from our next result.

Lemma 15. *There is a polynomial $\Phi : \mathbb{C} \rightarrow \mathbb{C}$ which satisfy $\operatorname{Re} \Phi(e^{ix}) = (1 - \cos x)^k$, for any $k \in \mathbb{N}$.*

Proof. Fix N with $k \leq 2N$, and for real numbers a_n and b_n consider

$$\Phi(z) = \sum_{n=1}^N \frac{(-1)^{n-1}}{2^n} (a_n(1-z)^{2n-1} - b_n(1-z)^{2n}).$$

Our first goal is to expand the real part of $\Phi(e^{ix})$ as degree $2N$ polynomial in $(1 - \cos x)$ with no constant term. To this end, we compute

$$(1 - e^{ix})^{2n-1} = e^{i(2n-1)x/2} (e^{-ix/2} - e^{ix/2})^{2n-1} = e^{i(2n-1)x/2} 2^{2n-1} (-1)^n i \sin^{2n-1}\left(\frac{x}{2}\right).$$

We use $2 \sin^2(x/2) = 1 - \cos x$, and obtain
(22)

$$\operatorname{Re}(1 - e^{ix})^{2n-1} = 2^{2n-1}(-1)^{n-1} \sin^{2n-1}\left(\frac{x}{2}\right) \sin\left(nx - \frac{x}{2}\right) = (-1)^{n-1} 2^n (1 - \cos x)^n \frac{\sin\left(nx - \frac{x}{2}\right)}{2 \sin\left(\frac{x}{2}\right)}.$$

Similarly, we obtain

$$(23) \quad \operatorname{Re}(1 - e^{ix})^{2n} = 2^{2n}(-1)^n \sin^{2n}\left(\frac{x}{2}\right) \cos(nx) = (-1)^n 2^n (1 - \cos x)^n \cos(nx).$$

To continue the computations, we introduce the Chebyshev polynomials

$$U_n(y) = \sum_{j=0}^n (-2)^j \frac{(n+j+1)!}{(n-j)!(2j+1)!} (1-y)^j, \quad \text{and} \quad T_n(y) = n \sum_{j=0}^n (-2)^j \frac{(n+j-1)!}{(n-j)!(2j)!} (1-y)^j.$$

The Chebyshev polynomials are relevant due to the formulas $\sin nx = \sin(x)U_{n-1}(\cos x)$ and $\cos nx = T_n(\cos x)$. We record the following coefficients.

$$\begin{aligned} u_{n-2}^{(n)} &= (-2)^{n-1} \left(-\frac{(2n-1)(n-2)}{2} \right), & u_{n-1}^{(n)} &= (-2)^{n-1} (2n), & u_n^{(n)} &= (-2)^{n-1} (-2), \\ t_{n-1}^{(n)} &= (-2)^{n-1} (n), & t_n^{(n)} &= (-2)^{n-1} (-1). \end{aligned}$$

Now, we rewrite (23) as

$$\operatorname{Re}(1 - e^{ix})^{2n} = (-1)^n 2^n (1 - \cos x)^n T_n(\cos x),$$

which is then clearly a degree $2n$ polynomial in $(1 - \cos x)$ with no constant term. For (22) we have to work a bit more, so we first compute

$$\frac{\sin\left(nx - \frac{x}{2}\right)}{2 \sin\left(\frac{x}{2}\right)} = \frac{\sin(nx) \cos\left(\frac{x}{2}\right) - \cos(nx) \sin\left(\frac{x}{2}\right)}{2 \sin\left(\frac{x}{2}\right)} = \cos^2\left(\frac{x}{2}\right) U_{n-1}(\cos x) - \frac{T_n(\cos x)}{2},$$

which implies that we may rewrite (22) as

$$\operatorname{Re}(1 - e^{ix})^{2n-1} = (-1)^{n-1} 2^n (1 - \cos x)^n \left(\left(1 - \frac{1 - \cos x}{2}\right) U_{n-1}(\cos x) - \frac{T_n(\cos x)}{2} \right).$$

Again we observe that this is a polynomial of degree $2n$ in $(1 - \cos x)$ with no constant term. In total, we have

$$\operatorname{Re}\Phi(1 - e^{ix}) = \sum_{m=1}^{2N} c_m (1 - \cos x)^m = \sum_{n=1}^N (a_n P_n(1 - \cos x) + b_n Q_n(1 - \cos x)),$$

where

$$\begin{aligned} P_n(y) &= \sum_{j=0}^n d_j^{(n)} y^{n+j} = y^n \left(\left(1 - \frac{y}{2}\right) U_{n-1}(1-y) - \frac{T_n(1-y)}{2} \right), \\ Q_n(y) &= \sum_{j=0}^n e_j^{(n)} y^{n+j} = y^n T_n(1-y). \end{aligned}$$

Given any choice of c_m (for instance $c_m = 0$ for $m \neq k$ and $c_k = 1$), we now have $2N$ linear equations and $2N$ unknowns, a_n and b_n for $1 \leq n \leq N$. We will now show that this system can always be solved.

We first observe that a_n and b_n only have an effect on c_m when $n \leq m \leq 2n$. Ordering the unknowns as $a_N, b_N, a_{N-1}, b_{N-1}, \dots, a_1, b_1$ and the datas as $c_{2N}, c_{2N-1}, \dots, c_1$, this means that the matrix of our system can be written in upper triangular block form, where the blocks on the diagonal are

$$\begin{pmatrix} e_n^{(n)} & d_n^{(n)} \\ e_{n-1}^{(n)} & d_{n-1}^{(n)} \end{pmatrix}, \quad n = N, N-1, \dots, 1.$$

We know that $e_{n-1}^{(n)} = t_{n-1}^{(n)}$ and $e_n^{(n)} = t_n^{(n)}$, which we recorded above. It is now easy to verify that

$$d_{n-1}^{(n)} = u_{n-1}^{(n)} - \frac{u_{n-2}^{(n)}}{2} - \frac{t_{n-1}^{(n)}}{2} = (-2)^{n-1} \left(\frac{3n}{2} + \frac{(2n-1)(n-2)}{4} \right),$$

$$d_n^{(n)} = u_n^{(n)} - \frac{u_{n-1}^{(n)}}{2} - \frac{t_n^{(n)}}{2} = (-2)^{n-1} \left(-\frac{3}{2} - n \right).$$

Hence we are reduced to considering the equation

$$0 = \frac{d_{n-1}^{(n)} e_n^{(n)} - d_n^{(n)} e_{n-1}^{(n)}}{4^{n-1}} = \left(\frac{3n}{2} + \frac{(2n-1)(n-2)}{4} \right) (-1) - \left(-\frac{3}{2} - n \right) n = n^2 - \frac{(2n-1)(n-2)}{4},$$

which has no integer solutions, and we are done. \square

The construction of Φ with specific expansion facilitated by Lemma 15 will be used in the next section to prove Corollary 4.

8. APPROXIMATION NUMBERS

In this section, we consider only the case $c_0 = 0$. We intend to estimate the decay of $a_n(\mathcal{C}_\varphi)$ for maps φ which are, in a certain sense, regular at their boundary points. For this we need as previously a careful inspection of the behaviour of the Bohr lift Φ near these boundary points.

Definition. Suppose that $\varphi(s) = c_1 + \sum_{n=2}^N c_n n^{-s} \in \mathcal{G}$, and that φ has complex dimension d and unrestricted range. Let Φ be a minimal Bohr lift of φ and let $w \in \mathbb{T}^d$ be such that $\operatorname{Re} \Phi(w) = 0$. We say that φ is *boundary regular* at w if there exist independent linear forms ℓ_1, \dots, ℓ_d on \mathbb{C}^d , even integers $k_1 \geq k_2 \geq \dots \geq k_d$ and real numbers b_1, \dots, b_d, τ with $b_1 \neq 0$ such that

$$(24) \quad \operatorname{Re} \Phi(e^{i\theta_1} w_1, \dots, e^{i\theta_d} w_d) = \ell_1(\theta)^{k_1} + \dots + \ell_d(\theta)^{k_d} + \sum_{j=1}^d o\left(\ell_j^{k_j}(\theta)\right)$$

$$(25) \quad \operatorname{Im} \Phi(e^{i\theta_1} w_1, \dots, e^{i\theta_d} w_d) = \tau + b_1 \ell_1(\theta) + \dots + b_d \ell_d(\theta) + o\left(\sum_{j=1}^d |\ell_j(\theta)|\right).$$

We define the *compactness index* of φ at w as

$$\eta_{\varphi, w} = \left(\sum_{j=2}^d \frac{1}{k_j} \right) \times \frac{k_1}{2(k_1 - 1)}.$$

If every boundary point is boundary regular, we say that φ is *boundary regular*.

The proof of Theorem 2 then shows that given a boundary regular map φ , the composition operator \mathcal{C}_φ is compact if and only if $d \geq 2$. We shall now assume that there is only one point $w \in \mathbb{T}^d$ such that $\operatorname{Re} \Phi(z) = 0$. In this case, we let the *compactness index* of φ be $\eta_\varphi := \eta_{\varphi, w}$.

The main theorem of this section now reads.

Theorem 16. *Let $\varphi(s) = c_1 + \sum_{n=2}^N c_n n^{-s} \in \mathcal{G}$ have unrestricted range and complex dimension d . Let Φ be a minimal Bohr lift and assume that there exists a unique $w \in \mathbb{T}^d$ such that $\operatorname{Re} \Phi(w) = 0$. Suppose moreover that φ is boundary regular at w . Then*

$$\left(\frac{1}{n}\right)^{\eta_\varphi} \ll a_n(\mathcal{C}_\varphi) \ll \left(\frac{\log n}{n}\right)^{\eta_\varphi}.$$

This statement may be applied to several cases.

Corollary 17. *Let $\varphi(s) = c_1 + \sum_{n=2}^N c_n n^{-s} \in \mathcal{G}$ have unrestricted range and complex dimension d . Let Φ be a minimal Bohr lift of φ and assume that there exists a unique $w \in \mathbb{T}^d$ such that $\operatorname{Re} \Phi(w) = 0$ and that $J(\Phi, w) = d$. Then*

$$\left(\frac{1}{n}\right)^{(d-1)/2} \ll a_n(\mathcal{C}_\varphi) \ll \left(\frac{\log n}{n}\right)^{(d-1)/2}.$$

Proof. Under these assumptions, φ is boundary regular at w with $k_1 = \dots = k_d = 2$. \square

In particular, this corollary covers the result of Queffélec and Seip for linear symbols (3), as well as the map φ_1 given in (4). We may also apply Theorem 16 to the maps considered in Theorem 2. In this case, one has simply $\ell_j(\theta) = \theta_j$ (up to a reordering of the terms).

Another interesting application of Theorem 16 is that we may distinguish the Schatten classes of bounded linear operators on \mathcal{H}^2 using composition operators, as mentioned in the introduction.

Proof of Corollary 4. Let $p' < q'$ and $\varepsilon > 0$ be such that $p \leq p'/2$ and $(\frac{1}{2} + \varepsilon)q' \leq q$. Then let $d \geq 2$ and $k \geq 2$ even such that

$$p' < \frac{d-1}{k} < q' \text{ and } \frac{1}{2} < \frac{k}{2(k-1)} < \frac{1}{2} + \varepsilon.$$

By Lemma 15, we know that there exists a boundary regular polynomial $\Phi : \mathbb{T}^d \rightarrow \mathbb{C}_0$ such that $\operatorname{Re} \Phi(w) = 0$ if and only if $w = \mathbf{1}$ and

$$\Phi(e^{i\theta_1}, \dots, e^{i\theta_d}) = \theta_1^k + \dots + \theta_d^k + o(\theta_1^k) + \dots + o(\theta_d^k).$$

Letting $\varphi \in \mathcal{G}$ any map such that Φ is a minimal Bohr lift of φ , we immediately get

$$\left(\frac{1}{n}\right)^{\frac{d-1}{k} \times \frac{k}{2(k-1)}} \ll a_n(\mathcal{C}_\varphi) \ll \left(\frac{\log n}{n}\right)^{\frac{d-1}{k} \times \frac{k}{2(k-1)}},$$

which completes the proof. \square

Theorem 16 may be also applied to many other maps. We will consider here the map φ_2 given in (4). Its boundary regularity is different than that of φ_1 , and hence the degree of compactness is also different.

Example. Let $\varphi_2(s) = 13/2 - 4 \cdot 2^{-s} - 4 \cdot 3^{-s} + 2 \cdot 6^{-s}$ as in (4) and let Φ be its minimal Bohr lift. It can be shown that $\operatorname{Re} \Phi(w) = 0$ for $w \in \mathbb{T}^2$ if and only if $w = (1, 1)$, and

$$\begin{aligned} \operatorname{Re} \Phi(e^{i\theta_1}, e^{i\theta_2}) &= \ell_1(\theta)^4 + \ell_2(\theta)^2 + o(\ell_1^4(\theta)) + o(\ell_2^2(\theta)), \\ \operatorname{Im} \Phi(e^{i\theta_1}, e^{i\theta_2}) &= -2\ell_1(\theta) + o(|\ell_1(\theta)| + |\ell_2(\theta)|), \end{aligned}$$

where $\ell_1(\theta) = \theta_1 + \theta_2$ and $\ell_2(\theta) = \theta_1 - \theta_2$. Hence $\eta_{\varphi_2} = (1/2) \times (4/6) = 1/3$.

The remaining part of this section is devoted to the proof of Theorem 16. We use the scheme introduced by Queffélec and Seip in [11] in the context of Dirichlet series (see also [12] for similar works on the classical Hardy space of the disk). Their method is based on Carleson measures, interpolation sequences and model spaces. In Subsection 8.1, we survey these tools and give a couple of lemmas. Subsection 8.2 is devoted to the proof of the upper bound, in a more general context, whereas Subsection 8.3 will be devoted to the lower bound.

8.1. Tools.

The Hyperbolic Metric. The pseudo-hyperbolic metric on the half-plane \mathbb{C}_0 is defined by

$$\rho(z, w) = \left| \frac{z - w}{z + \overline{w}} \right| = \frac{1 - e^{-d(z, w)}}{1 + e^{d(z, w)}}$$

where $d(z, w)$ is the hyperbolic distance between two points z and w in \mathbb{C}_0 . The hyperbolic length of a curve $\Gamma \subset \mathbb{C}_0$ is given by the integral

$$L_p(\Gamma) = \int_{\Gamma} \frac{|dz|}{\operatorname{Re} z}.$$

Carleson Measures and Interpolating Sequences. Let H be a Hilbert space of functions defined on some measurable set Ω in \mathbb{C} . A non-negative Borel measure μ on Ω is a *Carleson measure* for H if there exists some constant $C > 0$ such that

$$\int_{\Omega} |f(z)|^2 d\mu(z) \leq C \|f\|_H^2,$$

for every f in H . The smallest possible C will be called the *Carleson norm* of μ with respect to H and will be denoted by $\|\mu\|_{\mathcal{C}, H}$.

We also assume that the linear point evaluation is bounded at any $z \in \Omega$. Then H admits a reproducing kernel $K_z^H \in H$ for any $z \in \Omega$ which satisfies $f(z) = \langle f, K_z^H \rangle$ for every $f \in H$. We then say that a sequence $Z = (z_m)$ of distinct points in Ω is a *Carleson sequence* for H if the measure

$$\mu_{Z, H} := \sum_m \|K_{z_m}^H\|_H^{-2} \delta_{z_m}$$

is a *Carleson measure* for H .

We say that a sequence $Z = (z_m)$ of distinct points in Ω is an *interpolating sequence* for H if the interpolation problem $f(z_m) = a_m$ has a solution $f \in H$ whenever the admissibility condition

$$\sum_m |a_m|^2 \|K_{z_m}^H\|_H^{-2} < \infty$$

is satisfied. By the open mapping theorem, if Z is an interpolating sequence for H , there is a constant $C > 0$ such that we can solve $f(z_m) = a_m$ with f satisfying

$$\|f\|_H \leq C \left(\sum_m |a_m|^2 \|K_{z_m}^H\|_H^{-2} \right)^{1/2}.$$

The smallest constant C with this property will be called the *constant of interpolation* of Z and will be denoted by $M_H(Z)$.

We shall consider the two spaces $H = \mathcal{H}^2$ and $H = H^2(\mathbb{T}^d)$. Then we have, respectively, $\Omega = \mathbb{C}_{1/2}$ and $\Omega = \mathbb{D}^d$, and moreover

$$\|K_s^{\mathcal{H}^2}\|^{-2} = [\zeta(2\operatorname{Re} s)]^{-1} \quad \text{and} \quad \|K_z^{H^2(\mathbb{T}^d)}\|^{-2} = \prod_{j=1}^d (1 - |z_j|^2).$$

We will need the three following lemmas.

Lemma 18. *Let μ be a Borel measure on $\overline{\mathbb{C}_0}$, let $\sigma \in (0, 1)$ and $R > 0$. Assume that μ is supported on the rectangle $0 \leq \operatorname{Re} s \leq \sigma$, $|\operatorname{Im} s| \leq R$. Then*

$$\|\mu\|_{\mathcal{C}, \mathcal{H}^2} \ll_R \sup_{\varepsilon > 0, \tau \in \mathbb{R}} \frac{\mu(Q(\tau, \varepsilon))}{\varepsilon} \leq 2 \sup_{\varepsilon \in (0, \sigma), \tau \in \mathbb{R}} \frac{\mu(Q(\tau, \varepsilon))}{\varepsilon}.$$

Proof. The first inequality is Lemma 2.3 in [11] (the involved constant does not depend on $\sigma \in (0, 1)$). The second follows from the inequality

$$\sup_{\tau \in \mathbb{R}} \frac{\mu(Q(\tau, 2^{k+1}\sigma))}{2^{k+1}\sigma} \leq \sup_{\tau \in \mathbb{R}} \frac{\mu(Q(\tau, 2^k\sigma))}{2^k\sigma},$$

valid for any $k \geq 0$. Indeed, for any $\tau \in \mathbb{R}$ and any $k \geq 0$, we may find $\tau_1, \tau_2 \in \mathbb{R}$ such that

$$\mu(Q(\tau, 2^{k+1}\sigma)) = \mu(Q(\tau_1, 2^k\sigma)) + \mu(Q(\tau_2, 2^k\sigma)),$$

since the support of μ is contained in $0 \leq \operatorname{Re} s \leq \sigma$. □

Lemma 19. *Let $\nu > 0$. There exists $C > 0$ such that, for any $\delta \in (0, 1/\nu)$, $M_{\mathcal{H}^2}(S_\delta) \leq C$ where $S_\delta = (s_m)_{m=1}^{1/\delta}$ with $s_m = \frac{1}{2} + \nu\delta + im\delta$.*

Proof. The proof of this lemma can be found in [11, Sec 8.2]. □

Lemma 20. *Let $C_1, C_2 > 0$. There exists $D > 0$ such that for any $\delta > 0$ and any (finite) sequence*

$$Z = (Z(\alpha)) = ((1 - \rho_1(\alpha))e^{i\theta_1(\alpha)}, \dots, (1 - \rho_d(\alpha))e^{i\theta_d(\alpha)})$$

in \mathbb{D}^d satisfying

- $\sup_{j=1, \dots, d} |\theta_j(\alpha) - \theta_j(\beta)| \geq C_1\delta$, when $\alpha \neq \beta$,
- $\rho_j(\alpha) \leq C_2\delta$, for any α and $j = 1, \dots, d$,

we have $\|\mu_{Z, H^2(\mathbb{T}^d)}\|_{\mathcal{C}, H^2(\mathbb{T}^d)} \leq D$.

Proof. To each point $Z(\alpha)$, we associate a rectangle R_α on the distinguish boundary \mathbb{T}^d centered at

$$\left(\frac{z_1(\alpha)}{|z_1(\alpha)|}, \dots, \frac{z_d(\alpha)}{|z_d(\alpha)|} \right),$$

with side lengths $2(1-|z_1(\alpha)|), \dots, 2(1-|z_d(\alpha)|)$. By Chang's characterization of Carleson measures on the polydisc (see [4] or [6]), it is enough to show that we for all open sets \mathcal{U} of \mathbb{T}^d have

$$\sum_{R_\alpha \subset \mathcal{U}} \mathbf{m}_d(R_\alpha) \leq D \mathbf{m}_d(\mathcal{U}).$$

If R is some rectangle in \mathbb{T}^d and $\lambda > 0$, denote by λR the rectangle with the same center and side lengths multiplied by λ . Then our assumptions on Z imply that there exists some $\lambda \in (0, 1)$ depending only on C_1 and C_2 such that the rectangles R_α are pairwise disjoint. Thus

$$\sum_{R_\alpha \subset \mathcal{U}} \mathbf{m}_d(\mathcal{U}) \leq \sum_{R_\alpha \subset \mathcal{U}} \frac{1}{\lambda^d} \mathbf{m}_d(\lambda R_\alpha) \leq \frac{1}{\lambda^d} \mathbf{m}_d \left(\bigcup_{R_\alpha \subset \mathcal{U}} \lambda R_\alpha \right) \leq \frac{1}{\lambda^d} \mathbf{m}_d(\mathcal{U}),$$

which completes the proof with $D = 1/\lambda^d$. \square

The Queffélec–Seip Method. We have to introduce additional conventions. For $\varphi \in \mathcal{G}$ and Ω a compact subset of \mathbb{C}_0 , we denote by $\mu_{\varphi, \Omega}$ the non-negative Borel measure on $\overline{\mathbb{C}_0}$ defined by

$$\mu_{\varphi, \Omega}(E) := \mathbf{m}_\infty(\{z \in \mathbb{T}^\infty : \Phi(z) \in E \setminus \Omega\}).$$

Next, assume that φ has complex dimension d and Bohr lift $\Phi : \mathbb{C}^d \rightarrow \mathbb{C}$. Let $S = (s_m)$ be a sequence of n points in $\mathbb{C}_{1/2}$ and let Z be a finite sequence of points in \mathbb{D}^d such that $\Phi(Z) = S - \frac{1}{2}$. We set

$$N_\Phi(s_m; Z) := \sum_{z \in Z \cap \Phi^{-1}(s_m - 1/2)} \|K_z^{H^2(\mathbb{T}^d)}\|^{-2}.$$

We state Theorem 4.1 of [11] as the forthcoming lemma (we have modified it slightly to take into account our normalization).

Lemma 21. *Let $\varphi(s) = \sum_{n=1}^\infty c_n n^{-s} \in \mathcal{G}$ such that $\varphi(\mathbb{C}_0)$ is bounded.*

- (a) *Let $\sigma > 0$ and Ω be a compact subset of $\overline{\mathbb{C}_\sigma}$. Let B be a Blaschke product of degree n on \mathbb{C}_0 whose zeros lie in Ω . Then*

$$a_n(\mathcal{E}_\varphi) \leq \left(\sup_{s \in \Omega} |B(s)|^2 \zeta(1 + 2\sigma) + \sup_{\varepsilon > 0, \tau \in \mathbb{R}} \frac{\mu_{\varphi, \Omega}(Q(\tau, \varepsilon))}{\varepsilon} \right)^{1/2}.$$

- (b) *Assume that φ has complex dimension d . Let S and Z be finite sets in respectively $\mathbb{C}_{1/2}$ and \mathbb{D}^d such that $\Phi(Z) = S - \frac{1}{2}$. Then*

$$a_n(\mathcal{E}_\varphi) \geq [M_{\mathcal{H}^2}(S)]^{-1} \|\mu_{Z, H^2(\mathbb{T}^d)}\|_{\mathcal{E}, H^2(\mathbb{T}^d)}^{-1/2} \inf_m [N_\Phi(s_m; Z) \zeta(2\operatorname{Re} s_m)]^{1/2}.$$

8.2. The Upper Bound.

Let $\varphi(s) = \sum_{n=1}^{\infty} c_n n^{-s} \in \mathcal{G}$ and suppose that $\varphi(\mathbb{C}_0)$ bounded. By Lemma 5, \mathcal{C}_φ is compact if and only if $\mu_\varphi(Q(\tau, \varepsilon)) = o(\varepsilon)$ uniformly in $\tau \in \mathbb{R}$. We are planning to get an upper bound of $a_n(\mathcal{C}_\varphi)$ depending on the behaviour of $\sup_{\tau \in \mathbb{R}} \mu_\varphi(Q(\tau, \varepsilon))$ with respect to ε and on the size of the image of φ near a boundary point.

Thus, let Φ be a Bohr lift of φ . We define κ_φ as the infimum of those $\kappa \geq 1$ such that there exists a constant $C > 0$ such that, for every $\tau \in \mathbb{R}$ and every $\varepsilon > 0$,

$$\mathbf{m}_\infty(\{z \in \mathbb{T}^\infty : \Phi(z) \in Q(\tau, \varepsilon)\}) \leq C\varepsilon^\kappa.$$

Assume now that there exists a unique $w \in \mathbb{T}^\infty$ such that $\operatorname{Re} \Phi(w) = 0$ and write $\Phi(w) = i\tau$. Let ω_φ be the infimum of the positive ω such that, for any $s \in \mathbb{C}_0$,

$$|\operatorname{Im} \varphi(s) - \tau|^\omega \leq C \left(\operatorname{Re} \varphi(s) - \frac{1}{2} \right).$$

Theorem 22. *Let $\varphi(s) = \sum_{n=1}^{\infty} c_n n^{-s} \in \mathcal{G}$ with $\varphi(\mathbb{C}_0)$ bounded, let Φ be a Bohr lift of φ and assume that there is a unique $w \in \mathbb{T}^\infty$ such that $\operatorname{Re} \Phi(w) = 0$. Then*

$$a_n(\mathcal{C}_\varphi) \ll \begin{cases} \exp(-\lambda n^{-1/2}) & \text{if } \omega_\varphi \leq 1, \\ \left(\frac{\log n}{n}\right)^{(\kappa_\varphi - 1) \times \frac{\omega_\varphi}{2(\omega_\varphi - 1)}} & \text{if } \omega_\varphi > 1. \end{cases}$$

Here λ is some positive constant depending on φ .

This theorem illustrates the following general principle for composition operators (valid beyond \mathcal{H}^2): The more restricted the image of the symbol is, the more compact the associated composition operator is. In particular, the case $\omega_\varphi = 1$ (the range of φ is contained in an angle) is reminiscent from [12, Thm. 1.2] where a similar result was obtained for composition operators on $H^2(\mathbb{D})$.

Before we embark upon the proof of Theorem 22, we first employ it to deduce the upper bound of Theorem 16.

Final part in the proof of the upper bound of Theorem 16. Suppose that $\varphi \in \mathcal{G}$ is a boundary regular Dirichlet polynomial, and assume that $\operatorname{Re} \Phi(\mathbf{1}) = 0$. We write

$$\begin{aligned} \operatorname{Re} \Phi(e^{i\theta_1}, \dots, e^{i\theta_d}) &= \ell_1(\theta)^{k_1} + \dots + \ell_d(\theta)^{k_d} + \sum_{j=1}^d o(\ell_j^{k_j}(\theta)), \\ \operatorname{Im} \Phi(e^{i\theta_1}, \dots, e^{i\theta_d}) &= \tau + b_1 \ell_1(\theta) + \dots + b_d \ell_d(\theta) + o\left(\sum_{j=1}^d |\ell_j(\theta)|\right), \end{aligned}$$

with $k_1 \geq \dots \geq k_d$ and $b_1 \neq 0$. The proof of Theorem 2 shows that we have $\kappa_\varphi \geq 1 + \sum_{j=2}^d 1/k_j$.

Now, let us write the Taylor expansion of $\operatorname{Re} \Phi$ and $\operatorname{Im} \Phi$ near $\mathbf{1}$, but also now for a point belonging to the unit polydisc. Writing

$$\Phi(z) = \sum_{j=1}^d a_j(1 - z_j) + o\left(\sum_{j=1}^d |1 - z_j|\right)$$

and $z = ((1 - \rho_1)e^{i\theta_1}, \dots, (1 - \rho_d)e^{i\theta_d})$, it is easy to get

$$\begin{aligned} \operatorname{Re}\Phi(z) &= a_1\rho_1 + \dots + a_d\rho_d + \ell_1(\theta)^{k_1} + \dots + \ell_d(\theta)^{k_d} + o\left(\sum_{j=1}^d (\rho_j + \ell_j^{k_j}(\theta))\right), \\ \operatorname{Im}\Phi(z) &= \tau + b_1\ell_1(\theta) + \dots + b_d\ell_d(\theta) + o\left(\sum_{j=1}^d |\ell_j(\theta)|\right). \end{aligned}$$

Recalling that $a_j \geq 0$ for $j = 1, \dots, d$, it is easy to conclude that there exists a neighbourhood $\mathcal{U} \ni \mathbf{1}$ in $\overline{\mathbb{D}^d}$ and $C > 0$ such that, for all $z \in \mathcal{U}$,

$$|\operatorname{Im}\Phi(z) - \tau|^{k_1} \leq C\operatorname{Re}\Phi(z).$$

Outside \mathcal{U} , $\operatorname{Re}\Phi(z)$ is bounded away from 0, and $|\operatorname{Im}\Phi(z) - \tau|$ is here trivially majorized. Hence, the upper bound of Theorem 16 follows from Theorem 22. \square

Let us now turn to the proof of Theorem 22. The proof will be preceded by two lemmas. The first one is inspired by Lemma 3.1 in [12].

Lemma 23. *Let Ω be a bounded domain in \mathbb{C}_0 whose boundary is a piecewise regular Jordan curve Γ , with $L_p(\Gamma) \geq 1$. Let s_1, \dots, s_n be points in Γ such that the hyperbolic length of the curve between any two points s_j and s_{j+1} is equal to $L_p(\Gamma)/n$, $1 \leq j \leq n$, where $s_{n+1} = s_1$. Let B be the Blaschke product of degree n whose zeros are precisely s_1, \dots, s_n . Then, for any $s \in \Omega$,*

$$|B(s)| \leq \exp\left(-C\frac{n}{L_p(\Gamma)}\right).$$

Proof. By the maximum principle, it is sufficient to prove this inequality for $s \in \Gamma$. In this case, we know that there exists some $j \in \{1, \dots, n\}$ such that $d(s, s_j) \leq L_p(\Gamma)/n$, from which we deduce that

$$d(s, s_k) \leq \frac{L_p(\Gamma)}{n}(1 + |k - j|)$$

for any $k = 1, \dots, n$. Using the link between the pseudo hyperbolic distance and the hyperbolic distance, we deduce that

$$|B(s)| \leq \prod_{j=1}^n \left(\frac{1 - e^{-j\frac{L_p(\Gamma)}{n}}}{1 + e^{-j\frac{L_p(\Gamma)}{n}}} \right).$$

By a Riemann sum argument, this means that

$$|B(s)| \leq \exp\left(-n \int_0^1 \ln\left(\frac{1 - e^{-xL_p(\Gamma)}}{1 + e^{-xL_p(\Gamma)}}\right) dx\right) \leq \exp\left(-\frac{n}{L_p(\Gamma)} \int_{e^{-L_p(\Gamma)}}^1 \frac{1}{y} \ln\left(\frac{1+y}{1-y}\right) dy\right),$$

and we get the desired conclusion, since by assumption $L_p(\Gamma) \geq 1$. \square

Hence, we require estimates of the hyperbolic length of some curves which are linked to the way that φ touches the boundary. Such estimates are contained in the following result.

Lemma 24. *Let $\omega \geq 1$, $\sigma \in (0, 1/2)$ and $C > 1$. Consider*

$$\Omega_{\omega, \sigma, C} = \{s \in \mathbb{C}_0 : |\operatorname{Im} s|^\omega \leq C \operatorname{Re}(s), \sigma \leq \operatorname{Re} s \leq C\}.$$

Let $\Gamma_{\omega, \sigma, C}$ denote the boundary of $\Omega_{\omega, \sigma, C}$. Then

$$L_p(\Gamma_{\omega, \sigma, C}) \ll_{\omega, C} \begin{cases} \left(\frac{1}{\sigma}\right)^{\frac{\omega-1}{\omega}} & \text{if } \omega > 1, \\ -\ln(\sigma) & \text{if } \omega = 1. \end{cases}$$

Proof. Consider the curves

$$\begin{aligned} \Gamma_1 &= \{s \in \mathbb{C}_0; \operatorname{Re} s = \sigma, |\operatorname{Im} s| \leq C^{1/\omega} (\operatorname{Re} s)^{1/\omega}\}, \\ \Gamma_2 &= \{s \in \mathbb{C}_0; \operatorname{Re} s = C, |\operatorname{Im} s| \leq C^{1/\omega} (\operatorname{Re} s)^{1/\omega}\}, \\ \Gamma_3 &= \{s \in \mathbb{C}_0; \sigma \leq \operatorname{Re} s \leq C, |\operatorname{Im} s| = C^{1/\omega} (\operatorname{Re} s)^{1/\omega}\}. \end{aligned}$$

Clearly, $\Gamma_{\omega, \sigma, C} \subset \Gamma_1 \cup \Gamma_2 \cup \Gamma_3$ and it is sufficient to prove the corresponding inequalities for Γ_j , $j = 1, 2, 3$. Firstly, $L_p(\Gamma_2) \ll_{\omega, C} 1$. Regarding Γ_1 ,

$$L_p(\Gamma_1) = \int_{-C^{1/\omega} \sigma^{1/\omega}}^{C^{1/\omega} \sigma^{1/\omega}} \frac{dy}{\sigma} \ll_{\omega, C} \left(\frac{1}{\sigma}\right)^{\frac{\omega-1}{\omega}}$$

which is even a stronger inequality than required when $\omega = 1$. Finally,

$$L_p(\Gamma_3) \ll_{\omega, C} \int_{\sigma}^C \frac{\sqrt{1+x^{\frac{2}{\omega}-1}}}{x} dx \ll_{\omega, C} \begin{cases} -\ln(\sigma) & \text{if } \omega = 1, \\ \left(\frac{1}{\sigma}\right)^{\frac{\omega-1}{\omega}} & \text{if } \omega > 1. \end{cases}$$

The last estimate follows from inspecting the integrand near $x = 0$, since $\sigma \in (0, 1)$. \square

Proof of Theorem 22. Let $\sigma \in (0, 1)$ and $n \geq 1$. Without loss of generality, we may assume that $\Phi(\mathbf{1}) = 0$. Keeping the notations of Lemma 24, there exists $C > 0$ such that

$$\varphi(\mathbb{C}_0) - \frac{1}{2} \subset \{s \in \mathbb{C}_0 : 0 \leq \operatorname{Re} s \leq \sigma\} \cup \Omega_{\omega, \varphi, \sigma, C}.$$

Let B be a Blaschke product of degree n defined as in Lemma 23 with $\Omega_{\omega, \varphi, \sigma, C}$. Enlarging C if necessary, we may always assume that $L_p(\Gamma_{\omega, \varphi, \sigma, C}) \geq 1$, so that the assumptions of Lemma 23 are satisfied. The set

$$\Omega = \overline{\left\{ \varphi(s) - \frac{1}{2} : \operatorname{Re} \varphi(s) \geq \frac{1}{2} + \sigma \right\}}$$

is a compact subset of \mathbb{C}_0 , and we may apply part (a) of Lemma 21. Since $\Omega \subset \Omega_{\omega, \varphi, \sigma, C}$, we obtain

$$\sup_{s \in \Omega} |B(s)|^2 \leq \exp\left(-2C' \frac{n}{L_p(\Gamma_{\omega, \varphi, \sigma, C})}\right).$$

Moreover, $\zeta(1+2\sigma) \ll 1/\sigma$. Finally, using Lemma 18, we obtain

$$\|\mu_{\varphi, \Omega}\|_{\mathcal{C}, \mathcal{H}^2} \ll \sup_{\varepsilon \in (0, \sigma), \tau \in \mathbb{R}} \frac{\mu_{\varphi, \Omega}(Q(\tau, \varepsilon))}{\varepsilon} \ll \sigma^{\kappa_{\varphi}-1}.$$

We will now optimize the choice of σ with respect to n . When $\omega_\varphi > 1$, we set

$$\sigma = \rho \left(\frac{\log n}{n} \right)^{\frac{\omega_\varphi}{\omega_\varphi - 1}},$$

where ρ is some numerical parameter to be chosen later. Then

$$\sup_{s \in \Omega} |B(s)|^2 \zeta(1 + 2\sigma) \leq \exp \left(-2C' \rho^{\frac{\omega_\varphi - 1}{\omega_\varphi}} \log n \right) \cdot \frac{1}{\sigma} \ll \left(\frac{\log n}{n} \right)^{\frac{\omega_\varphi}{\omega_\varphi - 1} (\kappa_\varphi - 1)},$$

provided $\rho > 0$ is sufficiently large. When $\omega_\varphi \leq 1$, we set $\sigma = \exp(-\rho n^{-1/2})$, so that

$$\sup_{s \in \Omega} |B(s)|^2 \zeta(1 + 2\sigma) \leq \exp \left(-\frac{C''}{\rho} n^{1/2} + \rho n^{1/2} \right),$$

and the result is proved provided $\rho > 0$ is sufficiently small. \square

Remark. Our method of proof also shows, provided $\varphi(\mathbb{C}_0)$ is bounded and $\kappa_\varphi > 1$, that

$$a_n(\mathcal{C}_\varphi) \leq \left(\frac{\log n}{n} \right)^{\frac{\kappa_\varphi - 1}{2}}.$$

Indeed, we apply the same method with $\Omega_{\sigma, C} = \{s \in \mathbb{C}_0 : \sigma \leq \operatorname{Re} s \leq C, |\operatorname{Im} s| \leq C\}$ which satisfies $L_p(\Gamma) \ll_C \sigma^{-1}$. The rest of the proof remains unchanged.

8.3. The Lower Bound.

Let $\varphi \in \mathcal{G}$ satisfying the assumptions of Theorem 16 and let us assume that around $\mathbf{1}$, Φ satisfies (24) and (25). Let $\nu > 0$. For $\delta \in (0, 1/\nu)$, we consider the sequence $S_\delta = (s_m)$, given by

$$s_m = \frac{1}{2} + \nu\delta + im\delta, \quad \text{where } 1 \leq m \leq \left(\frac{1}{\delta} \right)^{1 - \frac{1}{k_1}}.$$

We intend to apply part (b) of Lemma 21. We will require the construction of preimages of $S_\delta - 1/2$ by Φ , and the inverse function theorem will provide the solution.

Lemma 25. *Let $\varphi \in \mathcal{G}$ satisfy the assumptions of Theorem 16. Then there exist $\nu_0, C_1, C_2 > 0$ such that for all $\nu \geq \nu_0$ and every $\delta \in (0, 1/\nu)$, there exists a finite sequence $Z_\delta = (Z(\alpha))$ in \mathbb{D}^d with*

$$Z(\alpha) = [(1 - \rho_1(\alpha))e^{i\theta_1(\alpha)}, \dots, (1 - \rho_d(\alpha))e^{i\theta_d(\alpha)}]$$

such that

- for any $\alpha \neq \beta$, we have $\sup_{j=1, \dots, d} |\theta_j(\alpha) - \theta_j(\beta)| \geq C_1 \delta$,
- for any α and any $j = 1, \dots, d$, we have $C_2^{-1} \delta \leq \rho_j(\alpha) \leq C_2 \delta$,
- $\Phi(Z_\delta) = S_\delta - 1/2$ and, for any $1 \leq m \leq \left(\frac{1}{\delta} \right)^{1 - \frac{1}{k_1}}$, the equation $\Phi(Z(\alpha)) = s_m - \frac{1}{2}$ has at least $\prod_{j=2}^d \left\lfloor \left(\frac{1}{\delta} \right)^{1 - \frac{1}{k_j}} \right\rfloor$ solutions.

Proof. We start as in the deduction of the upper bound in Theorem 16 from Theorem 22, writing

$$\begin{aligned}\operatorname{Re}\Phi(z) &= a_1\rho_1 + \cdots + a_d\rho_d + \ell_1(\theta)^{k_1} + \cdots + \ell_d(\theta)^{k_d} + o\left(\sum_{j=1}^d \left(\rho_j + \ell_j^{k_j}(\theta)\right)\right), \\ \operatorname{Im}\Phi(z) &= \tau + b_1\ell_1(\theta) + \cdots + b_d\ell_d(\theta) + o\left(\sum_{j=1}^d |\ell_j(\theta)|\right),\end{aligned}$$

for $z = ((1 - \rho_1)e^{i\theta_1}, \dots, (1 - \rho_d)e^{i\theta_d})$. To simplify the notations, we use the (linear) change of variables $u_j = \ell_j(\theta)$. We also set

$$\Lambda = \mathbb{N}^d \cap \prod_{j=1}^d \left[1, \left(\frac{1}{\delta}\right)^{1-\frac{1}{k_j}}\right]$$

and, for $\alpha \in \Lambda$ and $j = 2, \dots, d$ we let $\rho_j(\alpha) = \delta$ and $u_j(\alpha) = \alpha_j\delta$.

Setting $m = \alpha_1$, we want to find $Z(\alpha)$ such that $\operatorname{Re}\Phi(Z(\alpha)) = \nu\delta$ and $\operatorname{Im}\Phi(Z(\alpha)) = m\delta$. It remains to determine $\rho_1(\alpha)$ and $u_1(\alpha)$. We rewrite this system as

$$(26) \quad \begin{cases} f_\alpha(\rho_1, u_1)\rho_1 + g_\alpha(\rho_1, u_1)u_1^{k_1} &= \nu\delta + d_\alpha \\ h_\alpha(\rho_1, u_1)u_1 &= m\delta + e_\alpha \end{cases}$$

where f_α , g_α and h_α are smooth functions depending only on $\alpha_2, \dots, \alpha_d$ and there exists a neighbourhood $\mathcal{U} \ni (0, 0)$ so that for every $(\rho, u) \in \mathcal{U}$,

$$|f_\alpha(\rho, u) - a_1| \ll \delta, \quad |g_\alpha(\rho, u) - 1| \ll \delta \quad \text{and} \quad |h_\alpha(\rho, u) - 1| \ll \delta.$$

Here, the open set \mathcal{U} and the involved constants are uniform with respect to α , $\nu \geq 1$ and $\delta \in (0, 1/\nu)$. Moreover, the real numbers d_α and e_α satisfy

$$\begin{aligned}d_\alpha &\ll \sum_{j=2}^d \delta + \sum_{j=2}^d \left(\left(\frac{1}{\delta}\right)^{1-\frac{1}{k_j}}\right)^{k_j} \ll \delta \\ e_\alpha &\ll \delta \sum_{j=2}^d \left(\frac{1}{\delta}\right)^{1-\frac{1}{k_j}} \ll \delta^{\frac{1}{k_1}}.\end{aligned}$$

We now apply the inverse function theorem to solve the system (26). Provided ν is large enough, we get a solution $(\rho_1(\alpha), u_1(\alpha))$ satisfying $\sup(\rho_1(\alpha), |u_1(\alpha)|) \ll \delta^{1/k_1}$. In this case, the involved constant depends on ν , but it is uniform with respect to α and δ .

Now, a look at the first equation of (26) shows that we in fact have the more precise inequality $\delta \ll \rho_1(\alpha) \ll \delta$, provided ν is sufficiently large, and this is independent of α and $\delta \in (0, 1)$. Looking now at the second equation of (26), if $\alpha \neq \beta \in \Lambda$ satisfy $\alpha_j = \beta_j$ for $j \geq 2$, so that $e_\alpha = e_\beta$ and $h_\alpha = h_\beta$, then $|u_1(\alpha) - u_1(\beta)| \gg \delta$.

Hence, we have obtained $\prod_{j=2}^d \left[\left(\frac{1}{\delta}\right)^{1-\frac{1}{k_j}}\right]$ solutions to the equation $\Phi(Z(\alpha)) = s_m$, and they satisfy the conclusions of Lemma 25 since the inequalities on $u_j(\alpha)$ are also valid for $\theta_j(\alpha)$ up to a constant depending only of Φ . \square

Final part in the proof of the lower bound of Theorem 16. We apply Lemma 21 to S_δ and Z_δ given by the previous lemma, for

$$\delta = \left(\frac{1}{n}\right)^{\frac{k_1}{k_1-1}},$$

so that S_δ has cardinal number equal to n . Since $M_{\mathcal{H}^2}(S_\delta) \ll 1$ and $\|\mu_{Z, H^2(\mathbb{T}^d)}\|_{\mathcal{C}, H^2(\mathbb{T}^d)} \ll 1$ by Lemma 19 and Lemma 20, it remains to estimate the sum $N_\Phi(s_m; Z)\zeta(2\operatorname{Re} s_m)$ for any m . Using the fact that $\rho_j(\alpha) \gg \delta$ for any $j = 1, \dots, d$ and any α , we obtain

$$N_\Phi(s_m; Z)\zeta(2\operatorname{Re} s_m) \gg \left(\frac{1}{\delta}\right)^{\sum_{j=2}^d \left(1 - \frac{1}{k_j}\right)} \cdot \delta^d \cdot \delta^{-1} \gg \delta^{\sum_{j=2}^d \frac{1}{k_j}} \gg \left(\frac{1}{n}\right)^{\left(\sum_{j=2}^d \frac{1}{k_j}\right) \times \frac{k_1}{k_1-1}},$$

and we are done. □

Remark. We may modify the proof of Theorem 16 so that we do not assume that there exists a unique $w \in \mathbb{T}^d$ such that $\operatorname{Re} \Phi(w) = 0$. Suppose that φ is boundary regular at any $w \in \mathbb{T}^d$ such that $\operatorname{Re} \Phi(w) = 0$. Define now the *compactness index* of φ as the real number

$$\eta_\varphi(s) = \inf\{\eta_{\varphi, w} : \operatorname{Re} \Phi(w) = 0\}.$$

It should be observed that this infimum is in fact a minimum. Indeed, our assumptions imply that the points $w \in \mathbb{T}^d$ such that $\operatorname{Re} \Phi(w) = 0$ are isolated. Theorem 16 remains true with this new definition of η_φ .

REFERENCES

1. F. Bayart, *Hardy spaces of Dirichlet series and their composition operators*, Monatsh. Math. **136** (2002), 203–236.
2. ———, *Compact composition operators on a Hilbert space of Dirichlet series*, Illinois J. Math. **47** (2003), no. 3, 725–743.
3. F. Bayart and O. F. Brevig, *Xcas file for Lemma 11*, <http://bayart.perso.math.cnrs.fr/nonlin.xws>.
4. B. Berndtsson, A. Chang, and K-C. Lin, *Interpolating sequences in the polydisc*, Trans. of the AMS **302** (1987), 161–169.
5. J. W. Bruce and P. J. Giblin, *Curves and Singularities: A geometrical introduction to singularity theory*, Cambridge University Press, 1992.
6. A. Chang, *Carleson measure on the bi-disc*, Ann. of Math. **109** (1979), 613–620.
7. C. Finet and H. Queffélec, *Numerical range of composition operators on a Hilbert space of Dirichlet series*, Linear Algebra Appl. **377** (2004), 1–10.
8. C. Finet, H. Queffélec, and A. Volberg, *Compactness of composition operators on a Hilbert space of Dirichlet series*, J. Funct. Anal. **211** (2004), no. 2, 271–287.
9. J. Gordon and H. Hedenmalm, *The composition operators on the space of Dirichlet series with square summable coefficients*, Michigan Math. J. **46** (1999), no. 2, 313–329.
10. G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, Oxford University Press, 1979.
11. H. Queffélec and K. Seip, *Approximation numbers of composition operators on the H^2 space of Dirichlet series*, J. Funct. Anal. **268** (2015), 1612–1648.
12. H. Queffélec and K. Seip, *Decay rates for approximation numbers of composition operators*, J. Anal. Math. **125** (2015), 371–399.

CLERMONT UNIVERSITÉ, UNIVERSITÉ BLAISE PASCAL, LABORATOIRE DE MATHÉMATIQUES, UMR 6620 - CNRS,
CAMPUS DES CÉZEAUX, 3, PLACE VASARELY, TSA 60026, CS 60026, 63178 AUBIÈRE CEDEX

E-mail address: bayart@math.univ-bpclermont.fr

DEPARTMENT OF MATHEMATICAL SCIENCES, NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY (NTNU),
NO-7491 TRONDHEIM, NORWAY

E-mail address: ole.brevig@math.ntnu.no