

The Joint Association of Low Vitamin D and Vitamin K Status with Blood Pressure and Hypertension

Running title: vitamin D and K status with blood pressure

Adriana J. van Ballegooijen¹, Aivaras Cepelis¹, Marjolein Visser^{1,2}, Ingeborg A. Brouwer¹,
Natasja M. van Schoor³, Joline W. Beulens^{3,4}

1) Department of Health Sciences, and the EMGO⁺ Institute for health and care research, VU University, Amsterdam

2) Department of Internal Medicine, section Nutrition and Dietetics, VU University Medical Center, Amsterdam

3) Department of Epidemiology and Biostatistics, EMGO⁺ Institute for health and care research, VU University Medical Center, Amsterdam

4) Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht

Corresponding author:

A.J. van Ballegooijen, Department of Health Sciences, VU University Amsterdam, the Netherlands, De Boelelaan 1085, 1081 HV Amsterdam, phone: +31 20 5983291, email: hanne.van.ballegooijen@vu.nl

Total word count: 5682; Number of tables: 3, number of figures: 0

Abstract (words: 247)

Low vitamin D and K status are both associated with an increased cardiovascular risk. New evidence from experimental studies on bone health suggest an interaction between vitamin D and K, however a joint association with vascular health outcomes is largely unknown. To prospectively investigate whether the combination of low vitamin D and K status is associated with higher systolic and diastolic blood pressure in 402 participants and with incident hypertension in 231 participants free of hypertension at baseline. We used data from a subsample of the Longitudinal Aging Study Amsterdam, a population-based cohort of Dutch participants 55-65 years. Vitamin D and K status were assessed by 25-hydroxyvitamin D [25(OH)D] and dephosphorylated uncarboxylated matrix gla protein (dp-ucMGP) concentrations (high dp-ucMGP is indicative for low vitamin K status) in stored samples from 2002/03. Vitamin D and K status were categorized into: 25(OH)D $<50/\geq 50$ nmol/L and median dp-ucMGP $<323/\geq 323$ pmol/L. During a median follow-up of 6.4 years, 62% of the participants (n=143) developed hypertension. The combination of low vitamin D and K status was associated with increased systolic 4.8 mmHg (95% CI 0.1-9.5) and diastolic 3.1 mmHg (0.5-5.7) blood pressure compared to high vitamin D and K status (P-for interaction=0.013 for systolic blood pressure, 0.068 for diastolic blood pressure). A similar trend was seen for incident hypertension: hazard ratio 1.62 (95% CI 0.96-2.73) for the low vitamin D and K group. The combination of low vitamin D and K status was associated with increased blood pressure and a trend for greater hypertension risk.

Keywords: 25-hydroxyvitamin D, vitamin K status, blood pressure, incident hypertension, epidemiology

Introduction

Prospective studies have shown that both low vitamin D measured as low 25-hydroxyvitamin D concentrations and low vitamin K status measured as high dephosphorylated uncarboxylated matrix Gla protein (dp-ucMGP) concentrations are associated with an increased risk of vascular disease and adverse cardiovascular events.¹⁻³ Both vitamin D and K are involved in bone metabolism and potentially to mineral deposition in the vasculature, which in turn could lead to calcification in valves and vessels. Vascular calcification in different beds is linked with vascular stiffness, increased blood pressure^{4,5} and is an important predictor of coronary heart disease.⁵

Vitamin K is involved in vascular calcification through its role as a cofactor in the carboxylation, i.e. activation, of the calcification inhibitor matrix Gla protein (MGP). A loss-of-function mutation of the MGP gene has been shown to cause rapid calcification of arteries and, consequently, death in mice.⁶ Studies showed that vitamin K deficiency, induced by vitamin K antagonists, may lead to under-carboxylated MGP and increased vascular calcification.⁷ Under-carboxylated or inactive MGP, measured as high dp-ucMGP concentrations, is a marker of low vitamin K status and has a high affinity to bind calcium, thereby stimulating the incorporation into extracellular matrix of soft tissues.⁸ High levels of dp-ucMGP have been associated with increased vascular calcification² and cardiovascular disease risk⁹, but not in all populations.¹⁰

Previous prospective studies have indicated that lower vitamin D concentrations are associated with greater hypertension risk.¹¹ At physiological levels, vitamin D can upregulate the expression of MGP in vascular smooth muscle cells, which, in turn, could increase the demand for vitamin K and potential benefits of MGP on vascular health.¹²⁻¹⁶ A randomized controlled trial in postmenopausal women found that supplementation of vitamin K₁, D and minerals prevented a decrease in elastic properties of the carotid artery after 3 years compared to vitamin

D and minerals or placebo alone.¹⁷ This suggests that adequate levels of both vitamin D and vitamin K are needed for optimal vascular health . However the combined association of vitamin D and vitamin K on vascular health is currently unknown. Previous analysis in our study population have indicated that both 25(OH)D and dp-ucMGP were associated with cardiovascular disease risk.^{18,19}

We now investigated the joint association of serum 25(OH)D and plasma dp-ucMGP, with systolic and diastolic blood pressure, and the incidence of hypertension in older adults in the Netherlands.

For Hypertension Peer Review. Do not distribute.
Destroy after use.

Materials and Methods

Study Design and Population

The Longitudinal Aging Study Amsterdam started in 1992 with the goal to determine predictors and consequences of ageing focusing on physical, cognitive, emotional and social aspects in a Dutch population. A detailed description of the cohort sampling and data collection procedures has been provided elsewhere.²⁰ In short, LASA personnel used registers of 11 municipalities in three geographical areas in the Netherlands to recruit males and females aged 55 to 85 years. The sample was randomly selected from municipal registries in 1992, with an oversampling of the oldest old and older men. The initial response rate was 60 % (n=3805).

For the current study, we used data from second LASA cohort, which started in 2002/03 with adults aged 55 to 65 years. Altogether, 1002 males and females were recruited for the baseline examination using the same procedures as in the first LASA cohort. Follow-up took place in 3 cycles: 2005/06, 2008/09 and 2011/12.

For the present analysis, we excluded participants with a missing baseline blood sample (n=254), or vitamin D and vitamin K measurements (n=53), or using anti-hypertensive medication at baseline (n=188). For the blood pressure analysis, we excluded participants with missing information on all follow-up blood pressure measurements (n=64) or with missing information on medication use for all measurements (n=39) or using vitamin K antagonists (n=2) resulting in 402 participants for the final analysis. For participants who started using anti-hypertensive medication during follow-up, we only included blood pressure measurements up to the point of anti-hypertensive medication use.

For the of incident hypertension analysis, we further excluded participants with prevalent hypertension at baseline based on blood pressure (n=171) resulting in a sample of 231 participants.

All participants provided written informed consent. The study was conducted according to the Declaration of Helsinki and has been approved by the Medical Ethical Committee of the VU University Medical Center Amsterdam.

Measurements of vitamin D and K

LASA study personnel collected morning blood samples during a medical interview in a non-fasted state and samples were shipped to the VU University Medical Center in 2002-2003.

Blood samples were centrifuged and stored at -80° until analysis in 2010-2011. For vitamin D status, serum 25(OH)D concentrations were determined by the Endocrine Laboratory of the VU University Medical Center using a radioimmunoassay (DiaSorin, Stillwater, Minnesota, USA). The inter-assay coefficient of variation was 10.0%.

For vitamin K, plasma dp-ucMGP concentrations were determined using a sandwich ELISA with the capture of antibody directed against the non-phosphorylated MGP sequence 3-15 (mAb-dpMGP; VitaK BV, Maastricht, the Netherlands). High levels of dp-ucMGP reflect a low vitamin K status.²¹ The inter-assay coefficient of variation was 9.9%.

Ascertainment of blood pressure and incident hypertension

During baseline and 3 follow-up visits, systolic and diastolic blood pressure was taken after 5 minutes of rest at the upper left arm with participants in a seated position, using an oscillometric monitor for a total of 4 times taken 1-minute apart (model HEM-706; Omron Corporation,

Tokyo, Japan). We used the last 3 tests to calculate mean blood pressure. We defined hypertension as systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg or use of blood pressure lowering medication.²² Trained LASA interviewers assessed medication use during each medical interview by medication inventory of all medicines that were currently taken by the participant, including name, dose, frequency of intake, and duration of use. Medications were recorded into official international ATC-codes; anti-hypertensive drugs were classified with the codes C01 to C10 (cardiovascular system).

Other Measurements

LASA interviewers obtained comprehensive data on participants' demographics, anthropometrics and co-morbid conditions. Height was measured to the nearest 0.1 cm using a stadiometer. Weight was measured without clothes and shoes to the nearest 0.1 kg using a calibrated bathroom scales (Seca, model 100, Lameris, Utrecht, the Netherlands). Body mass index (BMI) was calculated by dividing body weight (kg) by height (m) squared kg/m^2 .

A self-administered questionnaire was used to assess participants self-reported education level, smoking status, alcohol use and physical activity level. Level of education was reported on a 9-category scale. We distinguished education into three categories: low (elementary school or less), medium (lower vocational or general intermediate education) and high (intermediate vocational education, general secondary school, higher vocational education, college or university). Smoking status was categorized as never, former and current smoker. Alcohol use, based on the number of days and the amount of alcohol used per week, was characterized into 3 categories: none/light (0-3 glasses/week), moderate (4-7 glasses/week), excessive or very excessive (≥ 8 glasses/week) according to the Garretsen index.²³ Physical activity was assessed

using the validated LASA Physical Activity Questionnaire in minutes per day by multiplying the frequency and duration of each activity in the previous two weeks, and summing these values across activities. Total physical activity was divided into tertiles: low, moderate and high.²⁴

Season of blood collection was calculated from baseline blood collection date using meteorological seasons and divided into: autumn, winter and spring/summer (combined due to few summer measurements (n=5)). Total cholesterol was determined from fasting serum samples using enzymatic colorimetry method with a Hitachi 747 analyzer (Roche diagnostics, Mannheim, Germany). Furthermore, we estimated glomerular filtrate rate (eGFR), derived from serum creatinine using the Chronic Kidney Disease (CKD) Epidemiology Collaboration 2009 equation (21), as an indicator of kidney function. Parathyroid hormone was measured with a second generation immunoradiometric assay that measures intact parathyroid hormone (Inctar Corp., Stillwater, MN, USA) with an inter-assay CV of 5%. All biochemical analyses were carried out by the Endocrine Laboratory of the VU University Medical Center Amsterdam.

Statistical Methods

We categorized serum 25(OH)D and dp-ucMGP into 4 categories based on the clinical cut-off value of 50nmol/L 25(OH)D and median dp-ucMGP (323 pmol/L) because there is currently no cut-off value defined for high dp-ucMGP. The reference group was the group with 25(OH)D \geq 50 nmol/L and dp-ucMGP <323 pmol/L considered high vitamin D status and high vitamin K status. Baseline characteristics are presented as mean and standard deviation for continuous variables and number and percentage for the categorical variables vitamin D and K combined categories. Skewed variables are presented as median and interquartile range.

To investigate the association between 25(OH)D and dp-ucMGP categories with systolic and diastolic blood pressure we used a generalized estimating equation (GEE) model to estimate regression coefficients and 95% confidence intervals. For the incident hypertension analyses, we reported unadjusted incidence rates of hypertension per 100 person-years. Risk time was calculated from the baseline examination until the examination at which hypertension was first diagnosed, the last examination before loss to follow-up or the last LASA cycle 2011-2012. We used Cox proportional hazard models to estimate hazard ratios (HRs) and 95% confidence intervals for 25(OH)D and dp-ucMGP categories with incident hypertension. Given the high incidence of hypertension in our study population (62%), a Cox regression model is preferred because it takes risk-time into account and other methods may present difficulties when the outcome is very prevalent, experience the risk of over-dispersion or require the estimation of additional parameters for bands of time.²⁵

We used nested models to adjust for potential confounders. A minimally adjusted model included age (year) and sex (model 1). The fully adjusted model included age and sex, as well as BMI (kg/m^2), alcohol consumption (none-light/moderate/excessive), smoking (never/former/current), education (low/medium/high), season of blood collection (3 categories), total cholesterol (nmol/L) concentrations and estimated glomerular filtration rate ($\text{ml}/\text{min}/1.73\text{m}^2$). Further, we also added PTH as a potential mediator (pmol/L). We imputed missing data for 16 respondents with at least 1 baseline characteristic missing using 5 iterations using Rubin's rule.²⁶

To determine the joint association between vitamin D and vitamin K with blood pressure and incident hypertension, we estimated the presence of interaction. For the association between vitamin D and vitamin K with blood pressure we added vitamin D (2 groups) and vitamin K (2

groups) to the GEE model plus the product terms for each category (3 dummies) calculating the Likelihood-ratio test. For the Cox regression, interaction between vitamin D and vitamin K status was estimated by adding the product terms for vitamin D and vitamin K. A P-value <0.10 was considered statistically significant. We also calculated interaction on a continuous scale. In linear regression (GEE), the regression coefficient of the product term reflects interaction as departure from additivity. In the Cox regression the product term reflects interaction as departure from multiplicativity (multiplicative scale vs. additive scale). However, It has been argued that interaction estimated as departure from additivity better reflects biologic interaction.²⁷

Interaction on an additive scale for relative risks from Cox regression can be estimated by calculating the Relative Excess Risk due to Interaction (RERI). $RERI = RR_{A+B+} - RRA+B- - RRA-B+ + 1$; A=vitamin D group, B=vitamin K group. A RERI of 0 reflects the absence of interaction, meaning no evidence of additive interaction.²⁷ We performed data analysis using IBM SPSS Statistics 23.0 for Windows 7 (SPSS Inc. Chicago, Illinois) and all P-values are 2-sided.

Results

Study population

Among the 402 included LASA participants at baseline, the mean age was 59.7 ± 2.9 years and 55% (n=222) were female. The mean body mass index (BMI) was 26.4 ± 3.7 kg/m² and 27% were current smokers and 42% had a lower education level.

Excluded participants with a vitamin D and K sample (n= 293) had a higher BMI (28.2 vs. 26.4 kg/m²), lower 25(OH)D (54 vs. 59 nmol/L), higher dp-ucMGP (442 vs. 349 pmol/L)

concentrations, used more medication (74% vs 45%) and had lower estimated glomerular filtration rate values (81 vs 84 ml/min/1.73m²) ($P < 0.05$).

Serum 25(OH)D and dp-ucMGP concentrations

Plasma 25(OH)D and dp-ucMGP levels were approximately normally distributed with mean concentrations of 59 ± 21 nmol/L and 349 ± 209 pmol/L, respectively.

Participants in the combined low vitamin D and low vitamin K status group (25(OH)D < 50 nmol/L and dp-ucMGP ≥ 323 pmol/L) represented 19% of the sample and had higher BMI, waist circumference and PTH concentrations compared to the reference group (25(OH)D ≥ 50 nmol/L and dp-ucMGP < 323 pmol/L) (Table 1). Characteristics for plasma 25(OH)D dp-ucMGP categories separately are available in Supplemental Table S1).

Associations of vitamin D and vitamin K with blood pressure

Systolic and diastolic blood pressure were normally distributed with mean measurements of 137 ± 23 mmHg and 83 ± 12 mmHg at baseline, respectively. Systolic blood pressure was strongly correlated with diastolic blood pressure (correlation coefficient=0.82). Both systolic and diastolic blood pressure were highest in the low vitamin D and low vitamin K status group (Table 2). The average increase in systolic and diastolic blood pressure during follow-up was 7.4 ± 17.3 and 2.5 ± 9.7 .

In the GEE model, after adjusting for age and sex, participants in the group of low vitamin D and low vitamin K status had a 6.5 mmHg (1.9-11.1) higher systolic blood pressure compared to the reference group with high vitamin D and high vitamin K status (25(OH)D ≥ 50 nmol/L and dp-ucMGP < 323 pmol/L). After further adjustment for potential confounders

(model 2) the results remained significant with a 4.8 mmHg (0.1-9.5) difference between the low vitamin D and low vitamin K status group compared with the reference group. Similarly, diastolic blood pressure was significantly higher 4.2 mmHg (1.5-6.8) in the low vitamin D and low vitamin K status group compared to the reference, which attenuated in the fully adjusted model to a 3.1 mmHg (0.5-5.7) difference, but remained statistically significant. Diastolic blood pressure was also significantly higher in the group high vitamin D and low vitamin K status: 2.8 mmHg (0.5-5.0) compared to the reference group. No other significant associations between vitamin D and K groups were observed in relation to blood pressure. For systolic blood pressure the P-value for interaction was 0.013 for the fully adjusted model and 0.068 for diastolic blood pressure. Adjusting for PTH as a potential mediator slightly attenuated the results 4.5(-0.3,9.2) and 2.6 (0.0, 5.3) for systolic and diastolic blood pressure, respectively, comparing the low vitamin D and K category compared to the reference group. We additionally adjusted for waist circumference as a measure of central obesity, however, the associations were numerically similar 4.9 (0.2, 9.7) and 3.2 (0.6, 5.7) for systolic and diastolic blood pressure, respectively.

Associations of vitamin D and vitamin K with incident hypertension

During a median follow-up time of 6.4 years (range 2.0-9.7) 143 participants (62%) developed hypertension (Table 3). The associations between vitamin D and vitamin K categories and incident hypertension showed similar patterns as for blood pressure. After adjustment for sex and age, participants in the low vitamin D and K status group had a 69% higher risk of developing hypertension compared to the participants in the high vitamin D and vitamin K status group: HR 1.69 (1.06-2.68). However, after further adjustments for potential confounders, the association remained similar, but was no longer statistically significant: HR 1.62 (0.96-2.73). The HRs for

the other categories were: 1.28 (0.77, 2.13) 25(OH)D <50/dp-ucMGP <323 and 1.29 (0.84, 1.99) 25(OH)D ≥50/dp-ucMGP ≥323. Adjusting for PTH as a potential mediator further attenuated the results to HR 1.59 (0.93, 2.73) for the 25(OH)D <50/dp-ucMGP <323 category.

The P-for interaction on a multiplicative scale was 0.226. For incident hypertension the relative excess risk due to interaction for model 1 is: $1.69 - 1.28 - 1.29 + 1 = 0.12$ and model 2: $1.62 - 1.42 - 1.37 + 1 = -0.17$.

No interaction was observed between 25(OH)D and dp-ucMGP as continuous variables for systolic and diastolic blood pressure as well as incident hypertension, as presented in Supplemental table S2 P-for interaction >0.262.

For Hypertension Peer Review. Do not distribute.
Destroy after use.

Discussion

In the current study, we reported associations of serum 25(OH)D and dp-ucMGP with blood pressure and incident hypertension in a population-based cohort with serial follow-up in older men and women. The combination of low vitamin D (25(OH)D <50 nmol/L) and low vitamin K (dp-ucMGP \geq 323 pmol/L) status was associated with increased systolic and diastolic blood pressure with a similar trend for incident hypertension, albeit not statistically significant. An additive effect of 25(OH)D and dp-ucMGP on blood pressure was evident based on the interaction tests, however, we did not find a statistically significant interaction between vitamin D and K for incident hypertension based on the relative excess risk due to interaction, which was approximately zero indicating no evidence of additive interaction. No interaction was observed between 25(OH)D and dp-ucMGP as continuous variables, meaning there might be a threshold association for both 25(OH)D and dp-ucMGP as observed in the categorical associations.

To our knowledge, the importance of optimal vitamin D and K concentrations on blood pressure and incident hypertension has thus far only been studied in isolation. Previous prospective studies have reported lower plasma vitamin D concentrations to be associated with higher hypertension risk¹¹ and higher dp-ucMGP concentrations to be associated with arterial stiffness²⁸ and higher risk of coronary artery calcification.²⁹ The first study that explored the combined effect of vitamin D and K supplementation on cardiovascular health was a double-blind placebo controlled trial of 181 postmenopausal women aged 50-60 years and revealed that supplementation of vitamin K₁, D and minerals was superior to vitamin D plus minerals alone in preventing a decrease of elastic properties of the carotid artery over 3 year follow-up.¹⁷ However, in this experiment¹⁷ vitamin D and K was combined with calcium, magnesium and zinc, making

it difficult to solely pinpoint the effect to vitamin D and K. Our results appear in line with this trial, since a combined low vitamin D and vitamin K status was associated with increased blood pressure compared to those with a high vitamin D and K status. We also provide evidence for an additive effect of vitamin D and vitamin K, since the interaction for blood pressure was significant. A similar trend was seen for low vitamin D and K status with incident hypertension, however, it should be noted that the power to test interaction is limited in small sample sizes (e.g., <250 subjects) and that type 1 error rates could be high, which might explain the difference in interaction tests between blood pressure and hypertension.³⁰

The hypothesized joint association between vitamin D and K on cardiovascular health is further supported by evidence on bone health outcomes. In a recent case-control study of 184 older Norwegian men and women, low phylloquinone and low 25(OH)D concentrations were synergistically associated with a higher risk of hip fractures.³¹ Further, multiple human trials have demonstrated a synergistic effect of vitamin D and K on bone health.^{32,33} Only the combination of vitamin D and K supplementation resulted in reduced bone loss. It should be noted that in the Postmenopausal Health Study II, all intervention groups significantly increased total-body bone mineral density compared to the control group, however, only the vitamin D and K combination groups increased lumbar spine bone mineral density. In the study of Ushiroyama et al, postmenopausal women with osteopenia and osteoporosis with low vitamin D values were included and no compliance markers of circulating measures of vitamin D and vitamin K were taken into account. Other studies using solely vitamin K supplementation did not find effects on bone density.³⁴⁻³⁶ These limited studies indicate that the combined effect of vitamin D and K might be superior than using either vitamin alone. Our study provides evidence that this

combined effect may also be present for vascular outcomes. This underscores the need for a concomitant adequate intake/supply of vitamin D and vitamin K.

The potential mechanism to explain the joint association of vitamin D and K is via stimulation of the γ -carboxylase system of vitamin K dependent proteins. Two of these proteins are Gla-containing proteins osteocalcin and MGP play a role in the development of coronary artery disease.^{8,37} Osteocalcin has been shown to have a regulatory role in bone mineralization and calcium ion homeostasis³⁸ with the potential to prevent calcium build-up in soft tissues, thereby preventing vascular calcification.³⁹ Similarly, MGP limits calcium incorporation into extracellular matrix of soft tissues thus acting as a potent inhibitor of soft tissue calcification.^{8,38} Activation of osteocalcin and MGP is dependent on vitamin K, which acts as a cofactor in the carboxylation cycle. Low expression or inactivation of vitamin K-dependent proteins can lead to vascular stiffness, which can increase blood pressure and incident hypertension risk.⁴ Circulating vitamin D has been shown to increase MGP mRNA expression in humans.⁴⁰ It has also been demonstrated that vitamin K promotes 1,25-dihydroxyvitamin D stimulated osteocalcin accumulation and mineralization.⁴¹ In addition, in an animal model the co-administration of vitamin K and 1,25-dihydroxyvitamin D, increased levels of bone anabolic markers including osteocalcin in a time-dependent manner⁴², while these effects were not visible with the separate vitamins, as demonstrated by randomized trials.^{43,44} In addition, the results attenuated further after adjusting for PTH, meaning that there might be potential mediation by PTH, however it remains unclear if dp-ucMGP is related to PTH. Overall, from experimental studies it is clear that vitamin D and K can mutually enhance each other, which could explain a joint association of these vitamins on cardiovascular health.

Our study has some limitation that should be addressed. Due to the high number of exclusions, the current study sample was relatively small varying between 231 and 402 participants. Excluded participants had poorer health as observed from higher BMI, lower vitamin status and more medication use, which could have impacted our results, although this would most likely dilute the observed associations. Furthermore, the sample size for the occurrence of hypertension was relatively small, which probably explains the borderline significant results for the combined low vitamin D and K status and non-significant interactions for incident hypertension. Future studies with larger sample sizes are needed to confirm our results. Furthermore, the use of data-driven cut-off points for vitamin K status can cause difficulties comparing results across studies. In our study, the median dp-ucMGP was 323 pmol/L, while other studies report higher median concentrations (466 to 525 pmol/L).^{45,46} Currently, there is no clinically accepted dp-ucMGP cut-off point for healthy populations. More research is needed to define optimal concentrations for vitamin K status using studies with a broad range of dp-ucMGP concentration with long-term follow-up of clinical outcomes.

Our study has also some strengths. We assessed joint associations of 25(OH)D and dp-ucMGP in relation to both blood pressure and incident hypertension in a population free of baseline hypertension and vitamin K antagonist use, minimalizing potential bias and misclassification. The LASA cohort is a nationally representative sample with a substantial follow-up time, repeated measured blood pressure and medication information, and a large number of covariates to adjust for potential confounding. Furthermore, we used blood samples to assess vitamin K status by measuring dp-ucMGP, which has been shown to be a valid biomarker of vitamin K status and reflects menaquinone intake in a dose-response manner.²¹ The majority

of the previous studies assessed vitamin K intake with a food frequency questionnaire, which has well known limitations.^{3,47}

In conclusion, the combination of low vitamin D and low vitamin K status was associated with increased systolic and diastolic blood pressure in a cohort of older men and women with a similar trend for incident hypertension risk compared with high vitamin D and vitamin K status.

Perspective

This is the first study investigating joint associations of vitamin D and vitamin K in relation to cardiovascular health. An additive effect of vitamin D and vitamin K status on blood pressure was evident based on the interaction tests, however, we did not find a statistically significant interaction between vitamin D and K for incident hypertension. The combination of low vitamin D and K status was associated with increased blood pressure and might play a role in development of hypertension. Further prospective studies are needed to better understand the clinical implications of this relationship to promote cardiovascular health.

Sources of funding

This study is based on data collected in the context of the LASA Study, which is largely funded by the Ministry of Health, Welfare, and Sports of the Netherlands. The dp-ucMGP measurements are funded by FrieslandCampina, a dairy company.

Disclosures

None

References

1. Wang L, Song Y, Manson JE, Pilz S, Marz W, Michaelsson K, Lundqvist A, Jassal SK, Barrett-Connor E, Zhang C, Eaton CB, May HT, Anderson JL, Sesso HD. Circulating 25-hydroxy-vitamin D and risk of cardiovascular disease: a meta-analysis of prospective studies. *Circ Cardiovasc Qual Outcomes*. 2012;5:819-829.
2. Dalmeijer GW, van der Schouw YT, Magdeleyns EJ, Vermeer C, Elias SG, Velthuis BK, de Jong PA, Beulens JW. Circulating species of matrix Gla protein and the risk of vascular calcification in healthy women. *Int J Cardiol*. 2013;168:e168-170.
3. Gast GC, de Roos NM, Sluijs I, Bots ML, Beulens JW, Geleijnse JM, Witteman JC, Grobbee DE, Peeters PH, van der Schouw YT. A high menaquinone intake reduces the incidence of coronary heart disease. *Nutr Metab Cardiovasc Dis*. 2009;19:504-510.
4. Jensky NE, Criqui MH, Wright MC, Wassel CL, Brody SA, Allison MA. Blood pressure and vascular calcification. *Hypertension*. 2010;55:990-997.
5. Writing Group M, Mozaffarian D, Benjamin EJ, et al. Executive Summary: Heart Disease and Stroke Statistics--2016 Update: A Report From the American Heart Association. *Circulation*. 2016;133:447-454.
6. Luo G, Ducy P, McKee MD, Pinero GJ, Loyer E, Behringer RR, Karsenty G. Spontaneous calcification of arteries and cartilage in mice lacking matrix GLA protein. *Nature*. 1997;386:78-81.
7. Caluwe R, Pyfferoen L, De Boeck K, De Vriese AS. The effects of vitamin K supplementation and vitamin K antagonists on progression of vascular calcification: ongoing randomized controlled trials. *Clin Kidney J*. 2016;9:273-279.

8. Schurgers LJ, Cranenburg EC, Vermeer C. Matrix Gla-protein: the calcification inhibitor in need of vitamin K. *Thromb Haemost.* 2008;100:593-603.
9. Dalmeijer GW, van der Schouw YT, Magdeleyns EJ, Vermeer C, Verschuren WM, Boer JM, Beulens JW. Matrix Gla protein species and risk of cardiovascular events in type 2 diabetic patients. *Diabetes Care.* 2013;36:3766-3771.
10. Dalmeijer GW, van der Schouw YT, Magdeleyns EJ, Vermeer C, Verschuren WM, Boer JM, Beulens JW. Circulating desphospho-uncarboxylated matrix gamma-carboxyglutamate protein and the risk of coronary heart disease and stroke. *J Thromb Haemost.* 2014;12:1028-1034.
11. Ke L, Mason RS, Kariuki M, Mpofo E, Brock KE. Vitamin D status and hypertension: a review. *Integr Blood Press Control.* 2015;8:13-35.
12. Masterjohn C. Vitamin D toxicity redefined: vitamin K and the molecular mechanism. *Med Hypotheses.* 2007;68:1026-1034.
13. Fu X, Wang XD, Mernitz H, Wallin R, Shea MK, Booth SL. 9-Cis retinoic acid reduces 1alpha,25-dihydroxycholecalciferol-induced renal calcification by altering vitamin K-dependent gamma-carboxylation of matrix gamma-carboxyglutamic acid protein in A/J male mice. *J Nutr.* 2008;138:2337-2341.
14. Fraser JD, Price PA. Induction of matrix Gla protein synthesis during prolonged 1,25-dihydroxyvitamin D3 treatment of osteosarcoma cells. *Calcif Tissue Int.* 1990;46:270-279.
15. Miyake N, Hoshi K, Sano Y, Kikuchi K, Tadano K, Koshihara Y. 1,25-Dihydroxyvitamin D3 promotes vitamin K2 metabolism in human osteoblasts. *Osteoporos Int.* 2001;12:680-687.

16. Seyama Y, Horiuchi M, Hayashi M, Kanke Y. Effect of vitamin K2 on experimental calcinosis induced by vitamin D2 in rat soft tissue. *Int J Vitam Nutr Res.* 1996;66:36-38.
17. Braam LA, Hoeks AP, Brouns F, Hamulyak K, Gerichhausen MJ, Vermeer C. Beneficial effects of vitamins D and K on the elastic properties of the vessel wall in postmenopausal women: a follow-up study. *Thromb Haemost.* 2004;91:373-380.
18. El Hilali J, de Koning EJ, van Ballegooijen AJ, Lips P, Sohl E, van Marwijk HW, Visser M, van Schoor NM. Vitamin D, PTH and the risk of overall and disease-specific mortality: Results of the Longitudinal Aging Study Amsterdam. *J Steroid Biochem Mol Biol.* 2016;164:386-394.
19. van den Heuvel EG, van Schoor NM, Lips P, Magdeleyns EJ, Deeg DJ, Vermeer C, den Heijer M. Circulating uncarboxylated matrix Gla protein, a marker of vitamin K status, as a risk factor of cardiovascular disease. *Maturitas.* 2014;77:137-141.
20. Hoogendijk EO, Deeg DJ, Poppelaars J, van der Horst M, Broese van Groenou MI, Comijs HC, Pasman HR, van Schoor NM, Suanet B, Thomese F, van Tilburg TG, Visser M, Huisman M. The Longitudinal Aging Study Amsterdam: cohort update 2016 and major findings. *Eur J Epidemiol.* 2016;31:927-945.
21. Dalmeijer GW, van der Schouw YT, Magdeleyns E, Ahmed N, Vermeer C, Beulens JW. The effect of menaquinone-7 supplementation on circulating species of matrix Gla protein. *Atherosclerosis.* 2012;225:397-402.
22. Poulter NR, Prabhakaran D, Caulfield M. Hypertension. *Lancet.* 2015;386:801-812.
23. Comijs HC, Aartsen MJ, Visser M, Deeg DJ. [Alcohol consumption among persons aged 55+ in The Netherlands]. *Tijdschr Gerontol Geriatr.* 2012;43:115-126.

24. Stel VS, Smit JH, Pluijm SM, Lips P. Consequences of falling in older men and women and risk factors for health service use and functional decline. *Age Ageing*. 2004;33:58-65.
25. Lee J, Tan CS, Chia KS. A practical guide for multivariate analysis of dichotomous outcomes. *Ann Acad Med Singapore*. 2009;38:714-719.
26. Rubin DB. Multiple Imputation After 18+ Years. *J Am Stat Ass*. 1996;91:473-489.
27. Knol MJ, van der Tweel I, Grobbee DE, Numans ME, Geerlings MI. Estimating interaction on an additive scale between continuous determinants in a logistic regression model. *Int J Epidemiol*. 2007;36:1111-1118.
28. Pivin E, Ponte B, Pruijm M, *et al*. Inactive Matrix Gla-Protein Is Associated With Arterial Stiffness in an Adult Population-Based Study. *Hypertension*. 2015;66:85-92.
29. Dalmeijer GW, van der Schouw YT, Vermeer C, Magdeleyns EJ, Schurgers LJ, Beulens JW. Circulating matrix Gla protein is associated with coronary artery calcification and vitamin K status in healthy women. *J Nutr Biochem*. 2013;24:624-628.
30. Schmidt AF, Groenwold RH, Knol MJ, Hoes AW, Nielen M, Roes KC, de Boer A, Klungel OH. Exploring interaction effects in small samples increases rates of false-positive and false-negative findings: results from a systematic review and simulation study. *J Clin Epidemiol*. 2014;67:821-829.
31. Torbergsen AC, Watne LO, Wyller TB, Frihagen F, Stromsoe K, Bohmer T, Mowe M. Vitamin K1 and 25(OH)D are independently and synergistically associated with a risk for hip fracture in an elderly population: a case control study. *Clin Nutr*. 2015;34:101-106.
32. Kanellakis S, Moschonis G, Tenta R, Schaafsma A, van den Heuvel EG, Papaioannou N, Lyrītis G, Manios Y. Changes in parameters of bone metabolism in postmenopausal women following a 12-month intervention period using dairy products enriched with

- calcium, vitamin D, and phylloquinone (vitamin K(1)) or menaquinone-7 (vitamin K (2)): the Postmenopausal Health Study II. *Calcified tissue international*. 2012;90:251-262.
33. Ushiroyama T, Ikeda A, Ueki M. Effect of continuous combined therapy with vitamin K(2) and vitamin D(3) on bone mineral density and coagulofibrinolysis function in postmenopausal women. *Maturitas*. 2002;41:211-221.
34. Katsuyama H, Ideguchi S, Fukunaga M, Fukunaga T, Saijoh K, Sunami S. Promotion of bone formation by fermented soybean (Natto) intake in premenopausal women. *Journal of nutritional science and vitaminology*. 2004;50:114-120.
35. Emaus N, Gjesdal CG, Almas B, Christensen M, Grimsgaard AS, Berntsen GK, Salomonsen L, Fonnebo V. Vitamin K2 supplementation does not influence bone loss in early menopausal women: a randomised double-blind placebo-controlled trial. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2010;21:1731-1740.
36. van Summeren MJ, Braam LA, Lilien MR, Schurgers LJ, Kuis W, Vermeer C. The effect of menaquinone-7 (vitamin K2) supplementation on osteocalcin carboxylation in healthy prepubertal children. *The British journal of nutrition*. 2009;102:1171-1178.
37. Kim KM, Lim S, Moon JH, Jin H, Jung KY, Shin CS, Park KS, Jang HC, Choi SH. Lower uncarboxylated osteocalcin and higher sclerostin levels are significantly associated with coronary artery disease. *Bone*. 2016;83:178-183.
38. Theuwissen E, Smit E, Vermeer C. The role of vitamin K in soft-tissue calcification. *Adv Nutr*. 2012;3:166-173.

39. Okura T, Kurata M, Enomoto D, Jotoku M, Nagao T, Desilva VR, Irita J, Miyoshi K, Higaki J. Undercarboxylated osteocalcin is a biomarker of carotid calcification in patients with essential hypertension. *Kidney Blood Press Res.* 2010;33:66-71.
40. Farzaneh-Far A, Weissberg PL, Proudfoot D, Shanahan CM. Transcriptional regulation of matrix gla protein. *Z Kardiol.* 2001;90 Suppl 3:38-42.
41. Koshihara Y, Hoshi K, Ishibashi H, Shiraki M. Vitamin K2 promotes 1alpha,25(OH)2 vitamin D3-induced mineralization in human periosteal osteoblasts. *Calcif Tissue Int.* 1996;59:466-473.
42. Poon CC, Li RW, Seto SW, Kong SK, Ho HP, Hoi MP, Lee SM, Ngai SM, Chan SW, Leung GP, Kwan YW. In vitro vitamin K(2) and 1alpha,25-dihydroxyvitamin D(3) combination enhances osteoblasts anabolism of diabetic mice. *Eur J Pharmacol.* 2015;767:30-40.
43. Pilz S, Verheyen N, Grubler MR, Tomaschitz A, Marz W. Vitamin D and cardiovascular disease prevention. *Nat Rev Cardiol.* 2016;13:404-417.
44. Fulton RL, McMurdo ME, Hill A, Abboud RJ, Arnold GP, Struthers AD, Khan F, Vermeer C, Knapen MH, Drummen NE, Witham MD. Effect of Vitamin K on Vascular Health and Physical Function in Older People with Vascular Disease--A Randomised Controlled Trial. *J Nutr Health Aging.* 2016;20:325-333.
45. Cranenburg EC, Koos R, Schurgers LJ, Magdeleyns EJ, Schoonbrood TH, Landewe RB, Brandenburg VM, Bekers O, Vermeer C. Characterisation and potential diagnostic value of circulating matrix Gla protein (MGP) species. *Thromb Haemost.* 2010;104:811-822.
46. Shea MK, O'Donnell CJ, Vermeer C, Magdeleyns EJ, Crosier MD, Gundberg CM, Ordovas JM, Kritchevsky SB, Booth SL. Circulating uncarboxylated matrix gla protein is

associated with vitamin K nutritional status, but not coronary artery calcium, in older adults. *J Nutr.* 2011;141:1529-1534.

47. Geleijnse JM, Vermeer C, Grobbee DE, Schurgers LJ, Knapen MH, van der Meer IM, Hofman A, Witteman JC. Dietary intake of menaquinone is associated with a reduced risk of coronary heart disease: the Rotterdam Study. *J Nutr.* 2004;134:3100-3105.

For Hypertension Peer Review. Do not distribute.
Destroy after use.

Novelty and Significance

What Is New?

- This is the first study that prospectively investigated whether the combination of low vitamin D and K status is associated with higher systolic and diastolic blood pressure in 402 participants and with incident hypertension in 231 participants free of hypertension at baseline in a population-based cohort with serial follow-up in older men and women.

What Is Relevant?

- The combination of high vitamin D and K status was associated with increased systolic 4.8 mmHg (95% CI 0.1-9.5) and diastolic 3.1 mmHg (0.5-5.7) blood pressure compared to low vitamin D and K status (P-for interaction=0.013 for systolic blood pressure, 0.068 for diastolic blood pressure).
- A similar trend was seen for incident hypertension: hazard ratio 1.62 (95% CI 0.96-2.73) for the low vitamin D and K group.

Summary

- During a median follow-up of 6.4 years, 62% of the participants (n=143) developed hypertension. The combination of low vitamin D and K status was associated with increased blood pressure and a trend for greater hypertension risk after adjustment for potential confounders.

Table 1. Baseline characteristics of 402 LASA participants stratified by combined 25-hydroxyvitamin D and dephosphorylated uncarboxylated matrix Gla protein categories.

Baseline characteristics	Categories of 25(OH)D (nmol/L) and dp-ucMGP status (pmol/L)			
	25(OH)D \geq 50/ dp-ucMGP<323	25(OH)D <50/ dp-ucMGP <323	25(OH)D \geq 50/ dp-ucMGP \geq 323	25(OH)D <50/ dp-ucMGP \geq 323
Demographics				
N	130	72	125	75
Age (y)	59.6 \pm 3.0	59.7 \pm 2.8	59.5 \pm 2.8	60.1 \pm 3.0
Female	68 (52%)	42 (58%)	73 (58%)	39 (52%)
Season				
Fall	40 (31%)	27 (38%)	41 (33%)	24 (32%)
Winter	59 (45%)	31 (43%)	57 (46%)	37 (49%)
Spring/Summer	31 (24%)	14 (19%)	27 (22%)	14 (19%)
Alcohol use				
Light	70 (54%)	39 (54%)	62 (50%)	40 (53%)
Moderate	49 (38%)	22 (31%)	55 (44%)	24 (32%)
Excessive	11 (9%)	11 (15%)	8 (6%)	11 (15%)
Smoking				
Never	33 (26%)	15 (23%)	37 (28%)	20 (24%)
Former	67 (52%)	22 (31%)	64 (51%)	36 (48%)
Current	28 (22%)	33 (46%)	26 (21%)	21 (28%)
Physical activity				
Low	40 (31%)	29 (40%)	35 (28%)	31 (41%)
Medium	44 (34%)	24 (33%)	46 (37%)	22 (29%)
High	46 (35%)	19 (26%)	44 (35%)	22 (29%)
Education				
Low	54 (42%)	30 (42%)	53 (42%)	32 (43%)
Middle	52 (40%)	23 (32%)	45 (36%)	18 (24%)

High	24 (19%)	19 (26%)	27 (22%)	25 (33%)
Adiposity				
BMI (kg/m ²)	25.5±3.3	25.8 ± 4.2	26.9 ± 3.4	27.7 ± 4.0
Waist circumference (cm)	92.0 ± 10.3	92.3 ± 11.4	95.3 ± 10.2	97.5 ± 11.2
Waist/Hip	0.91±0.07	0.91±0.08	0.92±0.07	0.93±0.09
Metabolic Variables				
Total cholesterol (nmol/L)	5.9±0.9	5.7 ± 1.0	6.0 ± 0.94	6.0±1.0
Systolic blood pressure (mmol/L)	133±20	133±24	135±21	139±21
Diastolic blood pressure (mmol/L)	76±8	77±8	77±9	75±10
eGFR (ml/min/1.73m ²)	83.8±15.0	86.7 ± 14.0	82.7 ± 15.5	84.6±16.8
Parathyroid hormone(pmol/L)	5.1±1.6	6.0 ± 1.7	5.7 ± 2.3	6.6±2.4
25(OH)D (nmol/L)	71±15.8	38±9.3	72±14.9	39±8.4
dp-ucMGP (pmol/L)	208±72	217±71	480±164	480±244
Medication use				
Diuretics	0 (0%)	3 (2%)	1 (1%)	3 (4%)
Lipid-Lowering	6 (5%)	6 (5%)	3 (4%)	5 (7%)

Values are mean ± SD or n (%)

Table 2. Regression coefficients of multiple generalized estimating equation analysis for diastolic and systolic blood pressure and 25-hydroxyvitamin D and dephosphorylated uncarboxylated matrix Gla protein categories during 6.4 years of follow-up in 402 LASA participants

Outcome measure	25(OH)D \geq 50/ dp-ucMGP<323 N=130	25(OH)D <50/ dp-ucMGP <323 N=72	25(OH)D \geq 50/ dp-ucMGP \geq 323 N=125	25(OH)D <50/ dp-ucMGP \geq 323 N=75
	Systolic blood pressure (mmHg)			
Model 1	Reference (0.0)	2.4 (-3.3, 8.2)	4.4 (0.1, 8.6)	6.5 (1.9, 11.1)
Model 2	Reference (0.0)	2.9 (-2.4, 8.2)	3.6 (-0.6, 7.8)	4.8 (0.1, 9.5)
Diastolic blood pressure (mmHg)				
Model 1	Reference (0.0)	1.6 (-1.5, 4.7)	3.1 (0.9, 5.3)	4.2 (1.5, 6.8)
Model 2	Reference (0.0)	1.4 (-1.4, 4.3)	2.8 (0.5, 5.0)	3.1 (0.5, 5.7)

Regression coefficients can be interpreted as difference in mmHg blood pressure expressed as: β and 95% confidence intervals

Low dp-ucMGP indicates high vitamin K status and high dp-ucMGP indicates low status

Model 1 adjusted for age and sex

Model 2 adjusted for age (y), sex, BMI (kg/m²), alcohol consumption (none-

light/moderate/excessive), smoking (never, former, current), education (low/medium/high),

season of blood collection (fall/winter/spring-summer), total cholesterol (nmol/L) and estimated glomerular filtration rate (ml/min/1.73m²)

Table 3. Associations of serum 25-hydroxyvitamin D and dephosphorylated uncarboxylated matrix Gla protein categories with risk of hypertension after 6.4 years of follow-up among 231 LASA participants

Outcome measure	25(OH)D \geq 50/ dp-ucMGP<323	25(OH)D <50/ dp-ucMGP <323	25(OH)D \geq 50/ dp-ucMGP \geq 323	25(OH)D <50/ dp-ucMGP \geq 323
	N=76	N=39	N=70	N=46
Person-years, y	476	244	423	276
Incidence rate per 100	8.4	10.3	10.4	12.3
No. of cases (%)	40 (52%)	25 (64%)	44 (63%)	34 (74%)
Model 1	Reference (1.0)	1.28 (0.77, 2.13)	1.29 (0.84, 1.99)	1.69 (1.06, 2.68)
Model 2	Reference (1.0)	1.42 (0.83, 2.41)	1.37 (0.86, 2.19)	1.62 (0.96, 2.73)

Hazard ratios and 95% confidence intervals derived from Cox proportional hazards models

*Low dp-ucMGP indicates high vitamin K status and high dp-ucMGP indicates low status

Model 1 adjusted for age and sex

Model 2 adjusted for age (y), sex, BMI (kg/m²), alcohol consumption (none-light/moderate/excessive), smoking (never, former, current), education (low/medium/high), season of blood collection (3 categories), total cholesterol (nmol/L) and estimated glomerular filtration rate (ml/min/1.73m²)