

Ole Johan Kemi

On the Cellular Basis of Aerobic Fitness

Intensity-Dependence and Time-
Course of Cardiomyocyte and
Endothelial Adaptations to
Exercise Training

Doctoral thesis
for the degree of Philosophiae Doctor

Trondheim 2005

Norwegian University of Science and Technology
Faculty of Medicine
Department of Circulation and Medical Imaging



NTNU

Norwegian University of Science and Technology
Doctoral thesis
for the degree of Philosophiae Doctor
Faculty of Medicine
Department of Circulation and Medical Imaging

©Ole Johan Kemi

ISBN 82-471-7277-1 (printed vers.)
ISBN 82-471-7276-3 (electronic vers.)
ISSN 1503-8181

Doctoral theses at NTNU, 2005:192

Printed by NTNU-trykk

CONTENTS

Acknowledgements	3
Preface	5
Definitions	6
Summary	7
Introduction	8
Aerobic Fitness is Closely Associated with Maximal Oxygen Uptake	8
Determinants of Maximal Oxygen Uptake	8
Intensity of Exercise Determines Outcome	9
Exercise Training Induces Cardiomyocyte Hypertrophy	9
Cardiomyocyte Contractile Function	10
Cardiomyocyte Contractile Machinery	10
Cardiomyocyte Excitation-Contraction Coupling and Calcium Handling	10
Intracellular Calcium Controls Myocyte Contractility	12
Myofilament Calcium Sensitivity	12
Arterial Endothelial Function	13
Acetylcholine Stimulates Nitric Oxide-Mediated Vasorelaxation	13
Objective	15
Methodological Considerations	16
Animal Models	16
Rat	16
Mouse	16
Exercise Training Program	17
Testing of Oxygen Uptake	17
Anesthesia	18
Echocardiography in Rats and Mice	18
Cardiomyocyte Isolation	19
Electrical Stimulation of Cardiomyocytes	19
Fluorescence and Edge-Detection Microscopy	20
Cardiomyocyte Calcium Sensitivity Index	21
Arterial Endothelial Function	21
Allometric Scaling	22
Statistical Analysis	23
Summary of Results	24
Discussion	28
Evaluation of Test Procedures for Oxygen Uptake	28
Exercise Training Improves Maximal Oxygen Uptake and Cardiac Function	28
Maximal Oxygen Uptake is Pliable to Different Exercise Intensities	29
Time Course of Adaptations in Maximal Oxygen Uptake	29
Cardiomyocyte Hypertrophy with Training	30
Hypertrophy Regression with Detraining	31
Cardiomyocyte Contribution to Myocardial Changes	31
Mechanisms of Cardiomyocyte Size Adaptation	32
Cardiomyocyte Contractility during Exercise Training and Detraining	33
Electrical versus Current Stimulation of Cardiomyocytes	33
Calcium Handling is Linked to Contraction-Relaxation Rates	34
Mechanisms of Altered Calcium Handling	35
Calcium Sensitivity Accounts for Fractional Shortening	35
Mechanisms of Calcium Sensitivity	36
Molecular Coupling Between Hypertrophy and Contractility	36
Endothelium-Dependent Arterial Relaxation	36
Intensity-Dependence of Endothelium-Dependent Relaxation	36
Regression of Improved Endothelium-Dependent Relaxation	37
Mechanisms of Endothelial Adaptation	37
Cellular Mechanisms of Maximal Oxygen Uptake	38
General Considerations and Clinical Implications	38
Conclusions	40
Errata	42
References	43

ACKNOWLEDGEMENTS

To quote one of my favorite writers; “No author writes alone”. The written words may have been mine, but the influence from the community to which I belong and from where my ideas, thoughts, and perspectives stem from is inescapably ever-present in shaping every word and phrase. I appreciate many fine colleagues who in different ways supported, challenged, and above all; inspired me to endure exigent scientific endeavors. Many are those that I deeply respect and by now prefer saluting friends. First and foremost, I wish to pay gratitude to my supervisor Professor Øyvind Ellingsen, whose genuine enthusiasm, continuous support, and constructive criticism has been of vital importance when guiding me not only through my efforts, but also into independency. Øyvind carries a wealth of formal and informal knowledge spanning from basic cardiology to the elements of style of delivering ones message legibly, yet detailed. Importantly, he has always been available, even when half a world away on sabbatical. The same gratitude goes to my co-supervisor Ulrik Wisløff, an excellent and resourceful applied experimentalist whose experience gave me a head start when I arrived to the laboratory. It has been a delight working in lab with him ever since; constantly discussing problems or enjoying lunch in our own think tank. Godfrey Smith from the University of Glasgow is also highly appraised; he advises the world, and us continuously on biophysical and optical techniques, and provides academic leadership in the field. Together, they introduced me to means of how to study and visualize the ways regular exercise can modify the inner workings of the heart, and even mend a broken heart. Moreover, they provided a nurturing ground on where to test my ideas.

Jan Pål Loennechen and Per Magnus Haram deserve sincere thanks for generous experimental help that expanded the studies; Per Magnus and I enjoyed many long nights in lab together, and rightfully, share dual first authorship on one of the papers. When we met pharmacological problems, Tor Skomedal and Jan-Bjørn Osnes from the University of Oslo provided much appreciated help that led us through confusing data. Ingerid Arbo, Sissel Skarra, Vidar Beisvåg, Morten A. Høydal, electrical and mechanical engineers Arnfinn Sira and Ketil Jensen, Vibeke Videm for valuable statistical guidance, my office-mates Tom Chr. Martinsen and Sveinung Sørhaug and all the unmentioned members of the Basic Physiology

Laboratory are all warmly acknowledged for their various contributions and making this a vivid and fun place to be at. The two Heads of Department whom I served with, Per Jynge and Stig A. Slørdahl, provided excellent working conditions; the entire staff at the animal facility, headed by Ingolf Hansen and Karin Bakkelund provided great help. Finally, I am indebted to Jan Hoff and Jan Helgerud for graciously directing me to my current laboratory when my research interests turned to basic biomedical concepts in exercise settings, whilst still maintaining exciting collaborative exercise physiology efforts.

It is also a pleasure to acknowledge the granted Research Fellowship from the Norwegian University of Science and Technology; and the financial support from the St. Olavs Hospital, the Norwegian Council on Cardiovascular Diseases, the Norwegian Society of Cardiology, Sintef Unimed, and the following charity foundations: Agnes Sars, Arild and Emilie Bachke, ESW, Ingeborg and Anders Nordheim, Lise and Arnfinn Heje, Randi and Hans Arnet, and Torstein Erbo.

Lastly, I appreciate the encouragement from family and friends. It is hard to know exactly what sparked an interest in advanced scientific studies, but I suspect my parents Margrethe and Anders had something to do with it, much the way they supported me.

Trondheim, August 2005

Ole Johan Kemi

PREFACE

The thesis is based on four papers listed below, referred to by roman numerals in the text.

The work was carried out in the Basic Physiology Laboratory at the Department of Circulation and Medical Imaging, and is presented to the Faculty of Medicine, the Norwegian University of Science and Technology, for the Doctoral Degree Ph.D. in Molecular Medicine.

Paper I

Ulrik Wisløff, Jan Helgerud, Ole Johan Kemi, Øyvind Ellingsen.

Intensity-controlled treadmill running in rats: VO_{2max} and cardiac hypertrophy. *American Journal of Physiology Heart and Circulatory Physiology* 2001;280:H1301-10.

Paper II

Ole Johan Kemi, Jan Pål Loennechen, Ulrik Wisløff, Øyvind Ellingsen.

Intensity-controlled treadmill running in mice: cardiac and skeletal muscle hypertrophy. *Journal of Applied Physiology* 2002;93:1301-9.

Paper III

Ole Johan Kemi, Per Magnus Haram, Jan Pål Loennechen, Jan-Bjørn Osnes, Tor Skomedal, Ulrik Wisløff, Øyvind Ellingsen.

Moderate vs. high intensity: differential effects on aerobic fitness, cardiomyocyte contractility, and endothelial function. *Cardiovascular Research* 2005;67:161-172.

Paper IV

Ole Johan Kemi, Per Magnus Haram, Ulrik Wisløff, Øyvind Ellingsen.

Aerobic fitness is associated with cardiomyocyte contractile capacity and endothelial function in exercise training and detraining. *Circulation* 2004;109:2897-904.

DEFINITIONS

Intracellular calcium concentration ($[Ca^{2+}]_i$) transient The transient increase and decay of $[Ca^{2+}]_i$ during a contraction-relaxation cycle of the cardiomyocyte; denotes the cytosolic Ca^{2+} changes that induce contraction and relaxation.

Ca^{2+} -induced Ca^{2+} release The subcellular events that initiate cardiomyocyte contraction. The inward L-type Ca^{2+} current stimulates the Ryanodine receptor to release Ca^{2+} from the sarcoplasmic reticulum. The released Ca^{2+} induces myofilament contraction.

Detraining Complete withdrawal of exercise after a history of regular exercise training.

Endothelial function Includes several faculties of arterial endothelium; is in this thesis indicated by acetylcholine-induced nitric oxide generation in the endothelium that relaxes the vessel wall smooth muscle and induces vasodilation.

Fractional shortening The decrease in cardiomyocyte length from end-diastole to end-systole divided by end-diastolic length; defines the degree of cell shortening.

Maximal oxygen uptake (VO_{2max}) The highest oxygen uptake achievable during dynamic exercise with large muscle groups; the best single physiological measure of aerobic fitness.

Myofilament Ca^{2+} sensitivity The myofilament contractile response to any given $[Ca^{2+}]_i$ in the cardiomyocyte.

SUMMARY

Beneficial effects of exercise are closely associated with fitness and maximal oxygen uptake (VO_{2max}). Capacity for oxygen transport increases mainly by improved cardiac function, including larger chamber volumes, myocardial hypertrophy, and enhanced diastolic and systolic function. Higher arterial conductance, capillarity, and oxygen utilization in skeletal muscle also contribute. The present thesis investigates the cellular basis for cardiac and arterial effects; how they correlate with changes in VO_{2max} during exercise training at high or moderate intensity, and during detraining.

The studies show that:

1. Regular aerobic exercise training increases running performance and VO_{2max} , induces cardiac hypertrophy, and improves cardiomyocyte contractility and arterial endothelial function.
2. High intensity exercise is more effective than moderate intensity exercise to increase aerobic fitness.
3. VO_{2max} correlates closely with cardiomyocytes size, contractility, and calcium handling during adaptation in the first weeks of exercise, during detraining, and with different training intensities.

We conclude that:

1. Aerobic fitness is intimately related to cardiomyocyte size and function.
2. The magnitude of adaptation to training depends on exercise intensity.
3. Intensity emerges as an important determinant for beneficial effects of exercise training.

INTRODUCTION

Physical activity reduces, but does not currently prevent the Western Society epidemic of cardiovascular disease from either reaching global proportions or taxing public health and economy (6,30). Lack of physical activity and lowered level of fitness contributes to the worldwide rise of life-style related diseases as cardiovascular disease, metabolic syndrome, and diabetes (118). Better, affordable prevention and treatment strategies to improve wide-scale health are called upon. Regular physical exercise represents such a strategy (17,20). It improves cardiovascular function, health and quality of life, and mortality in patients and in those at risk of developing disease (11,18,62,65,140,143). A Cochrane meta-analysis found that exercise training reduces mortality by 27% in coronary artery disease patients (91). However, recommendations on physical activity in primary and secondary prevention of cardiovascular disease are diffuse (5,52) despite indisputable evidence of aerobic fitness as an important clinical reference point and target (93,136). In order to develop optimal exercise protocols to improve health, a sound understanding of the responsible cellular biology of improved cardiac and vascular function after exercise training is required. Thus, this thesis studies the relationship between aerobic fitness and cardiomyocyte and arterial function.

Aerobic Fitness is Closely Associated with Maximal Oxygen Uptake

The single most important physiological measure of aerobic fitness in both health and disease is maximal oxygen uptake (VO_{2max}), which refers to the maximal rate at which oxygen can be transported from ambient air to peripheral working skeletal muscles, where it is utilized by intracellular metabolic processes (160,176,177). Work economy, i.e. oxygen cost of given work or ratio between power output and oxygen uptake (VO_2), and lactate threshold, i.e. highest VO_2 or power output without excess congestion of lactate, also contribute to aerobic fitness, but considerably less (77,126,145).

Determinants of Maximal Oxygen Uptake

Analytical modeling (26,47,176,177) and laboratory approaches (154,156) assert that cardiac output and arterial conductive and redistributive capacity are rate-limiting to aerobic exercise in every individual. A majority of studies indicate that VO_{2max} is by large determined by the

cardiac pumping capacity, as the main drop in oxygen partial pressure occurs between the pulmonary and skeletal muscle capillaries (160,176,177). This implies a regulatory role for arterial conductance. In contrast, pulmonary work and the problem of lung perfusion/diffusion mismatch is only rate-limiting to well-trained individuals, whereas skeletal muscle diffusion and metabolic flux appears rate-limiting only when convective supply surpasses peripheral demand, such as in unfit subjects or during work with only a small muscle mass (178). All things considered, the likelihood is that there is no sole determinant of VO_{2max} , but rather a complex chain of intertwined elements working in concert (154,176,177).

Intensity of Exercise Determines Outcome

Although defining studies of healthy individuals still are lacking, clinical trials point to the superiority of high aerobic intensity over low-to-moderate intensity to gain full effect of a training program (2,89,101,157,170). The emergence of VO_{2max} as a continuum from health to disease supports this notion (93,136). Cardiovascular adaptations may therefore rely on the exercise intensity during long-term regular training programs also in healthy individuals, especially since it was demonstrated that stroke volume in well-trained athletes increases continuously with increasing intensity up to maximal levels at or around peak aerobic exercise intensity (59,194).

Exercise Training Induces Cardiomyocyte Hypertrophy

Adaptive growth of the cell in response to regular exercise (physiological hypertrophy) usually involves proportional growth in length and width (83,128,129). This corresponds with increased ventricular weights (8) and the phenomenon of the athletes' heart, manifested as increased chamber volumes and mass, and increased thickness of the left ventricle wall (146,147,150). Both increased gene transcription and translation seem to underlie physiological hypertrophy (25,102). In contrast, abnormal growth in response to pressure and/or volume overload characterizes pathologic hypertrophy, and involves early longitudinal cell growth followed by a later stage of proportional length and width growth (38,83). Hypertrophy in heart failure is normally linked to reprogramming into a fetal gene program expressing embryonic growth cascades (28).

Cardiomyocyte Contractile Function

A central feature of regular exercise training is improved systolic and diastolic function and larger cardiac output (50,59,162,186). Exercise training over months usually improves cardiomyocyte shortening and rates of contraction-relaxation (129,187,188), providing a cellular basis for global function. In contrast, heart failure induces cardiac dysfunction with reduced ejection fraction and lower cardiac output, which also is associated with impaired cardiomyocyte contractile function in both man and rodent (22,38,41,66,109,110,148,165,174,182,187). Overall, the differences that occur between species seem to be quantitative, rather than qualitative (14,15,167). Experimental studies of cellular effects of regular exercise training appear therefore translatable to humans, also when exercise training is applied therapeutically in heart failure and cardiovascular disease. Such studies suggest restored contractility and attenuated pathological growth as important cellular mechanisms for the beneficial effects of physical activity in heart failure (187,190,192).

Cardiomyocyte Contractile Machinery

Cardiomyocyte sarcomere units composed of mainly actin and myosin filaments constitute the contractile protein machinery, together with tropomyosin and the troponin complex. When Ca^{2+} binds to troponin C, the complex changes conformation and troponin I moves to allow actin-myosin coupling, while troponin T attaches the complex to tropomyosin. Next, myosin heads bind to actin and flex to slide the filaments in opposite directions and cause shortening; hence, the process is termed the sliding filament model. Upon release of Ca^{2+} from troponin C, the machinery returns to its innate position and awaits the next contractile cycle. Titin, myosin binding protein C, M-protein, myomesin, nebulin and nebulin, and α - and β -actinin serve supporting and regulatory functions (14,15).

Cardiomyocyte Excitation-Contraction Coupling and Calcium Handling

The rapid transient increase in intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) evoke global cell shortening as described above. Conversely, the Ca^{2+} decay causes relaxation. Details of the subcellular events regulating Ca^{2+} induced Ca^{2+} release are illustrated in figure 1.




Figure 1. Excitation-contraction coupling in ventricular myocytes. DHPR: L-type Ca²⁺ channel; NCX; Na⁺/Ca²⁺ exchanger; T-Tubule: Transverse tubule; RyR2: Ryanodine receptor-2; SR: Sarcoplasmic reticulum; PLN: Phospholamban; SERCA2a: SR Ca²⁺ ATPase-2a. Modified from Bers (14).

Depolarization allows a small entry of Ca²⁺ through the dihydropyridine receptor (DHPR; L-type Ca²⁺ channel) and through reverse mode Na⁺/Ca²⁺ exchanger (NCX). The resulting [Ca²⁺]_i increase stimulates the ryanodine receptor-2 (RyR2) to release mM-order of Ca²⁺ from the Sarcoplasmic Reticulum (SR). The co-location of the inward Ca²⁺ channels in transverse tubules and the RyR2 complex in SR secures a fast and efficient coupling, whereas RyR2-embedded proteins as FKBP12.6 regulate the open probability of the RyR2 to prevent diastolic leakage. Localized Ca²⁺ release (Ca²⁺ sparks) coordinate and generate the [Ca²⁺]_i transient that induces contraction (60,104,112). Post-contraction, the main bulk of Ca²⁺ is re-sequestered (Ca²⁺ decay) by the SR Ca²⁺ ATPase-2a (SERCA2a), which activity is closely regulated by phospholamban (PLN). Thus, SERCA2a recharges the SR Ca²⁺ load, whereas normal mode NCX and plasma membrane Ca²⁺ ATPases translocate Ca²⁺ out of the cell to the extracellular space. This is the prevailing mechanism of cell contraction-relaxation

(14,15,153), although alternatives relying on other stimuli have been suggested (48,51,80), and rebutted (15,94,173).

Intracellular Calcium Controls Myocyte Contractility

Studies of excitation-contraction coupling and Ca^{2+} cycling after exercise training have demonstrated that increased levels of SERCA2a and PLN at least partly explain why Ca^{2+} handling improves and $[\text{Ca}^{2+}]_i$ transport rates increase with regular exercise (188). Studies of other subcellular factors that may improve cell contractility are sparse, but clues are provided from manipulative and mechanistic studies (27,35,76,111,119,121,151). Thus, exercise training improve subcellular Ca^{2+} events that account for larger cardiomyocyte shortening and higher contraction-relaxation rates, although a complete cellular perspective of the underlying biology has not yet been provided.

In contrast, abnormal excitation-contraction coupling and depressed Ca^{2+} handling provide mechanisms for cardiomyocyte contractile dysfunction (60,66,68,104,109,133,148,163,182). Reduced SERCA2a levels (54,76,84,149) and inhibited activity due to decreased phosphorylation of PLN (16,105) reduces the SR Ca^{2+} cycling ability to cope with higher heart rates, and may distort normal force-frequency relationships (38,68) and thus Frank-Starling mechanics. The importance of SERCA2a was further documented by adenoviral gene therapy; restoring SERCA2a in cardiomyocytes corrected abnormal Ca^{2+} handling and contractile dysfunction (41,42), whereas targeted overexpression enhanced contractility in transgenics (9). Reduced SERCA2a levels may be compensated by enhanced phosphorylation of PLN (36). However, close intrinsic control is needed to maintain balanced Ca^{2+} cycling (172).

Myofilament Calcium Sensitivity

Regular exercise training increases myofilament Ca^{2+} responsiveness (46,188). Thus, increased cell shortening may not always correspond to altered bio-availability of Ca^{2+} . The biological basis for altered Ca^{2+} sensitivity is not well defined yet, but modification in the contractile machinery, e.g. troponin subunits and myosin light chains have been proposed (7,

45,100,137). However, another plausible explanation to increased cell contractility may be changed cooperation of actin and myosin or other contractile proteins, independent of Ca^{2+} . After myocardial infarction and in heart failure, the role of myofilament Ca^{2+} sensitivity response is unsettled, as results range from decrease via no change to increase (67,69,70, 109,148,164), despite improved clinical outcome of Ca^{2+} sensitizers (75). It may be that cellular changes affecting Ca^{2+} sensitivity are adaptive in exercise training, and compensatory rather than causative in heart failure, in order to increase or preserve contractile capacity.

Arterial Endothelial Function

Improved arterial endothelial function with exercise training improves conductance of blood flow, and is associated with depressed atherosclerosis due to inhibition of leukocyte adhesion and invasion, less platelet aggregation and adhesion, and suppressed smooth muscle proliferation (3,23,73,97,179). Since these effects depend upon endothelial release of the vasoactive and anti-atherogenic agent nitric oxide (NO), it appears that measuring NO-mediated vasorelaxation is indicative of endothelial function (99,124,138,189). In this thesis, endothelial function is therefore indicated by acetylcholine-induced vasorelaxation (see below); higher artery relaxation is interpreted as improved endothelial function. Endothelial dysfunction represents abnormalities in the pathways that regulate the level of NO. Other vasoactive agents released from the endothelium are endothelium-derived hyperpolarizing factor, prostacyclin, angiotensin II, thromboxane, prostaglandin, and endothelin (57,87,139).

Acetylcholine Stimulates Nitric Oxide-Mediated Vasorelaxation

The bio-availability of NO is mechanistically linked to artery relaxation and endothelial function, as illustrated in figure 2. NO is a small uncharged radical compound produced by oxidation of the terminal guanidino nitrogen of the amino acid L-arginine. The process is catalyzed by the constitutive endothelial isoform of NO synthase (eNOS), after stimulation by acetylcholine-binding to muscarinic receptors. Next, NO enters smooth muscle cells and initiates the signal cascade that ultimately decreases $[\text{Ca}^{2+}]_i$ and induces vasorelaxation (23,138,139). Besides acetylcholine, shear stress, bradykinin, ATP, ischemia, and a large number of extra- and intracellular factors may also mediate NO production (23).




Figure 2. Acetylcholine-induced nitric oxide (NO)-dependent arterial relaxation pathway. eNOS: endothelial NO synthase; NO: nitric oxide; sGCi: inactivated soluble guanylate cyclase; sGCa: activated soluble guanylate cyclase; GTP: Guanosine triphosphate; cGMP: cyclic 3',5'-guanosine monophosphate; PKG: cGMP-dependent protein kinase. Modified from Walther et al (179).

Higher level of NO in the smooth muscle cell induces relaxation and is an important cause of improved endothelial function after exercise training (71,179). Although endothelial function is associated with aerobic fitness (97,189), conflicting results appear during exercise training in normal individuals, from increased to unchanged and decreased endothelial function (13,29, 95,99,117,123,124). In endothelial dysfunction, acetylcholine fails to induce normal relaxation, and may in fact induce a paradoxical vasoconstriction by binding to cholinergic receptors on the smooth muscle cell surface (115). Thus, endothelial (dys)function is linked to cardiovascular health. Recent studies link improved cardiovascular health in cardiovascular patients to corrected endothelial function due to regular exercise training (3,72,73,97,106), partly due to enhanced phosphorylation of eNOS to restore NO-levels (71). Hence, endothelial dysfunction reduces the levels of NO and has been identified as an independent risk factor and marker of morbidity and mortality (74,161).

OBJECTIVE

The main focus of the present study was to investigate the cellular basis for cardiac and vascular contributions to aerobic fitness (VO_{2max}).

The specific aims of the studies were to:

1. Develop valid and reproducible procedures to determine VO_{2max} during treadmill running in rats and mice.
2. Establish long-term intensity-controlled exercise training in rats and mice that yield robust cardiovascular effects mimicking adaptation in humans.
3. Determine how intensity of exercise influences VO_{2max} , cardiomyocyte hypertrophy, contractility and Ca^{2+} handling, and arterial endothelium-dependent relaxation.
4. Determine the time-course of VO_{2max} , cardiomyocyte hypertrophy, contractility and Ca^{2+} handling, and endothelial function when high-intensity exercise training is implemented and withdrawn.
5. Determine the relationship between cardiomyocyte contractile capacity and arterial endothelium-dependent relaxation on one side and VO_{2max} on the other side.

METHODOLOGICAL CONSIDERATIONS

Animal Models

The objective was to correlate VO_{2max} with cardiomyocyte and endothelial function. This requires access to viable cells and tissues and a close control of the intensity and time course of exercise training. Hence, procedures for exercise training and testing of VO_{2max} were developed and evaluated in animal models as detailed in papers I and II, since similar studies in humans are close to impossible. For both rodent models, availability and costs were favorable.

Rat

Adult female Sprague-Dawley rats were used in studies I, III, and IV, as they are previously well characterized in our lab (109,110,187,188) and elsewhere (43-46,190-193). Physiological body growth is less pronounced in females; thus, data related to body mass are less confounded than in males. As discussed below, this does not preclude gender-specificity on cardiac or vascular adaptations (63,142), which may affect cellular dynamics. As Sprague-Dawley rats come from an outbred strain, the genetic variability is preferably larger than in inbred animals and validates the results to a wider biological background. Moreover, the effects of anesthesia in this strain are well-known from previous experience in our laboratory (108-110).

Mouse

To expand future investigations to specific molecular and genetic targets via transgenic and gene-modified models, we extended our exercise model to adult mice. Thus, the first mouse training study from our laboratory is reported here (paper II). The C57BL/6J strain is inbred and more genetically homogenous than outbred animals. Hence, it is extensively used for pharmacological and physiological studies. Even though a recent report described it as inferior to other mouse strains with regard to running performance and exercise adaptability (122), responses in study II were fairly robust.

Exercise Training Program

To maximize cardiovascular adaptations, we chose the custom-made inclined treadmill running model that allows close control of exercise intensity, as our preferred model of exercise. The exercise sessions throughout the investigations were carried out at 85-90% of VO_{2max} , interspersed by short periods of milder intensity, except in paper III, in which moderate intensity at 65-70% of VO_{2max} also was employed to study cellular adaptations to different intensities. Uphill treadmill running is a full-body exercise that taxes the cardiovascular system and VO_2 maximally, and reduces running speed, which otherwise would have the potential to intrude reaching maximal work intensity (see data in papers I and II). Although this model may also cause stress reactions given the electrical grid (130), the physiological adaptations by large mimic those in humans (129,188). In comparison, other available exercise models as voluntary wheel running, fixed speed treadmill running or swim-training in a water tank (8,33,79,129), give little or no control of the amount or intensity of the work performed. In particular, forced swimming probably involves more stress.

The light-dark cycle of the animals was adjusted to accommodate exercise sessions during the animals' dark cycle, as normal behavior pattern summons activity at night. Finally, animals were rewarded chocolate after each exercise session as reinforcement to encourage treadmill running (158). To counteract bias, sedentary animals were also given the same amounts of chocolate.

Testing of Oxygen Uptake

To monitor training-induced adaptations and adjust treadmill running speed to maintain the desired relative exercise intensity, VO_{2max} was measured during treadmill running to exhaustion at the start of every training week. Running economy, i.e. VO_2 at a given submaximal running velocity was measured after the warm-up, but before testing VO_{2max} , to avoid accumulation of excess lactic acid. The protocols for measuring VO_2 in rats and mice are detailed in paper I and II, respectively. The methods have been found reproducible through several studies in our laboratory in both normal healthy and heart failure rats, and in rats genetically selected and bred for either high or low aerobic capacity (papers III and IV;

187-189). Since paper II, we have reproduced the test and training protocols for mice in later unpublished studies. The principles behind the VO_{2max} test resemble those from human subjects (181,195) and confirm that VO_{2max} is integral for evaluating physical capacity and integrative function.

Anesthesia

The normal method for sacrificing animals was with diethyl ether anesthesia, which provides a quick comatose, but maintains a stable cardiac function that is vital for harvesting cardiomyocytes. Intraperitoneal injections with ketamine and xylazine provided anesthesia for echocardiography; a mixture widely used for echocardiography recordings of both rats and mice because of the well-defined cardiovascular effects (88,107,169). Ketamine provides analgesia, sedation, and amnesia. It may affect hemodynamics by increasing blood pressure and cardiac output. Hence, the lowest possible dose was used to balance depressive and stimulatory effects to secure adequate sedation. Following echocardiography, the animals were observed until awakening.

Echocardiography in Rats and Mice

Global adaptations on cardiac morphology and work performance were evaluated by echocardiography, as described in papers II and III. 2-dimensional M-mode long-axis recordings providing 250 frames per second were used to evaluate left ventricular dimensions, whereas pulsed-wave Doppler recordings at 500 Hz were used to assess flow characteristics, according to recommendations from the American Society of Echocardiography (159). From each recording, 5-10 consecutive cardiac cycles were averaged. Although echocardiography is fairly well established as a valid tool to assess left ventricular dimensions and function in the rat myocardial infarction model (88,107,108,166), its feasibility to evaluate training-induced cardiac adaptations is less certain, probably due to a smaller window of changes. In our hands, echocardiography was less sensitive than single cell measurements to detect hypertrophy and different contractile function between sedentary and trained (papers II and III; 188). This is consistent with the large number of animals needed to detect biologically important changes by standard equipment; ~50-60 individuals

would be needed to detect significant differences of 15% (α 0.05, β 0.8) of left ventricular size between different mouse groups in standard M-mode (31).

Cardiomyocyte Isolation

Cardiomyocytes were isolated to study cell size and function, as detailed in papers I and II. The procedures are modified from previous protocols for rats (82) and mice (103). Due to discontinued delivery of Joklik's minimum essential medium used in I and II to isolate left ventricular myocytes, a slightly different solution replaced the isolation medium. Thus, studies III and IV proceeded with Krebs Henseleit Ca^{2+} free solution, but with similar concentrations of the proteolytic collagenase type II and similar introduction to Ca^{2+} to ensure Ca^{2+} tolerability after isolation, which was performed in absence of Ca^{2+} to separate cells from each other. Our laboratory has extensive experience with both Krebs Henseleit and Joklik's solutions, and no differences in cell yield and intrinsic properties were noted throughout the studies. However, the batch of collagenase was an important source of variation in cell yield. We also noted that different batches of laminin caused some variation in myocyte attachment to the cover-slips. The number of left ventricular myocyte nuclei of the rat is estimated to $\sim 33 \cdot 10^6$, whereas our average cell yield was $\sim 3\text{-}4 \cdot 10^6$ from healthy left ventricles; thus, a recovery rate of $\sim 10\%$, given that most cardiomyocytes are mononucleate; some may be binucleate (15). The percentage of rod-shaped cells was consistently $\sim 70\text{-}75\%$. Only viable rod-shaped cells without obvious cellular damage were selected for measurements of dimensions, and only undamaged cells that also responded to electrical stimulation were chosen for measurements of contractility and $[\text{Ca}^{2+}]_i$ transients. Cells that failed to complete the stimulation protocol during the experiment were excluded from analysis. Exclusion rates were similar in all groups. In every experiment, cells from two different groups were isolated simultaneously, to maintain a balanced design and avoid systematic variation.

Electrical Stimulation of Cardiomyocytes

Electrical field stimulation is a standard and widely used means to study contractile function and Ca^{2+} dynamics in isolated cardiomyocytes, as it effectively characterizes cellular adaptations of the heart to exercise training, heart failure, and other diseases (68,109,110,

129,148,151,172,187-189). As a measure of contractile function, fractional shortening was measured during continuous perfusion with HEPES solution at physiological $[Ca^{2+}]_i$ and 37° C. The magnitude of shortening was normalized to diastolic cell length, whereas changes in velocities of contraction and relaxation were indicated by time to half and peak shortening, or time from peak shortening to half-relaxation. $[Ca^{2+}]_i$ ratio was measured simultaneously.

Fluorescence and Edge-Detection Microscopy

In order to study contractility and intracellular Ca^{2+} handling and homeostasis (III and IV), cardiomyocytes were mounted on an inverted microscope with the optical train converted to epifluorescence, including a xenon arc 75 W excitation source, and fluorescence emission detection with a photometer and photomultiplier tube. Capture-and-analysis software handled the generated data. The extent of cell shortening and the rates of cell shortening and relaxation were measured by a sideport camera and video edge-detection from the light/dark contrast of the cell edge against background. Contractility and $[Ca^{2+}]_i$ transients were sampled simultaneously through an FI 40X/ 1.3 NA oil immersion objective, while cells were stimulated by 5 ms and 1-10 Hz electrical field bipolar pulses; bipolarity to avoid electrolysis. A 500 Hz rotating optical chopper and bandpass filters excited the Fura-2/AM loaded cells by alternatively near-ultraviolet 340 nm and 380 nm wavelengths, whereas 510 nm emission was counted to allow ratiometric evaluation of $[Ca^{2+}]_i$ transients. When excited at 340 nm, fluorescence emission increases with increasing $[Ca^{2+}]_i$, whereas at 380 nm excitation, emission decreases with increasing $[Ca^{2+}]_i$. Thus, ratioing minimizes possible aberrations of uneven dye loading, leakage, photobleaching and unwanted compartmentalization of Fura-2 in the cell (Grynkiewicz et al 1985). Because of uncertainty of calibration procedures and a non-linear relationship between the ratio and $[Ca^{2+}]_i$ (53), we used ratios without converting to $[Ca^{2+}]_i$ for further analyses.

For detection of cell size, a large number of cells were video taped through either a FI 20X/0.4 NA or FI 40X/0.7 NA objective, whereupon 100-150 random cells per rat and 170-250 random cells per mouse were measured for length and midpoint width. Cells without visible damage were selected and calibrated by a stage micrometer. The standard deviations for cell length

and width from a pool of sedentary rat cardiomyocytes were found to be $\pm 13 \mu\text{m}$ and $\pm 3 \mu\text{m}$, respectively, and $\pm 11 \mu\text{m}$ and $\pm 4 \mu\text{m}$ in mice.

Cardiomyocyte Calcium Sensitivity Index

To study how the contractile properties of the cell react to a given amount of available Ca^{2+} (67,70,148), a Ca^{2+} sensitivity index was calculated as the ratio between fractional shortening and the amplitude of the $[\text{Ca}^{2+}]_i$ ratio from diastole to peak systole. This is not a direct measure of the myofilament sensitivity to Ca^{2+} ; however, previous studies have established this as an adequate index (109,188). Direct evaluation would require permeabilization of the plasma membrane, which would compromise contractile function, or mounting the cell to force transducers to trace loaded shortening in order to determine the power production of the given Ca^{2+} (46). Moreover, indirect measurements of Ca^{2+} sensitivity imply that the intracellular environment in terms of e.g. pH is not controlled, which may affect myofilament Ca^{2+} sensitivity, whereas the time lag between peak $[\text{Ca}^{2+}]_i$, transient and peak shortening of the cell may to some degree distort the evaluation. Nonetheless, the Ca^{2+} sensitivity index results indicate that myofilament contractile dependence of Ca^{2+} in the isolated cardiomyocytes adapts to chronic exercise.

Arterial Endothelial Function

To assess endothelial function, acetylcholine-mediated endothelium-dependent vasorelaxation was measured by mounting the right common carotid artery onto a force transducer and a micrometer in a continuously equilibrated Krebs solution at 37°C , as detailed in papers III and IV. The protocol was adapted and modified from studies of piglet femoral arteries (168), and represented the first vascular assessments our laboratory performed in rat (paper III). After the first study, it became evident that, after pre-contractions with the α -agonist phenylephrine, higher concentrations of acetylcholine were required to ensure maximal relaxation in all artery segments (see paper III, figure 5). This was effected in the succeeding study (IV), whereas conventional curve-fitting methods were used to calculate maximal relaxation levels in the artery segments that did not level off in study III.

Acetylcholine induces endothelium-dependent vasodilatation via muscarin receptors. To ascertain endothelium-dependence in the tension measurements, arterial segments from each animal were also exposed to L-NAME, an L-Arginine competitor that inhibits the L-Arginine-NO pathway and thus blocks NO-production. Arterial segments were also exposed to Na⁺ nitroprusside; an endothelium-independent NO-donor. The prevailing NO synthase in the arterial wall is eNOS (71,74), which L-NAME inhibited effectively and Na⁺ nitroprusside bypassed by evoking maximal relaxation independent of endothelial NO. The interventions confirmed that arterial vasodilatation induced by acetylcholine were endothelium-dependent. Indomethacin was added to all experiments to exclude confounding by cyclooxygenase-mediated prostaglandin and thromboxane-conversion from arachidonic acid.

Allometric Scaling

Since studies I and II involved both sexes and every study involved long-term follow-up, body mass might influence the results. Thus, allometric dimensional scaling was applied to appropriately normalize VO_2 and cardiac and skeletal muscle weights to body size. Traditionally, data were usually divided by blank body mass. However, this may lead to underestimation in heavier subjects (10,37). Measuring lean body mass is difficult and would require body scanning to distinguish different tissues. At the time of our studies, scaling VO_2 to body mass raised to the power of 0.75 (exponent b) occurred as the best available method, in line with theoretical modeling and empirical studies over a wide range of body masses and species (171). Recent investigations published after the present studies, have estimated different mass exponential values, especially at maximal metabolic rate as indicated by VO_{2max} . By re-examining 34 eutherian mammalian species ranging 7 g to 500 kg from a large body of published data (31 original reports, including paper II), it was found that VO_{2max} should be reported in relation to a body mass with the exponent b 0.872 (\pm 0.029, 95% confidence limits 0.813-0.932; 184,185). However, the exponent b may differ between athletic and non-athletic phenotypes, as a tight connection was found between how VO_{2max} relates to body mass and aerobic capacity in the skeletal locomotor muscles. The allometric cascade was also examined with a different multiple-cause model, in which the relative contributions along a detailed cascade were assessed, from pulmonary ventilation and alveolar-to-arteriolar

oxygen diffusion, to skeletal muscle fiber, Ca^{2+} pump and myosin ATPase activities at the receiving end (81). With this approach, the exponent b was found to be 0.84-0.92 during maximal metabolic rate, and 0.74-0.77 during basal metabolic rate. However, when re-analyzing our $\text{VO}_{2\text{max}}$ data with the mass exponent 0.872, the conclusions remained the same, since body masses remained similar between trained and sedentary animals.

Following the best current approximation, cardiac weights were related to body mass with the exponent 0.78 (10). In paper II, skeletal muscle weights were also measured. To the best of our knowledge, no previous studies could advice on how to scale appropriately. Therefore, we calculated the exponent b from the present population by allometric equations. This approach yielded the exponent b 0.75, almost similar to the cardiac exponent. However, care should be granted, as defining a valid exponential power usually requires a much larger pool of data (90).

Statistical Analysis

As each study operates with a limited number of animals per group, data were analyzed by non-parametric procedures and complemented by one-way and repeated measures ANOVA, where appropriate. In papers III and IV, we investigated which cellular factors best corresponded to $\text{VO}_{2\text{max}}$, to identify the cellular determinants of aerobic fitness. Such attempts were performed by simple univariate and multivariate linear regression analyses. Although this approach identified several cellular features, variables that did not reach statistical significance may still be biologically important. The backward stepwise model was chosen to include all independent variables and then remove insignificant ones one at the time until a final model with significant cellular contributors was achieved. However, the same variables were identified using the forward stepwise model.

SUMMARY OF RESULTS

PAPER I Intensity-controlled treadmill running in rats: VO_{2max} and cardiac hypertrophy

The objectives of this study were to develop and evaluate experimental procedures for determining VO_{2max} during treadmill running in rats of both sexes, and to determine the effects of long-term high-intensity treadmill exercise training at 85-90% of VO_{2max} .

Evaluation of Test Procedures

1. Repeated measurements of submaximal running yielded a high reproducibility:
 VO_2 : $r = 0.96$, coefficient of variation = 4%.
Heart rate: $r = 0.98$, coefficient of variation = 1.5%.
2. Optimal treadmill inclination for measuring VO_{2max} : 25° (47%).
3. Respiratory exchange ratio > 1.05 and blood lactate > 6.0 mM confirmed exercise level at or close to exhaustion.
4. Linear relationship between VO_2 and heart rate was found during increasing treadmill running velocities; maximal heart rate required higher velocities than VO_{2max} .
5. VO_2 leveled off (VO_{2max}) during treadmill running, despite increased running velocity.

Effects of Regular Exercise Training

1. Exercise training induced an athletic phenotype resembling human cardiopulmonary adaptations to exercise.
2. VO_{2max} and maximal aerobic running velocity increased 60-70%.
3. Resting and maximal oxygen pulse increased 25-30%, indicating increased stroke volume.
4. Exercise training improved running economy during submaximal treadmill velocities:
Oxygen cost decreased $\sim 16\%$.
Heart rate decreased 11-13%.
Respiratory exchange ratio decreased $\sim 5\%$
5. Exercise training induced myocardial hypertrophy; right and left ventricular weights increased 23-34%, while left ventricular myocyte length increased 6-12%.

PAPER II Intensity-controlled treadmill running in mice: cardiac and skeletal muscle hypertrophy

The objective was to extend the experimental treadmill running protocols into a mouse model. Thus, the study established valid and reproducible protocols for determining aerobic capacity, and protocols for exercise-induced cardiopulmonary adaptations in mice of both sexes similar to those reported in rats and humans.

Evaluation of Test Procedures

1. Repeated measurements of submaximal VO_2 yielded a high reproducibility: $r = 0.98$, coefficient of variation = 9.1%.
2. Optimal treadmill inclination for measuring VO_{2max} was 15-35°; highest respiratory exchange ratios were obtained at 25°.
3. VO_2 increased linearly with increasing treadmill velocity, until leveling off at VO_{2max} , despite increasing running velocity. Respiratory exchange ratio > 1.0 confirmed exercise level at or close to exhaustion.

Regular Exercise Training

1. Regular exercise training in mice of both sexes induced a similar athletic phenotype as in paper I, and resembled human cardiopulmonary adaptations to exercise.
2. VO_{2max} increased 30-50%; maximal aerobic running velocity increased ~70%.
3. Exercise training improved running economy during submaximal treadmill velocities:
Oxygen cost decreased 19-25%.
Respiratory exchange ratio decreased ~8%
4. Exercise training induced myocardial hypertrophy; left and right ventricular weights increased 19-29% and 12-17% in females and males, respectively. Left ventricular myocyte length and width increased ~20-30% in both sexes.
5. Echocardiography indicated cardiac hypertrophy.
6. Extensor digitorum longus and soleus skeletal muscle weights increased 12-18%

PAPER III Moderate vs. high exercise intensity: differential effects on aerobic fitness, cardiomyocyte contractility, and endothelial function

The objective was to determine how regular exercise at either high (85-90%) or moderate (65-70%) intensity exercise affects VO_{2max} , cardiomyocyte contractile capacity and arterial endothelial function.

1. High and moderate intensity exercise increased VO_{2max} 71% and 28%, respectively.
2. High intensity exercise increased cardiomyocyte size 14% by proportional length and width growth, whereas moderate intensity increased cardiomyocyte length by 5% and cell width indicated growth. Ventricular weights and echocardiography indicated intensity-dependent biventricular hypertrophy.
3. Cell fractional shortening increased 45% and 23% with high and moderate intensity exercise, respectively. Intensity-dependent resting bradycardia was indicated.
4. Cardiomyocyte contraction rates increased 43% and 39%, whereas relaxation rates increased 20% and 10% with high and moderate intensity exercise, respectively. Rates of systolic and diastolic Ca^{2+} transient increase and decay paralleled contraction and relaxation rates.
5. Diastolic Ca^{2+} levels and Ca^{2+} transient amplitudes were unaffected by exercise training, whereas Ca^{2+} sensitivity index increased 40% and 30% after high and moderate intensity exercise, respectively.
6. Arterial endothelium-dependent relaxation improved with high and moderate intensity exercise; acetylcholine concentration evoking half-relaxation decreased 4.3-fold and 2.8-fold with high versus moderate intensity, but the difference between the exercise intensities was not statistically significant.
7. Differential integrative effects were paralleled by intensity-dependent adaptations on cardiomyocyte hypertrophy and myocyte contractile function, but not so clearly with arterial endothelial function. Multivariate regression analysis showed that rates of systolic Ca^{2+} increase and diastolic cell relaxation corresponded best with VO_{2max} , although also degree of cell hypertrophy and contraction, and vasorelaxation correlated to VO_{2max} with univariate analysis.

**PAPER IV Aerobic fitness is associated with cardiomyocyte contractile capacity
and endothelial function in exercise training and detraining**

The objective was to determine the time-course of VO_{2max} and associated cellular changes after regular high intensity exercise was implemented and withdrawn (detraining).

Time-Course of Adaptations to Implementation of Exercise Training (2-13 Weeks)

1. VO_{2max} and cardiomyocyte size, fractional shortening, and rates of myocyte relengthening and Ca^{2+} decay improved steadily up to ~8-10 weeks.
2. Multivariate regression analysis identified cardiomyocyte elongation and diastolic function as the main factors for determining VO_{2max} , although cell shortening and volume also correlated with VO_{2max} in univariate analysis.

Time-Course of Adaptations to Detraining (2-4 Weeks)

1. About 50% of the exercise-gained increase in VO_{2max} vanished within 2 weeks of detraining, and plateaued 5% above sedentary controls after 3-4 weeks.
2. Cardiomyocyte length and width increased 20-22% with exercise training. During detraining, cell width regressed completely within 2 weeks, whereas length remained 7% increased after 2 weeks and 5% after 4.
3. Myocyte fractional shortening increased 30% with exercise, but vanished almost completely within 2 weeks of detraining. The training-induced increase in contraction-relaxation rates and corresponding Ca^{2+} transient rates regressed within 2-4 weeks.
4. Diastolic Ca^{2+} levels and Ca^{2+} transient amplitudes were unaffected by exercise training and detraining, whereas Ca^{2+} sensitivity index increased with training, and regressed gradually within 4 weeks of detraining.
5. Endothelial function improved after exercise training, and regressed completely within 2 weeks of detraining.
6. Cardiomyocyte length and endothelial function appeared as the main cellular determinants for regressed VO_{2max} with detraining (multivariate regression analysis), whereas univariate analysis also showed that fractional shortening and rates of contraction-relaxation and Ca^{2+} transient correlated significantly with VO_{2max} .

DISCUSSION

The working hypothesis of the present thesis was that VO_{2max} is associated with cellular changes in the heart and in arteries. The magnitude of change depends on exercise intensity. Adaptations occur shortly after initiating regular exercise training, level off after ~2 months, and vanish almost completely within a month of detraining. Given that the level of VO_{2max} is closely related to clinical outcome, optimizing exercise programs is likely to yield significant health benefits.

Evaluation of Test Procedures for Oxygen Uptake

The procedures for determining VO_2 during treadmill running were found reproducible and valid both for rats and mice (I and II). To be strict, this validates measurements of submaximal VO_2 and not VO_{2max} . However, hallmarks of appropriate procedures for determining VO_{2max} are linear relationship between submaximal VO_2 and running speed, leveling off of VO_2 despite continued increase in running velocity (VO_{2max}), reproducible values from the same individual when duplicating tests with different inclinations, respiratory exchange ratios >1.0 , linear heart rate response up to maximal levels (I), and clinical signs of exhaustion. These results are indicative of appropriate experimental procedures for determining VO_{2max} in rats and mice (I and II). Our methods resemble clinical procedures in healthy and diseased human individuals (181). Thus, the procedures were valid to determine relative exercise intensity and evaluate adaptations to regular exercise training.

Exercise Training Improves Maximal Oxygen Uptake and Cardiac Function

As expected, VO_{2max} and maximal aerobic running velocity increased with regular exercise training (I-IV). Improvements on VO_{2max} of 50-70% in rats and 30-50% in mice are among the largest reported, which probably reflects the close control of relative exercise intensity (for comparisons, see Discussion sections in papers I and II). Although several factors influence aerobic fitness, VO_{2max} is by far the most important (77,145). Nonetheless, regular exercise also improved work economy considerably, as VO_2 and respiratory exchange ratio during submaximal running decreased. VO_{2max} is closely related to cardiac function (59,194). Thus, improved cardiac output accounts for a large component of increased VO_{2max} (176,177), as

confirmed by increased oxygen pulse (I), which indicates larger stroke volume. Both diastolic and systolic function and hypertrophy of the heart account for higher cardiac output. Multiple regression analysis identified hypertrophy, diastolic function, and Ca^{2+} handling of the cardiomyocyte as important cellular mechanisms of increased $\text{VO}_{2\text{max}}$, whereas single univariate analysis also found that myocyte systolic function correlated well with $\text{VO}_{2\text{max}}$ (III and IV).

Maximal Oxygen Uptake is Pliable to Different Exercise Intensities

High versus moderate exercise intensity during training sessions doubled $\text{VO}_{2\text{max}}$, even though moderate intensity induced a substantial increase (III). In fact, moderate exercise intensity increased $\text{VO}_{2\text{max}}$ to a degree previously regarded as high (134). Recently, intensity-dependence of training-induced $\text{VO}_{2\text{max}}$ has been confirmed in clinical trials of cardiovascular patients; high intensity exercise was superior for clinical outcome (see Introduction). This has important consequences since epidemiological data suggest that $\text{VO}_{2\text{max}}$ is closely linked to health and survival (93,136). Surprisingly though, no studies have thoroughly examined the effects of different exercise intensities on $\text{VO}_{2\text{max}}$ in healthy individuals during long-term exercise programs, although clues of intensity-dependence were provided early on also in healthy adults (39).

Time Course of Adaptations in Maximal Oxygen Uptake

Implementing regular high intensity exercise in previously inactive individuals imposes a rapid augmentation in $\text{VO}_{2\text{max}}$ up to ~8 weeks, whereupon it leveled off (IV). This probably reflects that at one point, exercise training shifts to maintaining fitness, leaving little room for improvement unless higher doses of exercise are implemented (195).

The time course of regressed $\text{VO}_{2\text{max}}$ is considerably shorter in rats than humans, as $\text{VO}_{2\text{max}}$ perished almost completely to sedentary levels in less than 1 month (IV). Ten days of detraining in well-trained men did not reduce $\text{VO}_{2\text{max}}$ (34), whereas 8 weeks detraining in previously active men and women only modestly decreased $\text{VO}_{2\text{max}}$ (152). However, the initial level of fitness in the latter study was much lower than in paper IV.

Cardiomyocyte Hypertrophy with Training

Cardiomyocyte hypertrophy was shortly induced after initiating a regular exercise training program, and continued steadily for ~2 months, when hypertrophy plateaued (IV). The magnitude of cardiomyocyte hypertrophy was dependent upon the intensity of regular exercise, as high intensity exercise training induced a substantially more pronounced response than moderate intensity, 14% vs. 5% longer cells, respectively (III). Cell width also increased after high intensity exercise (III and IV), but was only indicated with moderate intensity (III). Thus, high intensity exercise provides a stronger hypertrophy stimulus than moderate intensity. Moreover, proportional cardiomyocyte length and width hypertrophy was also induced by high intensity exercise in mice.

Physiological hypertrophy is considered beneficial to improve the cardiomyocyte contractile capacity. It provides a cellular basis for larger end-diastolic dimensions, and confirms with the concept of athletes' heart, i.e. enlarged left and right ventricular masses and volumes, and increased wall thicknesses (58,146,150). Global hypertrophy was indicated by increased left and right ventricular weights (I-IV), and by echocardiography indices of enlarged left ventricular posterior wall and septum thickness (II, III). Exercise-induced cardiomyocyte hypertrophy corresponds to previous studies (8,129,187,188), including previous studies reporting increased heart weights in both endurance-trained rats (49) and mice (4,92,125). In fact, the time course of cardiomyocyte hypertrophy (IV) corresponds well to a study finding heart weights to plateau after ~10 weeks of regular exercise, following a continuous increase from initiation of exercise (33). In contrast, there is a lack of well-controlled studies on intensity-dependence of cardiac hypertrophy; in fact, most experimental studies do not control exercise intensity but rely on fixed treadmill speeds, voluntary wheel running, or swim training. This is probably because controlling exercise intensity accurately in terms of VO_2 requires more sophisticated and time-consuming protocols. Intensity-dependence of hypertrophy may explain why only top endurance athletes develop hypertrophied, athletic hearts, whereas less intense programs usually do not (146,147). Cardiac factors account for the ability to increase stroke volume linearly with exercise intensity up to near-exhaustion in well-trained endurance athletes (59,194), and not just up to ~50-60% of VO_{2max} as previously suggested (195). This is

probably why the magnitude of cardiomyocyte adaptation depends on exercise-intensity beyond moderate levels.

Hypertrophy Regression with Detraining

Regression of cardiomyocyte hypertrophy with inactivity following a period of regular exercise (IV) is consistent with previous studies of deconditioning after regular swim training in rats. After 9-10 weeks of daily unloaded swimming that increased heart weights similar to the present studies (I-IV), 80% of the gained heart weight was lost after 2 weeks of detraining (56), and 100% after 3 weeks (33). In fact, 60% of training-induced gain in heart weight was lost one week after stopping regular exercise (79). In the latter study, the training period lasted only 3 weeks, rendering the degree of hypertrophy uncertain. Finally, it was shown that both training- and detraining-induced changes in heart weight are repeatable and susceptible to successive exercise programs (78). Jointly, our data and the previous reports suggest that cardiomyocyte size is closely regulated according to changes in physiological needs.

Regression of cardiomyocyte size is probably the mechanism of myocardial hypertrophy regression in humans, although time course may be slower. Ten days of detraining did not induce any regression in well-trained athletes (34), whereas hypertrophy regression was evident after 3 weeks of detraining in athletes with similar endurance level (120). Thus, the rate of global cardiac regression in humans is probably considerably longer. About ~50% of former elite athletes experienced complete regression of left ventricular mass, whereas almost all maintained enlarged chamber volume during 13 years of follow-up after ceasing competitive sports (147). Given the participants history of exercise training, it is possible that continued leisure activity may have maintained a certain degree of hypertrophy.

Cardiomyocyte Contribution to Myocardial Changes

Cardiac myocytes seem to have an ample potential of adaptation to regular exercise training and inactivity. Although the present results (I-IV) did not quantify contributions of water and connective tissue, other studies have. The roles of connective tissue and collagen were minimal for regulation of cardiac weights during exercise training and detraining (78,79),

whereas myocardial hydration did not contribute at all (33). However, the latter study indicated a minor role for non-myocyte protein components to changed heart weights. Thus, the combined data show that cardiomyocyte size account for the major part of training-induced cardiac hypertrophy and detraining-induced regression.

Mechanisms of Cardiomyocyte Size Adaptation

Physiologic hypertrophy has been associated with proportional increase in length and width of the cardiomyocyte (83), but this has not been demonstrated experimentally until now (II-IV). Other studies in our laboratory (I; 187,188) and elsewhere (129) have only reported increased cell length after regular training. The reason for this discrepancy is presently unclear, but may rely on the accuracy of measurements and number of cells studied. Proportional hypertrophy requires generation of new sarcomeres in series and parallel. Longer cardiomyocytes do not have longer sarcomeres (129), whereas tissue generation by differentiation, dedifferentiation, or stem cell proliferation remains dubious, although such traces have been found in infarcted hearts (12). Regression of cell length and width also occurs proportionally (IV).

Physiological and pathological hypertrophy is characterized by different molecular phenotypes and gene expression profiles (85,86). The underlying molecular mechanisms of adding and removing sarcomeres have not been determined, but several important mechanisms have been identified. Physiologic growth appears to rely on initiation of the phosphoinositol-3-kinase/AKT pathway to stimulate mTOR/p70 S6K signal transduction to increase rate of protein translation in the ribosomal machinery (19,25,125). Other suggested targets involve the Ca^{2+} -binding protein S100A1 (131) and the transcription factors GATA4 and GATA6 (102). Signal cascades normally associated with pathologic growth, such as atrial and brain natriuretic peptides (ANP and BNP; 108) and reprogramming into a fetal gene program (83) are usually depressed during physiological hypertrophy, although mildly increased mRNA levels of cardiac ANP and BNP also have been reported with mice voluntary running (4). Hypertrophy regression has been linked to depressed protein synthesis (56). Other mechanisms could also contribute, e.g. increased rates of protein metabolism or degradation to reduce redundant sarcomere mass.

Cardiomyocyte Contractility during Exercise Training and Detraining

Regular exercise training enhanced cardiomyocyte fractional shortening and rates of systolic contraction and diastolic relengthening (III and IV). The magnitude of improvement was larger with high intensity exercise than with moderate intensity (III). The adaptations increased steadily with regular exercise until a plateau was reached after ~2 months, whereas during detraining, adaptations subsided within 2-4 weeks (IV). No previous accounts exist of time course of cardiomyocyte contractile function after exercise training with different intensities or during detraining, but magnitude of improvements were in line with most data (e.g. 43,46,129, 187,188), but not with studies reporting no effect or decreased contractility (98,144,193). Different exercise training protocols and experimental conditions, e.g. ionic concentrations, pH, temperature, stimulation frequencies, and region of the heart from which cells were isolated, may explain some of the variation. Most studies did not stimulate cells up to physiological frequencies (7-10 Hz; e.g. 98); where our exercise-induced effects were most marked (III and IV). Together with hypertrophy, improved contractility effectuates a substantially greater contractile capacity of the cell, which contributes to greater systolic output and diastolic filling. Increased stroke volume is usually associated with resting bradycardia, as indicated in II and III.

Electrical versus Current Stimulation of Cardiomyocytes

Field stimulation does not completely resemble ordinary action potential-derived excitation-contraction coupling and should thus only cautiously be compared to patch-clamped cells stimulated with ion currents. For instance, field stimulation changes triggering mechanisms into dependence on L-type Ca^{2+} current and reverse mode NCX-current, instead of initiation by the Na^+ current, with the consequence that time to contraction is prolonged (24). In our hands, the extent of cell shortening with field stimulation has continuously been among the largest compared to other laboratories, with values regularly between ~15-25% at 7 Hz stimulation. Fractional shortening varies reportedly between ~3% and ~25%, in line with concordantly large variations in $[\text{Ca}^{2+}]_i$ amplitudes (15). Such differences probably result from experimental conditions, e.g. temperature, pH, concentrations of Ca^{2+} and other ions, stimulation frequencies, selection of cells, species and strains, and procedures for isolating

and culturing cardiomyocytes. These factors may also partially explain the controversy regarding a positive or negative force-frequency relationship (21,53). In our experimental conditions and procedures, we typically observed a positive staircase during 1-7 Hz stimulation, but negative >7 Hz (papers III and IV). This relationship is also maintained in heart failure cells (109). Furthermore, our approach studies intrinsic, unloaded contractile properties of the single cardiomyocyte, as mechanical influences by adjacent cells, tissues and impending in vivo load, as well as neurohormonal stimulation are removed.

Calcium Handling is Linked to Contraction-Relaxation Rates

The inter-dependency and the similar changes on $[Ca^{2+}]_i$ transients and contraction-relaxation velocities suggests that increase in rate of Ca^{2+} cycling explains faster shortening and relengthening of the trained cardiomyocytes (III and IV). Reduced rates of Ca^{2+} cycling with detraining also translate into slower contraction-relaxation rates (IV).

Altered Ca^{2+} handling and contractile capacity of the cardiomyocyte (III and IV) suggest that cellular adaptations contribute to improved global heart function in endurance-trained subjects, such as increased end-diastolic volume, stroke volume, and cardiac output (59,155, 162,194). Cell shortening has also been linked to myocardial contractility in heart failure (109). Regressed Ca^{2+} handling (IV) probably contributes to the detraining-induced reduction of stroke volume (32,120). Intensity-dependence of cardiac function has not been shown previously, but based upon paper III, we hypothesize that high intensity exercise is superior to moderate intensity to induce cardiac effects. Our results are of unloaded shortening with zero external work output, whereas in vivo loaded shortening includes external work due to tissue mass and blood pressure. However, single cell force- and power production also increase with regular exercise (43). Together, these results form a cellular basis for increased myocardial capability to perform external work, although effects of mechanical stress and neurohormonal stimulation are not accounted for (see Methods section for details).

Mechanisms of Altered Calcium Handling

Molecular changes that mechanistically explain the observed changes in Ca^{2+} handling or excitation-contraction coupling have not been determined, although clues are provided. More effective coupling of the L-type Ca^{2+} current to RyR2 Ca^{2+} release, but not more channels, has been shown with regular exercise (128,144). This suggests a mechanism for exercise-induced higher rate of systolic $[\text{Ca}^{2+}]_i$ transient increase (III and IV). Training-induced effects on isolated SR Ca^{2+} release *per se* have not been studied, but could potentially be interesting as the RyR2 complex is responsive to different stimuli (35,55,113,183). Faster $[\text{Ca}^{2+}]_i$ transients decay after exercise training (III and IV) seems to at least partly be explained by increased levels of SERCA2a, PLN and NCX (114,187,188). This improves Ca^{2+} re-sequestering and myocyte relaxation, and may secondarily improve contraction due to improved loading of the SR.

Calcium Sensitivity Accounts for Fractional Shortening

In rats, changed cardiomyocyte fractional shortening with exercise training or detraining could not be explained by systolic or diastolic levels of $[\text{Ca}^{2+}]_i$ or transient amplitude (III and IV). This suggests that the magnitude of cell shortening results from improved myofilament responsiveness to Ca^{2+} . Indeed, papers III and IV indicate that Ca^{2+} sensitivity improves with regular exercise training according to exercise intensity, and wanes with detraining. Adaptation of cardiac Ca^{2+} sensitivity to exercise training has previously been demonstrated thoroughly in rats (46) and has therefore been suggested as an important mechanism of intensity-dependency and time courses of increase and regression of cell fractional shortening (III and IV). Length-dependence of sarcomere Ca^{2+} sensitivity; stretched sarcomeres producing more force (44), implies that end-diastolic volume also may explain why myofilament Ca^{2+} responsiveness changes. These studies may also support the notion that Ca^{2+} sensitivity may be a compensatory mechanism to volume overload (148). Since the variation on Ca^{2+} sensitivity among different species is considerable (67,69,70,109,148,164) and no supportive data beyond rats exist (46, 188), the possibility remains that improved Ca^{2+} sensitivity is species-specific.

Mechanisms of Calcium Sensitivity

Likely mechanisms of changed Ca^{2+} sensitivity (III and IV) are found in the contractile machinery of the cell. The levels of atrial myosin light chain-1 in the heart (45), but also isoform shifting of troponin T (7), troponin I (100), and myosin heavy chains (137) have been associated with altered Ca^{2+} sensitivity. Regulation of pH may also affect Ca^{2+} sensitivity since protons may compete with Ca^{2+} and inhibit Ca^{2+} -binding to troponin C (141).

Molecular Coupling Between Hypertrophy and Contractility

Exercise training induces simultaneous adaptations on contractile function and size of the single cardiomyocyte (III and IV; see figure 2E paper IV). If the newly generated contractile machinery also contributes to improved contractility, then it suggests that intracellular regulatory mechanisms are activated, such as different protein kinases and phosphatases (e.g. 135). An interesting hypothesis is that the AKT pathway stimulates hypertrophy and facilitates Ca^{2+} cycling in concert (25).

Endothelium-Dependent Arterial Relaxation

Regular exercise training induced greater arterial endothelial function, but the effects were not significantly different between high and moderate exercise intensity (III), and vanished completely within days of detraining (less than 2 weeks; IV). Endothelium-mediated vasorelaxation measures the conductive ability of the artery (vasodilation), although endothelial function also includes mechanisms that protects against atherosclerosis (see Introduction). Exercise may therefore reverse endothelial dysfunction and exert anti-atherosclerotic effects on the endothelium. Thus, some effects of exercise training may be more marked in individuals with endothelial dysfunction.

Intensity-Dependence of Endothelium-Dependent Relaxation

Although endothelium-mediated vasorelaxation was indicated better after high intensity exercise training versus moderate, no statistical significance occurred between the exercise levels (III). No clear consensus has evolved on endothelial adaptations to exercise training in healthy individuals. Despite endothelium-dependent vasodilatation improving with moderate

intensity exercise (50% of VO_{2max}), it did not improve with high (75%) or low (25%) exercise intensities (61). Other reports show that endothelial function may improve (29,99,124), decrease (13) or remain unchanged (95,117,123) during regular exercise training. It was also reported that endurance-trained women do not have better endothelial-mediated vasodilatation than sedentary counterparts, but have structurally greater artery diameters (127). These discrepancies may be due to inadequate exercise programs, experimental procedures, selection of the specific artery to study (e.g. large conduit artery vs. smaller and more responsive arteries), initial status of the endothelium (normal vs. unhealthy), and inherent differences between human and animal biology.

In individuals with heart failure and cardiovascular disease, endothelial adaptation is associated with exercise-induced improvement of cardiovascular health and work capacity (3,71-73,97,106). This suggests that arterial adaptive potential may be larger when endothelial dysfunction is present.

Regression of Improved Endothelium-Dependent Relaxation

Regular exercise carried out for 10 weeks improved endothelial function substantially (III and IV). When training was stopped, the effects returned to normal levels within less than 2 weeks (IV). Given the rapid decline, one may assume that regular exercise is necessary to maintain, not just induce, improved endothelial function. The time course of regression appears longer in humans, as endothelial function returned to baseline ~2 months after ceasing regular exercise in patients with coronary artery disease and heart failure (117,180). How quickly regression progress in healthy humans is uncertain; whereas NO levels were maintained after 4 weeks of detraining, plasma levels of NO regress within 8 weeks (116).

Mechanisms of Endothelial Adaptation

Since dilatory responses to Na^+ nitroprusside and reaction to acetylcholine after eNOS inhibition with L-NAME were unaffected by training or detraining (III and IV), altered vasodilatory effects were probably caused by acetylcholine-mediated NO from the endothelium (see figure 2, Introduction). Besides increased eNOS enzyme activity (71),

regular exercise training also increases protein levels of eNOS if the tyrosine kinase c-Src is present (40). Also, increased shear stress and transmural pressure induces higher endothelial reactivity via AKT and vascular endothelial growth factor (VEGF; 1).

Although endothelial adaptation seemed to have a different time course than cardiomyocytes (III and IV), univariate correlation analysis established a weak association with VO_{2max} (III), whereas multivariate regression indicated that regressed endothelial function is associated with VO_{2max} (IV). However, further studies are required to investigate adaptations of the endothelium to exercise, especially in subjects with endothelial dysfunction.

Cellular Mechanisms of Maximal Oxygen Uptake

It takes a certain amount of flexibility and adaptability to respond to exercise training and withdrawal thereof. This thesis shows that changes in VO_{2max} during training and detraining are closely associated with adaptation in cardiac myocytes. It should be noted that statistical modeling to assign values to cellular determinants cannot provide causal proofs but indices, and does not take into account other variables not measured or controlled for. Nonetheless, it provides important targets for more defining studies. Although the scope of this thesis was on oxygen delivery and identifies cardiomyocyte contractile capacity as a main determinant of aerobic fitness, other biological factors are also important for integrative function after training and detraining, e.g. capillarization, mitochondrial mass, and oxidative and metabolic enzymes (96,132).

General Considerations and Clinical Implications

The data reported here are obtained in healthy rats and mice with no history of cardiovascular disease, and provide thus insight into processes occurring under normal physiological conditions. Thus, it remains uncertain how varying degree of disease would affect the results. Genetic background may also affect the outcome. Both young and old individuals, as well as different sexes may react differently to the interventions. Papers I and II indicate that sex differences mainly would appear on the magnitude of adaptations, not the nature. It also adds credibility to translating results from animal to man, that quantitative rather than qualitative

differences seem to separate excitation-contraction coupling between the species (14,15). In cases where human samples are available, a large degree of similarity between experimental and clinical studies seem to appear (e.g. 66-73).

The studies were conducted with the intention to identify mechanisms and form a basis for studies of exercise training as a therapeutic approach in heart failure and related conditions. The first such studies in our laboratory show comparable, beneficial effects of training in heart failure or in animals at risk of developing cardiovascular disease (187,189), in line with other studies reporting beneficial effects of exercise training in heart failure (134). Therefore, we studied intensity-dependence with a clinical approach by changing exercise intensity without changing exercise time. This is not a strict control of the sheer effects of exercise intensity per se, as volume is not adjusted with moderate intensity exercise, but provides evidence that clinical and cellular effects increase if exercise intensity is increased *within a given period of time “set” for exercise*. As described in paper III, if one were to appropriately adjust exercise time to match volume between groups, the exercise time would approach critical lengths in clinical settings. Further experimental and clinical studies of exercise training as a therapeutic tool, partly based upon the present data, are currently underway and show potentially large effects of increasing exercise intensity in patients, even when amount of work is the same.

Given the importance of VO_{2max} for predicting cardiovascular health and all-cause mortality in healthy and cardiovascular patients (93,136), our results should have consequences for designing appropriate exercise programs and studies. Nevertheless, we recognize that low-to-moderate exercise (175), and very intense short-bout sprint training (190-193), may also be valuable in both health and disease.

CONCLUSIONS

1. The experimental procedures for determining VO_{2max} and running economy were valid and reproducible in both rats and mice during inclined treadmill exercise. Thus, the protocols could be used to accurately evaluate adaptations to regular exercise training, and to determine the relative intensity of treadmill running.
2. Regular intensity-controlled exercise training induces robust integrative and cellular adaptations in rat and mouse. VO_{2max} and running economy improve. Cardiomyocyte hypertrophy is induced; and contractile function, Ca^{2+} handling and Ca^{2+} sensitivity increase. Arterial endothelium-dependent vasoreactivity to acetylcholine improved.
3. Regular exercise at high versus moderate intensity induces about two-fold greater adaptations on VO_{2max} and cardiomyocyte dimensions, fractional shortening, Ca^{2+} sensitivity, and velocities of contraction-relaxation and Ca^{2+} transient cycles. In contrast, arterial endothelial function was less dependent upon exercise intensity.
4. After implementing regular high-intensity exercise training, VO_{2max} increases rapidly until a peak level is reached after ~8 weeks. Cardiomyocyte adaptations including hypertrophy response, fractional shortening, and rates of contraction-relaxation and Ca^{2+} transients follow a similar pattern. When regular training is stopped, regression occurs about twice as fast; most of the effects vanished completely within 2-4 weeks, except cell size and VO_{2max} , which remained slightly elevated after 4 weeks of detraining. In contrast, training-induced arterial endothelium-dependent relaxation regressed completely within 2 weeks of detraining.
5. Aerobic fitness and VO_{2max} are closely associated with cellular function and adaptations to regular training at high and moderate exercise intensities and detraining. Multivariate correlation analysis identified cardiomyocyte hypertrophy, rate of Ca^{2+} increase and decay, myocyte relaxation, and endothelial function as the main variables linked to VO_{2max} . Thus, cardiac cell and endothelium features are dynamic and show great plasticity to exercise stimuli.




Figure 3. Summary of the main effects of exercise training at high or moderate intensity, and during detraining following regular exercise training.

The results indicate that exercise intensity is important for aerobic fitness and cellular adaptations that underlie health outcomes, and that cardiomyocyte contractile capacity emerges as an important determinant of VO_{2max} .

ERRATA

Paper IV

Page 2899, Figure 2 legend: references should read 6, 18.

Page 2899, last paragraph, line 4: "Figure 1" should read "Figure 2."

REFERENCES

1. Abid MR, Guo S, Minami T, Spokes KC, Ueki K, Skurk C, Walsh K, Aird WC. Vascular endothelial growth factor activates PI3K/Akt/forkhead signaling in endothelial cells. *Arterioscler Thromb Vasc Biol* 2004;24:294-300.
2. Adachi H, Koike A, Obayashi T, Umezawa S, Aonuma K, Inada M, Korenaga M, Niwa A, Marumo F, Hiroe M. Does appropriate endurance exercise training improve cardiac function in patients with prior myocardial infarction? *Eur Heart J* 1996;17:1511-21.
3. Adams V, Linke A, Krankel N, Erbs S, Gielen S, Mobius-Winkler S, Gummert JF, Mohr FW, Schuler G, Hambrecht R. Impact of regular physical activity on the NAD(P)H oxidase and angiotensin receptor system in patients with coronary artery disease. *Circulation* 2005;111:555-62.
4. Allen DL, Harrison BC, Maass A, Bell ML, Byrnes WC, Leinwand LA. Cardiac and skeletal muscle adaptations to voluntary wheel running in the mouse. *J Appl Physiol* 2001;90:1900-8
5. American College of Sports Medicine Position Stand. Exercise for patients with coronary artery disease. *Med Sci Sports Exerc* 1994;26:i-v.
6. American Heart Association. *Heart Disease and Stroke Statistics — 2003 Update*. American Heart Association, Dallas, TX. 2003.
7. Anderson PA, Greig A, Mark TM, Malouf NN, Oakeley AE, Ungerleider RM, Allen PD, Kay BK. Molecular basis of human cardiac troponin T isoforms expressed in the developing, adult, and failing heart. *Circ Res* 1995;76:681-6.
8. Anversa P, Beghi C, Levicky V, McDonald SL, Kikkawa Y. Morphometry of right ventricular hypertrophy induced by strenuous exercise in rat. *Am J Physiol* 1982;243:H856-61.
9. Baker DL, Hashimoto K, Grupp IL, Ji Y, Reed T, Loukianov E, Grupp G, Bhagwat A, Hoit B, Walsh R, Marban E, Periasamy M. Targeted overexpression of the sarcoplasmic reticulum Ca^{2+} -ATPase increases cardiac contractility in transgenic mouse hearts. *Circ Res* 1998;83:1205-14.
10. Batterham AM, George KP, Mullineaux DR. Allometric scaling of left ventricular mass by body dimensions in males and females. *Med Sci Sports Exerc* 1997;29:181-6.
11. Belardinelli R, Georgiou D, Cianci G, Purcaro A. Randomized, controlled trial of long-term moderate exercise training in chronic heart failure: effects on functional capacity, quality of life, and clinical outcome. *Circulation* 1999;99:1173-82.
12. Beltrami AP, Urbanek K, Kajstura J, Yan SM, Finato N, Bussani R, Nadal-Ginard B, Silvestri F, Leri A, Beltrami CA, Anversa P. Evidence that human cardiac myocytes divide after myocardial infarction. *N Engl J Med* 2001;344:1750-7.
13. Bergholm R, Makimattila S, Valkonen M, Liu ML, Lahdenpera S, Taskinen MR, Sovijarvi A, Malmberg P, Yki-Jarvinen H. Intense physical training decreases circulating antioxidants and endothelium-dependent vasodilatation in vivo. *Atherosclerosis* 1999;145:341-9.
14. Bers DM. Cardiac excitation-contraction coupling. *Nature* 2002;415:198-205.
15. Bers DM. *Excitation-Contraction Coupling and Cardiac Contractile Force*. Kluwer, Dordrecht, The Netherlands. 2001.

16. Bers DM, Eisner DA, Valdivia HH. Sarcoplasmic reticulum Ca^{2+} and heart failure: roles of diastolic leak and Ca^{2+} transport. *Circ Res* 2003;93:487-90.
17. Blair SN, Church TS. The fitness, obesity, and health equation: is physical activity the common denominator? *JAMA* 2004;292:1232-4.
18. Blair SN, Kohl HW 3rd, Barlow CE, Paffenbarger RS Jr, Gibbons LW, Macera CA. Changes in physical fitness and all-cause mortality. A prospective study of healthy and unhealthy men. *JAMA* 1995;273:1093-8.
19. Boluyt MO, Li ZB, Loyd AM, Scalia AF, Cirrincione GM, Jackson RR. The mTOR/p70S6K signal transduction pathway plays a role in cardiac hypertrophy and influences expression of myosin heavy chain genes in vivo. *Cardiovasc Drugs Ther* 2004;18:257-67.
20. Booth FW, Gordon SE, Carlson CJ, Hamilton MT. Waging war on modern chronic diseases: primary prevention through exercise biology. *J Appl Physiol* 2000;88:774-87.
21. Borzak S, Murphy S, Marsh JD. Mechanisms of rate staircase in rat ventricular cells. *Am J Physiol* 1991;260:H884-92.
22. Brixius K, Hoischen S, Reuter H, Lasek K, Schwinger RH. Force/shortening-frequency relationship in multicellular muscle strips and single cardiomyocytes of human failing and nonfailing hearts. *J Card Fail* 2001;7:335-41.
23. Bruckdorfer R. The basics about nitric oxide. *Mol Aspects Med* 2005;26:3-31.
24. Bøkenes J, Sjaastad I, Sejersted OM. Artfactual contractions triggered by field stimulation of cardiomyocytes. *J Appl Physiol* 2005;98:1712-9.
25. Ceci M, Ross J Jr, Condorelli G. Molecular determinants of the physiological adaptation to stress in the cardiomyocyte: a focus on AKT. *J Mol Cell Cardiol* 2004;37:905-12.
26. Cerretelli P, di Prampero PE. Gas exchange in exercise. In: Fishman AP et al (Eds). *Handbook of Physiology* Section 3, Chapter 16:297-339. American Physiological Society, Bethesda, MD. 1987.
27. Chen X, Piacentino V 3rd, Furukawa S, Goldman B, Margulies KB, Houser SR. L-type Ca^{2+} channel density and regulation are altered in failing human ventricular myocytes and recover after support with mechanical assist devices. *Circ Res* 2002;91:517-24.
28. Chien KR, Zhu H, Knowlton KU, Miller-Hance W, van-Bilsen M, O'Brien TX, Evans SM. Transcriptional regulation during cardiac growth and development. *Annu Rev Physiol* 1993;55:77-95.
29. Clarkson P, Montgomery HE, Mullen MJ, Donald AE, Powe AJ, Bull T, Jubb M, World M, Deanfield JE. Exercise training enhances endothelial function in young men. *J Am Coll Cardiol* 1999;33:1379-85.
30. Cleland JG, Khand A, Clark A. The heart failure epidemic: exactly how big is it? *Eur Heart J* 2001;22:623-26.
31. Collins KA, Korcarz CE, Shroff SG, Bednarz JE, Fentzke RC, Lin H, Leiden JM, Lang RM. Accuracy of echocardiographic estimates of left ventricular mass in mice. *Am J Physiol Heart Circ Physiol* 2001;280:H1954-62.
32. Coyle EF, Martin WH 3rd, Sinacore DR, Joyner MJ, Hagberg JM, Holloszy JO. Time course of loss of adaptations after stopping prolonged intense endurance training. *J Appl Physiol* 1984;57:1857-64.

33. Craig BW, Martin G, Betts J, Lungren M, Lambret V, Kaiserauer S. The influence of training-detraining upon the heart, muscle and adipose tissue of female rats. *Mech Ageing Dev* 1991;57:49-61.
34. Cullinane EM, Sady SP, Vadeboncoeur L, Burke M, Thompson PD. Cardiac size and VO_{2max} do not decrease after short-term exercise cessation. *Med Sci Sports Exerc* 1986;18:420-4.
35. Currie S, Quinn FR, Sayeed RA, Duncan AM, Kettlewell S, Smith GL. Selective down-regulation of sub-endocardial ryanodine receptor expression in a rabbit model of left ventricular dysfunction. *J Mol Cell Cardiol* 2005;39:309-17.
36. Currie S, Smith GL. Enhanced phosphorylation of phospholamban and downregulation of sarco/endoplasmic reticulum Ca^{2+} ATPase type 2 (SERCA 2) in cardiac sarcoplasmic reticulum from rabbits with heart failure. *Cardiovasc Res* 1999;41:135-46.
37. Darveau CA, Suarez RK, Andrews RD, Hochachka PW. Allometric cascade as a unifying principle of body mass effects on metabolism. *Nature* 2002;417:166-70.
38. Davies CH, Davia K, Bennett JG, Pepper JR, Poole-Wilson PA, Harding SE. Reduced contraction and altered frequency response of isolated ventricular myocytes from patients with heart failure. *Circulation* 1995;92:2540-9.
39. Davies CT, Knibbs AV. The training stimulus. The effects of intensity, duration and frequency of effort on maximum aerobic power output. *Int Z Angew Physiol* 1971;29:299-305.
40. Davis ME, Cai H, McCann L, Fukai T, Harrison DG. Role of c-Src in regulation of endothelial nitric oxide synthase expression during exercise training. *Am J Physiol Heart Circ Physiol* 2003;284:H1449-53.
41. del Monte F, Harding SE, Schmidt U, Matsui T, Kang ZB, Dec GW, Gwathmey JK, Rosenzweig A, Hajjar RJ. Restoration of contractile function in isolated cardiomyocytes from failing human hearts by gene transfer of SERCA2a. *Circulation* 1999;100:2308-11.
42. del Monte F, Williams E, Lebeche D, Schmidt U, Rosenzweig A, Gwathmey JK, Lewandowski ED, Hajjar RJ. Improvement in survival and cardiac metabolism after gene transfer of sarcoplasmic reticulum Ca^{2+} -ATPase in a rat model of heart failure. *Circulation* 2001;104:1424-9.
43. Diffeo GM, Chung E. Altered single cell force-velocity and power properties in exercise-trained rat myocardium. *J Appl Physiol* 2003;94:1941-8.
44. Diffeo GM, Nagle DF. Exercise training alters length dependence of contractile properties in rat myocardium. *J Appl Physiol* 2003;94:1137-44.
45. Diffeo GM, Seversen EA, Stein TD, Johnson JA. Microarray expression analysis of effects of exercise training: increase in atrial MLC-1 in rat ventricles. *Am J Physiol Heart Circ Physiol* 2003;284:H830-7.
46. Diffeo GM, Seversen EA, Titus MM. Exercise training increases the Ca^{2+} sensitivity of tension in rat cardiac myocytes. *J Appl Physiol* 2001;91:309-15.
47. di Prampero PE, Ferretti G. Factors limiting maximal oxygen consumption in humans. *Respir Physiol* 1990;80:113-27.
48. Fabiato A. Inositol (1,4,5)-triphosphate induced release of Ca^{2+} from the sarcoplasmic reticulum of skinned cardiac cells. *Biophys J* 1986;49:190. Abstract.
49. Fenning A, Harrison G, Dwyer D, Rose-Meyer R, Brown L. Cardiac adaptation to endurance exercise in rats. *Mol Cell Biochem* 2003;251:51-9.

50. Ferguson S, Gledhill N, Jamnik VK, Wiebe C, Payne N. Cardiac performance in endurance-trained and moderately active young women. *Med Sci Sports Exerc* 2001;33:1114-9.
51. Ferrier GR, Howlett SE. Cardiac excitation-contraction coupling: role of membrane potential in regulation of contraction. *Am J Physiol Heart Circ Physiol* 2001;280:H1928-44.
52. Fletcher GF, Balady GJ, Amsterdam EA, Chaitman B, Eckel R, Fleg J, Froelicher VF, Leon AS, Pina IL, Rodney R, Simons-Morton DA, Williams MA, Bazzarre T. Exercise standards for testing and training: a statement for healthcare professionals from the American Heart Association. *Circulation* 2001;104:1694-740.
53. Frampton JE, Orchard CH, Boyett MR. Diastolic, systolic and sarcoplasmic reticulum [Ca^{2+}] during inotropic interventions in isolated rat myocytes. *J Physiol* 1991;437:351-75.
54. Frank KF, Bolck B, Brixius K, Kranias EG, Schwinger RH. Modulation of SERCA: implications for the failing human heart. *Basic Res Cardiol* 2002;97:172-8.
55. Frank KF, Bolck B, Ding Z, Krause D, Hattebuhr N, Malik A, Brixius K, Hajjar RJ, Schrader J, Schwinger RH. Overexpression of sorcin enhances cardiac contractility in vivo and in vitro. *J Mol Cell Cardiol* 2005;38:607-15.
56. Frenzel H, Schwartzkopff B, Holtermann W, Schnurch HG, Novi A, Hort W. Regression of cardiac hypertrophy: morphometric and biochemical studies in rat heart after swimming training. *J Mol Cell Cardiol* 1988;20:737-51.
57. Furchgott RF, Zawadzki JV. The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. *Nature* 1980;288:373-6.
58. George KP, Wolfe LA, Burggraf GW, Norman R. Electrocardiographic and echocardiographic characteristics of female athletes. *Med Sci Sports Exerc* 1995;27:1362-70.
59. Gledhill N, Cox D, Jamnik R. Endurance athletes' stroke volume does not plateau: major advantage is diastolic function. *Med Sci Sports Exerc* 1994;26:1116-21.
60. Gomez AM, Valdivia HH, Cheng H, Lederer MR, Santana LF, Cannell MB, McCune SA, Altschuld RA, Lederer WJ. Defective excitation-contraction coupling in experimental cardiac hypertrophy and heart failure. *Science* 1997;276(5313):800-6.
61. Goto C, Higashi Y, Kimura M, Noma K, Hara K, Nakagawa K, Kawamura M, Chayama K, Yoshizumi M, Nara I. Effect of different intensities of exercise on endothelium-dependent vasodilation in humans: role of endothelium-dependent nitric oxide and oxidative stress. *Circulation* 2003;108:530-5.
62. Gregg EW, Cauley JA, Stone K, Thompson TJ, Bauer DC, Cummings SR, Ensrud KE; Study of Osteoporotic Fractures Research Group. Relationship of changes in physical activity and mortality among older women. *JAMA* 2003;289:2379-86.
63. Grohe C, Kahlert S, Lobbert K, Stimpel M, Karas RH, Vetter H, Neyses L. Cardiac myocytes and fibroblasts contain functional estrogen receptors. *FEBS Lett* 1997;416:107-12.
64. Grynkiewicz G, Poenie M, Tsien RY. A new generation of Ca^{2+} indicators with greatly improved fluorescence properties. *J Biol Chem* 1985;260:3440-50.
65. Gulati M, Pandey DK, Arnsdorf MF, Lauderdale DS, Thisted RA, Wicklund RH, Al-Hani AJ, Black HR. Exercise capacity and the risk of death in women. The St James women take heart project. *Circulation* 2003;108:1554-9.

66. Gwathmey JK, Copelas L, MacKinnon R, Schoen FJ, Feldman MD, Grossman W, Morgan JP. Abnormal intracellular calcium handling in myocardium from patients with end-stage heart failure. *Circ Res* 1987;61:70-6.
67. Gwathmey JK, Hajjar RJ. Relation between steady-state force and intracellular $[Ca^{2+}]$ in intact human myocardium. Index of myofibrillar responsiveness to Ca^{2+} . *Circulation* 1990;82:1266-78.
68. Gwathmey JK, Slawsky MT, Hajjar RJ, Briggs GM, Morgan JP. Role of intracellular calcium handling in force-interval relationships of human ventricular myocardium. *J Clin Invest* 1990;85:1599-613.
69. Hajjar RJ, Gwathmey JK, Briggs GM, Morgan JP. Differential effect of DPI 201-106 on the sensitivity of the myofilaments to Ca^{2+} in intact and skinned trabeculae from control and myopathic human hearts. *J Clin Invest* 1988;82:1578-84.
70. Hajjar RJ, Schwinger RH, Schmidt U, Kim CS, Lebeche D, Doye AA, Gwathmey JK. Myofilament calcium regulation in human myocardium. *Circulation* 2000;101:1679-85.
71. Hambrecht R, Adams V, Erbs S, Linke A, Krankel N, Shu Y, Baither Y, Gielen S, Thiele H, Gummert JF, Mohr FW, Schuler G. Regular physical activity improves endothelial function in patients with coronary artery disease by increasing phosphorylation of endothelial nitric oxide synthase. *Circulation* 2003;107:3152-8.
72. Hambrecht R, Walther C, Mobius-Winkler S, Gielen S, Linke A, Conradi K, Erbs S, Kluge R, Kendziorra K, Sabri O, Sick P, Schuler G. Percutaneous coronary angioplasty compared with exercise training in patients with stable coronary artery disease: a randomized trial. *Circulation* 2004;109:1371-8.
73. Hambrecht R, Wolf A, Gielen S, Linke A, Hofer J, Erbs S, Schoene N, Schuler G. Effect of exercise on coronary endothelial function in patients with coronary artery disease. *N Engl J Med* 2000;342:454-60.
74. Hare JM, Stamler JS. NO/redox disequilibrium in the failing heart and cardiovascular system. *J Clin Invest* 2005;115:509-17.
75. Hasenfuss G, Pieske B, Castell M, Kretschmann B, Maier LS, Just H. Influence of the novel inotropic agent levosimendan on isometric tension and calcium cycling in failing human myocardium. *Circulation* 1998;98:2141-7.
76. Hasenfuss G, Reinecke H, Studer R, Meyer M, Pieske B, Holtz J, Holubarsch C, Posival H, Just H, Drexler H. Relation between myocardial function and expression of sarcoplasmic reticulum Ca^{2+} -ATPase in failing and nonfailing human myocardium. *Circ Res* 1994;75:434-42.
77. Helgerud J. Maximal oxygen uptake, anaerobic threshold and running economy in women and men with similar performances level in marathons. *Eur J Appl Physiol Occup Physiol* 1994;68:155-61.
78. Hickson RC, Galassi TM, Dougherty KA. Repeated development and regression of exercise-induced cardiac hypertrophy in rats. *J Appl Physiol* 1983;54:794-7.
79. Hickson RC, Hammons GT, Holoszy JO. Development and regression of exercise-induced cardiac hypertrophy in rats. *Am J Physiol* 1979;236:H268-72.
80. Hirata M, Suematsu E, Hashimoto T, Hamachi T, Koga T. Release of Ca^{2+} from a non-mitochondrial store site in peritoneal macrophages treated with saponin by inositol 1,4,5-trisphosphate. *Biochem J* 1984;223:229-36.

81. Hochachka PW, Darveau CA, Andrews RD, Suarez RK. Allometric cascade: a model for resolving body mass effects on metabolism. *Comp Biochem Physiol A Mol Integr Physiol* 2003;134:675-91.
82. Holt E, Christensen G. Transient Ca²⁺ overload alters Ca²⁺ handling in rat cardiomyocytes: effects on shortening and relaxation. *Am J Physiol* 1997;273:H573-82.
83. Hunter JJ, Chien KR. Signaling pathways for cardiac hypertrophy and failure. *N Engl J Med* 1999;341:1276-83.
84. Hwang JJ, Allen PD, Tseng GC, Lam C-W, Fananapazir L, Dzau VJ, Liew C-C. Microarray gene expression profiles in dilated and hypertrophic cardiomyopathic end-stage heart failure. *Physiol Genomics* 2002;10:31-44.
85. Iemitsu M, Miyauchi T, Maeda S, Sakai S, Fujii N, Miyazaki H, Kakinuma Y, Matsuda M, Yamaguchi I. Cardiac hypertrophy by hypertension and exercise training exhibits different gene expression of enzymes in energy metabolism. *Hypertens Res* 2003;26:829-37.
86. Iemitsu M, Miyauchi T, Maeda S, Sakai S, Kobayashi T, Fujii N, Miyazaki H, Matsuda M, Yamaguchi I. Physiological and pathological cardiac hypertrophy induce different molecular phenotypes in the rat. *Am J Physiol Regul Integr Comp Physiol* 2001;281:R2029-36.
87. Ignarro LJ, Buga GM, Wood KS, Byrns RE, Chaudhuri G. Endothelium-derived relaxing factor produced and released from artery and vein is nitric oxide. *Proc Natl Acad Sci* 1987;84:9265-9.
88. Isgaard J, Kujacic V, Jennische E, Holmang A, Sun XY, Hedner T, Hjalmarson A, Bengtsson BA. Growth hormone improves cardiac function in rats with experimental myocardial infarction. *Eur J Clin Invest* 1997;27:517-25.
89. Jensen BE, Fletcher BJ, Rupp JC, Fletcher GF, Lee JY, Oberman A. Training level comparison study. Effect of high and low intensity exercise on ventilatory threshold in men with coronary artery disease. *J Cardiopulm Rehabil* 1996;16:227-32.
90. Jensen K, Johansen L, Secher NH. Influence of body mass on maximal oxygen uptake: effect of sample size. *Eur J Appl Physiol* 2001;84:201-5.
91. Jolliffe JA, Rees K, Taylor RS, Thompson D, Oldridge N, Ebrahim S. Exercise-based rehabilitation for coronary heart disease. *Cochrane Database Syst Rev* 2001;1:CD001800.
92. Kaplan ML, Cheslow Y, Vikstrom K, Malhotra A, Geenen DL, Nakouzi A, Leinwand LA, Buttrick PM. Cardiac adaptations to chronic exercise in mice. *Am J Physiol* 1994;267:H1167-73.
93. Kavanagh T, Mertens DJ, Hamm LF, Beyene J, Kennedy J, Corey P, Shephard RJ. Prediction of long-term prognosis in 12 169 men referred for cardiac rehabilitation. *Circulation* 2002;106:666-71.
94. Kentish JC, Barsotti RJ, Lea TJ, Mulligan IP, Patel JR, Ferenczi MA. Calcium release from cardiac sarcoplasmic reticulum induced by photorelease of calcium or Ins(1,4,5)P₃. *Am J Physiol* 1990;258:H610-5.
95. Kingwell BA, Arnold PJ, Jennings GL, Dart AM. Spontaneous running increases aortic compliance in Wistar-Kyoto rats. *Cardiovasc Res* 1997;35:132-7.
96. Klausen K, Andersen LB, Pelle I. Adaptive changes in work capacity, skeletal muscle capillarization and enzyme levels during training and detraining. *Acta Physiol Scand* 1981;113:9-16.

97. Laughlin MH, Joseph B. Wolfe Memorial lecture. Physical activity in prevention and treatment of coronary disease: the battle line is in exercise vascular cell biology. *Med Sci Sports Exerc* 2004;36:352-62.
98. Laughlin MH, Schaefer ME, Sturek M. Effect of exercise training on intracellular free Ca^{2+} transients in ventricular myocytes of rats. *J Appl Physiol* 1992;73:1441-8.
99. Laughlin MH, Woodman CR, Schrage WG, Gute D, Price EM. Interval sprint training enhances endothelial function and eNOS content in some arteries that perfuse white gastrocnemius muscle. *J Appl Physiol* 2004;96:233-44.
100. Layland J, Cave AC, Warren C, Grieve DJ, Sparks E, Kentish JC, Solaro RJ, Shah AM. Protection against endotoxemia-induced contractile dysfunction in mice with cardiac-specific expression of slow skeletal troponin I. *FASEB J* 2005;19:1137-9.
101. Lee IM, Sesso HD, Oguma Y, Paffenbarger RS Jr. Relative intensity of physical activity and risk of coronary heart disease. *Circulation* 2003;107:1110-6.
102. Liang Q, De Windt LJ, Witt SA, Kimball TR, Markham BE, Molkentin JD. The transcription factors GATA4 and GATA6 regulate cardiomyocyte hypertrophy in vitro and in vivo. *J Biol Chem* 2001;276:30245-53.
103. Lim CC, Apstein CS, Colucci WS, Liao R. Impaired cell shortening and relengthening with increased pacing frequency are intrinsic to the senescent mouse cardiomyocyte. *J Mol Cell Cardiol* 2000;32:2075-82.
104. Lindner M, Brandt MC, Sauer H, Hescheler J, Bohle T, Beuckelmann DJ. Calcium sparks in human ventricular cardiomyocytes from patients with terminal heart failure. *Cell Calcium* 2002;31:175-82.
105. Lindner M, Erdmann E, Beuckelmann DJ. Calcium content of the sarcoplasmic reticulum in isolated ventricular myocytes from patients with terminal heart failure. *J Mol Cell Cardiol* 1998;30:743-9.
106. Linke A, Schoene N, Gielen S, Hofer J, Erbs S, Schuler G, Hambrecht R. Endothelial dysfunction in patients with chronic heart failure: systemic effects of lower-limb exercise training. *J Am Coll Cardiol* 2001;37:392-7.
107. Litwin SE, Katz SE, Morgan JP, Douglas PS. Serial echocardiographic assessment of left ventricular geometry and function after large myocardial infarction in the rat. *Circulation* 1994;89:345-54.
108. Loennechen JP, Stoylen A, Beisvag V, Wisloff U, Ellingsen O. Regional expression of endothelin-1, ANP, IGF-1, and LV wall stress in the infarcted rat heart. *Am J Physiol Heart Circ Physiol* 2001;280:H2902-10.
109. Loennechen JP, Wisløff U, Falck G, Ellingsen Ø. Cardiomyocyte contractility and calcium handling partially recover after early deterioration during post-infarction failure in rat. *Acta Physiol Scand* 2002A;176:17-26.
110. Loennechen JP, Wisløff U, Falck G, Ellingsen Ø. Effects of carvedilol and losartan on hypertrophy, calcium transients, contractility, and gene expression in congestive heart failure. *Circulation* 2002B;105:1380-6.
111. Louch WE, Bito V, Heinzel FR, Macianskiene R, Vanhaecke J, Flameng W, Mubagwa K, Sipido KR. Reduced synchrony of Ca^{2+} release with loss of T-tubules—a comparison to Ca^{2+} release in human failing cardiomyocytes. *Cardiovasc Res* 2004;62:63-73.

112. Loughrey CM, MacEachern KE, Neary P, Smith GL. The relationship between intracellular $[Ca^{2+}]$ and Ca^{2+} wave characteristics in permeabilised cardiomyocytes from the rabbit. *J Physiol* 2002;543:859-70.
113. Loughrey CM, Seidler T, Miller SL, Prestle J, MacEachern KE, Reynolds DF, Hasenfuss G, Smith GL. Over-expression of FK506-binding protein FKBP12.6 alters excitation-contraction coupling in adult rabbit cardiomyocytes. *J Physiol* 2004;556:919-34.
114. Lu L, Mei DF, Gu AG, Wang S, Lentzner B, Gutstein DE, Zwas D, Homma S, Yi GH, Wang J. Exercise training normalizes altered calcium-handling proteins during development of heart failure. *J Appl Physiol* 2002;92:1524-30.
115. Ludmer PL, Selwyn AP, Shook TL, Wayne RR, Mudge GH, Alexander RW, Ganz P. Paradoxical vasoconstriction induced by acetylcholine in atherosclerotic coronary arteries. *N Engl J Med* 1986;315:1046-51.
116. Maeda S, Miyauchi T, Kakiyama T, Sugawara J, Iemitsu M, Irukayama-Tomobe Y, Murakami H, Kumagai Y, Kuno S, Matsuda M. Effects of exercise training of 8 weeks and detraining on plasma levels of endothelium-derived factors, endothelin-1 and nitric oxide, in healthy young humans. *Life Sci* 2001;69:1005-16.
117. Maiorana A, O'Driscoll G, Dembo L, Cheetham C, Goodman C, Taylor R, Green D. Effect of aerobic and resistance exercise training on vascular function in heart failure. *Am J Physiol Heart Circ Physiol* 2000;279:H1999-2005.
118. Malik S, Wong ND, Franklin SS, Kamath TV, L'Italien GJ, Pio JR, Williams GR. Impact of the metabolic syndrome on mortality from coronary heart disease, cardiovascular disease, and all causes in United States adults. *Circulation* 2004;110:1245-50.
119. Marks AR, Reiken S, Marx SO. Progression of heart failure: is protein kinase a hyperphosphorylation of the ryanodine receptor a contributing factor? *Circulation* 2002;105:272-5.
120. Martin WH 3rd, Coyle EF, Bloomfield SA, Ehsani AA. Effects of physical deconditioning after intense endurance training on left ventricular dimensions and stroke volume. *J Am Coll Cardiol* 1986;7:982-9.
121. Marx SO, Reiken S, Hisamatsu Y, Jayaraman T, Burkhoff D, Rosembliit N, Marks AR. PKA phosphorylation dissociates FKBP12.6 from the calcium release channel (ryanodine receptor): defective regulation in failing hearts. *Cell* 2000;101:365-76.
122. Massett MP, Berk BC. Strain-dependent differences in responses to exercise training in inbred and hybrid mice. *Am J Physiol Regul Integr Comp Physiol* 2005;288:R1006-13.
123. McAllister RM, Kimani JK, Webster JL, Parker JL, Laughlin MH. Effects of exercise training on responses of peripheral and visceral arteries in swine. *J Appl Physiol* 1996;80:216-25.
124. McAllister RM, Laughlin MH. Short-term exercise training alters responses of porcine femoral and brachial arteries. *J Appl Physiol* 1997;82:1438-44.
125. McMullen JR, Shioi T, Zhang L, Tarnavski O, Sherwood MC, Kang PM, Izumo S. Phosphoinositide 3-kinase(p110alpha) plays a critical role for the induction of physiological, but not pathological, cardiac hypertrophy. *Proc Natl Acad Sci* 2003;100:12355-60.
126. Meyer K, Gornandt L, Schwaibold M, Westbrook S, Hajric R, Peters K, Beneke R, Schnellbacher K, Roskamm H. Predictors of response to exercise training in severe chronic congestive heart failure. *Am J Cardiol* 1997;80:56-60.

127. Moe IT, Hoven H, Hetland EV, Rognmo Ø, Slørdahl SA. Endothelial function in highly endurance-trained and sedentary, healthy young women. *Vasc Med* 2005;10:97-102.
128. Mokolke EA, Palmer BM, Cheung JY, Moore RL. Endurance training does not affect intrinsic calcium current characteristics in rat myocardium. *Am J Physiol* 1997;273:H1193-7.
129. Moore RL, Musch TI, Yelamarty RV, Scaduto RC Jr, Semanchick AM, Elensky M, Cheung JY. Chronic exercise alters contractility and morphology of isolated rat cardiac myocytes. *Am J Physiol* 1993;264:C1180-9.
130. Moraska A, Deak T, Spencer RL, Roth D, Fleshner M. Treadmill running produces both positive and negative physiological adaptations in Sprague-Dawley rats. *Am J Physiol Regul Integr Comp Physiol* 2000;279:R1321-9.
131. Most P, Bernotat J, Ehlermann P, Pleger ST, Reppel M, Borries M, Niroomand F, Pieske B, Janssen PM, Eschenhagen T, Karczewski P, Smith GL, Koch WJ, Katus HA, Rempis A. S100A1: a regulator of myocardial contractility. *Proc Natl Acad Sci* 2001;98:13889-94.
132. Mujika I, Padilla S. Cardiorespiratory and metabolic characteristics of detraining in humans. *Med Sci Sports Exerc* 2001;33:413-21.
133. Mulieri LA, Hasenfuss G, Leavitt B, Allen PD, Alpert NR. Altered myocardial force-frequency relation in human heart failure. *Circulation* 1992;85:1743-50.
134. Musch TI, Moore RL, Smaldone PG, Riedy M, Zelis R. Cardiac adaptations to endurance training in rats with a chronic myocardial infarction. *J Appl Physiol* 1989;66:712-9.
135. Musi N, Hirshman MF, Arad M, Xing Y, Fujii N, Pomerleau J, Ahmad F, Berul CI, Seidman JG, Tian R, Goodyear LJ. Functional role of AMP-activated protein kinase in the heart during exercise. *FEBS Lett* 2005;579:2045-50.
136. Myers J, Prakash M, Froelicher V, Do D, Partington S, Atwood JE. Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med* 2002;346:793-801.
137. Nakao K, Minobe W, Roden R, Bristow MR, Leinwand LA. Myosin heavy chain gene expression in human heart failure. *J Clin Invest* 1997;100:2362-70.
138. Naseem KM. The role of nitric oxide in cardiovascular diseases. *Mol Aspects Med* 2005;26:33-65.
139. Ogut O, Brozovich FV. Regulation of force in vascular smooth muscle. *J Mol Cell Cardiol* 2003;35:347-55.
140. O'Neill JO, Young JB, Pothier CE, Lauer MS. Peak oxygen consumption as a predictor of death in patients with heart failure receiving beta-blockers. *Circulation* 2005;111:2313-8.
141. Orchard CH, Kentish JC. Effects of changes of pH on the contractile function of cardiac muscle. *Am J Physiol* 1990;258:C967-81.
142. Orshal JM, Khalil RA. Gender, sex hormones, and vascular tone. *Am J Physiol Regul Integr Comp Physiol* 2004;286:R233-49.
143. Paffenbarger RS Jr, Hyde RT, Wing AL, Lee IM, Jung DL, Kampert JB. The association of changes in physical-activity level and other lifestyle characteristics with mortality among men. *N Engl J Med* 1993;328:538-45.
144. Palmer BM, Thayer AM, Snyder SM, Moore RL. Shortening and $[Ca^{2+}]$ dynamics of left ventricular myocytes isolated from exercise-trained rats. *J Appl Physiol* 1998;85:2159-68.

145. Pate RR, Kriska A. Physiological basis of the sex difference in cardiorespiratory endurance. *Sports Med* 1984;1:87-98.
146. Pelliccia A, Maron BJ, Culasso F, Spataro A, Caselli G. Athlete's heart in women. Echocardiographic characterization of highly trained elite female athletes. *JAMA* 1996;276:211-5.
147. Pelliccia A, Maron BJ, De Luca R, Di Paolo FM, Spataro A, Culasso F. Remodeling of left ventricular hypertrophy in elite athletes after long-term deconditioning. *Circulation* 2002;105:944-9.
148. Perez NG, Hashimoto K, McCune S, Altschuld RA, Marban E. Origin of contractile dysfunction in heart failure; calcium cycling versus myofilaments. *Circulation* 1999;99:1077-83.
149. Pieske B, Maier LS, Bers DM, Hasenfuss G. Ca²⁺ handling and sarcoplasmic reticulum Ca²⁺ content in isolated failing and nonfailing human myocardium. *Circ Res* 1999;85:38-46.
150. Pluim BM, Zwinderman AH, van der Laarse A, van der Wall EE. The athlete's heart. A meta-analysis of cardiac structure and function. *Circulation* 2000;101:336-44.
151. Quinn FR, Currie S, Duncan AM, Miller S, Sayeed R, Cobbe SM, Smith GL. Myocardial infarction causes increased expression but decreased activity of the myocardial Na⁺-Ca²⁺ exchanger in the rabbit. *J Physiol* 2003;553:229-42.
152. Raven PB, Shi X. Cardiovascular adaptations following detraining. *J Gravit Physiol* 1995;21:P17-8.
153. Reuter H, Pott C, Goldhaber JI, Henderson SA, Philipson KD, Schwinger RH. Na⁺-Ca²⁺ exchange in the regulation of cardiac excitation-contraction coupling. *Cardiovasc Res* 2005;67:198-207.
154. Richardson RS. Oxygen transport: air to muscle cell. *Med Sci Sports Exerc* 1998;30:53-9.
155. Richardson RS. What governs skeletal muscle VO_{2max}? New evidence. *Med Sci Sports Exerc* 2000;32:100-7.
156. Richardson RS, Grassi B, Gavin TP, Haseler LJ, Tagore K, Roca J, Wagner PD. Evidence of O₂ supply-dependent VO_{2max} in the exercise-trained human quadriceps. *J Appl Physiol* 1999;86:1048-53.
157. Rognmo Ø, Hetland E, Helgerud J, Hoff J, Slørdahl SA. High intensity aerobic interval exercise is superior to moderate intensity exercise for increasing aerobic capacity in patients with coronary artery disease. *Eur J Cardiovasc Prev Rehabil* 2004;11:216-22.
158. Russell JC, Epling WF, Pierce D, Amy RM, Boer DP. Induction of voluntary prolonged running by rats. *J Appl Physiol* 1987;63:2549-53.
159. Sahn DJ, DeMaria A, Kisslo J, Weyman A. Recommendations regarding quantitation in M-mode echocardiography: results of a survey of echocardiographic measurements. *Circulation* 1978;58:1072-83.
160. Saltin B. Maximal oxygen uptake: limitation and malleability. In: Kane K and Ryan T (Eds). *International Symposium on Exercise Physiology* 26-40. Human Kinetics, Champaign, IL. 1990.

161. Schachinger V, Britten MB, Zeiher AM. Prognostic impact of coronary vasodilator dysfunction on adverse long-term outcome of coronary heart disease. *Circulation* 2000;101:1899-906.
162. Schaible TF, Scheuer J. Cardiac function in hypertrophied hearts from chronically exercised female rats. *J Appl Physiol* 1981;50:1140-5.
163. Schmidt U, Hajjar RJ, Helm PA, Kim CS, Doye AA, Gwathmey JK. Contribution of abnormal sarcoplasmic reticulum ATPase activity to systolic and diastolic dysfunction in human heart failure. *J Mol Cell Cardiol* 1998;30:1929-37.
164. Schwinger RH, Bohm M, Koch A, Schmidt U, Morano I, Eissner HJ, Uberfuhr P, Reichart B, Erdmann E. The failing human heart is unable to use the Frank-Starling mechanism. *Circ Res* 1994;74:959-69.
165. Sjaastad I, Birkeland JA, Ferrier G, Howlett S, Skomedal T, Bjørnerheim R, Wasserstrom JA, Sejersted OM. Defective excitation-contraction coupling in hearts of rats with congestive heart failure. *Acta Physiol Scand* 2005;184:45-58.
166. Sjaastad I, Sejersted OM, Ilebakk A, Bjørnerheim R. Echocardiographic criteria for detection of postinfarction congestive heart failure in rats. *J Appl Physiol* 2000;89:1445-54.
167. Stehle R, Kruger M, Scherer P, Brixius K, Schwinger RH, Pfitzer G. Isometric force kinetics upon rapid activation and relaxation of mouse, guinea pig and human heart muscle studied on the subcellular myofibrillar level. *Basic Res Cardiol* 2002;97:1127-35.
168. Støen R, Brubakk AM, Vik T, Lossius K, Jynge P, Karlsson JO. Postnatal changes in mechanisms mediating acetylcholine-induced relaxation in piglet femoral arteries. *Pediatr Res* 1997;41:702-7.
169. Tanaka N, Dalton N, Mao L, Rockman HA, Peterson KL, Gottshall KR, Hunter JJ, Chien KR, Ross J Jr. Transthoracic echocardiography in models of cardiac disease in the mouse. *Circulation* 1996;94:1109-17.
170. Tanasescu M, Leitzmann MF, Rimm EB, Willett WC, Stampfer MJ, Hu FB. Exercise type and intensity in relation to coronary heart disease in men. *JAMA* 2002;288:1994-2000.
171. Taylor CR, Maloij GM, Weibel ER, Langman VA, Kamau JM, Seeherman HJ, Heglund NC. Design of the mammalian respiratory system. III Scaling maximum aerobic capacity to body mass: wild and domestic mammals. *Respir Physiol* 1981;44:25-37.
172. Teucher N, Prestle J, Seidler T, Currie S, Elliott EB, Reynolds DF, Schott P, Wagner S, Kogler H, Inesi G, Bers DM, Hasenfuss G, Smith GL. Excessive sarcoplasmic/endoplasmic reticulum Ca^{2+} -ATPase expression causes increased sarcoplasmic reticulum Ca^{2+} uptake but decreases myocyte shortening. *Circulation* 2004;110:3553-9.
173. Trafford AW, Eisner DA. No role for a voltage sensitive release mechanism in cardiac muscle. *J Mol Cell Cardiol* 2003;35:145-51.
174. van der Velden J, Boontje NM, Papp Z, Klein LJ, Visser FC, de Jong JW, Owen VJ, Burton PB, Stienen GJ. Calcium sensitivity of force in human ventricular cardiomyocytes from donor and failing hearts. *Basic Res Cardiol* 2002;97:1118-26.
175. Veras-Silva AS, Mattos KC, Gava NS, Brum PC, Negrao CE, Krieger EM. Low-intensity exercise training decreases cardiac output and hypertension in spontaneously hypertensive rats. *Am J Physiol* 1997;273:H2627-31.
176. Wagner PD. Algebraic analysis of the determinants of VO_{2max} . *Respir Physiol* 1993;93:221-37.

177. Wagner PD. A theoretical analysis of factors determining VO_{2max} at sea level and altitude. *Respir Physiol* 1996;106:329-43.
178. Wagner PD. New ideas on limitations to VO_{2max} . *Exerc Sports Sci Rev* 2000;28:10-4.
179. Walther C, Gielen S, Hambrecht R. The effect of exercise training on endothelial function in cardiovascular disease in humans. *Exerc Sport Sci Rev* 2004;32:129-34.
180. Walsh JH, Bilsborough W, Maiorana A, Best M, O'Driscoll GJ, Taylor RR, Green DJ. Exercise training improves conduit vessel function in patients with coronary artery disease. *J Appl Physiol* 2003;95:20-5.
181. Wasserman K, Hansen JE, Sue DY, Casaburi R, Whipp BJ. *Principles of Exercise Testing and Interpretation*. Lippincott, Williams and Wilkins, Baltimore, MD. 1999.
182. Wasserstrom JA, Holt E, Sjaastad I, Lunde PK, Ødegaard A, Sejersted OM. Altered E-C coupling in rat ventricular myocytes from failing hearts 6 wk after MI. *Am J Physiol Heart Circ Physiol* 2000;279:H798-807.
183. Wehrens XH, Lehnart SE, Reiken S, van der Nagel R, Morales R, Sun J, Cheng Z, Deng SX, de Windt LJ, Landry DW, Marks AR. Enhancing calstabin binding to ryanodine receptors improves cardiac and skeletal muscle function in heart failure. *Proc Natl Acad Sci* 2005;102:9607-12.
184. Weibel ER, Bacigalupe LD, Schmitt B, Hoppeler H. Allometric scaling of maximal metabolic rate in mammals: muscle aerobic capacity as determinant factor. *Respir Physiol Neurobiol* 2004;140:115-32.
185. Weibel ER, Hoppeler H. Exercise-induced maximal metabolic rate scales with muscle aerobic capacity. *J Exp Biol* 2005;208:1635-44.
186. Wiebe CG, Gledhill N, Warburton DE, Jamnik VK, Ferguson S. Exercise cardiac function in endurance-trained males versus females. *Clin J Sport Med* 1998;8:272-9.
187. Wisløff U, Loennechen JP, Currie S, Smith GL, Ellingsen Ø. Aerobic exercise reduces cardiomyocyte hypertrophy and increases contractility, Ca^{2+} sensitivity and SERCA-2 in rat after myocardial infarction. *Cardiovasc Res* 2002;54:162-74.
188. Wisløff U, Loennechen JP, Falck G, Beisvåg V, Currie S, Smith GL, Ellingsen O. Increased contractility and calcium sensitivity in cardiac myocytes isolated from endurance trained rats. *Cardiovasc Res* 2001;50:495-508.
189. Wisløff U, Najjar SM, Ellingsen Ø, Haram PM, Swoap S, Al-Share Q, Fernstrom M, Rezaei K, Lee SJ, Koch LG, Britton SL. Cardiovascular risk factors emerge after artificial selection for low aerobic capacity. *Science* 2005;307:418-20.
190. Zhang LQ, Zhang XQ, Musch TI, Moore RL, Cheung JY. Sprint training restores normal contractility in postinfarction rat myocytes. *J Appl Physiol* 2000A;89:1099-105.
191. Zhang LQ, Zhang XQ, Ng YC, Rothblum LI, Musch TI, Moore RL, Cheung JY. Sprint training normalizes Ca^{2+} transients and SR function in postinfarction rat myocytes. *J Appl Physiol* 2000B;89:38-46.
192. Zhang XQ, Ng YC, Musch TI, Moore RL, Zelis R, Cheung JY. Sprint training attenuates myocyte hypertrophy and improves Ca^{2+} homeostasis in postinfarction myocytes. *J Appl Physiol* 1998;84:544-52.
193. Zhang XQ, Song J, Carl LL, Shi W, Qureshi A, Tian Q, Cheung JY. Effects of sprint training on contractility and $[Ca^{2+}]_i$ transients in adult rat myocytes. *J Appl Physiol* 2002;93:1310-7.

194. Zhou B, Conlee RK, Jensen R, Fellingham GW, George JD, Fisher AG. Stroke volume does not plateau during graded exercise in elite male distance runners. *Med Sci Sports Exerc* 2001;33:1849-54.

195. Åstrand PO, Rodahl K. *Textbook of Work Physiology*. McGraw-Hill, New York, NY. 1986.

Dissertations at the Faculty of Medicine, NTNU

1977

1. Knut Joachim Berg: EFFECT OF ACETYLSALICYLIC ACID ON RENAL FUNCTION
2. Karl Erik Viken and Arne Ødegaard: STUDIES ON HUMAN MONOCYTES CULTURED *IN VITRO*

1978

3. Karel Bjørn Cyvin: CONGENITAL DISLOCATION OF THE HIP JOINT.
4. Alf O. Brubakk: METHODS FOR STUDYING FLOW DYNAMICS IN THE LEFT VENTRICLE AND THE AORTA IN MAN.

1979

5. Geirmund Unsgaard: CYTOSTATIC AND IMMUNOREGULATORY ABILITIES OF HUMAN BLOOD MONOCYTES CULTURED *IN VITRO*

1980

6. Størker Jørstad: URAEMIC TOXINS
7. Arne Olav Jenssen: SOME RHEOLOGICAL, CHEMICAL AND STRUCTURAL PROPERTIES OF MUCOID SPUTUM FROM PATIENTS WITH CHRONIC OBSTRUCTIVE BRONCHITIS

1981

8. Jens Hammerstrøm: CYTOSTATIC AND CYTOLYTIC ACTIVITY OF HUMAN MONOCYTES AND EFFUSION MACROPHAGES AGAINST TUMOR CELLS *IN VITRO*

1983

9. Tore Syversen: EFFECTS OF METHYLMERCURY ON RAT BRAIN PROTEIN.
10. Torbjørn Iversen: SQUAMOUS CELL CARCINOMA OF THE VULVA.

1984

11. Tor-Erik Widerøe: ASPECTS OF CONTINUOUS AMBULATORY PERITONEAL DIALYSIS.
12. Anton Hole: ALTERATIONS OF MONOCYTE AND LYMPHOCYTE FUNCTIONS IN REACTION TO SURGERY UNDER EPIDURAL OR GENERAL ANAESTHESIA.
13. Terje Terjesen: FRACTURE HEALING AND STRESS-PROTECTION AFTER METAL PLATE FIXATION AND EXTERNAL FIXATION.
14. Carsten Saunte: CLUSTER HEADACHE SYNDROME.
15. Inggard Lereim: TRAFFIC ACCIDENTS AND THEIR CONSEQUENCES.
16. Bjørn Magne Eggen: STUDIES IN CYTOTOXICITY IN HUMAN ADHERENT MONONUCLEAR BLOOD CELLS.
17. Trond Haug: FACTORS REGULATING BEHAVIORAL EFFECTS OF DRUGS.

1985

18. Sven Erik Gisvold: RESUSCITATION AFTER COMPLETE GLOBAL BRAIN ISCHEMIA.
19. Terje Espevik: THE CYTOSKELETON OF HUMAN MONOCYTES.
20. Lars Bevanger: STUDIES OF THE Ibc (c) PROTEIN ANTIGENS OF GROUP B STREPTOCOCCI.
21. Ole-Jan Iversen: RETROVIRUS-LIKE PARTICLES IN THE PATHOGENESIS OF PSORIASIS.
22. Lasse Eriksen: EVALUATION AND TREATMENT OF ALCOHOL DEPENDENT BEHAVIOUR.
23. Per I. Lundmo: ANDROGEN METABOLISM IN THE PROSTATE.

1986

24. Dagfinn Berntzen: ANALYSIS AND MANAGEMENT OF EXPERIMENTAL AND CLINICAL PAIN.
25. Odd Arnold Kildahl-Andersen: PRODUCTION AND CHARACTERIZATION OF MONOCYTE-DERIVED CYTOTOXIN AND ITS ROLE IN MONOCYTE-MEDIATED CYTOTOXICITY.
26. Ola Dale: VOLATILE ANAESTHETICS.

1987

27. Per Martin Kleveland: STUDIES ON GASTRIN.
28. Audun N. Øksendal: THE CALCIUM PARADOX AND THE HEART.
29. Vilhjalmur R. Finsen: HIP FRACTURES

1988

30. Rigmor Austgulen: TUMOR NECROSIS FACTOR: A MONOCYTE-DERIVED REGULATOR OF CELLULAR GROWTH.
31. Tom-Harald Edna: HEAD INJURIES ADMITTED TO HOSPITAL.
32. Joseph D. Borsi: NEW ASPECTS OF THE CLINICAL PHARMACOKINETICS OF METHOTREXATE.
33. Olav F. M. Sellevold: GLUCOCORTICOIDS IN MYOCARDIAL PROTECTION.

34. Terje Skjærpe: NONINVASIVE QUANTITATION OF GLOBAL PARAMETERS ON LEFT VENTRICULAR FUNCTION: THE SYSTOLIC PULMONARY ARTERY PRESSURE AND CARDIAC OUTPUT.
 35. Eyvind Rødahl: STUDIES OF IMMUNE COMPLEXES AND RETROVIRUS-LIKE ANTIGENS IN PATIENTS WITH ANKYLOSING SPONDYLITIS.
 36. Ketil Thorstensen: STUDIES ON THE MECHANISMS OF CELLULAR UPTAKE OF IRON FROM TRANSFERRIN.
 37. Anna Midelfart: STUDIES OF THE MECHANISMS OF ION AND FLUID TRANSPORT IN THE BOVINE CORNEA.
 38. Eirik Helseth: GROWTH AND PLASMINOGEN ACTIVATOR ACTIVITY OF HUMAN GLIOMAS AND BRAIN METASTASES - WITH SPECIAL REFERENCE TO TRANSFORMING GROWTH FACTOR BETA AND THE EPIDERMAL GROWTH FACTOR RECEPTOR.
 39. Petter C. Borchgrevink: MAGNESIUM AND THE ISCHEMIC HEART.
 40. Kjell-Arne Rein: THE EFFECT OF EXTRACORPOREAL CIRCULATION ON SUBCUTANEOUS TRANSCAPILLARY FLUID BALANCE.
 41. Arne Kristian Sandvik: RAT GASTRIC HISTAMINE.
 42. Carl Bredo Dahl: ANIMAL MODELS IN PSYCHIATRY.
- 1989
43. Torbjørn A. Fredriksen: CERVICOGENIC HEADACHE.
 44. Rolf A. Walstad: CEFTAZIDIME.
 45. Rolf Salvesen: THE PUPIL IN CLUSTER HEADACHE.
 46. Nils Petter Jørgensen: DRUG EXPOSURE IN EARLY PREGNANCY.
 47. Johan C. Ræder: PREMEDICATION AND GENERAL ANAESTHESIA IN OUTPATIENT GYNECOLOGICAL SURGERY.
 48. M. R. Shalaby: IMMUNOREGULATORY PROPERTIES OF TNF- α AND THE RELATED CYTOKINES.
 49. Anders Waage: THE COMPLEX PATTERN OF CYTOKINES IN SEPTIC SHOCK.
 50. Bjarne Christian Eriksen: ELECTROSTIMULATION OF THE PELVIC FLOOR IN FEMALE URINARY INCONTINENCE.
 51. Tore B. Halvorsen: PROGNOSTIC FACTORS IN COLORECTAL CANCER.
- 1990
52. Asbjørn Nordby: CELLULAR TOXICITY OF ROENTGEN CONTRAST MEDIA.
 53. Kåre E. Tvedt: X-RAY MICROANALYSIS OF BIOLOGICAL MATERIAL.
 54. Tore C. Stiles: COGNITIVE VULNERABILITY FACTORS IN THE DEVELOPMENT AND MAINTENANCE OF DEPRESSION.
 55. Eva Hofslis: TUMOR NECROSIS FACTOR AND MULTIDRUG RESISTANCE.
 56. Helge S. Haarstad: TROPHIC EFFECTS OF CHOLECYSTOKININ AND SECRETIN ON THE RAT PANCREAS.
 57. Lars Engebretsen: TREATMENT OF ACUTE ANTERIOR CRUCIATE LIGAMENT INJURIES.
 58. Tarjei Rygnestad: DELIBERATE SELF-POISONING IN TRONDHEIM.
 59. Arne Z. Henriksen: STUDIES ON CONSERVED ANTIGENIC DOMAINS ON MAJOR OUTER MEMBRANE PROTEINS FROM ENTEROBACTERIA.
 60. Steinar Westin: UNEMPLOYMENT AND HEALTH: Medical and social consequences of a factory closure in a ten-year controlled follow-up study.
 61. Ylva Sahlin: INJURY REGISTRATION, a tool for accident preventive work.
 62. Helge Bjørnstad Pettersen: BIOSYNTHESIS OF COMPLEMENT BY HUMAN ALVEOLAR MACROPHAGES WITH SPECIAL REFERENCE TO SARCOIDOSIS.
 63. Berit Schei: TRAPPED IN PAINFUL LOVE.
 64. Lars J. Vatten: PROSPECTIVE STUDIES OF THE RISK OF BREAST CANCER IN A COHORT OF NORWEGIAN WOMAN.
- 1991
65. Kåre Bergh: APPLICATIONS OF ANTI-C5a SPECIFIC MONOCLONAL ANTIBODIES FOR THE ASSESSMENT OF COMPLEMENT ACTIVATION.
 66. Svein Svenningsen: THE CLINICAL SIGNIFICANCE OF INCREASED FEMORAL ANTEVERSION.
 67. Olbjørn Klepp: NONSEMINOMATOUS GERM CELL TESTIS CANCER: THERAPEUTIC OUTCOME AND PROGNOSTIC FACTORS.
 68. Trond Sand: THE EFFECTS OF CLICK POLARITY ON BRAINSTEM AUDITORY EVOKED POTENTIALS AMPLITUDE, DISPERSION, AND LATENCY VARIABLES.
 69. Kjetil B. Åsbakk: STUDIES OF A PROTEIN FROM PSORIATIC SCALE, PSO P27, WITH RESPECT TO ITS POTENTIAL ROLE IN IMMUNE REACTIONS IN PSORIASIS.

70. Arnulf Hestnes: STUDIES ON DOWN'S SYNDROME.
 71. Randi Nygaard: LONG-TERM SURVIVAL IN CHILDHOOD LEUKEMIA.
 72. Bjørn Hagen: THIO-TEPA.
 73. Svein Anda: EVALUATION OF THE HIP JOINT BY COMPUTED TOMOGRAPHY AND ULTRASONOGRAPHY.
- 1992
74. Martin Svartberg: AN INVESTIGATION OF PROCESS AND OUTCOME OF SHORT-TERM PSYCHODYNAMIC PSYCHOTHERAPY.
 75. Stig Arild Slørdahl: AORTIC REGURGITATION.
 76. Harold C Sexton: STUDIES RELATING TO THE TREATMENT OF SYMPTOMATIC NON-PSYCHOTIC PATIENTS.
 77. Maurice B. Vincent: VASOACTIVE PEPTIDES IN THE OCULAR/FOREHEAD AREA.
 78. Terje Johannessen: CONTROLLED TRIALS IN SINGLE SUBJECTS.
 79. Turid Nilsen: PYROPHOSPHATE IN HEPATOCYTE IRON METABOLISM.
 80. Olav Haraldseth: NMR SPECTROSCOPY OF CEREBRAL ISCHEMIA AND REPERFUSION IN RAT.
 81. Eiliv Brenna: REGULATION OF FUNCTION AND GROWTH OF THE OXYNTIC MUCOSA.
- 1993
82. Gunnar Bovim: CERVICOGENIC HEADACHE.
 83. Jarl Arne Kahn: ASSISTED PROCREATION.
 84. Bjørn Naume: IMMUNOREGULATORY EFFECTS OF CYTOKINES ON NK CELLS.
 85. Rune Wiseth: AORTIC VALVE REPLACEMENT.
 86. Jie Ming Shen: BLOOD FLOW VELOCITY AND RESPIRATORY STUDIES.
 87. Piotr Kruszewski: SUNCT SYNDROME WITH SPECIAL REFERENCE TO THE AUTONOMIC NERVOUS SYSTEM.
 88. Mette Haase Moen: ENDOMETRIOSIS.
 89. Anne Vik: VASCULAR GAS EMBOLISM DURING AIR INFUSION AND AFTER DECOMPRESSION IN PIGS.
 90. Lars Jacob Stovner: THE CHIARI TYPE I MALFORMATION.
 91. Kjell Å. Salvesen: ROUTINE ULTRASONOGRAPHY IN UTERO AND DEVELOPMENT IN CHILDHOOD.
- 1994
92. Nina-Beate Liabakk: DEVELOPMENT OF IMMUNOASSAYS FOR TNF AND ITS SOLUBLE RECEPTORS.
 93. Sverre Helge Torp: *erbB* ONCOGENES IN HUMAN GLIOMAS AND MENINGIOMAS.
 94. Olav M. Linaker: MENTAL RETARDATION AND PSYCHIATRY. Past and present.
 95. Per Oscar Feet: INCREASED ANTIDEPRESSANT AND ANTIPANIC EFFECT IN COMBINED TREATMENT WITH DIXYRAZINE AND TRICYCLIC ANTIDEPRESSANTS.
 96. Stein Olav Samstad: CROSS SECTIONAL FLOW VELOCITY PROFILES FROM TWO-DIMENSIONAL DOPPLER ULTRASOUND: Studies on early mitral blood flow.
 97. Bjørn Backe: STUDIES IN ANTENATAL CARE.
 98. Gerd Inger Ringdal: QUALITY OF LIFE IN CANCER PATIENTS.
 99. Torvid Kiserud: THE DUCTUS VENOSUS IN THE HUMAN FETUS.
 100. Hans E. Fjøsne: HORMONAL REGULATION OF PROSTATIC METABOLISM.
 101. Eylert Brodtkorb: CLINICAL ASPECTS OF EPILEPSY IN THE MENTALLY RETARDED.
 102. Roar Juul: PEPTIDERGIC MECHANISMS IN HUMAN SUBARACHNOID HEMORRHAGE.
 103. Unni Syversen: CHROMOGRANIN A. Physiological and Clinical Role.
- 1995
104. Odd Gunnar Brakstad: THERMOSTABLE NUCLEASE AND THE *nuc* GENE IN THE DIAGNOSIS OF *Staphylococcus aureus* INFECTIONS.
 105. Terje Engan: NUCLEAR MAGNETIC RESONANCE (NMR) SPECTROSCOPY OF PLASMA IN MALIGNANT DISEASE.
 106. Kirsten Rasmussen: VIOLENCE IN THE MENTALLY DISORDERED.
 107. Finn Egil Skjeldestad: INDUCED ABORTION: Timetrends and Determinants.
 108. Roar Stenseth: THORACIC EPIDURAL ANALGESIA IN AORTOCORONARY BYPASS SURGERY.
 109. Arild Faxvaag: STUDIES OF IMMUNE CELL FUNCTION *in mice infected with* MURINE RETROVIRUS.
- 1996
110. Svend Aakhus: NONINVASIVE COMPUTERIZED ASSESSMENT OF LEFT VENTRICULAR FUNCTION AND SYSTEMIC ARTERIAL PROPERTIES. Methodology and some clinical applications.

111. Klaus-Dieter Bolz: INTRAVASCULAR ULTRASONOGRAPHY.
 112. Petter Aadahl: CARDIOVASCULAR EFFECTS OF THORACIC AORTIC CROSS-CLAMPING.
 113. Sigurd Steinshamm: CYTOKINE MEDIATORS DURING GRANULOCYTOPENIC INFECTIONS.
 114. Hans Stifoss-Hanssen: SEEKING MEANING OR HAPPINESS?
 115. Anne Kvikstad: LIFE CHANGE EVENTS AND MARITAL STATUS IN RELATION TO RISK AND PROGNOSIS OF CANCER.
 116. Torbjørn Grøntvedt: TREATMENT OF ACUTE AND CHRONIC ANTERIOR CRUCIATE LIGAMENT INJURIES. A clinical and biomechanical study.
 117. Sigrid Hørven Wigert: CLINICAL STUDIES OF FIBROMYALGIA WITH FOCUS ON ETIOLOGY, TREATMENT AND OUTCOME.
 118. Jan Schjøtt: MYOCARDIAL PROTECTION: Functional and Metabolic Characteristics of Two Endogenous Protective Principles.
 119. Marit Martinussen: STUDIES OF INTESTINAL BLOOD FLOW AND ITS RELATION TO TRANSITIONAL CIRCULATORY ADAPATION IN NEWBORN INFANTS.
 120. Tomm B. Müller: MAGNETIC RESONANCE IMAGING IN FOCAL CEREBRAL ISCHEMIA.
 121. Rune Haaverstad: OEDEMA FORMATION OF THE LOWER EXTREMITIES.
 122. Magne Børset: THE ROLE OF CYTOKINES IN MULTIPLE MYELOMA, WITH SPECIAL REFERENCE TO HEPATOCYTE GROWTH FACTOR.
 123. Geir Smedslund: A THEORETICAL AND EMPIRICAL INVESTIGATION OF SMOKING, STRESS AND DISEASE: RESULTS FROM A POPULATION SURVEY.
- 1997
124. Torstein Vik: GROWTH, MORBIDITY, AND PSYCHOMOTOR DEVELOPMENT IN INFANTS WHO WERE GROWTH RETARDED *IN UTERO*.
 125. Siri Forsmo: ASPECTS AND CONSEQUENCES OF OPPORTUNISTIC SCREENING FOR CERVICAL CANCER. Results based on data from three Norwegian counties.
 126. Jon S. Skranes: CEREBRAL MRI AND NEURODEVELOPMENTAL OUTCOME IN VERY LOW BIRTH WEIGHT (VLBW) CHILDREN. A follow-up study of a geographically based year cohort of VLBW children at ages one and six years.
 127. Knut Bjørnstad: COMPUTERIZED ECHOCARDIOGRAPHY FOR EVALUATION OF CORONARY ARTERY DISEASE.
 128. Grethe Elisabeth Borchgrevink: DIAGNOSIS AND TREATMENT OF WHIPLASH/NECK SPRAIN INJURIES CAUSED BY CAR ACCIDENTS.
 129. Tor Elsås: NEUROPEPTIDES AND NITRIC OXIDE SYNTHASE IN OCULAR AUTONOMIC AND SENSORY NERVES.
 130. Rolf W. Gråwe: EPIDEMIOLOGICAL AND NEUROPSYCHOLOGICAL PERSPECTIVES ON SCHIZOPHRENIA.
 131. Tonje Strømholm: CEREBRAL HAEMODYNAMICS DURING THORACIC AORTIC CROSSCLAMPING. An experimental study in pigs.
- 1998
132. Martinus Bråten: STUDIES ON SOME PROBLEMS RELATED TO INTRAMEDULLARY NAILING OF FEMORAL FRACTURES.
 133. Ståle Nordgård: PROLIFERATIVE ACTIVITY AND DNA CONTENT AS PROGNOSTIC INDICATORS IN ADENOID CYSTIC CARCINOMA OF THE HEAD AND NECK.
 134. Egil Lien: SOLUBLE RECEPTORS FOR TNF AND LPS: RELEASE PATTERN AND POSSIBLE SIGNIFICANCE IN DISEASE.
 135. Marit Bjørngaas: HYPOGLYCAEMIA IN CHILDREN WITH DIABETES MELLITUS
 136. Frank Skorpen: GENETIC AND FUNCTIONAL ANALYSES OF DNA REPAIR IN HUMAN CELLS.
 137. Juan A. Pareja: SUNCT SYNDROME. ON THE CLINICAL PICTURE. ITS DISTINCTION FROM OTHER, SIMILAR HEADACHES.
 138. Anders Angelsen: NEUROENDOCRINE CELLS IN HUMAN PROSTATIC CARCINOMAS AND THE PROSTATIC COMPLEX OF RAT, GUINEA PIG, CAT AND DOG.
 139. Fabio Antonaci: CHRONIC PAROXYSMAL HEMICRANIA AND HEMICRANIA CONTINUA: TWO DIFFERENT ENTITIES?
 140. Sven M. Carlsen: ENDOCRINE AND METABOLIC EFFECTS OF METFORMIN WITH SPECIAL EMPHASIS ON CARDIOVASCULAR RISK FACTORES.
- 1999
141. Terje A. Murberg: DEPRESSIVE SYMPTOMS AND COPING AMONG PATIENTS WITH CONGESTIVE HEART FAILURE.

- 142.Harm-Gerd Karl Blaas: THE EMBRYONIC EXAMINATION. Ultrasound studies on the development of the human embryo.
- 143.Noëmi Becser Andersen:THE CEPHALIC SENSORY NERVES IN UNILATERAL HEADACHES. Anatomical background and neurophysiological evaluation.
- 144.Eli-Janne Fiskerstrand: LASER TREATMENT OF PORT WINE STAINS. A study of the efficacy and limitations of the pulsed dye laser. Clinical and morfological analyses aimed at improving the therapeutic outcome.
- 145.Bård Kulseng: A STUDY OF ALGINATE CAPSULE PROPERTIES AND CYTOKINES IN RELATION TO INSULIN DEPENDENT DIABETES MELLITUS.
- 146.Terje Haug: STRUCTURE AND REGULATION OF THE HUMAN UNG GENE ENCODING URACIL-DNA GLYCOSYLASE.
- 147.Heidi Brurok: MANGANESE AND THE HEART. A Magic Metal with Diagnostic and Therapeutic Possibilites.
- 148.Agnes Kathrine Lie: DIAGNOSIS AND PREVALENCE OF HUMAN PAPILLOMAVIRUS INFECTION IN CERVICAL INTRAEPITELIAL NEOPLASIA. Relationship to Cell Cycle Regulatory Proteins and HLA DQBI Genes.
- 149.Ronald Mårvik: PHARMACOLOGICAL, PHYSIOLOGICAL AND PATHOPHYSIOLOGICAL STUDIES ON ISOLATED STOMACS.
- 150.Ketil Jarl Hølen: THE ROLE OF ULTRASONOGRAPHY IN THE DIAGNOSIS AND TREATMENT OF HIP DYSPLASIA IN NEWBORNS.
- 151.Irene Hetlevik: THE ROLE OF CLINICAL GUIDELINES IN CARDIOVASCULAR RISK INTERVENTION IN GENERAL PRACTICE.
- 152.Katarina Tunøn: ULTRASOUND AND PREDICTION OF GESTATIONAL AGE.
- 153.Johannes Soma: INTERACTION BETWEEN THE LEFT VENTRICLE AND THE SYSTEMIC ARTERIES.
- 154.Arild Aamodt: DEVELOPMENT AND PRE-CLINICAL EVALUATION OF A CUSTOM-MADE FEMORAL STEM.
- 155.Agnar Tegnander: DIAGNOSIS AND FOLLOW-UP OF CHILDREN WITH SUSPECTED OR KNOWN HIP DYSPLASIA.
- 156.Bent Indredavik: STROKE UNIT TREATMENT: SHORT AND LONG-TERM EFFECTS
- 157.Jolanta Vanagaite Vingen: PHOTOPHOBIA AND PHONOPHOBIA IN PRIMARY HEADACHES
- 2000
- 158.Ola Dalsegg Sæther: PATHOPHYSIOLOGY DURING PROXIMAL AORTIC CROSS-CLAMPING
- 159.CLINICAL AND EXPERIMENTAL STUDIES
- 160.Christina Vogt Isaksen: PRENATAL ULTRASOUND AND POSTMORTEM FINDINGS – A TEN YEAR CORRELATIVE STUDY OF FETUSES AND INFANTS WITH DEVELOPMENTAL ANOMALIES.
- 161.Holger Seidel: HIGH-DOSE METHOTREXATE THERAPY IN CHILDREN WITH ACUTE LYMPHOCYTIC LEUKEMIA: DOSE, CONCENTRATION, AND EFFECT CONSIDERATIONS.
- 162.Stein Hallan: IMPLEMENTATION OF MODERN MEDICAL DECISION ANALYSIS INTO CLINICAL DIAGNOSIS AND TREATMENT.
- 163.Malcolm Sue-Chu: INVASIVE AND NON-INVASIVE STUDIES IN CROSS-COUNTRY SKIERS WITH ASTHMA-LIKE SYMPTOMS.
- 164.Ole-Lars Brekke: EFFECTS OF ANTIOXIDANTS AND FATTY ACIDS ON TUMOR NECROSIS FACTOR-INDUCED CYTOTOXICITY.
- 165.Jan Lundbom: AORTOCORONARY BYPASS SURGERY: CLINICAL ASPECTS, COST CONSIDERATIONS AND WORKING ABILITY.
- 166.John-Anker Zwart: LUMBAR NERVE ROOT COMPRESSION, BIOCHEMICAL AND NEUROPHYSIOLOGICAL ASPECTS.
- 167.Geir Falck: HYPEROSMOLALITY AND THE HEART.
- 168.Eirik Skogvoll: CARDIAC ARREST Incidence, Intervention and Outcome.
- 169.Dalius Bansevicius: SHOULDER-NECK REGION IN CERTAIN HEADACHES AND CHRONIC PAIN SYNDROMES.
- 170.Bettina Kinge: REFRACTIVE ERRORS AND BIOMETRIC CHANGES AMONG UNIVERSITY STUDENTS IN NORWAY.
- 171.Gunnar Qvigstad: CONSEQUENCES OF HYPERGASTRINEMIA IN MAN
- 172.Hanne Ellekjær: EPIDEMIOLOGICAL STUDIES OF STROKE IN A NORWEGIAN POPULATION. INCIDENCE, RISK FACTORS AND PROGNOSIS
- 173.Hilde Grimstad: VIOLENCE AGAINST WOMEN AND PREGNANCY OUTCOME.

174. Astrid Hjelde: SURFACE TENSION AND COMPLEMENT ACTIVATION: Factors influencing bubble formation and bubble effects after decompression.
175. Kjell A. Kvistad: MR IN BREAST CANCER – A CLINICAL STUDY.
176. Ivar Rossvoll: ELECTIVE ORTHOPAEDIC SURGERY IN A DEFINED POPULATION. Studies on demand, waiting time for treatment and incapacity for work.
177. Carina Seidel: PROGNOSTIC VALUE AND BIOLOGICAL EFFECTS OF HEPATOCYTE GROWTH FACTOR AND SYNDECAN-1 IN MULTIPLE MYELOMA.
- 2001
178. Alexander Wahba: THE INFLUENCE OF CARDIOPULMONARY BYPASS ON PLATELET FUNCTION AND BLOOD COAGULATION – DETERMINANTS AND CLINICAL CONSEQUENCES
179. Marcus Schmitt-Egenolf: THE RELEVANCE OF THE MAJOR HISTOCOMPATIBILITY COMPLEX FOR THE GENETICS OF PSORIASIS
180. Odrun Arna Gederaas: BIOLOGICAL MECHANISMS INVOLVED IN 5-AMINOLEVULINIC ACID BASED PHOTODYNAMIC THERAPY
181. Pål Richard Romundstad: CANCER INCIDENCE AMONG NORWEGIAN ALUMINIUM WORKERS
182. Henrik Hjorth-Hansen: NOVEL CYTOKINES IN GROWTH CONTROL AND BONE DISEASE OF MULTIPLE MYELOMA
183. Gunnar Morken: SEASONAL VARIATION OF HUMAN MOOD AND BEHAVIOUR
184. Bjørn Olav Haugen: MEASUREMENT OF CARDIAC OUTPUT AND STUDIES OF VELOCITY PROFILES IN AORTIC AND MITRAL FLOW USING TWO- AND THREE-DIMENSIONAL COLOUR FLOW IMAGING
185. Geir Bråthen: THE CLASSIFICATION AND CLINICAL DIAGNOSIS OF ALCOHOL-RELATED SEIZURES
186. Knut Ivar Aasarød: RENAL INVOLVEMENT IN INFLAMMATORY RHEUMATIC DISEASE. A Study of Renal Disease in Wegener's Granulomatosis and in Primary Sjögren's Syndrome
187. Trude Helen Flo: RECEPTORS INVOLVED IN CELL ACTIVATION BY DEFINED URONIC ACID POLYMERS AND BACTERIAL COMPONENTS
188. Bodil Kavli: HUMAN URACIL-DNA GLYCOSYLASES FROM THE UNG GENE: STRUCTURAL BASIS FOR SUBSTRATE SPECIFICITY AND REPAIR
189. Liv Thommesen: MOLECULAR MECHANISMS INVOLVED IN TNF- AND GASTRIN-MEDIATED GENE REGULATION
190. Turid Lingaas Holmen: SMOKING AND HEALTH IN ADOLESCENCE; THE NORD-TRØNDELAG HEALTH STUDY, 1995-97
191. Øyvind Hjertner: MULTIPLE MYELOMA: INTERACTIONS BETWEEN MALIGNANT PLASMA CELLS AND THE BONE MICROENVIRONMENT
192. Asbjørn Støylen: STRAIN RATE IMAGING OF THE LEFT VENTRICLE BY ULTRASOUND. FEASIBILITY, CLINICAL VALIDATION AND PHYSIOLOGICAL ASPECTS
193. Kristian Midthjell: DIABETES IN ADULTS IN NORD-TRØNDELAG. PUBLIC HEALTH ASPECTS OF DIABETES MELLITUS IN A LARGE, NON-SELECTED NORWEGIAN POPULATION.
194. Guanglin Cui: FUNCTIONAL ASPECTS OF THE ECL CELL IN RODENTS
195. Ulrik Wisløff: CARDIAC EFFECTS OF AEROBIC ENDURANCE TRAINING; HYPERTROPHY, CONTRACTILITY AND CALCIUM HANDLING IN NORMAL AND FAILING HEART
196. Øyvind Halaas: MECHANISMS OF IMMUNOMODULATION AND CELL-MEDIATED CYTOTOXICITY INDUCED BY BACTERIAL PRODUCTS
197. Tore Amundsen: PERFUSION MR IMAGING IN THE DIAGNOSIS OF PULMONARY EMBOLISM
198. Nanna Kurtze: THE SIGNIFICANCE OF ANXIETY AND DEPRESSION IN FATIGUE AND PATTERNS OF PAIN AMONG INDIVIDUALS DIAGNOSED WITH FIBROMYALGIA: RELATIONS WITH QUALITY OF LIFE, FUNCTIONAL DISABILITY, LIFESTYLE, EMPLOYMENT STATUS, CO-MORBIDITY AND GENDER
199. Tom Ivar Lund Nilsen: PROSPECTIVE STUDIES OF CANCER RISK IN NORD-TRØNDELAG: THE HUNT STUDY. Associations with anthropometric, socioeconomic, and lifestyle risk factors
200. Asta Kristine Häberg: A NEW APPROACH TO THE STUDY OF MIDDLE CEREBRAL ARTERY OCCLUSION IN THE RAT USING MAGNETIC RESONANCE TECHNIQUES
- 2002
201. Knut Jørgen Arntzen: PREGNANCY AND CYTOKINES
202. Henrik Døllner: INFLAMMATORY MEDIATORS IN PERINATAL INFECTIONS

203. Asta Bye: LOW FAT, LOW LACTOSE DIET USED AS PROPHYLACTIC TREATMENT OF ACUTE INTESTINAL REACTIONS DURING PELVIC RADIOTHERAPY. A PROSPECTIVE RANDOMISED STUDY.
204. Sylvester Moyo: STUDIES ON STREPTOCOCCUS AGALACTIAE (GROUP B STREPTOCOCCUS) SURFACE-ANCHORED MARKERS WITH EMPHASIS ON STRAINS AND HUMAN SERA FROM ZIMBABWE.
205. Knut Hagen: HEAD-HUNT: THE EPIDEMIOLOGY OF HEADACHE IN NORD-TRØNDELAG
206. Li Lixin: ON THE REGULATION AND ROLE OF UNCOUPLING PROTEIN-2 IN INSULIN PRODUCING β -CELLS
207. Anne Hildur Henriksen: SYMPTOMS OF ALLERGY AND ASTHMA VERSUS MARKERS OF LOWER AIRWAY INFLAMMATION AMONG ADOLESCENTS
208. Egil Andreas Fors: NON-MALIGNANT PAIN IN RELATION TO PSYCHOLOGICAL AND ENVIRONMENTAL FACTORS. EXPERIMENTAL AND CLINICAL STUDIES OF PAIN WITH FOCUS ON FIBROMYALGIA
209. Pål Klepstad: MORPHINE FOR CANCER PAIN
210. Ingunn Bakke: MECHANISMS AND CONSEQUENCES OF PEROXISOME PROLIFERATOR-INDUCED HYPERFUNCTION OF THE RAT GASTRIN PRODUCING CELL
211. Ingrid Susann Gribbestad: MAGNETIC RESONANCE IMAGING AND SPECTROSCOPY OF BREAST CANCER
212. Rønnaug Astri Ødegård: PREECLAMPSIA – MATERNAL RISK FACTORS AND FETAL GROWTH
213. Johan Haux: STUDIES ON CYTOTOXICITY INDUCED BY HUMAN NATURAL KILLER CELLS AND DIGITOXIN
214. Turid Suzanne Berg-Nielsen: PARENTING PRACTICES AND MENTALLY DISORDERED ADOLESCENTS
215. Astrid Rydning: BLOOD FLOW AS A PROTECTIVE FACTOR FOR THE STOMACH MUCOSA. AN EXPERIMENTAL STUDY ON THE ROLE OF MAST CELLS AND SENSORY AFFERENT NEURONS
- 2003
216. Jan Pål Loennechen: HEART FAILURE AFTER MYOCARDIAL INFARCTION. Regional Differences, Myocyte Function, Gene Expression, and Response to Cariporide, Losartan, and Exercise Training.
217. Elisabeth Qvigstad: EFFECTS OF FATTY ACIDS AND OVER-STIMULATION ON INSULIN SECRETION IN MAN
218. Arne Åsberg: EPIDEMIOLOGICAL STUDIES IN HEREDITARY HEMOCHROMATOSIS: PREVALENCE, MORBIDITY AND BENEFIT OF SCREENING.
219. Johan Fredrik Skomsvoll: REPRODUCTIVE OUTCOME IN WOMEN WITH RHEUMATIC DISEASE. A population registry based study of the effects of inflammatory rheumatic disease and connective tissue disease on reproductive outcome in Norwegian women in 1967-1995.
220. Siv Mørkved: URINARY INCONTINENCE DURING PREGNANCY AND AFTER DELIVERY: EFFECT OF PELVIC FLOOR MUSCLE TRAINING IN PREVENTION AND TREATMENT
221. Marit S. Jordhøy: THE IMPACT OF COMPREHENSIVE PALLIATIVE CARE
222. Tom Christian Martinsen: HYPERGASTRINEMIA AND HYPOACIDITY IN RODENTS – CAUSES AND CONSEQUENCES
223. Solveig Tingulstad: CENTRALIZATION OF PRIMARY SURGERY FOR OVARIAN CANCER. FEASIBILITY AND IMPACT ON SURVIVAL
224. Haytham Eloqayli: METABOLIC CHANGES IN THE BRAIN CAUSED BY EPILEPTIC SEIZURES
225. Torunn Bruland: STUDIES OF EARLY RETROVIRUS-HOST INTERACTIONS – VIRAL DETERMINANTS FOR PATHOGENESIS AND THE INFLUENCE OF SEX ON THE SUSCEPTIBILITY TO FRIEND MURINE LEUKAEMIA VIRUS INFECTION
226. Torstein Hole: DOPPLER ECHOCARDIOGRAPHIC EVALUATION OF LEFT VENTRICULAR FUNCTION IN PATIENTS WITH ACUTE MYOCARDIAL INFARCTION
227. Vibeke Nossun: THE EFFECT OF VASCULAR BUBBLES ON ENDOTHELIAL FUNCTION
228. Sigurd Fasting: ROUTINE BASED RECORDING OF ADVERSE EVENTS DURING ANAESTHESIA – APPLICATION IN QUALITY IMPROVEMENT AND SAFETY
229. Solfrid Romundstad: EPIDEMIOLOGICAL STUDIES OF MICROALBUMINURIA. THE NORD-TRØNDELAG HEALTH STUDY 1995-97 (HUNT 2)
230. Geir Torheim: PROCESSING OF DYNAMIC DATA SETS IN MAGNETIC RESONANCE IMAGING

231. Catrine Ahlén: SKIN INFECTIONS IN OCCUPATIONAL SATURATION DIVERS IN THE NORTH SEA AND THE IMPACT OF THE ENVIRONMENT
232. Arnulf Langhammer: RESPIRATORY SYMPTOMS, LUNG FUNCTION AND BONE MINERAL DENSITY IN A COMPREHENSIVE POPULATION SURVEY. THE NORD-TRØNDELAG HEALTH STUDY 1995-97. THE BRONCHIAL OBSTRUCTION IN NORD-TRØNDELAG STUDY
233. Einar Kjelsås: EATING DISORDERS AND PHYSICAL ACTIVITY IN NON-CLINICAL SAMPLES
234. Arne Wibe: RECTAL CANCER TREATMENT IN NORWAY – STANDARDISATION OF SURGERY AND QUALITY ASSURANCE
- 2004
235. Eivind Witsø: BONE GRAFT AS AN ANTIBIOTIC CARRIER
236. Anne Mari Sund: DEVELOPMENT OF DEPRESSIVE SYMPTOMS IN EARLY ADOLESCENCE
237. Hallvard Lærum: EVALUATION OF ELECTRONIC MEDICAL RECORDS – A CLINICAL TASK PERSPECTIVE
238. Gustav Mikkelsen: ACCESSIBILITY OF INFORMATION IN ELECTRONIC PATIENT RECORDS; AN EVALUATION OF THE ROLE OF DATA QUALITY
239. Steinar Krokstad: SOCIOECONOMIC INEQUALITIES IN HEALTH AND DISABILITY. SOCIAL EPIDEMIOLOGY IN THE NORD-TRØNDELAG HEALTH STUDY (HUNT), NORWAY
240. Arne Kristian Myhre: NORMAL VARIATION IN ANOGENITAL ANATOMY AND MICROBIOLOGY IN NON-ABUSED PRESCHOOL CHILDREN
241. Ingunn Dybedal: NEGATIVE REGULATORS OF HEMATOPOIETEC STEM AND PROGENITOR CELLS
242. Beate Sitter: TISSUE CHARACTERIZATION BY HIGH RESOLUTION MAGIC ANGLE SPINNING MR SPECTROSCOPY
243. Per Arne Aas: MACROMOLECULAR MAINTENANCE IN HUMAN CELLS – REPAIR OF URACIL IN DNA AND METHYLATIONS IN DNA AND RNA
244. Anna Bofin: FINE NEEDLE ASPIRATION CYTOLOGY IN THE PRIMARY INVESTIGATION OF BREAST TUMOURS AND IN THE DETERMINATION OF TREATMENT STRATEGIES
245. Jim Aage Nøttestad: DEINSTITUTIONALIZATION AND MENTAL HEALTH CHANGES AMONG PEOPLE WITH MENTAL RETARDATION
246. Reidar Fossmark: GASTRIC CANCER IN JAPANESE COTTON RATS
247. Wibeke Nordhøy: MANGANESE AND THE HEART, INTRACELLULAR MR RELAXATION AND WATER EXCHANGE ACROSS THE CARDIAC CELL MEMBRANE
- 2005
248. Sturla Molden: QUANTITATIVE ANALYSES OF SINGLE UNITS RECORDED FROM THE HIPPOCAMPUS AND ENTORHINAL CORTEX OF BEHAVING RATS
249. Wenche Brenne Drøyvold: EPIDEMIOLOGICAL STUDIES ON WEIGHT CHANGE AND HEALTH IN A LARGE POPULATION. THE NORD-TRØNDELAG HEALTH STUDY (HUNT)
250. Ragnhild Støen: ENDOTHELIUM-DEPENDENT VASODILATION IN THE FEMORAL ARTERY OF DEVELOPING PIGLETS
251. Aslak Steinsbekk: HOMEOPATHY IN THE PREVENTION OF UPPER RESPIRATORY TRACT INFECTIONS IN CHILDREN
252. Hill-Aina Steffenach: MEMORY IN HIPPOCAMPAL AND CORTICO-HIPPOCAMPAL CIRCUITS
253. Eystein Stordal: ASPECTS OF THE EPIDEMIOLOGY OF DEPRESSIONS BASED ON SELF-RATING IN A LARGE GENERAL HEALTH STUDY (THE HUNT-2 STUDY)
254. Viggo Pettersen: FROM MUSCLES TO SINGING: THE ACTIVITY OF ACCESSORY BREATHING MUSCLES AND THORAX MOVEMENT IN CLASSICAL SINGING
255. Marianne Fyhn: SPATIAL MAPS IN THE HIPPOCAMPUS AND ENTORHINAL CORTEX
256. Robert Valderhaug: OBSESSIVE-COMPULSIVE DISORDER AMONG CHILDREN AND ADOLESCENTS: CHARACTERISTICS AND PSYCHOLOGICAL MANAGEMENT OF PATIENTS IN OUTPATIENT PSYCHIATRIC CLINICS
257. Erik Skaasheim Haug: INFRARENAL ABDOMINAL AORTIC ANEURYSMS – COMORBIDITY AND RESULTS FOLLOWING OPEN SURGERY
258. Daniel Kondziella: GLIAL-NEURONAL INTERACTIONS IN EXPERIMENTAL BRAIN DISORDERS
259. Vegard Heimly Brun: ROUTES TO SPATIAL MEMORY IN HIPPOCAMPAL PLACE CELLS

260. Kenneth McMillan: PHYSIOLOGICAL ASSESSMENT AND TRAINING OF ENDURANCE AND STRENGTH IN PROFESSIONAL YOUTH SOCCER PLAYERS
261. Marit Sæbø Indredavik: MENTAL HEALTH AND CEREBRAL MAGNETIC RESONANCE IMAGING IN ADOLESCENTS WITH LOW BIRTH WEIGHT
262. Ole Johan Kemi: ON THE CELLULAR BASIS OF AEROBIC FITNESS, INTENSITY-DEPENDENCE AND TIME-COURSE OF CARDIOMYOCYTE AND ENDOTHELIAL ADAPTATIONS TO EXERCISE TRAINING