

ORIGINAL WORK



Extended Coagulation Profiling in Isolated Traumatic Brain Injury: A CENTER-TBI Analysis

Julia K. Böhm¹, Victoria Schaeben¹, Nadine Schäfer¹, Helge Güting¹, Rolf Lefering¹, Sophie Thorn², Herbert Schöchl^{3,4}, Johannes Zipperle⁴, Oliver Grottke⁵, Rolf Rossaint⁵, Simon Stanworth⁶, Nicola Curry⁷ and Marc Maegele^{1,8*} CENTER-TBI Participants and Investigators

© 2021 The Author(s)

Abstract

Background: Trauma-induced coagulopathy in traumatic brain injury (TBI) remains associated with high rates of complications, unfavorable outcomes, and mortality. The underlying mechanisms are largely unknown. Embedded in the prospective multinational Collaborative European Neurotrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI) study, coagulation profiles beyond standard conventional coagulation assays were assessed in patients with isolated TBI within the very early hours of injury.

Methods: Results from blood samples (citrate/EDTA) obtained on hospital admission were matched with clinical and routine laboratory data of patients with TBI captured in the CENTER-TBI central database. To minimize confounding factors, patients with strictly isolated TBI (iTBI) ($n = 88$) were selected and stratified for coagulopathy by routine international normalized ratio (INR): (1) $INR < 1.2$ and (2) $INR \geq 1.2$. An $INR > 1.2$ has been well adopted over time as a threshold to define trauma-related coagulopathy in general trauma populations. The following parameters were evaluated: quick's value, activated partial thromboplastin time, fibrinogen, thrombin time, antithrombin, coagulation factor activity of factors V, VIII, IX, and XIII, protein C and S, plasminogen, D-dimer, fibrinolysis-regulating parameters (thrombin activatable fibrinolysis inhibitor, plasminogen activator inhibitor 1, antiplasmin), thrombin generation, and fibrin monomers.

Results: Patients with iTBI with $INR \geq 1.2$ ($n = 16$) had a high incidence of progressive intracranial hemorrhage associated with increased mortality and unfavorable outcome compared with patients with $INR < 1.2$ ($n = 72$). Activity of coagulation factors V, VIII, IX, and XIII dropped on average by 15–20% between the groups whereas protein C and S levels dropped by 20%. With an elevated INR, thrombin generation decreased, as reflected by lower peak height and endogenous thrombin potential (ETP), whereas the amount of fibrin monomers increased. Plasminogen activity significantly decreased from 89% in patients with $INR < 1.2$ to 76% in patients with $INR \geq 1.2$. Moreover, D-dimer levels significantly increased from a mean of 943 mg/L in patients with $INR < 1.2$ to 1,301 mg/L in patients with $INR \geq 1.2$.

*Correspondence: Marc.Maegele@t-online.de

[†]Julia K. Böhm and Victoria Schaeben have contributed equally to this work.

¹Institute for Research in Operative Medicine, Faculty of Health, Department of Medicine, Witten/Herdecke University, Ostmerheimer Str. 200, Building 38, 51109 Cologne, Germany

Full list of author information is available at the end of the article

All CENTER-TBI participants and investigators are listed in the Appendix.

Conclusions: This more in-depth analysis beyond routine conventional coagulation assays suggests a counterbalanced regulation of coagulation and fibrinolysis in patients with iTBI with hemostatic abnormalities. We observed distinct patterns involving key pathways of the highly complex and dynamic coagulation system that offer windows of opportunity for further research. Whether the changes observed on factor levels may be relevant and explain the worse outcome or the more severe brain injuries by themselves remains speculative.

Keywords: CENTER-TBI, Coagulopathy, Fibrin monomers, Progressive intracranial hemorrhage, Thrombin generation, Traumatic brain injury

Introduction

Traumatic brain injury (TBI) remains a significant medical and socioeconomic burden for patients, relatives, and health care systems [1]. Current calculations estimate approximately 69 million cases of TBI to occur each year around the globe [2]. Apart from the initial injury, posttraumatic courses, including outcomes, may be complicated by preexisting hemostatic derangements or derangements that develop with TBI [3]. Recent observational data from the prospective multicentered Collaborative European Neurotrauma Effectiveness Research in Traumatic Brain Injury (CENTER-TBI) study have confirmed the presence of laboratory coagulopathy based on conventional coagulation assays (CCAs) in approximately 20% of all patients with isolated TBI (iTBI) on hospital admission [4]. Independent risk factors may include not only the degree of impact, shock, hypothermia, and hypotension but also preinjury use of anticoagulant and/or antiplatelet therapies [4]. Patients with preinjury anticoagulant and/or antiplatelet therapy were twice as likely to have an abnormal coagulation profile as those without premedication, and this was associated with greater expansion of hemorrhagic lesions and a higher risk of delayed traumatic intracranial hemorrhage [4, 5].

Apart from the physical impact, which typically leads to disruptions of the cerebral vasculature and the blood–brain barrier, along with hemorrhage, TBI-associated factors may further alter the overall hemostatic state, leading to disruptions in the balance between hypocoagulability and hypercoagulability, thereby exacerbating the initial injury [6]. The current definitions of post-TBI coagulopathy remain heterogeneous and are often based on CCAs, such as prothrombin time with its surrogates, prothrombin ratio and international normalized ratio (INR), and activated partial thromboplastin time (aPTT). In addition, the precise mechanisms that relate to the different phenotypes of coagulopathy seen with TBI are still incompletely understood. The aim of the present study was to further characterize alterations within the coagulation system of patients with iTBI by assessing concentration changes of selected proteins indicative for hypercoagulation and

hypocoagulation beyond CCAs within the early hours of injury. In this context, further interest was given to the assessment of hemorrhagic injuries, expansion, and outcome in the presence of hemostatic abnormalities after iTBI.

Methods

Patients and Blood Sampling

The present study was embedded into the longitudinal observational CENTER-TBI study, which had recruited patients from 60 selected centers across Europe and Israel between December 2014 and December 2017 [7]. The study was conducted in accordance with local ethical regulations and with all relevant laws of the European Union and the country of the recruiting site, including relevant laws and regulations on the use of human materials. Informed consent by the patients and/or the legal representative/next of kin was obtained according to the local legislations.

Patient inclusion criteria were clinical diagnosis of TBI, indication for computed tomography (CT) scanning, presentation to a study center within 24 h of injury, and informed consent according to local and national requirements [7]. Patients were excluded if they had any severe preexisting neurological disorder that could have confounded the outcome assessments.

Of the 60 recruiting CENTER-TBI study sites across Europe, the following centers had collected and submitted blood samples for the present analysis: (1) Antwerp University Hospital (Belgium), (2) University of Cambridge (UK), (3) Helsingin Yliopisto Finnish Institute for Molecular Medicine (Finland), (4) University Hospitals Leuven (Belgium), (5) Karolinska University Hospital Stockholm (Sweden), (6) University Hospital of Aachen (Germany), (7) Kaunas University of Technology (Lithuania), (8) Leids Universitair Medisch Centrum (Netherlands), and (9) Turku University Central Hospital (Finland). On site, EDTA and citrated blood tubes were centrifuged at $1.500 \times g$ for 10 min. Platelet poor plasma samples were aliquoted and frozen at -80°C directly or at -20°C for a maximum of 48 h before being transferred to the storage temperature of -80°C . Samples were sent

from the local recruiting sites to the central CENTER-TBI laboratory at the University of Pécs (Hungary) first and then sent on dry ice to the receiving Institute for Research in Operative Medicine (IFOM) in Cologne, Germany, for further analysis.

Sample Selection

On receipt, the samples were linked to the clinical and laboratory data comprehensively collected into the CENTER-TBI central database (INCF Neurobot 2.0; INCF, Stockholm, Sweden). To exclude any confounding effects by external factors (e.g., extracranial injuries and/or relevant interim anticoagulant or procoagulant therapies), further sample analysis was restricted to samples received from patients with iTBI ($AIS_{\text{Brain}} 2-5$) only and to those with available data on coagulation captured in the CENTER-TBI central database for the first 4 h after injury. Samples received from patients with documented preexisting neurological disorders, preinjury anticoagulant and/or antiplatelet therapy, presence of extracranial injuries ($AIS_{\text{Extracranial}} > 0$), and missing critical data points were excluded a priori. For further analysis, the first documented blood sample collected after local hospital admission was considered together with the clinical and laboratory data captured on the same timeline. The INR on admission captured in the CENTER-TBI central database served to distinguish two groups of samples and patients: those with (1) $INR < 1.2$ and those with (2) $INR \geq 1.2$. An $INR > 1.2$ has been well adopted over time as a threshold to define trauma-related coagulopathy in general trauma populations [8].

Clinical Data

The prospectively recorded clinical and laboratory data points in the scope of the CENTER-TBI study for patient characterization and captured in the CENTER-TBI central database included demographics, injury characteristics, medical presentation on admission (Glasgow Coma Scale score, systolic blood pressure, heart rate, temperature), scheduled emergency surgical intervention, neuroworsening, parameters for tissue injury and hypoperfusion (base excess and shock index), hemodilution (platelets, hemoglobin, and hematocrit), and outcome (death and Glasgow Outcome Score Extended [GOS-E]). Presence of progressive intracranial hemorrhage (PIH) was defined by the incidence of extradural or subdural hematoma or subarachnoid hemorrhage on the initial CT scan and increase in hemorrhage size/volume on the follow-up CT scan. Follow-up data on functional outcome, including mortality and GOS-E, were obtained 6 months post injury. A GOS-E between 1 and 4 (dead, vegetative state, low-severity disability, and high-severity disability) was considered unfavorable.

Extended Coagulation Testing

CCAs to evaluate the coagulation status at the time point of blood sampling were performed by the Institute of Transfusion Medicine (ITM) at the Cologne-Merheim Medical Center, Cologne (Germany), and included the following: Quick's value, INR, aPTT, thrombin time, fibrinogen, and antithrombin. The assessments beyond CCAs included coagulation factor V, VIII, IX, and XIII antigens; protein C; protein S; D-dimer; and plasminogen. Citrated plasma samples were analyzed by using HemosIL assays (Werfen, Bedford, MA) on the ACL TOP CTS 700 device (Werfen).

Enzyme-Linked Immunosorbent Assay

Soluble plasminogen activator inhibitor 1 (PAI-1) as a marker for fibrinolysis regulation was quantified via the commercial Human Serpin E1/PAI-1 DuoSet ELISA (R&D Systems, Minneapolis, MN) according to the manufacturer's recommendations.

Thrombin Generation and Fibrinolysis-Regulating Markers

Thrombin generation, thrombin activatable fibrinolysis inhibitor (TAFI), antiplasmin, and fibrin monomers were assessed at the Ludwig Boltzmann Institute (Vienna, Austria). By using a quantitative fluorogenic assay (STG-BleedScreen; Stago, Asnières-sur-Seine, France) on the thrombin generation analyzer (ST Genesia; Stago), the thrombin generation parameters (1) peak height and (2) endogenous thrombin potential (ETP) were determined. TAFI and the fibrinolysis regulator antiplasmin were quantified by a colorimetric assay using the STA-Stachrom TAFI and STA-Stachrom Antiplasmin kits (Stago). The level of fibrin monomers was measured with an immunoturbidimetric technology by using the STA-Liatest FM kit (Stago). For these assays, the frozen citrated plasma samples were thawed at 37 °C for 10 min in a water bath and stored at room temperature for up to 1 h.

Statistical Analysis

For the descriptive data analysis, metric data are presented as medians with interquartile ranges and categorical data are presented as percentages. Statistical differences in patient characteristics and coagulation factors/parameters were tested by using the Mann-Whitney *U*-test and χ^2 test, respectively. A *p* value < 0.05 was considered statistically significant. Statistical analyses were performed by using SPSS statistics version 25 for Windows (IBM Corp., Armonk, NY) and GraphPad Prism version 7.00 (GraphPad Software, La Jolla, CA).




Fig. 1 Overview of CENTER-TBI patients for selection into the present analysis to assess hemostatic abnormalities after iTBI. AIS Abbreviated Injury Scale, CENTER-TBI Collaborative European Neurotrauma Effectiveness Research in Traumatic Brain Injury, INR international normalized ratio, iTBI isolated traumatic brain injury, TBI traumatic brain injury

Results

Study Cohort Characteristics

Overall, 4,509 patients were included in the CENTER-TBI core study database; of these, 3,287 had to be excluded for coexisting extracranial injuries and 624 for missing data (Fig. 1). Of the 598 patients with iTBI who had available coagulation data within the acute

phase after TBI (< 4 h after injury), blood samples of 485 patients were missing, and to minimize interference and interfering factors, an additional 19 patients were excluded from the analysis (Fig. 1). Due to quality loss of samples during shipment and experimental settings, blood samples of another six patients had to be excluded.

Table 1 Characteristics of patients with iTBI stratified by INR (n = 88)

	INR <1.2 (n = 72)	INR ≥1.2 (n = 16)
Demographics		
Age, median (IQR) (years)	41.0 (30 to 57)	27.0 (22 to 54)
Age ≥ 75, n (%)	5 (6.9)	NA
Male sex, n (%)	48 (66.6)	12 (75)
Injury characteristics, n (%)		
Blunt TBI	65 (90.3)	15 (93.8)
AIS 2	6 (8.3)	1 (6.3)
AIS 3	27 (37.5)	1 (6.3)
AIS 4	21 (29.2)	6 (37.5)
AIS 5	18 (25)	8 (50)
Injuries identified on initial CT scan, n (%)		
Diffuse axonal injury	9 (12.5)	2 (12.5)
Extradural hematoma	15 (20.8)	4 (25)
Subdural hematoma	28 (38.8)	7 (43.7)
Subarachnoid hemorrhage	34 (47.2)	8 (50)
Midline shift	12 (16.7)	6 (37.5)*
Basal cistern compression	6 (8.3)	2 (12.5)
Depressed skull fracture	7 (9.7)	3 (18.8)
Severe contusion	2 (2.7)	2 (12.5)
Medical presentation at admission (ED)		
GCS, median (IQR)	14.0 (10.0 to 15.0)	12.0 (7.5 to 15.0)
SBP, median (IQR) (mm Hg)	134.0 (123 to 150)	135.5 (113 to 163)
Heart rate, median (IQR) (bpm)	84.0 (75 to 97)	82.0 (68 to 90)
Temperature, median (IQR) (°C)	36.4 (35.8 to 36.9)	36.0 (35.6 to 36.4)
Scheduled for emergency surgical intervention, n (%)	11 (15.3)	2 (12.5)
Neuroworsening, n (%)	8 (11.1)	3 (18.8)
Parameter for tissue injury and hypoperfusion, median (IQR)		
Base excess (mmol/L)	-1.1 (-2.9 to 0.55)	-2.7 (-3.7 to -0.9)
Shock index	0.61 (0.52 to 0.71)	0.64 (0.48 to 0.73)
Parameter for hemodilution, median (IQR)		
Platelet count (per nL)	220.5 (181.5 to 273.5)	211.5 (177.8 to 235.3)
Hemoglobin (g/dL)	13.9 (12.8 to 14.5)	13.4 (12.1 to 14.4)
Hematocrit	41.0 (38.0 to 44.0)	44.0 (37.7 to 45.7)*
ED therapy, n (%)		
Blood products (e.g., RBC, FFP, PC, WB)	NA	NA
Hemostatic agents		
TXA	2 (2.7)	1 (6.3)
Outcomes		
Death (overall), n (%)	7 (9.7)	3 (18.8)
GOS-E (6 months), median (IQR)	7 (6 to 8)	5 (2 to 7.5)

Neuroworsening was defined as follows: (1) a decrease in the GCS motor score of 2 or more points, (2) a new loss of pupillary reactivity or development of pupillary asymmetry ≥2 mm, or (3) deterioration in neurological or CT status sufficient to warrant immediate medical or surgical intervention. Shock index was calculated as follows: heart rate (bpm) divided by SBP (mm Hg). Patients with iTBI with AIS_{Brain 6} were not included.

AIS Abbreviated Injury Scale, aPTT activated partial thromboplastin time, CT computed tomography, ED emergency department, FFP fresh frozen plasma, GCS Glasgow Coma Scale, GOS-E Glasgow Outcome Score Extended, INR international normalized ratio, iTBI isolated traumatic brain injury, NA not available, PC platelet concentrate, PCC prothrombin complex concentrate, RBC red blood cells, SBP systolic blood pressure, TXA tranexamic acid, WB whole blood

*P < 0.05, **P < 0.001, ***P < 0.0001: statistically significant differences between patients with iTBI with INR <1.2 and patients with iTBI with INR ≥1.2.

The results from the blood sample analyses were matched with the clinical and routine laboratory data of corresponding patients with iTBI for a study cohort of 88 patients who were then stratified according to admission INR (Fig. 1). The basic demographics and injury characteristics of the patients included in the analysis are shown in Table 1. Almost all patients had sustained a blunt trauma mechanism, with high velocity trauma as the most common cause of injury (44%) in patients with $\text{INR} \geq 1.2$, followed by ground-level falls (34%) in patients with $\text{INR} < 1.2$ (Table 2). In more than 90% of cases, the iTBI was moderate to severe ($\text{AIS}_{\text{Brain}} \geq 3$) and closed (Table 1). Patients with $\text{INR} \geq 1.2$, in general, were younger, were more shocked, and had presented with a lower Glasgow Coma Scale score. PIH as detected by CT was observed more frequent with abnormal INR. Midline shifts on CT were seen twice as frequently in patients with $\text{INR} \geq 1.2$ (Table 1). Overall death increased with increasing INR as the median GOS-E at 6 months decreased (Table 1). In addition, 3.4% of the entire cohort of patients with iTBI had received hemostatic agents in the form of tranexamic acid (TXA), whereas just one patient with an $\text{INR} \geq 1.2$ had been treated with TXA upon emergency department admission (Table 1).

Hemostatic Changes with Increasing INR

The majority of the results from CCAs and coagulation factor concentrations, except for aPTT, thrombin time, and fibrinogen, decreased with increasing INR (Table 3). Whereas thrombin time and fibrinogen levels did not change with increasing INR, the aPTT was significantly prolonged in those patients (Table 3). The activity of coagulation factors V, VIII, IX, and XIII dropped between 10 and 25% with $\text{INR} \geq 1.2$, whereas protein C and S levels fell by approximately 20%. Coagulation factor V displayed the greatest decline with 25% with $\text{INR} \geq 1.2$. Also with increasing INR, thrombin generation potential became impaired, as reflected by lower peak height as well as significantly lower ETP (Fig. 2A, B). Conversely, the amount of fibrin monomers increased with increasing

INR, whereas D-dimer levels rose significantly from a mean of 943 $\mu\text{g/L}$ in patients with $\text{INR} < 1.2$ to 1,301 $\mu\text{g/L}$ in patients with $\text{INR} \geq 1.2$ (Fig. 2C, Table 3).

Fibrinolytic Markers with Increasing INR

Fibrinolysis inhibitor PAI-1 remained largely unaltered across the two groups (Fig. 3C), whereas TAFI tended to be lower with $\text{INR} \geq 1.2$ (Fig. 3A). Similarly, plasmin inhibitor antiplasmin decreased with increasing INR (Fig. 3B). In correspondence, plasminogen activity significantly decreased by 15% in patients with $\text{INR} \geq 1.2$ (Table 3).

Discussion

The results from the present study provide more detailed insights into the complex mechanisms behind the various hemostatic abnormalities occurring with and after TBI. Blood samples collected from patients with iTBI after admission to 9 of 60 recruiting CENTER-TBI study sites across Europe were submitted and used for more in-depth analyses of the coagulation system beyond those routinely used to characterize hemostatic abnormalities. The INR was used as a crude a priori way to stratify for the presence or absence of a coagulation deficit on emergency department admission [8].

Increasing INR was associated with a 1.2-fold decrease in thrombin formation peak height and ETP. This is most likely explained by the reduction of coagulation factors V and VIII in these patients because ETP is strongly dependent on these factors. The activity of coagulation factors V, VIII, and IX dropped by 10–25% of their activity with increasing INR. Coagulation factor V, in its active form, is part of the prothrombinase complex, which catalyzes the conversion of prothrombin (factor II) into thrombin (factor IIa), whereas coagulation factor IX is part of the intrinsic tenase complex, which activates prothrombinase. Also observed in the present study was a decrease in activity of coagulation factor XIII by 15% with increasing INR, which can most likely be linked to an increased activation and consumption of coagulation

Table 2 Mechanism of injury in patients with iTBI stratified by INR (n = 44)

Injury Mechanisms	INR < 1.2 (n = 35)	INR \geq 1.2 (n = 9)
High-velocity trauma, n (%)	6 (17)	4 (44)
Blow to head (direct impact), n (%)	5 (14)	NA
Head against object (direct impact), n (%)	2 (5.7)	1 (11)
Ground-level fall, n (%)	12 (34)	2 (22)
Fall from height (>1 m), n (%)	8 (23)	1 (11)
Other closed head injury, n (%)	2 (5.7)	1 (11)

Data on mechanism of injury were missing for $n = 44$ patients with iTBI.

INR international normalized ratio, iTBI isolated traumatic brain injury, NA not available.




Fig. 2 Analysis of thrombin activity and generation in plasma of patients with iTBI stratified by INR. Peak height (**A**), endogenous thrombin potential (**B**), and fibrin monomers (**C**) were measured after iTBI by using calibrated automated thrombography. Statistically significant differences between patients with iTBI with INR < 1.2 and patients with iTBI with INR ≥ 1.2 are marked with asterisks (* $P < 0.05$, ** $P < 0.001$, *** $P < 0.0001$). INR international normalized ratio, iTBI isolated traumatic brain injury


Table 3 Conventional coagulation parameters, coagulation factor activity and anticoagulant proteins of patients with iTBI stratified by INR (n = 88)

	INR <1.2 (n = 72)	INR ≥1.2 (n = 16)
Conventional coagulation parameters, median (IQR)		
Quick (%) (reference range 70–130)	101.0 (90.0–109.0)	74.0 (69.3–77.8)***
aPTT (seconds) (reference range 23–36)	28.1 (25.1–30.5)	32.0 (30.0–35.2)***
Thrombin time (seconds) (reference range 10–17)	14.0 (13.0–15.0)	14.0 (13.3–15.0)
Fibrinogen (mg/dL) (reference range 276–471)	371.0 (315.0–462.0)	389.0 (299.3–504.5)
Antithrombin (%) (reference range 83–128)	99.5 (89.3–113.0)	94.0 (84.8–103.3)*
Coagulation factors and anticoagulant proteins, median (IQR)		
Factor V (%) (reference range 62–139)	94.5 (79.3–110.8)	70.5 (51.3–78.8)***
Factor VIII (%) (reference range 50–150)	136.5 (111.8–175.5)	119.0 (100.3–169.8)
Factor IX (%) (reference range 65–150)	113.0 (99.5–128.8)	100.5 (85.3–122.3)
Factor XIII antigen (%) (reference range 75.2–154.8)	88.0 (75.0–104.0)	68.0 (61.5–83.0)**
Protein C (%) (reference range 70.0–140.0)	98.0 (86.0–112.0)	82.5 (69.3–92.0)**
Protein S (%) (reference range 63.5–149)	98.5 (81.8–118.3)	77.5 (67.2–96.3)**
D-dimer (µg/L) (reference range 0–232)	943.0 (418.0–1,300.5)	1,301.5 (966.0–6,260.5)*
Plasminogen (%) (reference range 80–133)	89.0 (79.0–97.3)	76.5 (60.0–83.8)**

Conventional coagulation parameters, coagulation factors, and anticoagulant proteins of patients with iTBI are represented as median and IQR

aPTT activated partial thromboplastin time, INR international normalized ratio, IQR interquartile range, iTBI isolated traumatic brain injury

* $P < 0.05$, ** $P < 0.001$, *** $P < 0.0001$: statistically significant differences between patients with iTBI with INR < 1.2 and those with INR ≥ 1.2



factors, thus coagulation factor XIII was depleted. Coagulation factor XIII is activated by thrombin to cross-link fibrin. A deficiency of coagulation factor XIII is known to worsen clot stability and increase bleeding tendency [9].

The reduced activity of thrombin-dependent factors, such as V, VIII, and XIII, may suggest impaired clot formation in iTBI with a bleeding phenotype. The authors of several studies have reported elevated thrombin generation and thrombin-generation-associated markers within 6 and 12 h following iTBI [10–12]. However, Massaro and co-workers [13] have shown a progressive and delayed procoagulant state after 48 h of injury in iTBI using thromboelastography. Thus, the decrease in thrombin activity observed in the present study indicates a consumption of thrombin, which may be caused by severe consumption of coagulation factors. Accordingly, the concentration of fibrin monomers increased with INR up to 3.0-fold with $\text{INR} \geq 1.2$, whereas the activity of antithrombin decreased. A recent prospective study demonstrated that high fibrin monomer levels on admission in severe iTBI was associated with progression of hemorrhage [14]. High concentrations of fibrin monomers were observed in patients with hemostatic abnormalities (e.g.,

disseminated intravascular coagulation and deep vein thrombosis), and it seems promising to further evaluate the procoagulant status in these patients [15].

In a prospective study in severely injured patients with iTBI, the authors reported a reduction of antithrombin levels, which was associated with increased clot strength due to high fibrinogen concentration in the subacute phase (between days 5 and 7) after iTBI [16]. In the present study, fibrinogen levels were not depleted with increasing INR. It was concluded that a procoagulant status may occur during different phases after severe TBI [16]. Moreover, clinical data indicated the presence of severe brain injuries ($\text{AIS}_{\text{Brain}} \geq 4$) in about 80% with PIH. An earlier study in critically injured trauma patients demonstrated a higher risk of thromboembolic complications with decreased antithrombin activity in those patients [17].

The balance between coagulation and fibrinolysis determines the stability of the fibrin clot. Apart from the aforementioned properties, thrombin is also involved in stabilizing the clot by activating TAFI. Alterations in thrombin generation and activation of TAFI will therefore directly affect the stabilization of clots against lysis

[18, 19]. In the present study, reduced thrombin generation was associated with decreased activation of TAFI in patients with $\text{INR} \geq 1.2$, which might have had a direct impact on clot protection against lysis.

In the present study, PAI-1 levels in plasma remained largely unaltered. PAI-1 is mainly produced by the endothelium and acts as a principal inhibitor of tissue plasminogen activator (tPA). According to the clinical data, patients suffered neither from hypoperfusion nor from shock, both of which are known triggers of the protein C pathway. This pathway inhibits PAI-1, leading to less inhibition of tPA, with the overall effect to promote lysis. In the experimental setting, TBI has been shown to initiate fibrinolysis, independent of shock and hypoperfusion, by releasing tPA and urokinase plasminogen activator from contused brain tissue [20]. It has also been shown that tPA activity is increased by approximately 30% within 1–3 h post TBI and returns to baseline levels by 24 h post TBI [21]. In the present study, plasminogen, which is cleaved by tPA to generate plasmin, decreased to 76% activity with increasing INR, potentially indicating enhanced conversion into plasmin. At the same time, antiplasmin activity also declined with increasing INR. These observations, together with the decreased activation of TAFI in $\text{INR} \geq 1.2$, are suggestive of increased fibrinolysis, as evidenced by increasing D-dimer levels with increasing INR. It may be precluded that TXA had an influence on fibrinolysis in patients with iTBI with increasing INR because of the restricted number of patients who actually had received the agent. Endogenous release of tPA from contused brain tissue may be a distinct trigger to promote coagulopathy in iTBI apart from hemostatic abnormalities seen in the general trauma population.

The present study also suggests an activation of the protein C pathway, independent of shock and hypoperfusion, to further promote lysis, as evidenced by reductions in pathway-associated proteins (e.g., protein S, coagulation factors V and X [indirect measurement of factor X], and protein C) itself with increasing INR. Low protein C levels have previously been reported in patients with iTBI and trauma patients with hypoperfusion and were associated with poor outcome [22, 23].

There is also retrospective evidence for patients with iTBI with high D-dimer levels in plasma being at higher risk for progressive hemorrhagic injury (PHI) and worse outcomes [24, 25]. In the present study, PHI, which included the enlargement of both extradural and subdural hematoma (25% and 43.7% in patients with $\text{INR} \geq 1.2$ vs. 20.8% and 38.8% in patients with $\text{INR} < 1.2$, respectively) and subarachnoid hemorrhage (50% in patients with $\text{INR} \geq 1.2$ vs. 47.2% in patients with $\text{INR} < 1.2$), could

be observed in patients with iTBI with elevated INR. In addition, mortality and unfavorable 6-month outcomes (GOS-E) increased with increasing INR. Several studies have previously confirmed D-dimer to predict PHI as well as correlations between elevated D-dimer levels and unfavorable outcomes, longer in-hospital stay, and higher mortality in patients with iTBI [25–28].

Although all efforts were undertaken to mirror early changes within the coagulation system in patients with iTBI, the major limitation of the present study remains the fairly large time window of patient inclusion into the CENTER-TBI core study. This precluded a stricter timeline for blood sampling after admission. In addition, blood sampling did not follow a timeline-guided protocol specifying exact time points, thereby causing variation between study centers that contributed to this study. The median time interval between injury and first blood sampling in the present study was 13.1 h (interquartile range 6.2–20.8 h). It is well accepted that hemostatic abnormalities after TBI occur quickly after the initial impact and follow dynamic patterns involving both hypercoagulative and hypocoagulative states with potential overlap [10]. Whether these findings may be interpreted as maladaptive/pathologic or represent a physiologic response to the impact remains unknown. The observation that thrombin generation and TAFI were both reduced in patients with $\text{INR} \geq 1.2$ could be interpreted as a protection against lysis. Hemostatic and/or resuscitation therapies in addition to emergency procedures during acute care might have had a theoretical impact on the present results, although the focus of the analysis was given to early acute and thus mostly undiluted blood samples. In addition, prehospital care was not documented in detail in the CENTER-TBI central database. Although the present work initially aimed to present a more detailed picture of the possible underlying mechanistic changes/exchanges in hemostasis after iTBI, including outcomes, the study remains rather descriptive. Nevertheless, we observed in the present study distinct patterns involving key pathways of the highly complex and dynamic coagulation system in patients with iTBI that offer windows of opportunity for further research.

Author details

¹ Institute for Research in Operative Medicine, Faculty of Health, Department of Medicine, Witten/Herdecke University, Ostmerheimer Str. 200, Building 38, 51109 Cologne, Germany. ² Emergency and Trauma Centre, Alfred Health, 55 Commercial Road, Melbourne, VIC 3004, Australia. ³ Department of Anesthesiology and Intensive Care, AUVA Trauma Hospital, Academic Teaching Hospital of the Paracelsus Medical University, Doktor-Franz-Rehrl-Platz 5, 5010 Salzburg, Austria. ⁴ Ludwig Boltzmann Institute for Experimental and Clinical Traumatology, AUVA Trauma Research Center, Donaueschingenstr. 13, 1200 Vienna,

Austria.⁵ Department of Anesthesiology, RWTH Aachen University Hospital, Pauwelsstraße 30, Aachen 52074, Germany.⁶ NHS Blood and Transplant, John Radcliffe Hospital, Oxford University Hospital NHS Foundation Trust, Headley Way, Headington, Oxford OX3 9DU, UK.⁷ Oxford Haemophilia and Thrombosis Centre, Churchill Hospital, Oxford University Hospitals NHS Foundation Trust and NIHR BRC Haematology Theme, Old Road, Headington, Oxford OX37LE, UK.⁸ Department of Traumatology, Orthopedic Surgery and Sports Traumatology, Cologne-Merheim Medical Centre, Witten/Herdecke University, Campus Cologne-Merheim, Ostmerheimer Str. 200, 51109 Cologne, Germany.

Acknowledgements

We would like to thank all CENTER-TBI centers, participants, and investigators for all their efforts realizing this project. We would like to thank the colleagues of the Institute of Transfusion Medicine at Cologne-Merheim Medical Center for the analysis of conventional coagulation parameters, coagulation factors, and anticoagulant proteins. We would further like to thank Sandra Spreizer and Gerhard Fritsch from the AUVA Lorenz Böhler Trauma Center, Vienna, for their support in the analysis of thrombin generation and fibrinolysis-regulating parameters.

Author contributions

Data were acquired, analyzed, and interpreted by JKB, VS, ST, HG, NS, and MM. Statistical expertise was provided by RL, HS, JZ, OG, RR, SS, and NC contributed to the conception of the study, provided scientific support, and critically revised the data. The manuscript was written by JKB and was critically reviewed by all authors. Supervision was provided by MM. All authors read and approved the final manuscript.

Source of support

Open Access funding enabled and organized by Projekt DEAL. The research described in this article was supported by the European Union's Seventh Framework Programme (FP7/2007–2013) under Grant agreement No. 602150 (CENTER-TBI). This research received further financial support for hemostatic analysis from the Hannelore Kohl Foundation under Grant agreement No. 2014014 (TBI study).

Conflict of interest

The authors declare that they have no conflict of interest regarding this article.

Ethics approval/informed consent

As part of the CENTER-TBI core study, the present analysis was performed in accordance with relevant local ethics and European law.

Open Access

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Appendix

The CENTER-TBI participants and investigators

Cecilia Åkerlund¹, Krisztina Amrein², Nada Andelic³, Lasse Andreassen⁴, Audny Anke⁵, Anna Antoni⁶, Gérard Audibert⁷, Philippe Azouvi⁸, Maria Luisa Azzolini⁹, Ronald Bartels¹⁰, Pál Barzó¹¹, Romuald Beauvais¹², Ronny Beer¹³, Bo-Michael Bellander¹⁴, Antonio Belli¹⁵, Habib Benali¹⁶, Maurizio Berardino¹⁷, Luigi Beretta⁹, Morten

Blaabjerg¹⁸, Peter Bragge¹⁹, Alexandra Brazinova²⁰, Vibeke Brinck²¹, Joanne Brooker²², Camilla Brorsson²³, Andras Buki²⁴, Monika Bullinger²⁵, Manuel Cabeleira²⁶, Alessio Caccioppola²⁷, Emiliana Calappi²⁷, Maria Rosa Calvi⁹, Peter Cameron²⁸, Guillermo Carbayo Lozano²⁹, Marco Carbonara²⁷, Simona Cavallo¹⁷, Giorgio Chevallard³⁰, Arturo Chiericato³⁰, Giuseppe Citerio^{31, 32}, Iris Ceyisakar³³, Hans Clusmann³⁴, Mark Coburn³⁵, Jonathan Coles³⁶, Jamie D. Cooper³⁷, Marta Correia³⁸, Amra Čović³⁹, Nicola Curry⁴⁰, Endre Czeiter²⁴, Marek Czosnyka²⁶, Claire Dahyot-Fizelier⁴¹, Paul Dark⁴², Helen Dawes⁴³, Véronique De Keyser⁴⁴, Vincent Degos¹⁶, Francesco Della Corte⁴⁵, Hugo den Boogert¹⁰, Bart Depreitere⁴⁶, Đula Đilvesi⁴⁷, Abhishek Dixit⁴⁸, Emma Donoghue²², Jens Dreier⁴⁹, Guy-Loup Dulière⁵⁰, Ari Ercole⁴⁸, Patrick Esser⁴³, Erzsébet Ezer⁵¹, Martin Fabricius⁵², Valery L. Feigin⁵³, Kelly Foks⁵⁴, Shirin Frisvold⁵⁵, Alex Furmanov⁵⁶, Pablo Gagliardo⁵⁷, Damien Galanaud¹⁶, Dashiell Gantner²⁸, Guoyi Gao⁵⁸, Pradeep George⁵⁹, Alexandre Ghuysen⁶⁰, Lelde Giga⁶¹, Ben Glocker⁶², Jagoš Golubovic⁴⁷, Pedro A. Gomez⁶³, Johannes Gratz⁶⁴, Benjamin Gravesteijn³³, Francesca Grossi⁴⁵, Russell L. Gruen⁶⁵, Deepak Gupta⁶⁶, Juanita A. Haagsma³³, Iain Haitsma⁶⁷, Raimund Helbok¹³, Eirik Helseth⁶⁸, Lindsay Horton⁶⁹, Jilske Huijben³³, Peter J. Hutchinson⁷⁰, Bram Jacobs⁷¹, Stefan Jankowski⁷², Mike Jarrett²¹, Ji-yao Jiang⁵⁸, Faye Johnson⁷³, Kelly Jones⁵³, Mladen Karan⁴⁷, Angelos G. Kolias⁷⁰, Erwin Kompanje⁷⁴, Daniel Kondziella⁵², Evgenios Koraropoulos⁴⁸, Lars-Owe Koskinen⁷⁵, Noémi Kovács⁷⁶, Ana Kowark³⁵, Alfonso Lagares⁶³, Linda Lanyon⁵⁹, Steven Laureys⁷⁷, Fiona Lecky^{78, 79}, Didier Ledoux⁷⁷, Rolf Lefering⁸⁰, Valerie Legrand⁸¹, Aurelie Lejeune⁸², Leon Levi⁸³, Roger Lightfoot⁸⁴, Hester Lingsma³³, Andrew I.R. Maas⁴⁴, Ana M. Castaño-León⁶³, Marc Maegele⁸⁵, Marek Majdan²⁰, Alex Manara⁸⁶, Geoffrey Manley⁸⁷, Costanza Martino⁸⁸, Hugues Maréchal⁵⁰, Julia Mattern⁸⁹, Catherine McMahon⁹⁰, Béla Melegh⁹¹, David Menon⁴⁸, Tomas Menovsky⁴⁴, Ana Mikolic³³, Benoit Misset⁷⁷, Visakh Muraleedharan⁵⁹, Lynnette Murray²⁸, Ancuta Negru⁹², David Nelson¹, Virginia Newcombe⁴⁸, Daan Nieboer³³, József Nyirádi², Otesile Olubukola⁷⁸, Matej Oresic⁹³, Fabrizio Ortolano²⁷, Aarno Palotie^{94, 95, 96}, Paul M. Parizel⁹⁷, Jean-François Payen⁹⁸, Natascha Perera¹², Vincent Perlbarg¹⁶, Paolo Persona⁹⁹, Wilco Peul¹⁰⁰, Anna Piippo-Karjalainen¹⁰¹, Matti Pirinen⁹⁴, Horia Ples⁹², Suzanne Polinder³³, Inigo Pomposo²⁹, Jussi P. Posti¹⁰², Louis Puybasset¹⁰³, Andreea Radoi¹⁰⁴, Arminas Ragauskas¹⁰⁵, Rahul Raj¹⁰¹, Malinka Rambadagalla¹⁰⁶, Jonathan Rhodes¹⁰⁷, Sylvia Richardson¹⁰⁸, Sophie Richter⁴⁸, Samuli Ripatti⁹⁴, Saulius Rocka¹⁰⁵, Cecilie Roe¹⁰⁹, Olav Roise^{110, 111}, Jonathan Rosand¹¹², Jeffrey V. Rosenfeld¹¹³, Christina Rosenlund¹¹⁴,

Guy Rosenthal⁵⁶, Rolf Rossaint³⁵, Sandra Rossi⁹⁹, Daniel Rueckert⁶², Martin Rusnák¹¹⁵, Juan Sahuquillo¹⁰⁴, Oliver Sakowitz^{89, 116}, Renan Sanchez-Porras¹¹⁶, Janos Sandor¹¹⁷, Nadine Schäfer⁸⁰, Silke Schmidt¹¹⁸, Herbert Schoechl¹¹⁹, Guus Schoonman¹²⁰, Rico Fredrik Schou¹²¹, Elisabeth Schwendenwein⁶, Charlie Sewalt³³, Toril Skandsen^{122, 123}, Peter Smielewski²⁶, Abayomi Sorinola¹²⁴, Emmanuel Stamatakis⁴⁸, Simon Stanworth⁴⁰, Robert Stevens¹²⁵, William Stewart¹²⁶, Ewout W. Steyerberg^{33, 127}, Nino Stocchetti¹²⁸, Nina Sundström¹²⁹, Anneliese Synnot^{22, 130}, Riikka Takala¹³¹, Viktória Tamás¹²⁴, Tomas Tamosuities¹³², Mark Steven Taylor²⁰, Braden Te Ao⁵³, Olli Tenovuo¹⁰², Alice Theadom⁵³, Matt Thomas⁸⁶, Dick Tibboel¹³³, Marjolein Timmers⁷⁴, Christos Tolia¹³⁴, Tony Trapani²⁸, Cristina Maria Tudora⁹², Andreas Unterberg⁸⁹, Peter Vajkoczy¹³⁵, Shirley Vallance²⁸, Egils Valeinis⁶¹, Zoltán Vámos⁵¹, Mathieu van der Jagt¹³⁶, Gregory Van der Steen⁴⁴, Joukje van der Naalt⁷¹, Jeroen T.J.M. van Dijk¹⁰⁰, Thomas A. van Essen¹⁰⁰, Wim Van Hecke¹³⁷, Caroline van Heugten¹³⁸, Dominique Van Praag¹³⁹, Thijs Vande Vyvere¹³⁷, RoelP. J. van Wijk¹⁰⁰, Alesia Vargiolu³², Emmanuel Vega⁸², Kimberley Velt³³, Jan Verheyden¹³⁷, Paul M. Vespa¹⁴⁰, Anne Vik^{122, 141}, Rimantas Vilcinis¹³², Victor Volovic⁶⁷, Nicole von Steinbüchel³⁹, Daphne Voormolen³³, Petar Vulekovic⁴⁷, Kevin K.W. Wang¹⁴², Eveline Wieggers³³, Guy Williams⁴⁸, Lindsay Wilson⁶⁹, Stefan Winzeck⁴⁸, Stefan Wolf¹⁴³, Zhihui Yang¹⁴², Peter Ylén¹⁴⁴, Alexander Younsi⁸⁹, Frederick A. Zeiler^{48, 145}, Veronika Zelinkova²⁰, Agate Ziverte⁶¹, Tommaso Zoerle²⁷.

¹Department of Physiology and Pharmacology, Section of Perioperative Medicine and Intensive Care, Karolinska Institutet, Stockholm, Sweden

²János Szentágothai Research Centre, University of Pécs, Pécs, Hungary

³Division of Surgery and Clinical Neuroscience, Department of Physical Medicine and Rehabilitation, Oslo University Hospital and University of Oslo, Oslo, Norway

⁴Department of Neurosurgery, University Hospital Northern Norway, Tromsø, Norway

⁵Department of Physical Medicine and Rehabilitation, University Hospital Northern Norway, Tromsø, Norway

⁶Trauma Surgery, Medical University Vienna, Vienna, Austria

⁷Department of Anesthesiology & Intensive Care, University Hospital Nancy, Nancy, France

⁸Raymond Poincaré hospital, Assistance Publique – Hôpitaux de Paris, Paris, France

⁹Department of Anesthesiology & Intensive Care, S Raffaele University Hospital, Milan, Italy

¹⁰Department of Neurosurgery, Radboud University Medical Center, Nijmegen, The Netherlands

¹¹Department of Neurosurgery, University of Szeged, Szeged, Hungary

¹²International Projects Management, ARTTIC, Munchen, Germany

¹³Department of Neurology, Neurological Intensive Care Unit, Medical University of Innsbruck, Innsbruck, Austria

¹⁴Department of Neurosurgery & Anesthesia & intensive care medicine, Karolinska University Hospital, Stockholm, Sweden

¹⁵NIHR Surgical Reconstruction and Microbiology Research Centre, Birmingham, UK

¹⁶Anesthésie-Réanimation, Assistance Publique – Hôpitaux de Paris, Paris, France

¹⁷Department of Anesthesia & ICU, AOU Città della Salute e della Scienza di Torino - Orthopedic and Trauma Center, Torino, Italy

¹⁸Department of Neurology, Odense University Hospital, Odense, Denmark

¹⁹BehaviourWorks Australia, Monash Sustainability Institute, Monash University, Victoria, Australia

²⁰Department of Public Health, Faculty of Health Sciences and Social Work, Trnava University, Trnava, Slovakia

²¹Quesgen Systems Inc., Burlingame, California, USA

²²Australian & New Zealand Intensive Care Research Centre, Department of Epidemiology and Preventive Medicine, School of Public Health and Preventive Medicine, Monash University, Melbourne, Australia

²³Department of Surgery and Perioperative Science, Umeå University, Umeå, Sweden.

²⁴Department of Neurosurgery, Medical School, University of Pécs, Hungary and Neurotrauma Research Group, János Szentágothai Research Centre, University of Pécs, Hungary

²⁵Department of Medical Psychology, Universitätsklinikum Hamburg-Eppendorf, Hamburg, Germany

²⁶Brain Physics Lab, Division of Neurosurgery, Dept of Clinical Neurosciences, University of Cambridge, Addenbrooke's Hospital, Cambridge, UK

²⁷Neuro ICU, Fondazione IRCCS Cà Granda Ospedale Maggiore Policlinico, Milan, Italy

²⁸ANZIC Research Centre, Monash University, Department of Epidemiology and Preventive Medicine, Melbourne, Victoria, Australia

²⁹Department of Neurosurgery, Hospital of Cruces, Bilbao, Spain

³⁰NeuroIntensive Care, Niguarda Hospital, Milan, Italy

³¹School of Medicine and Surgery, Università Milano Bicocca, Milano, Italy

³²NeuroIntensive Care, ASST di Monza, Monza, Italy

-
- ³³Department of Public Health, Erasmus Medical Center-University Medical Center, Rotterdam, The Netherlands
- ³⁴Department of Neurosurgery, Medical Faculty RWTH Aachen University, Aachen, Germany.
- ³⁵Department of Anaesthesiology, University Hospital of Aachen, Aachen, Germany
- ³⁶Department of Anesthesia & Neurointensive Care, Cambridge University Hospital NHS Foundation Trust, Cambridge, UK.
- ³⁷School of Public Health & PM, Monash University and The Alfred Hospital, Melbourne, Victoria, Australia
- ³⁸Radiology/MRI department, MRC Cognition and Brain Sciences Unit, Cambridge, UK
- ³⁹Institute of Medical Psychology and Medical Sociology, Universitätsmedizin Göttingen, Göttingen, Germany
- ⁴⁰Oxford University Hospitals NHS Trust, Oxford, UK
- ⁴¹Intensive Care Unit, CHU Poitiers, Poitiers, France
- ⁴²University of Manchester NIHR Biomedical Research Centre, Critical Care Directorate, Salford Royal Hospital NHS Foundation Trust, Salford, UK
- ⁴³Movement Science Group, Faculty of Health and Life Sciences, Oxford Brookes University, Oxford, UK
- ⁴⁴Department of Neurosurgery, Antwerp University Hospital and University of Antwerp, Edegem, Belgium
- ⁴⁵Department of Anesthesia & Intensive Care, Maggiore Della Carità Hospital, Novara, Italy
- ⁴⁶Department of Neurosurgery, University Hospitals Leuven, Leuven, Belgium
- ⁴⁷Department of Neurosurgery, Clinical centre of Vojvodina, Faculty of Medicine, University of Novi Sad, Novi Sad, Serbia
- ⁴⁸Division of Anaesthesia, University of Cambridge, Addenbrooke's Hospital, Cambridge, UK
- ⁴⁹Center for Stroke Research Berlin, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin, Germany
- ⁵⁰Intensive Care Unit, CHR Citadelle, Liège, Belgium
- ⁵¹Department of Anaesthesiology and Intensive Therapy, University of Pécs, Pécs, Hungary
- ⁵²Departments of Neurology, Clinical Neurophysiology and Neuroanesthesiology, Region Hovedstaden Rigshospitalet, Copenhagen, Denmark
- ⁵³National Institute for Stroke and Applied Neurosciences, Faculty of Health and Environmental Studies, Auckland University of Technology, Auckland, New Zealand
- ⁵⁴Department of Neurology, Erasmus MC, Rotterdam, the Netherlands
- ⁵⁵Department of Anesthesiology and Intensive care, University Hospital Northern Norway, Tromsø, Norway
- ⁵⁶Department of Neurosurgery, Hadassah-hebrew University Medical center, Jerusalem, Israel
- ⁵⁷Fundación Instituto Valenciano de Neurorrehabilitación (FIVAN), Valencia, Spain
- ⁵⁸Department of Neurosurgery, Shanghai Renji hospital, Shanghai Jiaotong University/school of medicine, Shanghai, China
- ⁵⁹Karolinska Institutet, INCF International Neuroinformatics Coordinating Facility, Stockholm, Sweden
- ⁶⁰Emergency Department, CHU, Liège, Belgium
- ⁶¹Neurosurgery clinic, Pauls Stradins Clinical University Hospital, Riga, Latvia
- ⁶²Department of Computing, Imperial College London, London, UK
- ⁶³Department of Neurosurgery, Hospital Universitario 12 de Octubre, Madrid, Spain
- ⁶⁴Department of Anesthesia, Critical Care and Pain Medicine, Medical University of Vienna, Austria
- ⁶⁵College of Health and Medicine, Australian National University, Canberra, Australia
- ⁶⁶Department of Neurosurgery, Neurosciences Centre & JPN Apex trauma centre, All India Institute of Medical Sciences, New Delhi-110029, India
- ⁶⁷Department of Neurosurgery, Erasmus MC, Rotterdam, the Netherlands
- ⁶⁸Department of Neurosurgery, Oslo University Hospital, Oslo, Norway
- ⁶⁹Division of Psychology, University of Stirling, Stirling, UK
- ⁷⁰Division of Neurosurgery, Department of Clinical Neurosciences, Addenbrooke's Hospital & University of Cambridge, Cambridge, UK.
- ⁷¹Department of Neurology, University of Groningen, University Medical Center Groningen, Groningen, Netherlands
- ⁷²Neurointensive Care, Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, UK
- ⁷³Salford Royal Hospital NHS Foundation Trust Acute Research Delivery Team, Salford, UK
- ⁷⁴Department of Intensive Care and Department of Ethics and Philosophy of Medicine, Erasmus Medical Center, Rotterdam, The Netherlands
- ⁷⁵Department of Clinical Neuroscience, Neurosurgery, Umeå University, Umeå, Sweden.
- ⁷⁶Hungarian Brain Research Program - Grant No. KTIA_13_NAP-A-II/8, University of Pécs, Pécs, Hungary
- ⁷⁷Cyclotron Research Center, University of Liège, Liège, Belgium
- ⁷⁸Centre for Urgent and Emergency Care Research (CURE), Health Services Research Section, School of Health and Related Research (SchARR), University of Sheffield, Sheffield, UK

-
- ⁷⁹Emergency Department, Salford Royal Hospital, Salford UK
- ⁸⁰Institute of Research in Operative Medicine (IFOM), Witten/Herdecke University, Cologne, Germany
- ⁸¹VP Global Project Management CNS, ICON, Paris, France
- ⁸²Department of Anesthesiology-Intensive Care, Lille University Hospital, Lille, France
- ⁸³Department of Neurosurgery, Rambam Medical Center, Haifa, Israel
- ⁸⁴Department of Anesthesiology & Intensive Care, University Hospitals Southampton NHS Trust, Southampton, UK
- ⁸⁵Cologne-Merheim Medical Center (CMMC), Department of Traumatology, Orthopedic Surgery and Sportmedicine, Witten/Herdecke University, Cologne, Germany
- ⁸⁶Intensive Care Unit, Southmead Hospital, Bristol, Bristol, UK
- ⁸⁷Department of Neurological Surgery, University of California, San Francisco, California, USA
- ⁸⁸Department of Anesthesia & Intensive Care, M. Bufalini Hospital, Cesena, Italy
- ⁸⁹Department of Neurosurgery, University Hospital Heidelberg, Heidelberg, Germany
- ⁹⁰Department of Neurosurgery, The Walton centre NHS Foundation Trust, Liverpool, UK
- ⁹¹Department of Medical Genetics, University of Pécs, Pécs, Hungary
- ⁹²Department of Neurosurgery, Emergency County Hospital Timisoara, Timisoara, Romania
- ⁹³School of Medical Sciences, Örebro University, Örebro, Sweden.
- ⁹⁴Institute for Molecular Medicine Finland, University of Helsinki, Helsinki, Finland
- ⁹⁵Analytic and Translational Genetics Unit, Department of Medicine; Psychiatric & Neurodevelopmental Genetics Unit, Department of Psychiatry; Department of Neurology, Massachusetts General Hospital, Boston, MA, USA
- ⁹⁶Program in Medical and Population Genetics; The Stanley Center for Psychiatric Research, The Broad Institute of MIT and Harvard, Cambridge, MA, USA
- ⁹⁷Department of Radiology, University of Antwerp, Edegem, Belgium
- ⁹⁸Department of Anesthesiology & Intensive Care, University Hospital of Grenoble, Grenoble, France
- ⁹⁹Department of Anesthesia & Intensive Care, Azienda Ospedaliera Università di Padova, Padova, Italy
- ¹⁰⁰Dept. of Neurosurgery, Leiden University Medical Center, Leiden, The Netherlands and Dept. of Neurosurgery, Medical Center Haaglanden, The Hague, The Netherlands
- ¹⁰¹Department of Neurosurgery, Helsinki University Central Hospital
- ¹⁰²Division of Clinical Neurosciences, Department of Neurosurgery and Turku Brain Injury Centre, Turku University Hospital and University of Turku, Turku, Finland
- ¹⁰³Department of Anesthesiology and Critical Care, Pitié-Salpêtrière Teaching Hospital, Assistance Publique, Hôpitaux de Paris and University Pierre et Marie Curie, Paris, France
- ¹⁰⁴Neurotraumatology and Neurosurgery Research Unit (UNINN), Vall d'Hebron Research Institute, Barcelona, Spain
- ¹⁰⁵Department of Neurosurgery, Kaunas University of Technology and Vilnius University, Vilnius, Lithuania
- ¹⁰⁶Department of Neurosurgery, Rezekne Hospital, Latvia
- ¹⁰⁷Department of Anaesthesia, Critical Care & Pain Medicine NHS Lothian & University of Edinburgh, Edinburgh, UK
- ¹⁰⁸Director, MRC Biostatistics Unit, Cambridge Institute of Public Health, Cambridge, UK
- ¹⁰⁹Department of Physical Medicine and Rehabilitation, Oslo University Hospital/University of Oslo, Oslo, Norway
- ¹¹⁰Division of Orthopedics, Oslo University Hospital, Oslo, Norway
- ¹¹¹Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway
- ¹¹²Broad Institute, Cambridge MA Harvard Medical School, Boston MA, Massachusetts General Hospital, Boston MA, USA
- ¹¹³National Trauma Research Institute, The Alfred Hospital, Monash University, Melbourne, Victoria, Australia
- ¹¹⁴Department of Neurosurgery, Odense University Hospital, Odense, Denmark
- ¹¹⁵International Neurotrauma Research Organisation, Vienna, Austria
- ¹¹⁶Klinik für Neurochirurgie, Klinikum Ludwigsburg, Ludwigsburg, Germany
- ¹¹⁷Division of Biostatistics and Epidemiology, Department of Preventive Medicine, University of Debrecen, Debrecen, Hungary
- ¹¹⁸Department Health and Prevention, University Greifswald, Greifswald, Germany
- ¹¹⁹Department of Anaesthesiology and Intensive Care, AUA Trauma Hospital, Salzburg, Austria
- ¹²⁰Department of Neurology, Elisabeth-TweeSteden Ziekenhuis, Tilburg, the Netherlands
- ¹²¹Department of Neuroanesthesia and Neurointensive Care, Odense University Hospital, Odense, Denmark
- ¹²²Department of Neuromedicine and Movement Science, Norwegian University of Science and Technology, NTNU, Trondheim, Norway

¹²³Department of Physical Medicine and Rehabilitation, St.Olavs Hospital, Trondheim University Hospital, Trondheim, Norway

¹²⁴Department of Neurosurgery, University of Pécs, Pécs, Hungary

¹²⁵Division of Neuroscience Critical Care, John Hopkins University School of Medicine, Baltimore, USA

¹²⁶Department of Neuropathology, Queen Elizabeth University Hospital and University of Glasgow, Glasgow, UK

¹²⁷Dept. of Department of Biomedical Data Sciences, Leiden University Medical Center, Leiden, The Netherlands

¹²⁸Department of Pathophysiology and Transplantation, Milan University, and Neuroscience ICU, Fondazione IRCCS Cà Granda Ospedale Maggiore Policlinico, Milano, Italy

¹²⁹Department of Radiation Sciences, Biomedical Engineering, Umeå University, Umeå, Sweden.

¹³⁰Cochrane Consumers and Communication Review Group, Centre for Health Communication and Participation, School of Psychology and Public Health, La Trobe University, Melbourne, Australia

¹³¹Perioperative Services, Intensive Care Medicine and Pain Management, Turku University Hospital and University of Turku, Turku, Finland

¹³²Department of Neurosurgery, Kaunas University of Health Sciences, Kaunas, Lithuania

¹³³Intensive Care and Department of Pediatric Surgery, Erasmus Medical Center, Sophia Children's Hospital, Rotterdam, The Netherlands

¹³⁴Department of Neurosurgery, Kings college London, London, UK

¹³⁵Neurologie, Neurochirurgie und Psychiatrie, Charité – Universitätsmedizin Berlin, Berlin, Germany

¹³⁶Department of Intensive Care Adults, Erasmus MC– University Medical Center Rotterdam, Rotterdam, the Netherlands.

¹³⁷icoMetrix NV, Leuven, Belgium

¹³⁸Movement Science Group, Faculty of Health and Life Sciences, Oxford Brookes University, Oxford, UK

¹³⁹Psychology Department, Antwerp University Hospital, Edegem, Belgium

¹⁴⁰Director of Neurocritical Care, University of California, Los Angeles, USA

¹⁴¹Department of Neurosurgery, St.Olavs Hospital, Trondheim University Hospital, Trondheim, Norway

¹⁴²Department of Emergency Medicine, University of Florida, Gainesville, Florida, USA

¹⁴³Department of Neurosurgery, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin, Germany.

¹⁴⁴VTT Technical Research Centre, Tampere, Finland.

¹⁴⁵Section of Neurosurgery, Department of Surgery, Rady Faculty of Health Sciences, University of Manitoba, Winnipeg, MB, Canada

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 14 June 2021 Accepted: 11 November 2021

Published online: 16 December 2021

References

1. Maas AIR, Menon DK, David Adelson PD, et al. Traumatic brain injury: Integrated approaches to improve prevention, clinical care, and research. *Lancet Neurol.* 2017;16(12):987–1048.
2. Dewan MC, Rattani A, Gupta S, et al. Estimating the global incidence of traumatic brain injury. *J Neurosurg.* 2019;130(4):1080–97.
3. Maegele M, Schöchl H, Menovsky T, et al. Coagulopathy and haemorrhagic progression in traumatic brain injury: advances in mechanisms, diagnosis, and management. *Lancet Neurol.* 2017;16(8):630–47.
4. Böhm JK, Güting H, Thorn S, et al. Global characterisation of coagulopathy in isolated traumatic brain injury (ITBI): a CENTER-TBI analysis. *Neurocrit Care.* 2020;2:1029.
5. Mathieu F, Güting H, Gravesteyn B, et al. Impact of antithrombotic agents on radiological lesion progression in acute traumatic brain injury: a CENTER-TBI propensity-matched cohort analysis. *J Neurotrauma.* 2020;37(19):2069–80.
6. Maegele M, Aversa J, Marsee MK, et al. Changes in coagulation following brain injury. *Semin Thromb Hemost.* 2020;46(2):155–66.
7. Steyerberg EW, Wieggers E, Sewalt C, et al. Case-mix, care pathways, and outcomes in patients with traumatic brain injury in CENTER-TBI: a European prospective, multicentre, longitudinal, cohort study. *Lancet Neurol.* 2019;18(10):923–34.
8. Peltan ID, Vande Vusse LK, Maier RV, Watkins TR. An international normalized ratio-based definition of acute traumatic coagulopathy is associated with mortality, venous thromboembolism, and multiple organ failure after injury. *Crit Care Med.* 2015;43(7):1029.
9. Levy JH, Greenberg C. Biology of Factor XIII and clinical manifestations of Factor XIII deficiency. *Transfusion.* 2013;53(5):1009.
10. Fletcher-Sandersjö A, Thelin EP, Maegele M, Svensson M, Bellander BM. Time course of hemostatic disruptions after traumatic brain injury: a systematic review of the literature. *Neurocrit Care.* 2020;20:1798.
11. Scherer RU, Spangenberg P. Procoagulant activity in patients with isolated severe head trauma. *Crit Care Med.* 1998;26(1):149–56.
12. Albert V, Arulsevi S, Agrawal D, Pati HP, Pandey RM. Early posttraumatic changes in coagulation and fibrinolysis systems in isolated severe traumatic brain injury patients and its influence on immediate outcome. *Hematol Oncol Stem Cell Ther.* 2019;12(1):32–43.
13. Massaro AM, Doerfler S, Nawalinski K, et al. Thromboelastography defines late hypercoagulability after TBI: a pilot study. *Neurocrit Care.* 2015;22(1):45–51.
14. Esnault P, Mathais Q, D'Aranda E, et al. Ability of fibrin monomers to predict progressive hemorrhagic injury in patients with severe traumatic brain injury. *Neurocrit Care.* 2020;33(1):182–95.
15. Refaai MA, Riley P, Mardovina T, Bell PD. The clinical significance of fibrin monomers. *Thromb Haemost.* 2018;118(11):1856–66.
16. Kalgudi S, Ho KM. Incidence of antithrombin deficiency and anti-cardiolipin antibodies after severe traumatic brain injury: a prospective cohort study. *Neurocrit Care.* 2021;34(1):227–35.
17. Owings JT, Bagley M, Gosselin R, Romac D, Disbrow E. Effect of critical injury on plasma antithrombin activity: low antithrombin levels are associated with thromboembolic complications. *J Trauma Injury Infect Crit Care.* 1996;93:396–406.

-
18. Mosnier LO, Bouma BN. Regulation of fibrinolysis by Thrombin Activatable Fibrinolysis Inhibitor, an unstable carboxypeptidase B that unites the pathways of coagulation and fibrinolysis. *Arterioscler Thromb Vasc Biol.* 2006;26(11):2445–53.
 19. Bouma BN, Mosnier LO. Thrombin activatable fibrinolysis inhibitor (TAFI)—How does thrombin regulate fibrinolysis? *Ann Med.* 2006;38(6):378–88.
 20. Hijazi N, Fanne RA, Abramovitch R, et al. Endogenous plasminogen activators mediate progressive intracerebral hemorrhage after traumatic brain injury in mice. *Blood.* 2015;125(16):2558–67.
 21. Sashindranath M, Samson AL, Downes CE, et al. Compartment- and context-specific changes in tissue-type plasminogen activator (tPA) activity following brain injury and pharmacological stimulation. *Lab Invest.* 2011;91(7):1079–91.
 22. Cohen MJ, Brohi K, Ganter MT, Manley GT, Mackersie RC, Pittet JF. Early coagulopathy after traumatic brain injury: the role of hypoperfusion and the protein c pathway. *J Trauma.* 2007;63(6):1254–62.
 23. Brohi K, Cohen MJ, Ganter MT, Matthay MA, MacKersie RC, Pittet JF. Acute traumatic coagulopathy: initiated by hypoperfusion: modulated through the protein C pathway? *Ann Surg.* 2007;245(5):812–8.
 24. Karri J, Cardenas JC, Matijevic N, et al. Early fibrinolysis associated with hemorrhagic progression following traumatic brain injury. *Shock.* 2017;48(6):644–50.
 25. Zhang J, He M, Song Y, Xu J. Prognostic role of D-dimer level upon admission in patients with traumatic brain injury. *Med (United States).* 2018;97(31):e11774.
 26. Saggat V, Mittal RS, Vyas MC. Hemostatic abnormalities in patients with closed head injuries and their role in predicting early mortality. *J Neurotrauma.* 2009;26(10):1665–8.
 27. Kuo JR, Lin KC, Lu CL, Lin HJ, Wang CC, Chang CH. Correlation of a high D-dimer level with poor outcome in traumatic intracranial hemorrhage. *Eur J Neurol.* 2007;14(10):1073–8.
 28. Subedi PP, Shilpakar SK. Correlation of D-Dimer level with outcome in traumatic brain injury. *J Soc Surg Nepal.* 2016;17(1):21–30.