






# Quantifying Europe's biodiversity footprints and the role of urbanization and income

Maximilian Koslowski<sup>1,2</sup> , Daniel D. Moran<sup>1</sup> , Alexandre Tisserant<sup>1</sup> ,  
Francesca Veronesi<sup>1</sup>  and Richard Wood<sup>1</sup> 

## Long Form Research Paper

**Cite this article:** Koslowski M, Moran DD, Tisserant A, Veronesi F, Wood R (2020). Quantifying Europe's biodiversity footprints and the role of urbanization and income. *Global Sustainability* 3, e1, 1–12. <https://doi.org/10.1017/sus.2019.23>

Received: 23 November 2018  
Revised: 29 October 2019  
Accepted: 29 October 2019

### Keywords:

Biodiversity footprint; MRIO; household consumption; footprint

**Author for correspondence:** Maximilian Koslowski, E-mail: [maximilian.koslowski@inatech.uni-freiburg.de](mailto:maximilian.koslowski@inatech.uni-freiburg.de)

<sup>1</sup>Programme for Industrial Ecology, Department of Energy and Process Technology, Norwegian University of Science and Technology, Trondheim 7010 Norway and <sup>2</sup>Institute for Sustainable Systems Engineering, Albert Ludwig University of Freiburg, 79110 Freiburg i. Br., Germany

### Non-technical abstract

Biodiversity footprinting links consumers to the biodiversity pressure their consumption induces, thereby informing choices and enabling participation in remediation measures. In order for countries, cities and households to reduce their impacts it is useful to know more precisely what the various drivers of their footprints are. Here we ask: do urban or rural areas in Europe exert higher biodiversity footprints? And how strongly coupled are income and biodiversity losses? Studying urban versus rural households at the country level in Europe, we found both have generally similar footprints, but that higher income households clearly drive higher footprints.

### Technical abstract

We examined the role of selected socio-economic variables regarding biodiversity-related impacts associated with European household consumption in 2005 and 2010, asking: does urbanization alone drive higher biodiversity footprints, or what are the relative contributions of income and urbanity? We applied a multi-regional input-output (MRIO) model, supplemented with data from consumer expenditure surveys and extended by life cycle impact assessment methodologies to account for biodiversity losses. We find that urbanization and higher income are important sources of higher absolute biodiversity footprints. On a per capita basis, results are mixed, though a slight trend of higher impacts from city residents in most countries, as well as a general positive correlation with income, can be observed. Also, while wealthy European countries are accountable for the largest species losses overall, it is the ones with a high gross domestic product per capita and those bordering the Mediterranean Sea that have the highest per capita biodiversity footprints. Additionally, most European countries and Europe as a whole are net importers of biodiversity losses, with land use generally being the dominating impact category.

## Social media summary

Wealthy urban areas most responsible for Europe's global biodiversity footprint.

### 1. Introduction

The International Union for Conservation of Nature and Natural Resources (IUCN) estimates that more than 28,000 species are presently threatened with extinction (IUCN 2019). The primary causes of biodiversity loss include habitat loss, climate change, pollution, overexploitation and invasive species (Millennium Ecosystem Assessment 2005; WWF 2016). While the production of goods like coffee or beef is a direct domestic driver for such environmental pressures, the demand for these goods often lies in other regions of the world (Lenzen *et al.* 2012; Chaudhary & Kastner 2016; Moran & Kanemoto 2017; Veronesi *et al.* 2017; Wilting *et al.* 2017; Wiedmann & Lenzen 2018). European countries in particular are responsible for a large share of environmental pressures originating abroad (Tukker *et al.* 2016; Wood *et al.* 2018).

Household consumption is an endpoint of global supply chains and can thus be considered to be a major driver of environmental burdens (Ivanova *et al.* 2016). What shapes household consumption will therefore have an impact on global biodiversity loss. Additionally, the composition of household demand is neither static, nor equal across different groups, but rather depends on multiple socio-cultural and economic factors (Froemelt *et al.* 2018). As countries and the world as a whole become increasingly urban, it is important to understand how this urbanization process could lead to overall changes in the aforementioned composition of household demand. Do urban, semi-urban and rural households have similar consumption preferences? How much does urbanization alone, rather than changes in income or household

© The Author(s) 2020. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

size, influence changes in consumption preferences? Whilst the role of urbanization in influencing household carbon footprints has been well studied (Jones & Kammen 2014; Kanemoto *et al.* 2014; Wiedmann *et al.* 2016; Moran *et al.* 2018), there is no assessment for biodiversity footprints, which are associated with different consumption categories. That is, biodiversity footprints are associated majorly with food and fibre, whereas carbon footprints are more associated with metals, complex products and services. And while urbanites in wealthy countries tend to have lower emissions, urbanization in developing countries is often linked to increases in wealth and thus higher footprints compared with rural households. Hence, findings on the relationship between urbanization and emissions might differ from those between urbanization and biodiversity footprints.

This paper proceeds as follows: we first discuss the study methodology, after which we present European biodiversity footprints on regional, national and sub-national levels, with a focus on urban vs rural consumption patterns. We outline the influence of selected socio-economic drivers, give detailed accounts of species losses embodied in trade, and describe the role of different product sectors and impact categories. Finally, we discuss these results with regard to current socio-economic developments, and future research.

## 2. Materials and methods

Much traditional environmental impact analysis studies impacts associated with production. The footprint concept reallocates, through trade and transformation steps, those impacts of primary producers to the households whose final demand ultimately induces them. Thus, for instance, a consumer buying palm oil, or products containing palm oil, may be considered responsible for forest losses in Indonesia and other environmental burdens that occurred along the supply chain of the palm oil product.

To implement this in the present study, we establish a standard environmentally extended Leontief demand-pull model (Leontief 1936, 1970; see also SI6) using a global multi-regional input-output (MRIO) table to estimate consumption footprints. The MRIO database includes detail on average household consumption patterns in each country, but for this study we disaggregated that by incorporating data from consumer expenditure surveys (CESs) in order to obtain detail on consumption patterns by various socio-economic variables. Additionally, the MRIO database natively describes only the physical resources required to satisfy consumption, e.g. hectares of wheat cultivated, cubic metres of roundwood, etc. To evaluate not just the resource demands of consumption but estimate the full ecosystem impact of that consumption, we extend the MRIO model with a life cycle impact assessment (LCIA) model (Veronesi *et al.* 2017). LCIA is commonly used within life cycle assessments (LCA) as a modelling step to convert physical resource demands into estimates of how impactful those pressures are on ecosystems. Extending pressure footprints from MRIO with an LCIA model allows us to model the potential consequences on biodiversity and thus helps to highlight the potential impacts. The analyses were conducted for the years 2005 and 2010.

### 2.1. Environmentally extended MRIO analysis

Various methods exist for analysing environmental impacts. Although bottom-up approaches such as LCA allow for comparatively higher detail, namely analyses on product or technology

level, the top-down accounting method MRIO is preferable for national or regional assessments because of the complete coverage of supply chains and the avoidance of truncation errors (Majeau-Bettez *et al.* 2011; Wiedmann & Barrett 2013; Steen-Olsen *et al.* 2016). MRIO analysis covers both inter-industry and final demand in multiple regions and their bilateral trade interlinkages. By including environmental extensions, MRIO can track environmental pressures associated with trade flows of goods and services.

Environmentally extended MRIO (EE-MRIO) is an established and well-described approach for analysing footprints and environmental burdens associated with supply chains and consumption (Leontief 1936, 1970; Miller & Blair 2009; Kitzes 2013; Wiedmann & Lenzen 2018; see also SI6 for a description of a basic input-output model). EE-MRIO reattributes emissions and resource uses within a nation (production-based account) through multiple trade and transformation steps to final consumers (consumption-based account; Wiebe & Yamano 2016; see SI1 for an introduction into both variations). EE-MRIO analysis has been used in numerous previous studies for calculating footprints regarding environmental pressures (see SI2 for a detailed overview of MRIO applications), e.g. greenhouse gas emissions (Kanemoto *et al.* 2016; Wiedmann *et al.*, 2016; Moran *et al.*, 2018), land use (Ewing *et al.*, 2012), material requirements (Wiedmann *et al.* 2015; Giljum *et al.* 2016; Tukker *et al.* 2016), water stress (Lenzen *et al.* 2013; Lutter *et al.* 2016; Mekonnen & Hoekstra 2011), or threatened biodiversity (Lenzen *et al.* 2012; Moran & Kanemoto 2017).

For our analytical EE-MRIO model, we used data from the EXIOBASE MRIO database (Tukker *et al.*, 2013; Wood *et al.*, 2015; Stadler *et al.*, 2018) to calculate consumption-based accounts. EXIOBASE v3.4 represents the world economy for the period 1995–2011, distinguishing between 28 EU member countries, 16 major economies, and five rest of the world (RoW) regions, each of the above with a sectoral detail of 163 industries by 200 products. Moreover, EXIOBASE includes an extensive account of environmental pressures, covering a variety of emissions and resource uses (see section 2.4). The rationale for choosing EXIOBASE (over other MRIO databases such as Eora, WIOD or GTAP; see SI3 for a comparative overview) was twofold: first, it provides the highest sector resolution in comparison with other MRIO databases, and second, its individual representation of European countries allowed for a satisfying CES-MRIO data fit.

### 2.2. Consumer expenditure survey data reconciliation

While the applied EE-MRIO model in its original form provides information on household final demand per country, no further disaggregation of it with respect to socio-economic variables is provided. CESs contain this information, so that complementing the EE-MRIO model with CES data allows the analysis of household final demand on a virtual sub-national level for a multi-region (Steen-Olsen *et al.* 2016; Ivanova *et al.* 2017).

Eurostat's CES data (Eurostat 2018b) stems from annual large-scale household budget surveys conducted by National Statistical Institutes (Eurostat 2018a). These have been run every 5–6 years since 1988 with 2010 being the newest comprehensive dataset available (Eurostat 2015). 2005 data also had good country coverage.

The extension of the present EE-MRIO model with CES data required several reconciliation steps, the procedure of which followed to a large extent the approach taken by Steen-Olsen *et al.*

(2016) and Ivanova *et al.* (2017), and is described in detail in SI4. Essential stages were the upscaling of country-specific mean consumption expenditure per household to the national level, structured by socio-economic parameters and sectors according to the COICOP classification (COICOP is the acronym of 'Classification of Individual Consumption according to Purpose'; United Nations Statistics Division 2018); the accounting for underreporting, i.e. including detail on the difference between household expenditures reported in CES and MRIO data (Bee *et al.* 2012; Steen-Olsen *et al.*, 2016; Ivanova *et al.* 2017); and the COICOP-EXIOBASE sector bridging using country-specific, weighted concordance matrices. In the course of this adjustment, the valuation in purchaser prices had to be converted into basic prices.

CES data were structured according to various socio-economic variables: complete data were available for degrees of urbanization (note that we use the terms 'degrees of urbanization' and 'area types' interchangeably), income quintiles, age groups and types of households (for most countries for the selected years; Tables S3–S7). While degrees of urbanization are the central point in this study for examining urban vs rural consumption patterns, the other socio-economic variables were examined additionally to account for differences in lifestyle and wealth. As defined by Eurostat (2018b, 2019), the degrees of urbanization classification distinguishes between three parameters (cities/urban, towns/suburban, rural), based on population densities, whereas income is divided into quintiles. Types of households describe the household size, i.e. the number of people living there. As the category of age groups is only accounting for the age of the main income earner per household, resulting footprints were deemed to be less relevant and are therefore only shown in the spreadsheet appendix and SI10.

### 2.3. Impact assessment methods

For the conversion of environmental pressures to biodiversity impacts, the LCIA methods LC-IMPACT (Verones *et al.* 2019) and ReCiPe (Huijbregts *et al.* 2016) were chosen due to their level of detail and ability to expand the impact assessment to endpoint levels (also sometimes called damage levels) such as potentially lost fractions of species as proxy for biodiversity loss via characterization factors. These endpoint characterisation factors quantify the consequence of an anthropogenic intervention on ecosystem quality, human health or resource use (Verones *et al.* 2019), which are the three most commonly used *Areas of Protection* and are used both in ReCiPe and LC-IMPACT. In comparison, midpoint characterization factors quantify impacts at an intermediary step in the cause-effect chain and are usually indicated in emission or resource use equivalents. Here, we focus on endpoint impacts related to ecosystem quality only, scilicet biodiversity losses associated with household final demand. The term biodiversity being ambiguous (Curran *et al.* 2011), it is species richness (or rather, losses of it) that the above impact assessment methods measure via distinct metrics.

Despite their similar applications and foundations, LC-IMPACT and ReCiPe differ considerably. While LC-IMPACT offers spatially explicit endpoint characterization factors of fine granularity, ReCiPe provides only country-specific as well as weighted, globally averaged endpoint characterization factors. Moreover, LC-IMPACT accounts for potential *global* species losses including the vulnerability of species through corresponding scores, whereas ReCiPe covers potential *local* species losses. The metrics used for measuring biodiversity loss

are the globally potentially disappeared fraction of species over time (PDF; LC-IMPACT) and the time-integrated number of species lost (species.yr; ReCiPe), respectively. As for the actual factors, these are given per emission or resource use, for instance, PDF.yr/m<sup>2</sup> for land use (Tables S10–S13).

Both PDF and species.yr are addressing potential biodiversity loss in terms of a loss in species richness. The included taxonomic groups vary for both ReCiPe and LC-IMPACT between the impact categories. For example, while toxicity is based on lab tests and a range of species, terrestrial acidification takes into account vascular plants. In LC-IMPACT, land use takes birds, amphibians, reptiles and mammals into account, while water use is based on birds, amphibians, reptiles, water-dependent mammals and vascular plants (Verones *et al.* 2019). In ReCiPe, water use looks at fish and vascular plants and land use includes plants, mammals, birds and arthropods (Huijbregts *et al.* 2016).

The PDF is a relative measure, based on the fraction of species richness lost in relation to the total number of species (in the considered taxonomic groups). The global PDF used in LC-IMPACT considers the global loss of these fractions in a spatially differentiated way by considering the individual species' vulnerability. This was done by including range area and IUCN threat levels, as explained in Verones *et al.* (2013) and Chaudhary *et al.* (2015). As a simple and extreme example, an endemic species occurring in one region only, would be globally extinct if it is extinct in this region, thus the PDF of that species would be 1.


Species.yr, on the other hand, is an absolute measure of the potential loss of species, which is used in ReCiPe to be able to aggregate impacts across ecosystem types. To arrive at this absolute metric, based on previously calculated local, relative species losses, global species densities per ecosystem type are used (Huijbregts *et al.*, 2017).

For calculating biodiversity impact footprints in the present analysis, spatially explicit LC-IMPACT characterization factors and weighted globally averaged ReCiPe mid- to endpoint conversion factors were applied (the latter were applied to the midpoint factors in EXIOBASE). Time constraints prevented the application of ReCiPe's country-specific factors. We used LC-IMPACT's characterization factors for 100-year time horizons and certain effects, and ReCiPe's hierarchist perspective. Footprint results based on LC-IMPACT's extended characterization factors are included in the corresponding Excel spreadsheet in the digital SI. More details on the applied impact assessment methods, as well as the preparation and derivation of factors can be found in SI9.

### 2.4. Accounts of emissions and resource uses

The environmental satellite account of EXIOBASE covers 1338 stressors of different domains and is given in intensities, i.e. as environmental loads in respective units per monetary unit (Stadler *et al.* 2018). In the present study, multiple impact categories were considered, each of which relied on the aggregation of relevant stressors, when calculating four different footprint types (see SI8).

LC-IMPACT and pressure footprints comprise the impact categories land occupation, blue water consumption, global warming, photochemical ozone formation, freshwater eutrophication, and terrestrial acidification. ReCiPe and characterized pressure footprints encompass land use, water consumption, global warming, terrestrial acidification, and toxicity. Photochemical ozone formation was not assessed for ReCiPe footprints, although a mid- to endpoint conversion factor is available; this is due to the lack of corresponding midpoint characterisation factors in EXIOBASE.



**Fig. 1.** LC-IMPACT biodiversity footprints of EU28 + Norway for 2010. (a) shows absolute national biodiversity footprints and (b) shows per capita biodiversity footprints against the per capita GDP per country. See Table S2 for country abbreviations. Countries other than EU28 + Norway are grey-shaded in (a). The dotted lines in (b) represent the per capita footprint and GDP averages (see Tables S14–S16).

For a comparison of the footprints, only impact categories covered by all respective accounts were considered (unless otherwise stated).

While pressure accounts aggregate stressor intensities where applicable and are classified according to the above categories, LC-IMPACT accounts convert these aggregated stressor intensities into endpoint intensities, i.e. environmental impacts per monetary unit, before the actual footprint calculation. For both pressure and LC-IMPACT footprints, the stressors were selected manually. In comparison, characterised pressure and ReCiPe footprints applied both the same selection of EXIOBASE midpoint characterization factors to the stressor intensity matrix, thus retrieving emission and resource use equivalents per monetary unit; that is, no stressors had to be selected manually. In the case of ReCiPe footprints, the stressor equivalents per monetary unit were then multiplied by the corresponding mid- to endpoint conversion factors, thus yielding, similar to LC-IMPACT accounts, environmental impacts per monetary unit. As a modelling choice and due to data limitations, direct household impacts were not considered in either case.

### 2.5. Methodological limitations


Our approach requires certain considerations. For one, we only examine the biodiversity impacts due to production processes, and not the direct household impacts nor the direct impact of urban expansion in Europe which may potentially fragment existing ecosystems. Furthermore, an allocation of environmental impacts to illegal activities such as illegal logging or poaching was not possible due to data unavailability – although the impacts were potentially accounted for through the impact assessment characterization factors; thus, ecosystem damage due to illegal activities was falsely allocated to legal economic trade flows. Furthermore, the impacts of production were allocated based on expenditure – rather than for example mass – meaning the inter-industry transactions and household consumption were measured in monetary terms (Tukker *et al.* 2016).

In terms of years studied, we were highly limited to available data for 2005 and 2010 – two years that bridge the global financial

crisis and consequent economic restructuring. Moreover, weighted country-specific bridge matrices were only available for 2010. These were applied in the CES data reconciliations for both reference years, as changes were assumed to be negligible.

A temporal mismatch also existed between the MRIO model and the characterization factors of both impact assessment methods, as the characterization factors were mainly valid for 2016 and later. However, it was assumed that over the years the characterization factors would not have changed structurally, but only in their magnitude. In addition, LC-IMPACT and ReCiPe differ considerably in their methodologies and assumptions; hence, a direct comparison between the two respective impact footprints must be treated with caution. Although both impact assessment methods account for biodiversity loss in their respective units, neither of them allows an interpretation of ecosystem functioning. That is, implications of impact footprints depend not only on the presently assessed species richness, but also on ecosystem resilience. However, LC-IMPACT's vulnerability scores already point in this direction. Moreover, taxa coverage differs between the methodologies and even across impact categories. Further, the high spatial detail of the original LC-IMPACT characterization factors gets lost by averaging them per country when preparing for the MRIO calculus, whereas the globally averaged ReCiPe characterization factors provide too little spatial detail. The absence of spatial detail on production is another limitation. In this study it is assumed that all goods exported from a country have national-average biodiversity intensity; in other words, lacking information about the true spatial pattern of production within a country, it is not possible to make use of the spatially explicit characterization factors that are available, so a national average is used instead. Other studies have worked to improve the spatial resolution of production, including Moran & Kanemoto (2017) and Godar *et al.* (2015). For details on limitations of the applied LCIA methods, the reader is referred to the respective publications.

Uncertainty and sensitivity analyses would add additional detail but would require more information on the source data.



**Fig. 2.** Total global biodiversity losses due to European household consumption in 2010, measured in PDF. The panel in the bottom-left corner shows the biodiversity losses in Europe due to land use for annual crops (cf. Figure 4), associated with European household consumption in 2010. The countries are grouped according to the EXIOBASE-classification.

Crucially, the level of uncertainty itself is an unknown, so most sensitivity analysis approaches would make an assumption about the degree of confidence (i.e. the relative standard deviation for values) and thus the net information gain from conducting an exercise with a sensitivity can be quite small. There is no uncertainty information available for the LC-IMPACT, ReCiPe or CES data, so making assumptions about the uncertainty of those would not add information about the reliability of the results. In this study, all values reported should be treated as a mean estimate. As for the choice of the MRIO model, other studies on the reliability of MRIO-based results (Inomata & Owen 2014; Moran & Wood 2014; Rodrigues *et al.* 2018) indicate that normal consumption-based footprints converge fairly well. The largest variation in carbon footprints across MRIO databases actually comes from the absolute volume of emissions in each country. Excluding this factor, the footprints calculated across different MRIOs agree to approximately within  $\pm 10\%$ . In our experience, the largest source of uncertainty is likely to be the quantification of the biodiversity impact, rather than the pressure, trade or demand portion of the model. Measuring and quantifying biodiversity impact and change is extremely difficult, even for dedicated studies. This will be the overwhelmingly dominant source of uncertainty in the numerical results, yet no better information to quantify these characterization factors exists.

In future research on the present topic, the multi-impact assessment character as presented here must be preserved to avoid potential problem shifting. Additionally, as a next step of the parallel analysis of socio-economic variables undertaken in this study, a simultaneous disaggregation of income and urbanity


would deepen the understanding of a combination of drivers and enable adequate policy responses.

### 3. Results

#### 3.1. European trends

According to either impact methodology, the European biodiversity footprint associated with household final demand in 2010 was in total  $2.8E-01$  PDF or  $4.6E+04$  species.yr, and  $5.7E-10$  PDF or  $9.1E-05$  species.yr on the per capita level, respectively, from LC-IMPACT (PDF) and ReCiPe (species.yr).<sup>1</sup> While a decline of the European average by about 10% and the European total by about 4% can be observed between 2005 and 2010, some countries show increased footprints in one or multiple impact categories, both on national average and per capita (see Figure S5 as well as Tables S14–S19). Particularly countries with low to medium footprints such as Croatia, Lithuania or Poland express this behaviour, but also Italy as a high-impact country does. Normalizing these national footprints against the respective gross domestic product (GDP) reveals that between 2005 and 2010 only biodiversity impacts associated with Italian household consumption increased and that Bulgaria and Romania experienced the largest decreases (Figure S6).

<sup>1</sup>Note that these values only account for the impact categories that are covered by both LC-IMPACT and ReCiPe footprints, i.e. land occupation, water stress, greenhouse gas emissions, and terrestrial acidification. For values on all impact categories per footprint type, please see the SI dataset.



**Fig. 3.** LC-IMPACT biodiversity footprints disaggregated by degrees of urbanisation for 2010. The axes show the biodiversity footprints and balanced consumer expenditure (BCE) per household per country; circle sizes indicate the total balanced consumer expenditure (small – low, big – high); colouring denotes the total biodiversity footprint (blue – low, red – high). The dotted lines are linear trend lines. See also Figure S11 and Tables S24–S25.

In absolute terms, the countries responsible for the highest absolute biodiversity losses are by far Europe's large economies, i.e. Germany, Italy, Spain, France and the UK (Figure 1a). On a per capita basis, however, results are less unequivocal – which is also linked to the choice of impact methodology (Figures S7 and S8). More specifically, LC-IMPACT footprints exhibit a two-fold pattern: Mediterranean countries such as Greece and Spain, and those with high per-capita GDP such as Luxembourg have the highest per capita biodiversity impacts (Figure 1b). Following the ReCiPe-methodology, however, no clear pattern can be identified for the most impactful countries on the per-capita scale, except for the two top GDP-per-capita nations Norway and Luxembourg leading the list, while it is East-European countries that have the lowest impacts.

The major driver of biodiversity losses is found to be land use. On average across Europe and fairly stable over the years with not more than 4% deviation (the per-country fluctuations have a wider range), land use as a mechanism is responsible for 63–81% maximum (from ReCiPe and LC-IMPACT, respectively; cf. country levels in Tables S20–S21) of the total biodiversity impact footprint in 2010. The next largest mechanisms are global warming and terrestrial acidification, at roughly 20% and 13% for ReCiPe and 9% and 8% for LC-IMPACT, all in 2010.

### 3.2. Trade related biodiversity losses


European countries have a significant share of environmental pressures and impacts embodied in trade. For brevity, the following results only refer to the LC-IMPACT footprints. Only about 31% of species losses attributable to total European household consumption are sourced from inside Europe, and all countries are net importers of biodiversity footprints, i.e. the species losses associated with the import of goods to the respective country are greater than those linked to its exports (Table S22). Around 40% of the exports of European countries stay within European boundaries. With the exceptions of Bulgaria, Spain, Greece and

Croatia, all of which have a higher domestic impact, all European countries have more imported biodiversity losses than domestically sourced ones (Table S23). The import shares are highest for countries with high GDP per capita, namely Norway (95%), Belgium (97%), the Netherlands (98%) and Luxembourg (99%). For the remaining countries, import shares range typically from 50 to 70%. While most countries had even higher import shares in 2005, a few countries, most prominently eastern European ones, showed decreases in their domestic share from 2005 to 2010. The most noteworthy change during this period is, however, Italy's drop in import shares from 74 to 57%. A country's characteristic of being a net-importer of species losses is, with slight fluctuations, also reflected in the individual impact categories.

European countries clearly differ as to where the sources of their imported biodiversity losses lie. For instance, RoW Africa is the largest contributor to France's footprint with about 23%, whereas only 5% of species losses attributable to Bulgaria's household consumption are sourced from RoW Africa. Overall, the largest sources for biodiversity footprints embodied in international trade for European consumption are RoW Africa (12.1%), RoW America (14.2%) and Spain (12.7%), followed by Italy (8.8%), RoW Asia and Pacific (6.7%) and France (5.4%) (Figure 2). The largest absolute domestic flows of species losses are in Spain and Italy (Figures S9 and S10).

### 3.3. The effect of urbanization

Both LC-IMPACT and ReCiPe biodiversity footprints in cities are, in absolute terms, higher than in towns or rural areas. If not indicated differently, the following results refer to the LC-IMPACT footprints. On the European level, more than 50% of total ecosystem damage is caused by household consumption of urban populations. Towns are accountable for about 27%, and the remainder is due to final demand in rural areas. This same pattern can, however, be observed only across a few individual countries, mainly the ones with high absolute footprints, i.e. Italy, the UK and



**Fig. 4.** Differences across urbanization degrees and the importance of land use. a) shows the absolute biodiversity footprint per area type, indicating the share of land use; (b) shows land use pressure and impact disaggregated by land use type and the share of each degree of urbanization. For Ireland, Malta, the Netherlands and Sweden, no disaggregated 2005 data were available; therefore, 2010 data were used for these countries. Romania is not included due to a lack of data in both years.


Germany. While urban areas have the highest share in total biodiversity losses in most countries, e.g. Cyprus (60%), Finland (65%) or the UK (61%), rural areas are often the ones with the second highest contribution, e.g. in Austria (37%), France (37%) or Ireland (34%). But there are also countries where the highest biodiversity footprints are borne by people living in the countryside, for instance, in Hungary, Sweden and Slovakia with between 40 and 60%. The strongest signal of urban biodiversity footprints is in Malta with 93%. In Latvia, cities and rural areas are responsible for about 50% of the national biodiversity footprint each.

Both on the per capita as well as per household levels, differences are, however less distinct. A city resident in Europe is accountable for about 5% higher biodiversity losses than the average European citizen, whereas suburban residents have biodiversity footprints that are only 2% higher, and rural residents have footprints that are 12% lower than average.<sup>ii</sup> While this is also the case in most countries, some exceptions do exist: in smaller economies such as Bulgaria, Croatia, Hungary or Ireland, the per capita biodiversity footprint distribution shares of urban populations are relatively higher than the European city

<sup>ii</sup> For ReCiPe biodiversity footprints, the differences are +3% (city), -3% (town) and -3% (rural). This pattern can be found across most European countries when using the ReCiPe method.

distribution share. Conversely, rural populations in, amongst others, France and the UK have a stronger per capita impact on biodiversity compared with the average European city resident. While the biodiversity footprints of Norwegians living in any area type are highest in direct comparison across Europe according to the ReCiPe assessment, this applies to Luxembourg citizens in cities and towns as well as Greek rural residents when applying the LC-IMPACT methodology. For a comparison on the per household level, see Figure 3 as well as Figure S11. In either case, biodiversity losses disaggregated by degree of urbanization correspond to the balanced consumer expenditure, i.e. the higher the expenditure, the higher the footprint.

The relative breakdown of impact categories *per* degree of urbanization in a country differs only slightly, i.e. max. 3%, compared with national averages, both in absolute and per capita terms for all countries. That is, for instance, Germany's share of land use in all area types of around 78% in 2010 is about the same as in its national average. However, comparing the contributions per impact category *across* the degrees of urbanization shows that in most countries and impact categories city residents are more accountable than people living in towns or in the countryside. Scaling this to the European level, urban citizens carry higher weights in the categories of land occupation and water stress (up to 7% higher than the average per capita footprint per impact



**Fig. 5.** LC-IMPACT biodiversity footprint per sector for 2010. The left axis describes the absolute European footprint disaggregated by types of households. The sector grouping was established using a concordance matrix (EXIOBASE sector to sector group) from Ivanova *et al.* (2017). Colouring denotes the distinct types of households. The full sector group names can be found in Table S30.

category), whereas people living in towns have higher footprints than the average European citizen regarding ecosystem damage related to emissions of, for example, fossil methane and sulphur hexafluoride (Tables S26–S28).

Land use prevails as the highest contributor to the overall biodiversity footprint across cities, towns and rural areas. While the absolute land use for annual crops, intensive forestry and pastures is of similar magnitude, it is annual crops that have the greatest impact (Figures 4 and S12). The ratios of urbanization degrees within each land use type are comparable to the total.


Since land use is the most important mechanism driving biodiversity impact, land-intensive products are seen to be the most intensive product-level drivers (Figure 5). Food, services (including restaurants, and other services, which buy food as part of their business), clothing, and to a lesser extent, building materials, are the largest product drivers. We also observe impacts through water stress, chemical pollution (including from manufacturing and burning gasoline), and eutrophication impacts (including from the production of meat and leather), though these categories are minor (see spreadsheet appendix for full

detail). The relative contributions from the various product groups do not change significantly across household types (Figures 5 and S13–S15). In per capita terms, households consisting of three or more adults without dependents are the ones with the highest overall footprints. In absolute terms, the major contribution per sector comes from households with two adults (see also Table S29).

### 3.4. Income as a determinant for impact

While we find that both per capita and absolute biodiversity footprints increased with increasing income in 2010, no such development can be observed for 2005 (Figure 6 as well as Figure S16). More specifically, the absolute footprint still follows the former pattern, although less pronounced than in 2010, but per capita biodiversity impacts associated with final demand in low income quintiles are considerably higher in 2005. The per capita footprints attributable to Europe's second income quintile even exceed the footprint attributable to the fifth income quintile in 2005.





**Fig. 6.** LC-IMPACT biodiversity footprints disaggregated by income quintiles. a) shows 2005 footprints, (b) shows 2010 footprints. The primary axes describe per capita footprints, whereas the secondary axes scale absolute footprints (black line). Colouring denotes different impact categories. Note that the 2005 footprints do not include the contribution of Ireland and Sweden due to missing data; replacing these with 2010 data as in Figure 3 is not possible due to the nature of the data. No data on income quintiles in Norway were available in either year.

However, when normalizing the biodiversity footprints against the respective balanced consumer expenditure, the patterns for both years resemble one another (Figure S17). Across both reference years and all European countries, these normalized footprints decrease, the higher the income. Independent of the normalization procedure, land use accounts in both years and across all income quintiles for most of the biodiversity footprint, ranging from around 80% (62% for ReCiPe) for quintile five in 2010 to slightly above 84% (65% for ReCiPe) for the first quintile in 2005.

#### 4. Discussion and conclusion

A decrease in both absolute and per capita footprints in the period 2005–2010 was identified for the European total and most individual countries. Because the GDPs of most European countries increased at the same time, a decoupling of biodiversity impacts from affluence is indicated. With only two reference years that cover the financial crisis, more evidence is needed to state that this is a longer-term trend. Differences within each country exist across impact categories as well as within each impact category across countries.

The ranking of national biodiversity footprints depends largely on the chosen impact assessment methodology. That is, although ReCiPe and LC-IMPACT footprints show a similar overall pattern, they differ in detail across European countries, with location and per-capita wealth being the dominating factors for both

footprint types. Already Verones *et al.* (2017) showed the inter-linkage between the size of biodiversity footprints and income levels and that biodiversity losses are influenced by levels of endemism, species richness and their vulnerability in richer countries. Deviations between the footprint types can be attributed to their different nature and the underlying methodology. A brief discussion on fundamental implications regarding the choice of impact methodology is provided in SI11.

While urbanization has a major influence on the absolute biodiversity footprints of countries, it is less pronounced on the per capita and per household levels. The high share of species losses attributable to city residents on a national level is mainly due to the size of urban populations in Europe. In times of urban sprawl, related social and demographic changes, as well as society's high impact on the planet, sustainable urban development and regional planning become more important than ever before. Moreover, it can be observed that, with few exceptions, absolute biodiversity footprints in each degree of urbanization follow the magnitude of the total GDP and the population size across all countries and all impact categories. The variation of national per capita footprints across all area types and all impact categories, however, has a less clear signal. Nonetheless, we find that both absolute and per household footprints are correlated to GDP and balanced consumer expenditure across all degrees of urbanization.

In relation to that, it was shown that income is a major driver of biodiversity losses due to household final demand on absolute

national averages and for the whole of Europe. That is, the higher the income, the higher the footprint. This is in alignment with studies explaining the magnitude of environmental pressures with both expenditure and income (Jones & Kammen 2014; Chancel & Piketty 2015; Ivanova *et al.* 2016, 2017; Steen-Olsen *et al.* 2016). While that holds also for the 2010 per capita level, per capita footprints in 2005 appear to be decoupled from income. Such variation can, however, be explained by expenditure patterns. That is, per capita footprints in proportion to per capita expenditure decrease from low to high income for both reference years. While the raw results of the income-footprint nexus extend the finding of non-saturation regarding environmental pressures with increasing wealth by Hertwich & Peters (2009) and others, the normalized results rather corroborate the controversial hypothesis of the environmental Kuznets curve (Stern 2004). A definitive, generalized answer on the role of income across both, absolute and per capita, levels is therefore not possible, but differentiation is necessary.

In line with other biodiversity footprint studies (Verones *et al.* 2017; Wilting *et al.* 2017), land use was found to be the major contributor to biodiversity losses. Land use, in turn, was shown to be mainly driven by the demand for agricultural food products, which is in accordance with Kitzes *et al.* (2017) and Wilting *et al.* (2017). The role of food commodities was also recently highlighted by Castellani *et al.* (2019), Crenna *et al.* (2019) and Marques *et al.* (2019). The impacts of this demand and other European consumption are, however, to a large extent imposed on countries in other regions of the world, whereas domestically caused species losses are considerably lower. Additionally, it must be noted that climate change, on both a local and global level, could take a larger toll on biodiversity in the future given the dramatically increasing concentrations of greenhouse gases in the atmosphere. Moreover, we observe that Europe is a net-importer of biodiversity losses, a result that aligns with earlier studies on both environmental pressures and impacts (Lenzen *et al.* 2012; Giljum *et al.* 2016; Lutter *et al.* 2016; Kitzes *et al.* 2017; Wilting *et al.* 2017; Wood *et al.* 2018). Such an imbalance may be a source for ethical demer and raises the question of producer vs consumer responsibility (Lenzen *et al.* 2007) – an answer to which shall not be attempted here.

Additionally, the impact distribution is sector dependent. Therefore, directed intervention via policy instruments such as taxes on certain goods for curbing further ecosystem damage associated with household consumption is not straightforward. With the highest biodiversity losses embodied in agricultural products, the discourse on the role of food and non-food commodities must be widened to reach multiple stakeholders including the public, the scientific community and policymakers. Differences regarding the environmental performance of producers of agricultural products as well as the effectiveness of environmental policies and regulations could not be analysed here but should be addressed in future research. Additionally, our model could be combined with the approach by Froemelt *et al.* (2018) for gaining more insights into consumption patterns of individual countries.

Given that urbanization is expected to increase and that more countries strive for ever more wealth, stronger impacts on the environment would be the consequence, both domestically and, even more so, abroad. Therefore, political action is crucial – yet, not only that: the responsibility of the individual is also asked for to avert further negative environmental corollaries of household consumption. Be it environmental laws and regulations, or just the decision of the individual to abstain from, for instance,

meat consumption and do without the extra cup of coffee in the morning, both behavioural and structural changes could reduce humanity's footprint on the planet.

**Supplementary material.** To view supplementary material for this article, please visit <https://doi.org/10.1017/sus.2019.23>

**Acknowledgements.** The authors would like to thank the editor Xuemei Bai and two anonymous reviewers for their constructive suggestions improving the quality of the manuscript.

**Author contributions.** MK led the work. RW and DM designed the research and assisted with the paper and implementation. AT assisted with the consumer expenditure data. FV contributed the LCIA models and biodiversity metric expertise.

**Financial support.** This work was supported by the Norwegian Research Council (grant # 255483/E50).

**Conflict of interest.** The authors declare they have no conflict of interest.

## References

- Bee, A., Meyer, B., & Sullivan, J. (2012). *The Validity of Consumption Data: Are the Consumer Expenditure Interview and Diary Surveys Informative?* Cambridge, MA: National Bureau of Economic Research. Retrieved from <https://doi.org/10.3386/w18308>.
- Castellani, V., Beylot, A., & Sala, S. (2019). Environmental impacts of household consumption in Europe: comparing process-based LCA and environmentally extended input-output analysis. *Journal of Cleaner Production*, 117966. Retrieved from <https://doi.org/10.1016/j.jclepro.2019.117966>
- Chancel, L. & Piketty, T. (2015). *Carbon and Inequality: From Kyoto to Paris*. Paris School of Economics. Retrieved from <http://piketty.pse.ens.fr/files/ChancelPiketty2015.pdf>.
- Chaudhary, A. & Kastner, T. (2016). Land use biodiversity impacts embodied in international food trade. *Global Environmental Change*, 38, 195–204.
- Chaudhary, A., Verones, F., Baan, L. de, & Hellweg, S. (2015). Quantifying land use impacts on biodiversity: combining species-area models and vulnerability indicators. *Environmental Science & Technology*, 49(16), 9987–9995.
- Crenna, E., Sinkko, T., & Sala, S. (2019). Biodiversity impacts due to food consumption in Europe. *Journal of Cleaner Production*, 227, 378–391.
- Curran, M., Baan, L. de, Schryver, A. M. de, van Zelm, R., Hellweg, S., Koellner, T., ... Huijbregts, M. A. J. (2011). Toward meaningful end points of biodiversity in life cycle assessment. *Environmental Science & Technology*, 45(1), 70–79.
- Eurostat (2015). *Household Budget Survey: 2010 Wave*. Retrieved from <http://ec.europa.eu/eurostat/web/household-budget-surveys/publication>.
- Eurostat (2018a). Consumption expenditure of private households (hbs) – Reference Metadata in Euro SDMX Metadata Structure (ESMS). Retrieved from [http://ec.europa.eu/eurostat/cache/metadata/en/hbs\\_esms.htm](http://ec.europa.eu/eurostat/cache/metadata/en/hbs_esms.htm).
- Eurostat (2018b). Eurostat Database: European Commission. Retrieved from <http://ec.europa.eu/eurostat/data/database>.
- Eurostat (2019). Degree of Urbanisation Classification – 2011 Revision. Retrieved from [https://ec.europa.eu/eurostat/statistics-explained/index.php/Degree\\_of\\_urbanisation\\_classification\\_-\\_2011\\_revision#Degree\\_of\\_urbanisation\\_classification](https://ec.europa.eu/eurostat/statistics-explained/index.php/Degree_of_urbanisation_classification_-_2011_revision#Degree_of_urbanisation_classification).
- Ewing, B. R., Hawkins, T. R., Wiedmann, T. O., Galli, A., Ertug Erinc, A., Weinzettel, J., & Steen-Olsen, K. (2012). Integrating ecological and water footprint accounting in a multi-regional input-output framework. *Ecological Indicators*, 23, 1–8.
- Froemelt, A., Dürrenmatt, D. J., & Hellweg, S. (2018). Using data mining to assess environmental impacts of household consumption behaviors. *Environmental Science & Technology*, 52(15), 8467–8478.
- Giljum, S., Wieland, H., Lutter, S., Bruckner, M., Wood, R., Tukker, A., & Stadler, K. (2016). Identifying priority areas for European resource policies: a MRIO-based material footprint assessment. *Journal of Economic Structures*, 5(1), 99.

- Godar, J., Persson, U. M., Tizado, E. J., & Meyfroidt, P. (2015). Towards more accurate and policy relevant footprint analyses: tracing fine-scale socio-environmental impacts of production to consumption. *Ecological Economics*, 112, 25–35.
- Hertwich, E. G. & Peters, G. P. (2009). Carbon footprint of nations: a global, trade-linked analysis. *Environmental Science & Technology*, 43(16), 6414–6420.
- Huijbregts, M. A. J., Steinmann, Z. J. N., Elshout, P. M. F., Stam, G., Verones, F., Vieira, M. D. M., & van Zelm, R. (2016). *ReCiPe 2016: A Harmonized Life Cycle Impact Assessment Method at Midpoint and Endpoint Level. Report I: Characterization*. Bilthoven: National Institute for Public Health and the Environment. Retrieved from [www.rivm.nl/en/Topics/L/Life\\_Cycle\\_Assessment\\_LCA/Download](http://www.rivm.nl/en/Topics/L/Life_Cycle_Assessment_LCA/Download).
- Huijbregts, M. A. J., Steinmann, Z. J. N., Elshout, P. M. F., Stam, G., Verones, F., Vieira, M. D. M., ... van Zelm, R. (2017). ReCiPe2016: a harmonised life cycle impact assessment method at midpoint and endpoint level. *International Journal of Life Cycle Assessment*, 22(2), 138–147.
- Inomata, S. & Owen, A. (2014). Comparative evaluation of MRIO databases. *Economic Systems Research*, 26(3), 239–244.
- IUCN (2019). *The IUCN Red List of Threatened Species*. Retrieved from <https://www.iucnredlist.org/>.
- Ivanova, D., Stadler, K., Steen-Olsen, K., Wood, R., Vita, G., Tukker, A., & Hertwich, E. G. (2016). Environmental impact assessment of household consumption. *Journal of Industrial Ecology*, 20(3), 526–536.
- Ivanova, D., Vita, G., Steen-Olsen, K., Stadler, K., Melo, P. C., Wood, R., & Hertwich, E. G. (2017). Mapping the carbon footprint of EU regions. *Environmental Research Letters*, 12(5), 54013.
- Jones, C. & Kammen, D. M. (2014). Spatial distribution of U.S. household carbon footprints reveals suburbanization undermines greenhouse gas benefits of urban population density. *Environmental Science & Technology*, 48(2), 895–902.
- Kanemoto, K., Moran, D., Lenzen, M., & Geschke, A. (2014). International trade undermines national emission reduction targets: new evidence from air pollution. *Global Environmental Change*, 24, 52–59.
- Kanemoto, K., Moran, D., & Hertwich, E. G. (2016). Mapping the carbon footprint of nations. *Environmental Science & Technology*, 50(19), 10512–10517.
- Kitzes, J. (2013). An introduction to environmentally-extended input-output analysis. *Resources*, 2(4), 489–503.
- Kitzes, J., Berlow, E., Conlisk, E., Erb, K.-H., Iha, K., Martinez, N., ... Harte, J. (2017). Consumption-based conservation targeting: linking biodiversity loss to upstream demand through a global wildlife footprint. *Conservation Letters*, 10(5), 531–538.
- Lenzen, M., Murray, J., Sack, F., & Wiedmann, T. O. (2007). Shared producer and consumer responsibility – theory and practice. *Ecological Economics*, 61(1), 27–42.
- Lenzen, M., Moran, D., Kanemoto, K., Foran, B., Lobefaro, L., & Geschke, A. (2012). International trade drives biodiversity threats in developing nations. *Nature*, 486(7401), 109–112.
- Lenzen, M., Moran, D., Bhaduri, A., Kanemoto, K., Bekchanov, M., Geschke, A., & Foran, B. (2013). International trade of scarce water. *Ecological Economics*, 94, 78–85.
- Leontief, W. W. (1936). Quantitative input and output relations in the economic systems of the United States. *Review of Economics and Statistics*, 18(3), 105.
- Leontief, W. W. (1970). Environmental repercussions and the economic structure: an input-output approach. *Review of Economics and Statistics*, 52(3), 262.
- Lutter, S., Pfister, S., Giljum, S., Wieland, H., & Mutel, C. L. (2016). Spatially explicit assessment of water embodied in European trade: a product-level multi-regional input-output analysis. *Global Environmental Change*, 38, 171–182.
- Majeau-Bettez, G., Strømman, A. H., & Hertwich, E. G. (2011). Evaluation of process- and input-output-based life cycle inventory data with regard to truncation and aggregation issues. *Environmental Science & Technology*, 45(23), 10170–10177.
- Marques, A., Martins, I. S., Kastner, T., Plutzer, C., Theurl, M. C., Eisenmenger, N., ... Pereira, H. M. (2019). Increasing impacts of land use on biodiversity and carbon sequestration driven by population and economic growth. *Nature Ecology & Evolution*, 3(4), 628–637.
- Mekonnen, M. & Hoekstra, A. (2011). *National Water Footprint Accounts: The Green, Blue and Grey Water Footprint of Production and Consumption. Value of Water Research: Vol. 50*. UNESCO-IHE Institute for Water Education.
- Millennium Ecosystem Assessment (2005). *Ecosystems and Human Well-being: Biodiversity Synthesis*. Island Press.
- Miller, R. E. & Blair, P. D. (2009). *Input-output Analysis: Foundations and Extensions* (Second edition). Cambridge University Press.
- Moran, D. & Kanemoto, K. (2017). Identifying species threat hotspots from global supply chains. *Nature Ecology & Evolution*, 1(1), 109.
- Moran, D. & Wood, R. (2014). Convergence between the Eora, WIOD, EXIOBASE, and Open EU's consumption-based carbon accounts. *Economic Systems Research*, 26(3), 245–261.
- Moran, D., Kanemoto, K., Jiborn, M., Wood, R., Többen, J., & Seto, K. C. (2018). Carbon footprints of 13 000 cities. *Environmental Research Letters*, 13(6), 64041.
- Rodrigues, J. F. D., Moran, D., Wood, R., & Behrens, P. (2018). Uncertainty of consumption-based carbon accounts. *Environmental Science & Technology*, 52(13), 7577–7586.
- Stadler, K., Wood, R., Bulavskaya, T., Södersten, C.-J., Simas, M., Schmidt, S., ... Tukker, A. (2018). EXIOBASE 3: developing a time series of detailed environmentally extended multi-regional input-output tables. *Journal of Industrial Ecology*, 22(3), 502–515.
- Steen-Olsen, K., Wood, R., & Hertwich, E. G. (2016). The carbon footprint of Norwegian household consumption 1999–2012. *Journal of Industrial Ecology*, 20(3), 582–592.
- Stern, D. I. (2004). The rise and fall of the environmental Kuznets curve. *World Development*, 32(8), 1419–1439.
- Tukker, A., Koning, A. de, Wood, R., Hawkins, T. R., Lutter, S., Acosta-Fernández, J., ... Kuenen, J. (2013). EXIOPOL – development and illustrative analyses of a detailed global MR EE SUT/IOT. *Economic Systems Research*, 25(1), 50–70.
- Tukker, A., Bulavskaya, T., Giljum, S., Koning, A. de, Lutter, S., Simas, M., ... Wood, R. (2016). Environmental and resource footprints in a global context: Europe's structural deficit in resource endowments. *Global Environmental Change*, 40, 171–181.
- United Nations Statistics Division (2018). Classification of Individual Consumption According to Purpose. Retrieved from <https://unstats.un.org/unsd/publications/catalogue?selectID=145>.
- Verones, F., Saner, D., Pfister, S., Baisero, D., Rondinini, C., & Hellweg, S. (2013). Effects of consumptive water use on biodiversity in wetlands of international importance. *Environmental Science & Technology*, 47(21), 12248–12257.
- Verones, F., Moran, D., Stadler, K., Kanemoto, K., & Wood, R. (2017). Resource footprints and their ecosystem consequences. *Scientific Reports*, 7(1), 171.
- Verones, F., Huijbregts, M. A. J., Azevedo, L. B., Chaudhary, A., Cosme, N., Baan, L. de, ... Hellweg, S. (2019). LC-Impact Version 1.0 – a spatially differentiated life cycle impact assessment approach. Retrieved from <http://lc-impact.eu>.
- Wiebe, K. & Yamano, N. (2016). *Estimating CO2 Emissions Embodied in Final Demand and Trade Using the OECD ICIO 2015: Methodology and Results. OECD Science, Technology and Industry Working Papers: 2016/05*. Paris: OECD Publishing.
- Wiedmann, T. O. & Barrett, J. (2013). Policy-relevant applications of environmentally extended MRIO databases – experiences from the UK. *Economic Systems Research*, 25(1), 143–156.
- Wiedmann, T. O. & Lenzen, M. (2018). Environmental and social footprints of international trade. *Nature Geoscience*, 11(5), 314–321.
- Wiedmann, T. O., Schandl, H., Lenzen, M., Moran, D., Suh, S., West, J., & Kanemoto, K. (2015). The material footprint of nations. *Proceedings of the National Academy of Sciences USA*, 112(20), 6271–6276.
- Wiedmann, T. O., Chen, G., & Barrett, J. (2016). The concept of city carbon maps: a case study of Melbourne, Australia. *Journal of Industrial Ecology*, 20(4), 676–691.
- Wiling, H. C., Schipper, A. M., Bakkenes, M., Meijer, J. R., & Huijbregts, M. A. J. (2017). Quantifying biodiversity losses due to human consumption: a global-scale footprint analysis. *Environmental Science & Technology*, 51(6), 3298–3306.
- Wood, R., Stadler, K., Bulavskaya, T., Lutter, S., Giljum, S., Koning, A. de, ... Tukker, A. (2015). Global sustainability accounting – developing

- EXIOBASE for multi-regional footprint analysis. *Sustainability*, 7(1), 138–163.
- Wood, R., Stadler, K., Simas, M., Bulavskaya, T., Giljum, S., Lutter, S., & Tukker, A. (2018). Growth in environmental footprints and environmental impacts embodied in trade: resource efficiency indicators from EXIOBASE3. *Journal of Industrial Ecology*, 22(3), 553–564.
- WWF (2016). *Living Planet Report 2016: Risk and Resilience in a New Era*. World Wildlife Fund For Nature.