

ISUOG Safety Committee updated recommendation on use of respirators by practitioners undertaking obstetric and gynecological ultrasound in context of SARS-CoV-2 Omicron variant of concern

K. SALVESEN, G. TER HAAR, P. MILORO, E. SINKOVSKAYA, C. LEES, T. BOURNE, K. MARŠÁL and A. DALL'ASTA, on behalf of the Bioeffects and Safety Committee and the Board of the International Society of Ultrasound in Obstetrics and Gynecology (ISUOG)

In view of the rapid spread of the Omicron variant of concern (VOC) of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)¹, the International Society of Ultrasound in Obstetrics and Gynecology (ISUOG) has decided to revise its position statements on use of personal protective equipment for the safe performance of obstetric and gynecological ultrasound^{2,3}, particularly in relation to the protection of healthcare workers.

The World Health Organization (WHO) recommendation on mask use by healthcare workers, issued on 22 December 2021, states that ‘... the Omicron variant is spreading significantly faster than the Delta variant in countries with documented community transmission, with a doubling time between 1.5–3 days. Preliminary, unpublished data show a reduction in neutralizing titres against Omicron suggesting a level of immune evasion and an important reduction in vaccine effectiveness against infection and symptomatic disease for Omicron compared to Delta’⁴.

ISUOG previously recommended the use of fit-approved masks or respirators (FFP2, FFP3, NIOSH-approved N95 or other equivalent or higher-level certified respirator) for the ultrasound examination of patients with suspected or confirmed SARS-CoV-2 infection³ or in the event of widespread community transmission². The latest WHO recommendation on mask use by healthcare workers¹ states that ‘A respirator... should be worn by health workers along with other personal protective equipment (PPE) – a gown, gloves and eye protection – before entering a room where there is a patient with suspected or confirmed COVID-19... in care settings where ventilation is known to be poor or cannot be assessed or the ventilation system is not properly maintained...’. This applies to the rooms in which ultrasound scans are typically performed.

In light of the growing number of SARS-CoV-2 infections and in particular the highly infectious Omicron VOC, the USA Centers for Disease Control and Prevention has predicted and provided strategies to limit the effects of healthcare staff shortages⁵. Patients do not undergo routinely polymerase chain reaction or lateral flow testing prior to ultrasound assessment for obstetric or gynecological indications, nonetheless, a

high number of asymptomatic cases are to be expected in the context of the SARS-CoV-2 epidemic caused by the Omicron VOC⁶. On this basis, ISUOG now recommends that fit-approved masks or respirators should be worn routinely in ultrasound clinics and departments to limit the risk of infection for practitioners performing obstetric and gynecological ultrasound scans, in addition to preventive measures including hand hygiene, patient mask-wearing^{2–4} and vaccination^{7,8}. The recommendation can be adapted locally based on community prevalence and transmissibility of the current and future variants of SARS-CoV-2.

ACKNOWLEDGMENT

Open Access Funding provided by Università degli Studi di Parma within the CRUI-CARE Agreement. [Correction added on May 20, 2022, after first online publication: CRUI funding statement has been added.]

References

1. Enhancing Readiness for Omicron (B.1.1.529): Technical Brief and Priority Actions for Member States. Geneva: World Health Organization, 17 December 2021. [https://www.who.int/publications/m/item/enhancing-readiness-for-omicron-\(b-1.1.529\)-technical-brief-and-priority-actions-for-member-states](https://www.who.int/publications/m/item/enhancing-readiness-for-omicron-(b-1.1.529)-technical-brief-and-priority-actions-for-member-states)
2. Abramowicz JS, Basseal JM, Brezinka C, Dall'Asta A, Deng J, Harrison G, Lee JCS, Lim A, Maršal K, Miloro P, Poon LC, Salvesen KÅ, Sande R, Ter Haar G, Westerway SC, Xie MX, Lees C. ISUOG Safety Committee Position Statement on use of personal protective equipment and hazard mitigation in relation to SARS-CoV-2 for practitioners undertaking obstetric and gynecological ultrasound. *Ultrasound Obstet Gynecol* 2020; 55: 886–891.
3. Poon LC, Abramowicz JS, Dall'Asta A, Sande R, Ter Haar G, Maršal K, Brezinka C, Miloro P, Basseal J, Westerway SC, Abu-Rustum RS, Lees C. ISUOG Safety Committee Position Statement on safe performance of obstetric and gynecological scans and equipment cleaning in context of COVID-19. *Ultrasound Obstet Gynecol* 2020; 55: 709–712.
4. WHO recommendations on mask use by health workers, in light of the Omicron variant of concern. World Health Organization, 22 December 2021. https://www.who.int/publications/i/item/WHO-2019-nCoV-IPC_Masks-Health_Workers-Omicron_variant-2021.1
5. CDC Releases Emergency Guidance for Healthcare Facilities to Prepare for Potential Omicron Surge. 23 December 2021. <https://www.cdc.gov/media/releases/2021/s1223-emergency-guidance-prepare-for-omicron.html>
6. Sutton D, Fuchs K, D'Alton M, Goffman D. Universal Screening for SARS-CoV-2 in Women Admitted for Delivery. *N Engl J Med* 2020; 382: 2163–2164.
7. Pilishvili T, Gierke R, Fleming-Dutra KE, Farrar JL, Mohr NM, Talan DA, Krishnadasan A, Harland KK, Smithline HA, Hou PC, et al. Effectiveness of mRNA Covid-19 Vaccine among U.S. Health Care Personnel. *N Engl J Med* 2021; 385: e90.
8. Bar-On YM, Goldberg Y, Mandel M, Bodenheimer O, Freedman L, Alroy-Preis S, Ash N, Huppert A, Milo R. Protection against Covid-19 by BNT162b2 Booster across Age Groups. *N Engl J Med* 2021; 385: 2421–2430.