

# SCIENTIFIC REPORTS



OPEN

## A CRISPR/Cas9 system adapted for gene editing in marine algae

Marianne Nymark, Amit Kumar Sharma, Torfinn Sparstad, Atle M. Bones & Per Winge

Received: 08 February 2016

Accepted: 07 April 2016

Published: 25 April 2016

Here we report that the CRISPR/Cas9 technology can be used to efficiently generate stable targeted gene mutations in microalgae, using the marine diatom *Phaeodactylum tricorutum* as a model species. Our vector design opens for rapid and easy adaption of the construct to the target chosen. To screen for CRISPR/Cas9 mutants we employed high resolution melting based PCR assays, mutants were confirmed by sequencing and further validated by functional analyses.


Marine microalgae are responsible for 40% of global primary production and capture more CO<sub>2</sub> than all the rain forests combined<sup>1</sup>. Diatoms (Bacillariophyceae) are the most important group of eukaryotic microalgae, and have enormous ecological significance<sup>2–4</sup>. This group of microalgae can also be used as a source of lipids for biofuels, as well as other applications in bio-, nano- and environmental technology<sup>5,6</sup>. Studies of microalgae have been hampered by inefficient genome editing methods restricting functional analyses. An improved system for generation of mutants is important for 1) functional analyses of genes/proteins to improve our understanding of diatom biology and 2) for strain improvement for biotechnological applications. In previous years it has only been possible to create non-targeted knock-down (RNAi) or overexpression lines in diatoms, but recently two independent groups demonstrated for the first time precise targeted genome editing of the *Phaeodactylum tricorutum* genome<sup>7,8</sup>. The modifications of the diatom genome were achieved using protein-based systems involving meganucleases and transcription activator-like effector nucleases (TALENs) that have specific DNA binding activities. These systems are labour intensive and costly, so we set out to establish the much simpler and inexpensive CRISPR/Cas9 technology for genetic engineering of diatom genomes<sup>9</sup>.

We developed a highly efficient CRISPR/Cas9 based system optimized for creating stable targeted gene knock-outs in the marine diatom *P. tricorutum*. The two components necessary for this system to function, the Cas9 nuclease and a guide RNA directing the nuclease to a specific DNA sequence, were expressed from the same vector. The diatom codon-optimized Cas9 was driven by a promoter for a fucoxanthin chlorophyll a/c-binding protein gene (*LHCF2*), and the single guide RNA (sgRNA) was under control of the *P. tricorutum* U6 promoter. The Cas9 and the sgRNA modules were cloned into the same vector (pKS diaCas9\_sgRNA; Supplementary Fig. 1) to increase the probability of co-delivery during biolistic transformation of the *P. tricorutum* cells. By exploiting two BsaI cutting sites located within the sgRNA cassette (Supplementary Fig. 1), a small adapter can easily be inserted into the sgRNA, making it target-specific to the gene that it is desired to disrupt.


As a proof of concept we targeted the chloroplast signal recognition particle 54 (CpSRP54; Draft ID Phatr2\_35185), a member of the chloroplast signal recognition particle (CpSRP) pathway, for CRISPR/Cas9-based disruption. By using a combination of high resolution melting (HRM) analyses and Sanger sequencing we detected a variety of indels at the target site (located in the conserved SRP54 N-terminal domain (Fig. 1A)) in 8 out of 26 transformants (Fig. 1B,C), giving a mutation frequency of 31%. In addition to the CRISPR/Cas9-mediated mutagenesis of the *CpSRP54* gene, we have successfully knocked out two other genes, using two different sgRNAs (targets) per gene. The mutation frequencies in these experiments were also high, ranging between 25–63% (unpublished data). All transformants cannot be expected to contain mutations because: 1) The non-selectable pKS diaCas9\_sgRNA plasmid is co-transformed with the pAF6 plasmid to confer resistance to zeocin<sup>10</sup>. According to literature 30–40% of the transformants are expected to contain only the pAF6 plasmid<sup>10</sup>, and no targeted gene mutations will occur. 2) Biolistic-mediated transformation is known to cause fragmentation and incomplete incorporation of the Cas9 gene<sup>11</sup>. Intact Cas9 and sgRNA is a necessity for CRISPR mutagenesis to take place. In plants, the incomplete incorporation of Cas9 has been suggested as a likely explanation for the much lower mutation efficiencies (5–12%) reported when using biolistic bombardment for introduction of the Cas9 and sgRNA<sup>11–13</sup>, compared to *Agrobacterium*-mediated transformation (20–90%)<sup>14–17</sup>.

The detected indel mutations in the *CpSRP54* gene included single nucleotide (nt) insertions, a variety of mostly short deletions and in one case a 212 bp insertion (Fig. 1C). As reported for large insertions generated

Department of Biology, Norwegian University of Science and Technology, N-7491 Trondheim, Norway. Correspondence and requests for materials should be addressed to P.W. (email: per.winge@bio.ntnu.no)



**Figure 1. Genome editing and screening of *CpSRP54* gene mutants.** (A) Schematic presentation of the *CpSRP54* protein. The area of the protein corresponding to the target region for CRISPR/Cas9 based gene editing is located within the conserved SRP54\_N domain (red highlighting). CLS: Chloroplast localization signal; SRP54\_N: SRP54 N-terminal helical bundle domain; SRP: SRP GTPase containing domain; SRP\_SPB: SRP54 signal peptide binding domain (B) Left side: Sequence chromatograms produced by direct sequencing of PCR product from WT cells and colonies containing cells with targeted mutations. Chromatograms produced by Chromas Lite version 2.0 software. Right side: High resolution melting analyses (HRM) data presented as normalized melting peaks generated from the corresponding PCR product. WT profiles (purple lines) are included in each plot to visualize the differences in melting behaviour between WT amplification products and products containing indels. (C) Overview of indels generated by transformation of *P. tricornutum* with the pKS diaCas9\_sgrNA construct targeting the *CpSRP54* gene. Sequences represent cloned PCR products from colonies exhibiting targeted mutations. The number of times identical sequences were detected within the same colony is indicated in brackets to the right. Colour coding: Blue: WT target sequence; Orange: PAM sequence; Red dashes: Deleted bases; Red letters: Inserted bases.



**Figure 2. Photosynthetic efficiency after high intensity blue light exposure and recovery in dim white light.** The photosynthetic efficiency ( $F_v/F_m$ ) was determined in wild type cultures (WT;  $n = 6$ ) and cultures containing CpSRP54 CRISPR/Cas9 mutants (M1, M4, M5 and M8;  $n = 3$ ) after a 3 min dark acclimation period. Prior to the measurements the cells were exposed to blue high light (BHL;  $500 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) for 5 min, 0.5 h and 1 h with a following recovery in dim white light (WL;  $65 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) for 0.5 h, 1 h and 3 h. Values are presented with  $\pm$ SD bars.

by repair of TALEN induced double strand breaks<sup>7</sup>, the 212 bp insertion found here also consisted of fragments of the vectors used for transformation. HRM and sequencing data indicate that in colonies where indels are detected, cells with mutations completely dominate over cells with wild type (WT) alleles (Fig. 1B,C). Amplified genomic DNA from colonies with targeted mutations was TOPO-cloned and 7–8 clones from each colony were sequenced. Only three out of a total of 62 clones contained the WT allele.

Chromatograms produced from direct sequencing of PCR product confirming insertions in colony M1, M2 and M8 showed clear peaks with low background levels also after the site of the insertion (Fig. 1B). The 21 bp deletion detected in CpSRP54 from colony M5 could be identified by direct sequencing of the PCR product (Fig. 1B). The data imply that the vast majority of the cells in these colonies are mutants having identical biallelic mutations, suggesting a double-strand break (DSB) induced gene conversion mechanism<sup>18</sup>. Sequencing of the cloned PCR products from M1, M2, M5 and M8 detected only one type of mutation in each colony, supporting this assumption (Fig. 1C). The results for the M1, M2, M5 and M8 mutants suggest that whereas the DSB in one allele is repaired by the error-prone Non-homologous end joining (NHEJ) pathway, the DSB in the other allele might be repaired by homologous recombination (HR) using the newly mutated allele as template for repair. The limited number of different indels in cells derived from individual transformants suggests that the DSBs and genome editing start already within the initially transformed cell. When several different mutated alleles are found within a colony (Fig. 1C), implying that a few cell divisions have occurred before the first mutagenic event, the high efficiency of the CRISPR/Cas9 system still causes an almost complete elimination of the WT sequence. In contrast, the percentage of mutant *P. tricornutum* cells within colonies shown to have targeted gene modifications induced by meganucleases was only 1–15%<sup>7</sup>. For TALEN the equivalent number was 6–80%<sup>7</sup>.

Proteins of the CpSRP pathway are involved in assembly and integration of proteins into the thylakoid membrane<sup>19</sup>. To the best of our knowledge, no CpSRP54 knockout mutant has previously been reported in microalgae, but based on information from higher plants we expected such a mutant to have an easily recognizable phenotype connected to alterations of the photosynthetic apparatus<sup>20,21</sup>. Sensitivity to high light has also been reported in CpSRP54 mutants in higher plants<sup>21</sup>. Since high light sensitivity would be an easy way to show a functional effect of the disruption of the CpSRP54 gene in *P. tricornutum*, we decided to provoke light-induced damage to photosystem II (PSII) by exposing mutants to high intensity blue light (BHL). The photosynthetic efficiency (measured as the maximum quantum yield,  $F_v/F_m$ ) was monitored during the period of BHL stress, and through the following recovery period in dim white light (WL) conditions (Fig. 2). Exposure to BHL for 1 h caused a greater decrease in  $F_v/F_m$  in the CpSRP54 mutant cultures than in WT cultures. Both mutants and WT cells showed an almost complete recovery after 3 h of dim WL conditions (Fig. 2). The increased sensitivity to high intensity light in mutant cultures is in support of the HRM and sequencing data indicating that cells with mutations completely dominate over wild type within the culture.

“Clean” CpSRP54 mutant lines originating from single cells with biallelic mutations have also been isolated by spreading out highly diluted cell suspensions from the original cultures on agar plates, and repeating the screening process on the resulting colonies. These cultures will be used for future characterization of the CpSRP54 mutants. Gene expression analyses detected high levels of sgRNA in all of the clean CpSRP54 mutant lines, and similar high levels of Cas9 mRNA in all but one of the same cultures (Supplementary Table 1). The transcript levels of Cas9 and sgRNA were within the same range as the endogenous Light Harvesting Complex gene *LHCX1* (XM\_002179724) (Supplementary Table 1), known to be one of the most abundant mRNAs in *P. tricornutum*<sup>22</sup>. The transcriptional analyses confirm high gene expression and stable integration of the Cas9 and sgRNA genes into the *P. tricornutum* genome. Constitutively expressed Cas9 protein has been reported to be toxic for *Chlamydomonas reinhardtii*<sup>23</sup>. Clean *P. tricornutum* CpSRP54 mutant lines grow at a similar or slightly slower rate (Div. per day:  $\sim 1.5$ – $2.0 \text{ day}^{-1}$ ; Supplementary Table 2) than WT ( $\sim 2.0 \text{ day}^{-1}$ ) in dim WL conditions.

Transcriptional analyses of the clean mutant culture M8.1, originating from M8, showed that the sgRNA, but not the *Cas9* gene is expressed in this culture. Still, the M8.1 culture has the lowest growth rate of all the mutant lines ( $\sim 1.5 \text{ day}^{-1}$ ). The above described results indicate that the observed growth differences are not likely to be connected to a toxic effect of constitutive *Cas9* expression.

Diatoms are important as the major group of primary producers of the oceans and as a promising and basically unexploited group of eukaryotic organisms for bio-production and biotechnological exploitation. We have developed a very efficient genome editing and selection method for the diatom *P. tricornutum*, which can be potentially adapted for use in other microalgae. We anticipate our method to be useful for all kinds of functional studies in microalgae, and also for engineering of specific traits like modified light-harvesting antennas and lipid profile.

## Methods

**Growth conditions.** Axenic *P. tricornutum* cells (CCMP2561) were grown as described previously<sup>24</sup> in a controlled *in vitro* growth room under a 16 h photoperiod at 22 °C, 65  $\mu\text{mol photon m}^{-2} \text{ s}^{-1}$ .

**Vector construction.** A codon-optimized *Cas9* protein (dia*Cas9*) modelled after hSp*Cas9*<sup>25</sup>, containing N- and C-terminal nuclear localization signals (NLS) with 3  $\times$  FLAG repeats at the N-terminal part, was designed using a codon usage table derived from available gene models of *P. tricornutum* in GenBank, NCBI. The dia*Cas9* gene was placed under the control of the *P. tricornutum* *LHCF2* promoter and has a *LHCF1* terminator sequence (Z24761.1). The whole module, promoter, dia*Cas9* and terminator, was synthesized by GeneArt<sup>®</sup> Services Thermo Fisher Scientific Inc. and SacI and PstI sites were included at 5' and 3' end of the module respectively, to facilitate cloning in a pBlueScript KS+ vector (Stratagene) where the BsaI restriction site in the ampicillin gene had been removed.

The *P. tricornutum* U6 snRNA promoter, located on Chr8 (complement pos. 239707–239986) acc: CM000611.1, was used for sgRNA expression. The RNA pol III transcription initiation site was estimated based on partially conserved 5' DNA sequence motifs in the U6 gene, high-throughput DNA sequencing data, and a conserved guanine located 27 bp downstream of the TATA box. To enable easy generation of sgRNAs with customized target sites, the vector contains two BsaI restriction sites located at the 5'-end of the guide RNA, which allow ligation of adapters (target sites) with 5' TCGA and AAAC overhangs<sup>26</sup>. The adapter for targeting the *CpSRP54* gene was made by annealing complementary oligos (1  $\mu\text{g}$  of each oligo) with the above described overhangs (Supplementary Table 3) in an annealing reaction containing 1  $\times$  T4 Ligase Buffer (NEB) in a total volume of 50  $\mu\text{l}$ . The annealing reaction was incubated in a heating block for 10 min at 85 °C, followed by slow cooling to room temperature (approx. 60 min). The resulting adapter was ligated into the pKS dia*Cas9*\_sgRNA plasmid using a molar vector to insert ratio of 1:20, and a T4 DNA ligase (Fermentas). A poly-T termination signal and 3' sequences from the U6 gene was added at the 3' end of the sgRNA module Chr8 (complement pos. 239266–239605). The module containing U6 snRNA promoter, sgRNA and U6 terminator was synthesized by Eurofins Genomic Services Ltd, with PstI and HindIII sites at 5' and 3' respectively to facilitate cloning. See Supplementary Fig. 2 for full DNA sequence of dia*Cas9* and sgRNA module. The pKS dia*Cas9*\_sgRNA plasmid vector will be made available through AddGene (Addgene ID: 74923).

**PAM-target site selection.** A Perl-based script and a local Blast server (Ncbi-blast 2.2.27) were used to identify target sites with low homology to other genomic loci. PAM-target sites with high homology to other loci identified through Blast searches (blastN-short) were excluded. Due to the limitations of blastN-short to find sequences with mismatches, using a seed sequence of 12 to 14 bp (proximal to the PAM site), an additional string-based search was performed that screen for single bp mismatches located in the proximal seed region. Based on the string searches the PAM sites were rated according to their “uniqueness”.

**Biolistic transformation.** *P. tricornutum* cell culture (500  $\mu\text{l}$ ) adjusted to 1  $\times 10^8$  cells/ml was plated on 1/2 f/2-Si, 1% agar plates and grown for 1 day. The pKS dia*Cas9*\_sgRNA vector was introduced to the cells by biolistic bombardment using the Biolistic PDS-1000/He Particle Delivery System (Bio-Rad, Hercules, CA, USA) as previously described<sup>10</sup>. The cells were co-transfected with the pAF6 vector<sup>10</sup> to enable selection of transformants on 1/2 f/2-Si, 1% (w/v) agar plates containing 100  $\mu\text{g/ml}$  zeocin (InvivoGen). Tungsten M17 microcarriers (Bio-Rad) were coated with 2.5  $\mu\text{g}$  of each vector following the manufacturer's instructions (Bio-Rad). Cells were transferred to selection plates 1 day after bombardment. Resistant colonies appearing 3–4 weeks after bombardment were transferred to liquid f/2 containing 50  $\mu\text{g/ml}$  zeocin (InvivoGen).

**Screening for gene targeted mutations.** Transformants were screened for DNA sequence mutations in the target region of the *CpSRP54* gene (Supplementary Fig. 3) by performing high-resolution melting (HRM) based PCR assays on the region spanning the target site of the *Cas9*-sgRNA complex. A small amount of cells derived from each transformant were lysed<sup>7</sup>. An amplicon with a size of approx. 500 bp surrounding the target site were amplified by PCR using the lysate as template and ExTaq DNA polymerase and Buffer system (TaKaRa). The PCR products were purified using Wizard<sup>®</sup> SV Gel and PCR Clean-Up kit (Promega). The concentration of the purified PCR products were adjusted to 5 ng/ $\mu\text{l}$ , diluted 1:10<sup>6</sup> and used as DNA template in following HRM analyses. An amplicon of approx. 100 bp surrounding the target site was amplified using the LightCycler 480 High Resolution Melting Master kit (Roche) following the manufacturer's instructions. The PCR reactions and HRM curve analyses were performed on a LightCycler 96 instrument (Roche). Mutations were confirmed by Sanger sequencing. The 500 bp PCR product was cloned into pCR<sup>™</sup>4-TOPO<sup>®</sup> TA vector (Invitrogen), and re-sequenced



to get an overview of the different indels present within a colony. Primers used are presented in Supplementary Table 3.

**High blue light experiment.** Six biological replicates of wild type (WT) cells and three biological replicates of each of the four different CpSRP54 mutant lines (M1, M4, M5 and M8) were subjected to 1 h of high intensity blue light (BHL;  $500 \mu\text{mol photon m}^{-2} \text{s}^{-1}$ ) provided by a SL3500 LED light source (Photon System Instruments) at  $15^\circ\text{C}$ . The PSII efficiency (maximum quantum yield;  $F_v/F_m$ ) was measured at the end of a 3 min dark acclimation period using an AquaPen-C (Photon System Instruments) after 5 min, 0.5 h and 1 h of BHL exposure. After 1 h of BHL, cell cultures were returned to the previous growth conditions at  $65 \mu\text{mol photon m}^{-2} \text{s}^{-1}$  white light (WL) at  $22^\circ\text{C}$ . The PSII efficiency was measured 0.5 h, 1 h and 3 h into the recovery period.

**Isolation of “clean” CpSRP54 mutant lines.** Highly diluted cell suspensions ( $\sim 200$  cells/ml;  $500 \mu\text{l}$ ) from five (M1, M2, M4, M5, M8) of the eight original mutant cultures were plated on  $\frac{1}{2}$  f/2-Si, 1% agar plates containing  $100 \mu\text{g/ml}$  zeocin (InvivoGen). Clean CpSRP54 mutant lines with biallelic mutations originating from single cells, were isolated and identified by repeating the screening process described above on the resulting colonies.

**RNA isolation and quantitative real-time PCR.** Harvesting of three biological replicates of WT and clean CpSRP54 mutant cultures (M1.1, M2.1, M4.1, M5.1, M8.1) grown at  $22^\circ\text{C}$ ,  $65 \mu\text{mol photon m}^{-2} \text{s}^{-1}$ , and subsequent RNA isolation, quantification and verification of RNA integrity were performed as described in Nymark *et al.*<sup>24</sup>. Reverse transcription of  $1 \mu\text{g}$  of total RNA from all samples was performed with QuantiTect Reverse Transcription kit (Qiagen) following the recommended protocol. Reactions where the reverse transcriptase had been omitted were included for all samples to be used as genomic DNA controls during the quantitative real-time PCR (qRT-PCR) analyses. The qRT-PCR reactions were performed on a LightCycler 96 instrument (Roche) as described in Nymark *et al.*<sup>24</sup>, using cDNA diluted 1:10 as template. Forward and reverse primers used are listed in Supplementary Table 3. Quantification cycle (Cq) – values were determined using the LightCycler<sup>®</sup> 96 Software.

**Cell divisions per day.** Algal cells were counted by a NovoCyte<sup>™</sup> flow cytometer (ACEA Biosciences) to calculate the cell division per day in three biological replicates of WT and clean CpSRP54 mutant cultures (M1.1, M2.1, M4.1, M5.1, M8.1) grown under dim WL conditions (16 h photoperiod,  $22^\circ\text{C}$ ,  $65 \mu\text{mol photon m}^{-2} \text{s}^{-1}$ ). Samples were excited by a 488 nm laser and chlorophyll fluorescence emission collected on a detector with a 675/30 nm bandpass filter.

## References

- Field, C. B., Behrenfeld, M. J., Randerson, J. T. & Falkowski, P. Primary production of the biosphere: integrating terrestrial and oceanic components. *Science* **281**, 237–240 (1998).
- Armbrust, E. V. Identification of a new gene family expressed during the onset of sexual reproduction in the centric diatom *Thalassiosira weissflogii*. *Appl Environ Microbiol* **65**, 3121–3128 (1999).
- Nelson, D. M., Treguer, P., Brzezinski, M. A., Leynaert, A. & Queguiner, B. Production and dissolution of biogenic silica in the ocean - Revised global estimates, comparison with regional data and relationship to biogenic sedimentation. *Global Biogeochem Cy* **9**, 359–372, doi: 10.1029/95gb01070 (1995).
- Smetacek, V. Diatoms and the ocean carbon cycle. *Protist* **150**, 25–32, doi: 10.1016/S1434-4610(99)70006-4 (1999).
- Lebeau, T. & Robert, J. M. Diatom cultivation and biotechnologically relevant products. Part II: current and putative products. *Appl Microbiol Biotechnol* **60**, 624–632, doi: 10.1007/s00253-002-1177-3 (2003).
- Bozarth, A., Maier, U. G. & Zauner, S. Diatoms in biotechnology: modern tools and applications. *Appl Microbiol Biotechnol* **82**, 195–201, doi: 10.1007/s00253-008-1804-8 (2009).
- Daboussi, F. *et al.* Genome engineering empowers the diatom *Phaeodactylum tricornutum* for biotechnology. *Nat Commun* **5**, 3831, doi: 10.1038/ncomms4831 (2014).
- Weyman, P. D. *et al.* Inactivation of *Phaeodactylum tricornutum* urease gene using transcription activator-like effector nuclease-based targeted mutagenesis. *Plant Biotechnol J* **13**, 460–470 (2015).
- Sander, J. D. & Joung, J. K. CRISPR-Cas systems for editing, regulating and targeting genomes. *Nat Biotechnol* **32**, 347–355, doi: 10.1038/nbt.2842 (2014).
- Falciatore, A., Casotti, R., Leblanc, C., Abrescia, C. & Bowler, C. Transformation of nonselectable reporter genes in marine diatoms. *Mar Biotechnol* **1**, 239–251, doi: MBT30 [pii] (1999).
- Jacobs, T. B., LaFayette, P. R., Schmitz, R. J. & Parront, W. A. Targeted genome modifications in soybean with CRISPR/Cas9. *BMC Biotechnol* **15**, 16, doi: 10.1186/s12896-015-0131-2 (2015).
- Shan, Q. *et al.* Targeted genome modification of crop plants using a CRISPR-Cas system. *Nat Biotechnol* **31**, 686–688, doi: 10.1038/nbt.2650 (2013).
- Wang, Y. P. *et al.* Simultaneous editing of three homoeoalleles in hexaploid bread wheat confers heritable resistance to powdery mildew. *Nat Biotechnol* **32**, 947–951, doi: 10.1038/nbt.2969 (2014).
- Feng, Z. Y. *et al.* Efficient genome editing in plants using a CRISPR/Cas system. *Cell Res* **23**, 1229–1232, doi: 10.1038/cr.2013.114 (2013).
- Miao, J. *et al.* Targeted mutagenesis in rice using CRISPR-Cas system. *Cell Res* **23**, 1233–1236, doi: 10.1038/cr.2013.123 (2013).
- Xu, R. F. *et al.* Gene targeting using the *Agrobacterium tumefaciens*-mediated CRISPR-Cas system in rice. *Rice* **7**, doi: ARTN 5 10.1186/s12284-014-0005-6 (2014).
- Zhang, H. *et al.* The CRISPR/Cas9 system produces specific and homozygous targeted gene editing in rice in one generation. *Plant Biotechnol J* **12**, 797–807, doi: 10.1111/pbi.12200 (2014).
- Pâques, F. & Haber, J. E. Multiple pathways of recombination induced by double-strand breaks in *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev* **63**, 349–404 (1999).
- Richter, C. V., Bals, T. & Schunemann, D. Component interactions, regulation and mechanisms of chloroplast signal recognition particle-dependent protein transport. *Eur J Cell Biol* **89**, 965–973, doi: 10.1016/j.ejcb.2010.06.020 (2010).
- Amin, P. *et al.* *Arabidopsis* mutants lacking the 43- and 54-kilodalton subunits of the chloroplast signal recognition particle have distinct phenotypes. *Plant Physiol* **121**, 61–70 (1999).
- Hutin, C. *et al.* Double mutation cpSRP43<sup>-</sup>/cpSRP54<sup>-</sup> is necessary to abolish the cpSRP pathway required for thylakoid targeting of the light-harvesting chlorophyll proteins. *Plant J* **29**, 531–543 (2002).

22. Nymark, M. *et al.* Molecular and photosynthetic responses to prolonged darkness and subsequent acclimation to re-illumination in the diatom *Phaeodactylum tricornutum*. *PLoS One* **8**, e58722, doi: 10.1371/journal.pone.0058722 (2013).
23. Jiang, W., Brueggeman, A. J., Horken, K. M., Plucinak, T. M. & Weeks, D. P. Successful transient expression of Cas9 and single guide RNA genes in *Chlamydomonas reinhardtii*. *Eukaryot Cell* **13**, 1465–1469, doi: 10.1128/EC.00213-14 (2014).
24. Nymark, M. *et al.* An integrated analysis of molecular acclimation to high light in the marine diatom *Phaeodactylum tricornutum*. *PLoS One* **4**, e7743, doi: 10.1371/journal.pone.0007743 (2009).
25. Cong, L. *et al.* Multiplex Genome Engineering Using CRISPR/Cas Systems. *Science* **339**, 819–823, doi: 10.1126/science.1231143 (2013).
26. Hwang, W. Y. *et al.* Efficient genome editing in zebrafish using a CRISPR-Cas system. *Nat Biotechnol* **31**, 227–229, doi: 10.1038/nbt.2501 (2013).

### Acknowledgements

This work was funded by a grant from the Gordon and Betty Moore Foundation GBMF 4966 to Atle Bones and the NTNU enabling technologies program.

### Author Contributions

M.N. and P.W. conceived and designed the experiments. M.N., A.K.S. and T.S. performed the experiments. M.N., P.W. and T.S. analysed the data. A.M.B. and P.W. contributed reagents and materials. M.N., A.M.B. and P.W. wrote the paper.

### Additional Information

**Supplementary information** accompanies this paper at <http://www.nature.com/srep>

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Nymark, M. *et al.* A CRISPR/Cas9 system adapted for gene editing in marine algae. *Sci. Rep.* **6**, 24951; doi: 10.1038/srep24951 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>