

Julius William Nyahongo

**Depredation of Livestock
by Wild Carnivores and
Illegal Utilization of Natural
Resources by Humans in the
Western Serengeti, Tanzania**

Thesis for the degree philosophiae doctor

Trondheim, November 2007

Norwegian University of Science and Technology
Faculty of Natural Sciences and Technology
Department of Biology



NTNU

Norwegian University of Science and Technology

Thesis for the degree philosophiae doctor

Faculty of Natural Sciences and Technology
Department of Biology

© Julius William Nyahongo

ISBN 978-82-471-5272-0 (printed version)
ISBN 978-82-471-5286-7 (electronic version)
ISSN 1503-8181

Doctoral theses at NTNU, 2007:238

Printed by NTNU-trykk

PREFACE

This work is an output of the Biodiversity and Human Wildlife Interface (BHWI) in the western Serengeti, a collaborative project between the Tanzania Wildlife Research Institute (TAWIRI), the Norwegian Institute for Nature Research (NINA) and the Norwegian University of Science and Technology (NTNU). The Messerli Foundation of Switzerland, Institute for Zoology and Wildlife Research (IZW) of Germany and the Norwegian Agency for Development Cooperation (NORAD) funded the initial stage of the project between 2001 and 2004. The final stage of this work (2005 and 2007) was financially supported by the Norwegian Peace Corps and the Quota Programme Scheme (NORAD). Among objectives of the BHWI has been to build research capacity at TAWIRI.

I thank the management of TAWIRI, in particular Charles Mlingwa, (the former Director General), Simon Mduma (the Acting Director General), George Sabuni and Julius Keyyu for granting me a three years study leave and assisting me logistically during the data collection. I also thank the Department of Biology, NTNU for hosting me as a student during my study. My deepest appreciation is extended to my academic supervisors; Eivin Røskoft and Bjørn Kaltenborn who guided me through academic life throughout my stay in Norway. The following staff members at the Department of Biology were always available when I needed any one kind of assistance: Lisbeth, Jasenka, Inger, Hans Jacob, Anne Beate, Tove, Ingunn and Claudia Melis. Bård Stokke and Tomas Holmern offered invaluable advice in data analyses. I am grateful to Marion East, Fatima Mturi and Heribert Hofer who supervises my field work during the initial stage of this project

(2001-2003). From the Norwegian Institute of Nature Research (NINA), the assistance offered by Graciella Rusch, Sigbjørn Stokke, Bjørn Kalternborn and Jørn Thomassen has been appreciated. The logistical support provided by The International Office, in particular Ms Ragnhild Brakstad played an important role in ensuring my peaceful stay in Norway. I thank Grumeti Reserve Fund, especially, Brian Harris and Ed Sayer who devoted their valuable time to organize and finally carrying out the hunting activities for this research against their will. I acknowledge the assistance offered by Grumeti and Ikorongo Game Reserve, especially Benjamin Kijika and the Serengeti Regional Conservation Programme (SRCP), especially John Muya.

I would also like to thank my field assistants, Ally Kashindye, Richard Ndaskoi and Herry Lema who endured the long trips each month to and from Serengeti. I also appreciate the assistance of livestock enumerators; Maganga Chama (Nattambiso), Mangora Samson (Nyamakendo), George Nyahongo (Kowak) and John Wambura (Robanda). The Serengeti Wildlife Research Centre staff members are all acknowledged, especially Robert Fyumagwa who shared his apartment with me when I was in Serengeti. All village leaders, Robanda, Nyamakendo, Nattambiso, Nyatwali, Butiama, Rwamkoma, Busegwe, Kowak, Chereche and Omuga are acknowledged for their valuable support. I also appreciate the good time with my fellow students at NTNU, Shombe Hassan, Jafari Kideghesho, Flora Magige, Jarle Inge Holten, Johan Vikan, Anne Skjetne Motensen, Wilfred Marealle, Vedasto Ndibalema, and Iddi Mfunda for their academic and social companion.

In addition to academic life, we enjoyed social gatherings that always were organized in Eivin Røskaft's house. 'Mama' Berit Røskaft is thanked for her valuable time devoted preparing food for all of us from Tanzania, Bangladesh, Bulgaria and Norway. I also enjoyed good time with Bjørn Kaltenborn's family in Lillehammer and Jørn Thomassen's family in Trondheim. Thank you very much. Ragnvald Larsen and Synne Rudsar gave me a map of Trondheim in Arusha, Tanzania and explained to me how to survive the winter in Norway when I was planning to come to Norway for my study.

I also extend my acknowledgment to my beloved mother, Beldine and my sisters, Helen, Penina, Jenipha, Rose and Mary who kept on praying for me during my study. I also thank my brother, George who is looking after *old mama* and the inherited livestock there at the home village-Kowak. Keeping inherited livestock link the living family to the dead great-great grand parents.

Last but no by means least, I thank my beloved wife; Lucy who cared children alone during my stay in Norway. That allowed me peace of mind to focus on the job at hands. My beloved sons, Frank and Innocent who missed the fatherly love for several months each year and our beloved *brand new* daughter, Dinna whose first ten months in this wonderful world passed without getting the fatherly kiss.

Table of Contents

PREFACE..... 1

TABLE OF CONTENTS 4

THIS THESIS CONSISTS OF THE FOLLOWING FIVE PAPERS..... 6

SUMMARY 7

INTRODUCTION..... 10

Carnivore-human conflicts 11

Effects of illegal hunting 14

Dietary contribution of bushmeat to local people..... 16

AIMS OF THE THESIS..... 18

METHODS 19

Study area 19

Illegal bushmeat hunting and law-enforcement..... 21

Off-take levels..... 23

Hunting techniques 24

Study species..... 25

Data collection 26

MAIN RESULTS 26

Paper I..... 26

Paper II 27

Paper III..... 27

Paper IV 28

Paper V 29

DISCUSSION	29
Natural resources utilization and wildlife experience	29
CONFLICTS OVER LIVESTOCK-LIVESTOCK DEPREDATION	32
MANAGEMENT RECOMMENDATIONS	35
FUTURE RESEARCH NEEDS	37
REFERENCES.....	39

THIS THESIS CONSISTS OF THE FOLLOWING FIVE PAPERS.

- I Nyahongo, J.W., East, M.L., Mturi, F. & Hofer, H. (2006) Benefits and costs of illegal grazing and illegal hunting in the Serengeti ecosystem. *Environmental Conservation*, 32 (4), 326-332
- II Holmern, T., Nyahongo, J. & Røskaft, E. (2007) Livestock loss caused by predators outside the Serengeti National Park, Tanzania. *Biological Conservation*, 135, 534-542.
- III Nyahongo, J.W., Holmern, T., Stokke, B.G., Keyyu, J.D., Kaltenborn, B.P. & Røskaft, E. (Submitted) Disease is a major cause of livestock loss in villages surrounding western Serengeti, Tanzania.
- IV Nyahongo, J.W., Holmern, T., Kaltenborn, B. P. & Røskaft, E. (Submitted) Spatial-temporal variation in meat and fish consumption among humans in the western Serengeti, Tanzania: the importance of migratory herbivores.
- V Nyahongo, J.W., Holmern, T., Stokke, B.G., Kaltenborn, B. P. & Røskaft, E. (Submitted) Bushmeat preference and species recognition based on meat taste by humans in the western Serengeti, Tanzania.

Declaration of contribution

EML, MF and HH contributed with ideas, planning for fieldwork, data collection and commenting on paper I. ER contributed with ideas, planning for fieldwork, data collection and commenting on the II, III, IV and V ms throughout. Contribution of the co-authors: Paper II, I provided some data, exchanged some notes, discussed some results and commented on the ms. Paper III, HT, SBG, KJD and KBP commented on the ms. Paper IV, HT and KBP commented on the ms. Paper V, HT, SBG and KBP commented on the ms.

SUMMARY

Human-wildlife interactions play an important role in shaping perceptions and conservation paradigms and the livelihoods in villages neighbouring protected areas. These interactions also determine the future survival of the wildlife in the face of increasing pressure due to high human population increase characterising most countries in sub-Saharan Africa. Most rural people in sub-Saharan Africa are agropastoral, combining small scale farming with animal husbandry, or they are purely agropastoralists or farming who relies on natural resources for sustenance.

The negative impacts from wildlife to humans may include crop damage, attacking and killing livestock and humans, competing for game species or acting as diseases reservoirs. Humans may affect wildlife through a wide range of lethal methods such as shooting, poisoning, trapping or snaring, habitat modification, encroachment or diseases exchange between wildlife and livestock.

Illegal hunting using traditional weapons is wide spread in communities surrounding areas rich of wildlife where in some countries in Africa (i.e. Liberia) up to 75% meat protein is derived from wildlife. The main factors attributing to high consumption of bushmeat is local availability, easy catch-ability (wire snares, pitfall traps), affordability and the consequent household savings.

This thesis evaluates the conflicts between human and wildlife in the human-wildlife interface using the western Serengeti as a case study. The first part of the thesis focuses

on the conflict related to utilization of natural resources and livestock depredation whereas the second part focuses on the dietary contribution of bushmeat to local people, bushmeat experience and utilization.

Local people living close to protected areas are rational when it comes to the illegal utilization of natural resources because they consider the benefits and cost implications. The bushmeat hunters, especially, know in advance which areas in the protected areas are profitable at the same time consider the cost of being arrested and the distance they need to walk to the profitable areas. While illegal hunting can take place far in the park, livestock keepers avoid grazing inside the park because they know the consequences (penalties and fines) of utilizing the pasture inside the protected areas illegally.

The local people living close to protected areas consume more meat meals during the period when the wildebeest are in the village proximities than when the herds are far in the southern plains. This further proves the rationality of illegal bushmeat hunters when planning for hunting trips (the benefits versus cost). In contrast, the fish meals in the villages located close to protected areas but far from Lake Victoria decrease with influx of migratory herbivores, which suggest that fish and meat complement each other when the distance from the sources fluctuates. This was proved true when test-persons from villages close, intermediate and distant from the nearest national park boundary were given pieces of meat in a combination of wild ungulates and beef to rank the meat and species recognition according to the perceived taste. While the test-persons from distant villages preferred beef to all, the test-person from villages close to national park

boundary prefer topi and those in the intermediate villages prefer impala. This suggests long term experience with beef to distant test-persons as no other source of meat is locally available in the area other than livestock meat and fish.

Wild carnivores are considered to be responsible for livestock losses in the villages surrounding the protected areas. The results from the current study in the villages surrounding the western Serengeti show that among the wild carnivores reported to kill livestock, 97.7% of all reported claims was spotted hyena, being responsible for 98.2%. Spotted hyenas are nocturnal animals capable of commuting up to 80 km from their territory areas and are the most numerous large carnivore species in the Serengeti ecosystem, mainly targeting goats and sheep. To evaluate the level of conflicts between carnivores and human on livestock depredation, enumeration of livestock loss causes was conducted for subsequent comparison. In all villages, diseases were responsible for major loss of livestock.

Based on the findings the current study recommends better education on wildlife conservation, livestock husbandry practices and extension. A change in wildlife policy in favour of compensation would reduce the retaliatory killing of carnivores in the villages. Livestock keepers should improve the night holding enclosures to reduce livestock depredation by nocturnal predators. The findings recommend further study on the alternative sources of meat protein to local communities living close to protected areas. Last but not least, I recommend a special conservation attention to resident herbivore population close to village proximities.

INTRODUCTION

Human-wildlife interactions play an important role in shaping perceptions and conservation paradigms and the livelihoods in villages neighbouring protected areas. These interactions also determine the future survival of the wildlife in the face of increasing pressure due to high human population increase characterising most countries in sub-Saharan Africa (Ceballos and Ehrlich 2002). Biodiversity is being depleted at a rate that is causing concern among conservation interests worldwide.

The human population in Africa has increased from 224 million in 1950 to 960 million in 2005, and is predicted to reach more than 1.2 billion people by 2030 (UNDP, 2002). Most of this increase will happen in the rural areas, which currently hold 65-85% of the African population (UNDP, 2002). This inevitably will affect the conservation in the future because most rural people in sub-Saharan Africa are agropastoral, combining small scale farming with animal husbandry, or are purely agropastoralists or farming. The future reliance on natural resources (i.e. water, firewood, rangeland for livestock, fish and bushmeat together with mining) for sustenance means exhaustion of their resources base that not only affects conservation and biodiversity but is also a threat to human welfare.

Impact from human-wildlife interactions may be either positive or negative on the parties affected (Conover, 2002). The negative impacts from wildlife to humans may include crop damage (Dey, 1991; Naughton-Treves, 1998), attacking and killing livestock (Mishra, 1997; Ogada et al., 2003), competing for game species (Gasaway et al., 1992; Thirgood et al., 2000) attacking and killing humans (Herero, 1985; Saberwal et al., 1994;

Nowell and Jackson, 1996; Løe and Røskaft, 2004; Packer et al., 2005) or acting as diseases reservoirs (Jenknis et al., 1998; Hudson et al., 2002; Kock, 2003). On the other hand, technological advancements have lead to a wide range of lethal methods for controlling wildlife, such as shooting, poisoning, trapping or snaring (Hofer et al., 1996; Brand and Nel, 1997; Treves and Naughton-Treves, 2005). Indirectly, humans may affect wildlife through habitat modification, encroachment or diseases exchange between wildlife and domestic stock all of which intend to satisfy the humans needs (Kock, 2003).

Carnivore-human conflicts

The common conflicts between humans and wild animals in different parts of the world involve livestock depredation and crop damages. Although a remarkable range of species cause conflicts with humans, from rodents such as prairie dogs (*Cynomys ludovicianus*) to mega-herbivores like African elephants (*Loxodonta africana*; Hoare, 1999), large carnivores are of particular interest in this conflict. This is due to their obligate instinctive carnivorous behaviour, which put them into direct competition with humans for both livestock and wild game species or their ability to kill humans, which create more fear, intensifying the conflicts (Sillero-Zuberi, and Laurenson, 2001; Baldus, 2004; Løe and Røskaft, 2004; Packer, 2005). These perceptions are always compounded by an innate fear of large predators and long term negative attitudes that have developed among humans towards large predators due to the past experiences they had or were told even if carnivores do not pose any threat in present time (Quammen, 2003, Røskaft, 2003; Dickman, 2005). While studies show that large carnivores are not responsible for as much damage as local people commonly perceive (Rasmussen, 1999), this perception of

severe conflict is the important factor, as negative attitudes are strongly linked to retaliatory killing of carnivores (Gittleman et al., 2001, Paper II). This has resulted in persecution of wild carnivores in most parts of the world. For instance, angry farmers in Norway were reported to kill wolf (*Canis lupus*) to reduce sheep predation (Kaltenborn et al., 1999), and even today is the conflict between sheep farmers and wolves at a serious level elsewhere several places outside Africa (Røskaft et al., 2003).

Large-scale predator control programs have historically been employed to reduce predator conflicts with humans (Kellert, 1985; Woodroffe, 2000). All predators have suffered persecution with the result that they have been exterminated over most of their former ranges, particularly in Europe, North America and parts of Asia (Ginsberg and Macdonald, 1990; Saberwal et al.; 1994; Nowell and Jackson, 1996; Mills and Hofer, 1998). Lethal control of carnivores has resulted in extinction of several species of carnivores. For instance, a combination of trapping for fur and poisons to protect sheep led to the extinction of the Falkland's wolf or Malvinas zorro (*Dusicyon australis*) in 1876 (Sillero-Zubiri et al., 2004). Similarly, conflict with humans was identified as a key factor behind the extinction of the Carolina parakeet (*Conuropsis carolinensis*) in 1904 and that of the thylacine or marsupial wolf (*Thylacinus cynocephalus*) in 1930 (IUCN, 2006; Woodroffe et al., 2005).

In Africa, killing of wild carnivores over livestock depredation has been reported. Berry (1990) reported the killing of at least 320 lions (*Panthera leo*) between 1980 and 1989 on farms bordering the Etosha National Park, Namibia. Stuart et al. (1985) reported the

killing of leopards (*Panthera pardus*) by farmers due to predation on livestock in the Cape Province, South Africa. Holekamp and Smale (1992) reported that the growing human population around Maasai Mara National Reserves in Kenya poisoned at least 14 spotted hyenas (*Crocuta Crocuta*) in a single incidence in June 1991 to reduce livestock predation.

Even where the carnivore-human conflicts does not result in extinction, it may have a devastating impact on species' population size and geographical range, often leading to local extirpation (Johnson et al., 2001; Treves and Naughton-Treves, 2005). For example, cheetahs (*Acinonyx jubatus*) historically ranged across Africa, Asia and into the Indian sub-continent, with numbers estimated at ca. 100 000 individuals in 1900 (Marker, 1998). However, the population of cheetah has declined to less than 15 000 individuals globally for the past 50 years, and a complete disappearance in at least 13 countries where cheetah has been recorded (Marker, 1998). Similarly, African wild dogs (*Lycaon pictus*) have suffered a severe eradication from 25 of the 39 countries they used to occupy and are now one of the world's most endangered carnivores, with total number estimated to fewer than 5000 individuals, with only six packs thought to hold over 100 individual dogs (Fanshawe et al., 1991; Woodroffe et al., 1997).

The carnivore species causing most conflicts are also those who are most important in ecological maintenance. Large carnivores fulfil many important ecological functions, such as regulating prey numbers (many of them crop pests or water pollutants), controlling number of mesopredators through competition or are maintaining a functional

balance of biodiversity in local communities (Krebs et al., 1995; Logan and Sweanor, 2001). Removing top predators from habitat patches often results in marked changes in biodiversity and community structure, which may have severe ecological effects (Terborgh et al., 2002).

Effects of illegal hunting

Illegal hunting using traditional weapons such as snares, bow and arrows is widespread in communities surrounding areas rich of wildlife. Legal hunting on the other hand require the possession of a license and the demand for a license include among others, a fire arm which the majority of people in poor countries rich of biodiversity cannot afford. Illegal hunting is motivated by the need for protein, income and sometimes acts as food especially during prolonged droughts. These and other factors magnify the hunting pressure resulting on the park-people conflicts (Holmern et al., 2004).

Estimated off-take as a percentage of total population has been indicated to diverge widely between species. Among several resident species in Africa, including giraffe (*Giraffa camelopardalis*), impala (*Aepyceros melampus*) and topi (*Damaliscus korrigum*), off-take must be considered high. For instance, past exploitation for bush meat in Serengeti has significantly reduced the Cape buffalo (*Syncerus caffer*) by 50-90% in parts of their range and local declines in waterbuck (*Kobus ellipsiprymnus*) and giraffe population (Campbell, 1989; Dublin et al., 1990). Furthermore, roan antelope *Hippotragus equines* might have never been common in Serengeti due to over hunting (Turner, 1987; McNaughton, 1989). Hippo (*Hippopotamus amphibious*) populations are

not known in many ecosystems, thus the effect of off-take is difficult to assess (Hofer et al., 1996). Furthermore, Rusch et al. (2005) reported a drastic decline in the topi population in Serengeti, while populations of other herbivores either remained steady or increased which raises concern that the topi is particularly targeted by illegal hunters and exploited at unsustainable level. In Zambia, a comparison between the 1960s and 1994, animal sighting close to the villages suggest a drop of 50% of hippo populations (Marks, 1994). Moreover, a long term study from Ghana suggests that bushmeat hunting caused a decline of about 41 species of mammals by 76% between 1970 and 1998 and a local extinction of between 16 to 45% of the same species (Brasheres et al., 2001).

Some previous studies suggest that illegal hunting, which is the major source of bush meat supply, has been more detrimental to animal population not only for bush meat species but also for trophy animals. For instance, unchecked illegal hunting between 1975 and 1986 drove the black rhino (*Diceros bicornis*) populations to factual extinction and significantly reduced the elephant population size in Tanzania (Hofer et al., 1996). In Kenya, black rhino population decreased by 90% between 1969 and 1979 whereas in Zambia the black rhino population decreased from 12 000 individuals in 1972 to a few hundred in the 1980s (Leader-Williams & Albon, 1988). Untargeted species has also been caught in snares set for bush meat. In the Serengeti for instance, 8% of spotted hyenas (*Crocuta crocuta*) from the study population of 423 individuals are killed each year by snares that are set for bush meat (East and Hofer, 2000).

Dietary contribution of bushmeat to local people

Generally, the main factors attributing to high consumption of bushmeat is local availability, easy catch-ability (wire snares, pitfall traps), affordability and the consequent household savings. Snares and pitfall traps are easy to set and very efficient, while it is easy to conceal (Paper I). Moreover, bushmeat is usually cheaper than meat from livestock. For example, in Kitui district, amounts of bush meat consumed equate to 34% of household monthly income and from 15.7% to 39.2% in Kweneng and Kgalagadi in Botswana, respectively (Barnett, 2000).

In Tanzania for example, illegal utilization of bushmeat represents a larger economic value of wildlife than legalized trophy hunting or photographic tourism (Barnett, 2000).

A current study in Serengeti indicates that 83% of households buy illegal bushmeat (Holmern et al., 2004) the majority of them are subsistence farmers (Loibooki et al., 2002). The estimated mean number of people per household in the western Serengeti is seven (Hofer et al., 1996). Considering the number of households within 45 km from the park boundary, i.e. 137,750 households (roughly 964,250 people) depend on bushmeat as their main animal protein in the area with about 1.37 million people (URT, 2002). Annual off-take from this part of Serengeti alone has been estimated as 159,811 wild animals including resident (28%) and migratory species (72%) (Hofer et al., 1996). This is equivalent to 11,950 tons of meat per year or 230.6 g of meat per person per day.

In other African countries like Zambia, a similar study shows that a total of 27.4 tons of meat was made available to 466 local residents during the course of a year (Marks, 1973).

This is equivalent to 162 g of meat per person per day. Consumption of bush meat from these two regions surveyed is much higher than the minimum Food and Agriculture Organizational (FAO) recommendation of 60 g meat per person per day (Barnett, 2000). However, in Zambia, bush meat off-take for commercial purposes was considered to replace trophy poaching as the main impact on wildlife populations in many areas (Marks, 1973; Marks, 1994).

A study of bushmeat utilization in Kenya suggests that bushmeat represents the bulk of all meat protein consumed by Kitui communities. The study observed domestic meat playing a reduced role in meeting protein requirement as the meat from livestock was expensive (Barnett, 2000). Furthermore, study indicates that 80% of households consume 14.1 kg of bushmeat each month. In addition, FitzGibbon et al. (1996) report that traditional hunter or gatherer forest dwelling people rely heavily on bush meat as protein and potential income generating activity. In Botswana, 18.2 kg of bush meat is consumed per household per month by 46% of the Kweneng local people and the meat was the only viable source of meat protein for many rural inhabitants living in the semi-arid range land of the country (Barnett, 2000). Assuming an average of 7 people per household (Hofer et al., 1996), the average amount of meat consumed per person per day would be 86.7 g. This, however, is still higher than the minimum FAO recommendation. In Maputo Mozambique, the study indicates that more than 50 tons of bushmeat is traded per month. This has attributed directly to a severe decline in wildlife populations in the area. In Malawi the mini-fauna species are presently the source of meat protein to the majority of people (Barnett, 2000).

In contrast to countries within the Congo basin, the bush meat intake per day in Gabon and Congo were 180g and 89g, respectively (John et al., 2003). Likewise, these intakes were still higher than the minimum amount recommended by FAO. However, within the region, the study indicates that Cameroon and Democratic Republic of Congo (DRC) had much lower bush meat intake per person per day of 26g and 28g, respectively (John et al., 2003) suggesting depletion of the resource in these areas.

Aims of the thesis

This thesis evaluates the conflicts between human and wildlife in the human-wildlife interface using the western section of the Serengeti ecosystem in Tanzania as a case study. The first part of the thesis focuses on the conflict related to utilization of natural resources and livestock depredation (Papers I-III) whereas the second part focuses on the dietary contribution of bushmeat to local people, bushmeat experience and utilization (Papers IV-V).

METHODS

Study area

The study area is located in the north-eastern corner of Tanzania (Fig 1) on the north-western part of Serengeti National Park (SNP) (14 763 km²). The SNP is the central part of the greater Serengeti Ecosystem in the northern Tanzanian highlands. Serengeti was declared a national park in 1951 and a World Heritage Site in 1981 when the bordering Ngorongoro Conservation Area became a Biosphere Reserve. The park is approximately one-half of the entire ecosystem, which includes the Ngorongoro Conservation Area, Maswa, Ikorongo, Grumeti Game Reserves, Loliondo Game Controlled Area, and Masai Mara Game Reserve in Kenya (Fig 2).

The Serengeti ecosystem is a highland savannah region with thorn tree woodlands and plains from approximately 900–1500 meters above sea level. Annual precipitation ranges from about 800 millimetres in the east to 1000 millimetres in the northwest (Norton-Griffiths et al., 1975). The world largest populations of herbivores and carnivore are found in this ecosystem and the majority of the species of the East African savannah are found there too. Serengeti is famous for the large scale herbivore migrations (wildebeest, Thomson's gazelle, zebra and eland, Fig 2) as well as for the large populations of resident herbivores (African buffalo, giraffe, Grant's gazelle, impala, topi, warthog (*Phacochoerus aethiopicus*), and waterbuck). Sizeable populations of large carnivores like lion, leopard, cheetah and hyenas also roam these areas (Sinclair, 1995).

The people inhabiting this region are either agro-pastoralists or pastoralists. The areas north and west of SNP are densely populated (> 70 people/km², human population in Mara Region was about 1.37 million growing at a rate of 2.9% per annum (URT 2002)) by a diversity of tribes and ethnic groups. The main tribes are Ikizu, Zanaki, Sukuma, Jita, Taturu, Ikoma, Kuryia, Natta, Issenye and Luo. In earlier years, the cultural and ethnic differences were much more distinct than they are today. Largely due to the rapid population growth and significant transmigration from other areas far from the park boundaries, most of the communities along western Serengeti are currently multiethnic. The communities are organised just as much around available space and agricultural land and the search for economic opportunities, as traditional culture.

The average annual cash income of local people living in the study area is low (i.e. US\$ 140 in 2001, Borge, 2003). Overall, Tanzania is a poor country with a per capita income of US \$ 280 (World Bank 2006). By most conventional standards the villagers residing around north-western Serengeti are impoverished, and a great number of them qualify as poor by the UN standard. The main economic activities include farming and livestock production. Farming is mostly based on crops like cassava, sorghum, millet, maize (food) and cotton (cash crop). The crops cycle follows the rain-pattern with long rain lasting from March-May and short rain October-December. January-February and June-September are always dry. Maize, sorghum and millet are planted twice a year; in February-March and August-October and harvesting period is between June and July and between January and February, respectively. Livestock includes cattle, goats, sheep and poultry, although few households keep pigs and donkeys. Hunting varies in importance

among these tribes. The wildebeest migration is a central part of the annual life cycle for tribes like Ikoma and Kurya where hunting has traditionally been a part of culture and life patterns. The estimated number of illegal bushmeat hunters within 45 km of SNP and adjacent protected areas is 23,294 and 31,655, respectively (Campbell and Hofer, 1995). A more recent estimate (Campbell et al. 2001) puts the number of illegal bushmeat hunters at approximately 60,000, i.e. an increase of 90% in ten years (from 1988 to 1998). In contrast, the population to the east of the park is dominated by pastoralists (Maasai, who supposedly do not hunt), and there is very little farming here.

Illegal bushmeat hunting and law-enforcement

In the western Serengeti illegal hunting has increasingly become a coping strategy for a major part of the population as legal access to resources has been restricted (Campbell et al., 2001). According to Loibooki et al. (2002) people of the western Serengeti participate in illegal hunting in order to offset food shortage and generate cash income. Participation in illegal bushmeat hunting decreased with increasing numbers of livestock owned, and people with access to alternative income means were also less likely to engage in illegal hunting. Furthermore, involvement in illegal hunting was not reduced by participation in community-based conservation programmes.

In Serengeti, anti-poaching patrols have been an important task for park staff since the inception as a national park (Arcese et al., 1995; Loibooki et al., 2002). Arcese et al. (1995) report a possible six-fold increase in arrests from 1957 to 1991. However, they also point out that the ranger force has doubled since 1963, and in order to understand the

changes one must know how many people actually enter the park to hunt. Given the contentious nature of the issue, it may be impossible to arrive at an accurate estimate of this figure, at least if it is based on observation and self-reports.




Figure 1: Map of the western Serengeti showing the location of some study villages (Robanda, Nyamakendo, Nyatwali, Rwamkoma and Kowak) and the surrounding villages (Black dots).

Off-take levels

The illegal hunting activity has been spatially modelled. Campbell and Hofer (1995) estimated that in and around Serengeti 210 000 herbivores are hunted illegally each year. Of this, wildebeest comprises 57 per cent (118 922 animals). However, Mduma et al. (1998) estimated a much lower number of 40 000 wildebeest illegally hunted each year, and predicted that a harvest of 80 000 animals per year is unsustainable and could cause a total collapse of the wildebeest population by 2018. Given the fact that the current wildebeest population is reasonably stable at approximately 1.3 million animals could indicate that the Campbell and Hofer (1995) estimates may be too high.




Figure 2. Map of the Serengeti ecosystem. The core area is Serengeti National Park (SNP), surrounded by Maswa Game Reserve (MGR), Grumeti Game Reserve (GGR), Ikorongo Game Reserve (IGR), Maasai Mara National Reserve (MMNR), Loliendo Game Controlled Area (LGCR) and Ikoma Open Area (IOA). The arrows show the movement of wildebeest around the Serengeti Ecosystem. The wildebeest usually carves in the southern short grass plain in December-February each year. The northern part of the ecosystem is the refuge of migratory herbivores during dry season (August-November). On their northward migration (May-July), the wildebeest herds use parts of the western corridor, as well as the adjacent game reserves and open village lands, depending upon the rainfall pattern (adapted from Thirgood et al., 2004 and Rusch et al., 2005).

Hunting techniques

Hunting is conducted in a number of ways. Few people own firearms so most illegal hunting is accomplished by setting snares and pitfall traps (Plates 1 and Plate 2). Snares and pitfall traps are unselective hunting methods and can injure or kill a wide range of animals from large carnivores to small and large herbivores. They are often inefficient in killing and animals may suffer for a long time before they are dealt with by hunters and/or sometimes die and scavenged by predators (Plate 3). In some cases an animal escapes with a snare wire deeply cut through the neck (Plate 4). In some cases hunting involves well organised parties on several week long expeditions into the bush where the hunters set up a secluded camp, butcher and sun dry the meat before they depart. Much of the meat is then preserved in a form (swahili: 'kimoro') that permits storage and selling or trading in markets locally or far away (Kaltenborn et al., 2005). Alternatively, smaller groups and individuals take what they can find in their immediate surroundings, and mostly for subsistence use. During the wildebeest migrations huge herds of animals roam through villages and agricultural lands and great numbers of animals are slaughtered literally at the doorstep. A few authors have attempted to quantify the economy linked to wildlife harvesting (Campbell et al. 2001; Borge 2003), but there is as yet no comprehensive picture or consensus neither on the magnitude of the harvest, nor on the contribution to rural household economies due to the delicacy of the subject to local communities.



Plate 1: Snare wire set ready waiting for the victim in the park section of Ndabaka plain (Photo: J. Nyahongo)



Plate 2. Pitfall trap in Serengeti National Park (Ndabaka plain): big enough to swallow a mature buffalo (Photo: J. Nyahongo)



Plate 3. The scavenged victim of wire snare (Photo: J. Nyahongo)



Plate 4: A lioness with a wound around neck that has been inflicted by a wire snare (Kirawira pride) (Photo: J. Nyahongo)

Study species

In this study, we include those wild animals that influence human livelihood (bushmeat) and livestock (depredation). In the western Serengeti, the wildebeest, zebra and Thomson's gazelle migration has large impact on the livelihood of local communities adjacent to park boundary. However, resident animals such as topi, giraffe, impala, buffalo, warthogs and waterbuck are important sources of meat for both human and large carnivores (lions, spotted hyena and leopard) in the area. Spotted hyena is the most numerous carnivores in the area and the species is also found outside the protected areas. Thus, this is the carnivore being mostly involved in livestock depredation.

Data collection

Data for this thesis were collected during several field trips. Data on the benefits and costs of illegal grazing and hunting (Paper I) was collected between May 2001 and March 2002 for livestock depredations (Paper II) was collected between September and November 2004. Data for livestock depredations (Paper III) and for bushmeat utilization (Paper IV and Paper V) was collected between January and December 2006 (See the respective papers for detailed complete descriptions of methods).

MAIN RESULTS

The following papers (I, II & III) focuses on the conflict related to utilization of natural resources and livestock depredation.

Paper I

The levels of illegal use of natural resources by local communities surrounding the western Serengeti were influenced by the likely value of the resources acquired and the probable costs associated with their acquisition. Evidence of hunting was found in the national park section closer to a ranger post, suggesting that benefits (bushmeat) of hunting mostly outweighed costs (chances of being arrested) in these areas. However, the level of illegal hunting was observed to decrease with the distance hunters have to travel on foot to hunting areas. Travel cost is likely to be assessed not only in terms of distance travelled but also in terms of time that could be devoted to other activities (opportunity cost). In contrast, despite the high densities of livestock close to the boundary of the

protected areas, livestock was rarely illegally present inside these protected areas. This may indicate that livestock owners considered the chance of detection (as it is difficult to conceal grazing livestock) and likely financial penalties (when livestock are confiscated) too high in relation to the benefit gained from illegally acquired forage and the use of watering areas inside the areas.

Paper II

Livestock depredation in the villages surrounding the western Serengeti is mostly caused by spotted hyena, followed by leopard, baboon (*Papio cynocephalus*), lion and jackal. Economically, the livestock depredation contributed to two-thirds of the annual cash income for the households in the study area. This does not only intensify the human-carnivore conflicts but also may be a serious obstacle to both human and livestock development. Depredation events were not only reported to villages close to the protected areas but also affected households in distant villages where only the spotted hyena was reported to be involved in livestock killings. Lion and leopards only killed livestock in the households that were close to protected areas. Tolerance of livestock depredation was low and the majority of livestock owners accept retaliatory killing as a way to reduce loss. Level of education, the number of livestock previously lost and the perceived effective protective measures had influence on acceptance of retaliatory killing.

Paper III

Among the recorded causes of livestock losses such as disease, depredation, theft, and loss in bush while grazing, the results suggest that diseases are responsible for the highest loss. Death of livestock due to diseases affected household similarly. Overall, diseases cost was 59.6% of average annual household cash income. In comparison, the

contribution of diseases to livestock loss was four times higher than depredation in the household located far away from the park boundary, and was about 10 times higher than the cost of depredation in the households that were close to the park boundary. More sheep were killed by spotted hyena in the households located farther away than those located close to the park boundary. Overall, spotted hyena killed more sheep than goats or cattle.

The following papers (IV & V) focuses on the dietary contribution of bushmeat to local people, bushmeat experience and utilization.

Paper IV

Meat and fish meals per household were studied in villages that were located close, intermediate and/or farther away from the boundary of the protected areas (Serengeti National Park and Grumeti-Ikorongo Game Reserves) and/or the Lake Victoria. Generally households that were close to the protected areas consume more meat during the migration than those located farther away where the peak meat consumption in the villages close to protected areas corresponded to the peak influx of migratory herbivores. Similarly, households located close to Lake Victoria eat more fish than those located farther away. The consumption of fish meals is not affected by the influx of migratory herbivores close to the villages located close to the lake. Fish consumption in villages that were close to the protected areas but far from the Lake Victoria declined with the influx of migration. The household income significantly influenced the meat consumption in the villages that were far from the protected areas but not in the villages that were located close or intermediate distance from the boundary of the protected areas.

Understanding human species preference and ability to recognize species by meat taste may be employed to explore how some group of people along the gradient of distance from the park have experience with different species of wild ungulates and beef. This can be an indirect method to evaluate the levels of the past and current bushmeat utilization.

Paper V

Our overall results show that test-persons favoured beef, followed by topi and impala. The preference patterns and the ranking position of beef, topi and impala alternated along the gradient of distance from the park suggesting high preference and acceptability of the three species by test-persons from different villages along the gradient of distance from the park boundary. Moreover, it was possible to predict the preferences of beef, topi, impala and wildebeest along the gradient of distance from the park. In contrast, the results indicated that most test-persons were not able to identify the species based on the meat test. Generally, the most correctly identified meat was beef while the least identified species was impala. Age and gender did not have a significant effect on meat preference for all species in the pooled data. Distance from the park had a negative effect on the preference of topi and the similar effect on the identification of all five species studied suggesting a different level or type of experience with topi in the immediate villages.

DISCUSSION

Natural resources utilization and wildlife experience

Generally the findings from this study show that local people living close to protected areas are rational when it comes to the illegal utilization of natural resources. They consider the benefits and cost of illegal utilization of natural resources (Hofer et al., 2000;

Paper I). In this context, local people living close to protected area are able to plan and carefully follow the laid plans during the hunting operation. They know in advance which areas in the protected areas are profitable (high herbivore densities), at the same time they consider the cost (chances of being arrested and the distance to walk). While illegal hunting can take place far inside the park, livestock keepers avoid grazing deep inside the national park because they know the consequences (penalties and fines) of utilizing the pasture in the park illegally. Illegal hunting can be easily concealed and often takes place at night while grazing take place during the day and involves large herds of cattle, which is easier to see from long distances. In the western Serengeti previous studies indicate that most arrested illegal bushmeat hunters are poor uneducated people who own few or no livestock (Loibooki et al., 2002), which suggest that bushmeat hunting has been and will continue to be (unless the economy and social services such as better education, employment opportunities, health and water sanitation are improved in the villages) a coping strategy for survival in the areas with relatively abundant wild ungulates (Kaltenborn et al., 2005).

In order to investigate the dietary contribution of bushmeat to local people and their experience with wildlife as a result of long term human-wildlife interactions (bushmeat utilization), one method we can use is direct and indirect observation of what local people consume during the period spanning several months or years and compare that with the seasonal movements of migratory herbivores through and around the Serengeti ecosystem. On their northward migration, wildebeest moves close or direct in village

areas during the period covering about three months (May-July) each year. During this period, substantial numbers of wildebeest and zebra are slaughtered for bushmeat.

The findings of the current study (Paper IV) shows that local people living close to protected areas consume more meat meals during the period when the wildebeest are in the village proximities than when the herds are far into the southern plains. This further proves the rationality of illegal bushmeat hunters when planning for hunting trips (the benefits versus cost). In contrast, the fish meals in the villages located close to protected areas but far from the Lake Victoria decrease with influx of migratory herbivores, which suggest that fish and meat complement each other when the distance from the sources fluctuate (because fish must be bought and expensive when compared to bushmeat that may be obtained freely or very cheap from the illegal bushmeat hunters during the influx of migratory herbivores).

Local people switch to more available and inexpensive sources of protein whenever the opportunity comes (Barnett, 2000; Brashares et al., 2004; Rowcliffe et al., 2005). In the distant villages (> 80 km), the meat and fish meals depended on the household income and not the movement of migratory herbivores around the Serengeti ecosystem. This was proved when test-persons from villages close, intermediate and distant from the nearest park boundary were given pieces of meat in a combination of wild ungulates and beef to rank the meat according to the perceived taste. In addition, the test-persons were requested to recognize the species of animal whose meat they tasted. Villages that were close to park boundaries as well as those in the intermediate (< 43 km) areas preferred

topi and impala, respectively. The test-persons from distant villages preferred beef to all other species, which suggests long term experience with beef as no other source of meat is locally available in the area other than livestock meat. In contrast, test-persons from the villages that are located at short or intermediate distance from the park may have different experience with different species of wild ungulates and preferred some meat over others although overall preferred meat was beef.

Conflicts over livestock depredation

Human-carnivore conflict over livestock depredation is a serious management issue often causing opposition towards conservation at a worldwide scale. The results from the current study in the villages surrounding the western Serengeti show that among the wild carnivores reported to kill livestock, 97.7% of reported species was spotted hyena, being responsible for 98.2% of total livestock loss (US \$ 12,846 in 2003) (Paper II). The killing was not only restricted to villages in the proximity of the park but also as far away as 80 km (Paper III). The most numerous large carnivore species in the Serengeti ecosystem is the spotted hyena (Mill and Hofer, 1998). Thus, it is not surprising that this is the species of carnivore villagers in the study area report to attack livestock most frequently. Given that this species is mostly a nocturnal predator, it is also to be expected that most attacks by this species occur at night. In the study area, livestock is taken out from the village to graze during the day, and then kept in enclosures or bomas, usually close to houses, at night. In addition, the nocturnal and opportunistic foraging behaviour, together with the ability of spotted hyena to take long distance commuting trips, make them particularly adaptable to anthropogenic environments (Kruuk, 1972; Hofer and East, 1993; Mills and

Hofer, 1998). Livestock keepers show no tolerance towards carnivores that kill livestock. The analysis shows that level of education associates with higher levels of tolerance, while for livestock keepers higher depredation rates is linked to approval of lethal retaliation and effective protection measures is associated with a reduced desire of retaliation. However, the negative attitudes towards large carnivores may be due to human safety as well (Kalterborn et al., 2005; Packer et al., 2005; Røskaft et al., 2007). Our study that included households in the villages that are close and distant from the park boundary indicates that goat and sheep are targeted by hyena and that more killings were recorded in the households farther away from the boundary of the park than in the households in the park proximity (Paper III). This suggests an existence of the spotted hyena in the area with high anthropogenic activities. However, the data collection period was not sufficiently long to warrant the fair comparison and because I only had one village far away from the park boundary, the results may also be a result of pseudoreplication. Moreover, the sample size in distant villages was not large and extensive enough for representing the conclusive picture on the livestock depredation. This is because the depredations recorded may involve one animal or a single group of animals.

When the level of loss due to diseases was compared to loss due to predation, diseases caused higher livestock loss in households than depredation, theft or loss while grazing. This observation is in agreement with other studies conducted elsewhere (Ogada et al. 2003; Frank et al., 2005; Kolowski and Holekamp, 2006; Holmern et al., 2007). Disease, although farmers in Africa do not consider it seriously (Mwangi, 1997), were responsible

for 3.5-7.0% livestock loss per household during the period of nine months, costing them US\$ 83.5 (Paper III). This loss is equivalent to 59.6% of the average annual household income (Borge, 2003). The cost per household of theft and poor management in the grazing field was US \$ 3.0 and US \$ 11.4, respectively. The cost of depredation recorded was higher in the household located far away from the park boundary. The depredation cost per household in the four villages was US\$ 16.5. Livestock keepers may not observe the direct effect of diseases to their livestock production due to the fact that the sick animals may be slaughtered and used as food or sold to neighbours while carnivores often consume all edible parts of a kill; leaving nothing to human consumption. Moreover, diseases often kill larger number of new born calves than adults (personal observation, 2006).

Due to poor livestock husbandry skill (records), livestock keepers may not observe this as an important loss because the capital investment in terms of veterinary service, feeding or grazing time and/or output in terms of meat or money (when sold) is relatively much lower for new born calves than for adults, although the new born calves are the future mature animals. The problem with carnivores is the level of economic loss caused in a single attack. This is because when a carnivore breaks into the livestock enclosures, usually at night (Nyahongo, 2004; Kolowski and Holekamp, 2006; Holmern et al., 2007), it may kill several adult livestock. However, since the compensation scheme that may offset some of the costs are always lacking in Tanzania, negative attitudes towards carnivores may have developed among farmers and which have resulted in retaliatory killing of carnivores in or close to village proximities (Holekamp and Smale, 1992;

Ogada et al., 2003; Dickman, 2005; Frank et al., 2005; Graham et al., 2005; Holmern et al., 2007).

MANAGEMENT RECOMMENDATIONS

Based on my findings and the experiences from other studies in similar ecosystems, I recommend the following:

- i) Improvement of primary and secondary education in village schools in rural areas with the emphasis on wildlife conservation programs in the areas adjacent to protected areas. Education may improve attitudes of local communities towards carnivores (Lindsey et al., 2005; Woodroffe et al., 2005; Røskaft et al., 2007).
- ii) This study suggest that local people would benefit from better education on animal husbandry practices and extension service to help them maintain the health of their livestock and to prevent theft and loss of livestock while grazing. I recommend that diseases control and management should be integral part of regional and national programs to limit disease transmission between livestock and wildlife and even among livestock in the villages.
- iii) To conserve the carnivores outside the protected areas, a change in wildlife policy to allow compensation when livestock are killed by wild predator may be required this could contribute to the changing of the negative attitude that exists among livestock keepers towards wild carnivore. That may reduce retaliatory killings of such carnivores commuting from protected areas or that taking refuge in the few remaining thickets, kopjes, hills and/or mountains that are located within the village areas.

- iv) I suggest that night enclosures for livestock should be improved to reduce the conflict due to livestock losses between local people and wild carnivores. However this would require considerable effort in terms of hours of work, and might require some financial investments.
- v) In order to reduce the dependence on bushmeat, alternative sources of meat protein like aquaculture together with some income generating projects such as poultry and horticulture need to be considered in both general local as well as national development planning.
- vi) The contribution of fish to household diet as an alternative to bushmeat should be emphasized so that the limitation on processing fresh fish and transportation to local market is solved. Industrial harvesting of fish from Lake Victoria need to be coordinated as it may reduce availability of fish to the local markets in both villages located close to the lake and the distant villages from the lake and thus increasing pressure and reliance on bushmeat around the lake region.
- vii) The increased reliance on bushmeat may have negative impacts on the resident herbivore populations. Thus, the policy makers need to understand the link and the need for coordinated management between these two ecologically very different resources; the bushmeat species and the fish.
- viii) The price of beef should be reduced and wildlife management somehow should manage to limit bushmeat supply (preferably by cooperating actively with communities) so that many people may choose to eat more beef rather

than wild ungulates. This will inevitably reduce the hunting pressure on resident herbivores.

- ix) The findings from this thesis suggest the need for special conservation attention to resident herbivore populations close to village proximities. Otherwise the long term harvest and uncontrolled illegal bushmeat hunting based on current meat preferences and habitat location may seriously deplete the resident herbivore species from their key habitats.

FUTURE RESEARCH NEEDS

Future studies on the effect of bushmeat processes before transportation to the market place is recommended. This is important because the bushmeat consumers may be used to sun-dried meat that may influence the fatty aroma and the texture in different levels among the different species.

Further research on the coping strategies for supplementing the low meat protein consumption in the villages located farther from the park needs to be carried out. This is important for advising local communities living close to the park boundaries on such alternative sources of protein so that they reduce reliance on bushmeat

The findings from this study suggest further research on the population ecology of wild carnivores outside the protected areas in order to establish the current coping strategy of these carnivores and possible current spatial-temporal conflicts resulting from the interactions between human and the carnivores.

Further research on the nutritional contribution of bushmeat to local communities in the western Serengeti and the consequences on both bushmeat species and humans in the future depending on the increasing human population is recommended.

Study on the alternative income generating projects that are socio-culturally acceptable and environmentally friendly such as beekeeping, poultry, aquaculture and horticulture is recommended in order to help in alleviating poverty among local people in the western Serengeti.

REFERENCES

- Arcese, P., Hando J., Campbell, K., 1995. Historical and Present – Day Anti-poaching Efforts in Serengeti In: Serengeti II – Dynamics, Management and Conservation of an Ecosystem (Eds. A.R.E, Sinclair and P. Arcese). The University of Chicago Press, 506-533.
- Baldus, R.D., 2004. Lion conservation in Tanzania leads to serious human-lion conflicts with case study of a man-eating lion killing 35 people. Tanzania Wildlife Discussion Paper, 4, 1-63.
- Barnett, R., 2000. Food for Thought. The utilization of wild meat in Eastern and Southern Africa. TRAFFIC East/Southern Africa, pp. 264. Nairobi, Kenya.
- Brand, D.J., Nel, J.A.J., 1997. Avoidance of cynide gums by black-backed jackals. Applied Animal Behaviour Science, 55, 177-182.
- Brashares, J.S., Arcese, P., Sam, M.K., 2001. Human demography and reserve size predict wildlife extinction in West Africa. Proceedings of Royal Society of London, 268, 2473-2478.
- Brashares, J.S., Arcese, P., Sam, M.K., Coppolillo, P.B., Sinclair, A.R.E., Balmford, A., 2004. Bushmeat hunting, wildlife decline and fish supply in West Africa. Science, 306, 1180-1183.
- Borge, A., 2003. Essays on the economics of African wildlife and utilization and management. Dr. Polit. Thesis, University of Trondheim, NTNU, 187 pp.
- Campbell, K. L. I, Hofer, H., 1995. People and Wildlife: Spatial Dynamics and Zones of Interaction. In: Serengeti II – Dynamics, Management and Conservation

- of an Ecosystem (Eds. A.R.E., Sinclair and P. Arcese). The University of Chicago Press, 534-570.
- Campbell, K.L.M., Nelson, V., Loibooki, M., 2001. Sustainable use of wildlife resources: ecological, economics and social interactions. Natural Resources Institute, University of Greenwich, Main Report.
- Ceballos, G., Ehrlich, P.R., 2005. Global mammal distribution, biodiversity hotspots and conservation. (<http://www.pnas.org/cgi/reprint/103/51/19374> (accessed on 16-09-2007)).
- Conover, M., 2002. Resolving human-wildlife conflicts: the science of wildlife damage management. CRC Press, Boca Raton, Florida, 418 pp.
- Dickman, A.J., 2005. An assessment of pastoralist attitudes and wildlife conflict in the Rungwa-Ruaha region, Tanzania, with particular reference to large carnivores. Msc. thesis. University of Oxford.
- Dey, S.C., 1991. Depredation by wildlife in the fringe areas of North Bengal forests with special reference to elephant damage. *Indian Forester*, 117, 901-908.
- East, M.L., Hofer, H., 2000. The impact of bush-meat hunting on non-target species: Commuting spotted hyenas, *Crocuta crocuta*, in the Serengeti. African Wildlife Management in the New Millenium Conference. College of African Wildlife Management, Mweka, Tanzania 13–15 December 2000
- Fanshawe, J.H., Frame, L.H., Ginsberg, J.R., 1991. The wild dog – Africa's vanishing carnivores. *Oryx*, 25, 137-146.

- FitzGibbon, C.D., Mogaka, H., Fanshawe, J.H., 1996. Subsistence hunting and mammal conservation in Kenyan coastal forest: resolving a conflict. The exploitation of mammal population. Chapman & Hall London. U.K.
- Frank, L.G., Woodroffe, R., Ogada, M.O., 2005. People and predators in Laikipia District, Kenya. In: people and wildlife conflicts or coexistence? (Eds. R. Woodroffe, S. Thirgood, A. Rabinowitz), Cambridge University Press, Cambridge, pp. 286-304.
- Gasagway, W.C., Boertje, R.D., Grangaard, D.V., Kelleyhouse, D.G., Stephenson, R.O., Larsen, D.G., 1992. The role of predation in limiting moose at low densities in Alaska and Yukon and implications for conservation. *Wildlife Monographs*, 120, 1-59.
- Gifford-Gonzalez, D. 2000. Animal disease challenges to the emergence of pastoralism in sub-Saharan Africa. *African Archaeological Review*, 17 (3), 95-139.
- Ginsberg, J. R., Macdonald, D. W., 1990. *Foxes, Wolves, Jackals and Dogs: An Action Plan for the Conservation of Canids*. IUCN, Gland, Switzerland.
- Gittleman, J.L., Funk, S.M., Macdonald, D.W., Wayne, R.K (eds), 2001. *Carnivore conservation*. Conservation Biology Series. Cambridge University Press, Cambridge, U.K.
- Graham, K., Beckerman, A.P., Thirgood, S., 2005. Human-predator-prey conflicts: ecological correlates, prey losses and patterns of management. *Biological Conservation*, 122, 159-171.
- Herero, S., 1985. *Bear attack: their causes and avoidance*. Nick Lyons Book, New York.

- Hoare, R., 1999. Determinants of human-elephant conflict in a land use mosaic. *Journal of Applied Ecology*, 36, 689-700.
- Hudson, P.J., Rizzoli, A., Grenfell, B.T., Heesterbeek, H., Dobson, A.P., 2002. *The ecology of wildlife diseases*. Oxford University Press, Oxford, U.K.
- Hofer, H., Campbell, K. L. I., East, L.M., Huish, S.A., 1996. The impact of game meat hunting on target and non - target species in the Serengeti. In: *the exploitation of mammal population*. Chapman & Hall London. U.K.
- Hofer, H., Campbell, K.L.I., East, M.L., Huish, S.A., 2000. Modelling the spatial distribution of the economic costs and benefits of illegal game meat hunting in the Serengeti. *Natural Resource Modelling*, 13, 151 – 177.
- Hofer, H., East, M.L., 1993. The commuting system of Serengeti spotted hyenas: how a predator copes with migratory prey. I. Social organization. *Animal Behaviour* 46, 547-557.
- Holekamp, K. E., Smale, L., 1992. Human-hyena relations in and around the Maasai Mara National Reserve, Kenya. *IUCN SSC Hyena Specialist Group Newsletter* 5: 19-20.
- Holmern, T., Johannesen, A.B., Mbaruka, J., Mkama, S., Muya, J., Røskaft, E., 2004. *Human-Wildlife Conflicts and Hunting in the Western Serengeti, Tanzania*,
- Holmern, T., Nyahongo, J., Røskaft, E., 2007. Livestock loss caused by predators outside the Serengeti National Park, Tanzania. *Biological Conservation*, 135, 534-542.

- IUCN The World Conservation Union, 2006. IUCN Red List of Threatened Species: A Global Assessment. Gland, Switzerland.
- Jenkins, S.R, Perry, B.D., Winkler, W.G., 1998. Ecology and epidemiology of raccoon rabies. *Reviews of Infectious Diseases*, 10, 620-625.
- John, E.F.A., Currie, D., Meeuwig, J., 2003. Bushmeat and food security in the Congo Basin: linkages between wildlife and people's future. *Environmental Conservation*, 30, 71-78.
- Johnson, W.E., Eizirik, E., Lento, G.M., 2001. The control, exploitation and conservation of carnivores. In: *Carnivore Conservation*. (Eds. J.L. Gittleman, S.M. Funk, D.W. Macdonald, R.K. Wayne). Cambridge University Press, Cambridge, U.K., pp. 196-216.
- Kaltenborn, B. P., Bjerke, T., Vittersø, J., 1999. Attitudes towards large carnivores among sheep farmers, wildlife managers and research biologists in Norway. *Human Dimensions of Wildlife*, 4, 57-73.
- Kaltenborn, B.P., Nyahongo, J.W., Tingstad, M.K, 2005. The nature of hunting around the western corridor of Serengeti National Park, Tanzania. *European Journal of Wildlife Research*, 51, 213-222.
- Kellert, S. R., 1985. Public Perception of Predators, Particularly the Wolf and Coyote. *Biological Conservation*, 31: 167-189.
- Kock, R.A., 2003. What is this infamous "wildlife/livestock diseases interface?" A review of current knowledge for the African continent. PACE Epidemiology Unit, AU/IBAR, Nairobi, Kenya. <http://wcs-ahead.org/book/chapter01.pdf> (accessed 03-09-2007).

- Kolowski, J.M., Holekamp, K.E., 2006. Spatial, temporal, and physical characteristics of livestock depredations by large carnivores along a Kenyan reserve border. *Biological Conservation* 128, 529-541.
- Krebs, C.J., Boutin, S., Boonstra, R., Sinclair, A.R.E., Smith, J.N.M., Dale, M.N.R. Martin, K., Turkington, R. 1995. Impact of food and predation on the snowshoe hare cycle. *Science*, 269, 1112-1118.
- Kruuk, H., 1972. *The spotted hyena. A study of Predation and Social Behavior.* University of Chicago Press. 335pp.
- Leader-Williams, N., Albon, S.D, 1988. Allocation of resources for conservation. *Nature*, 336, 533-535.
- Lindsey, P.A., du Toit, J.T., Mills, M.G.L., 2005. Attitudes of ranchers towards African wild dogs *Lycaon pictus*: Conservation implications on private land. *Biological Conservation* 125, 113-121.
- Logan, K.A. and Sweanor, L.L., 2001. *Desert Puma: evolutionary ecology and conservation of an enduring carnivore.* Island Press, Washington D.C.
- Loibooki, M.T., Hofer, H., Campbell, K.L.I., East, M.L., 2002. Bush meat Hunting by communities adjacent to the Serengeti National Park, Tanzania: the importance of livestock ownership and alternative sources of protein and income. *Environmental Conservation*, 29, 391-398
- Løe, J., E. Røskoft, 2004. Large carnivores and human safety: A review. *Ambio*, 33, 283-288.
- Marker, L., 1998. Current status of cheetah (*Acinonyx jubatus*). In: *A symposium on cheetahs as game ranch animals.* (Eds. B.L. Penzhorn). Wildlife group of

- South African Veterinary Association, Onderstepoort, South Africa, pp. 1-17.
- Marks, S.A., 1973. Prey selection and annual harvest of game in a rural Zambian community. *East African Wildlife Journal*, 11, 113-128.
- Marks, S.A., 1994. Local Hunters and Wildlife survey: a design to enhance participation. *African Journal of Ecology*, 32, 233-254.
- Mills, G., Hofer, H., 1998. Hyena: Status Survey and Conservation Action Plan. IUCN/SSC Hyena Specialist Group. Information Press, Oxford, UK.
- Mishra, C., 1997. Livestock depredation by large carnivores in the India trans-Himalaya: conflict perceptions and conservation prospects. *Environmental Conservation*, 24, 338-343.
- Mwangi, N.S., 1997. A cost-benefit analysis of livestock predation on the group ranches in the dispersal areas of Maasai Mara National Reserve, Kenya. Mphil. thesis, Moi University.
- Naughton-Treves, L., 1998. Predicting patterns of crop damage by wildlife around Kibale National Park, Uganda. *Conservation Biology*, 12, 156-168.
- Norton-Griffiths, M., Herlocker, D., Pennycuik, L., 1975. The pattern of rainfall in the Serengeti ecosystem, Tanzania. *East African Wildlife Journal*. 13, 347-374.
- Nowell, K., Jackson, K., 1996. Wild Cats: Status Survey and Conservation Action Plan. Burlington Press, Cambridge.

- Nyahongo, J., W., 2004. Impact of human activities on the carnivore populations in the western corridor of the Serengeti Ecosystem. MSc. Thesis, University of Dar es Salaam, Tanzania. 147 pp.
- Ogada, M.O., Woodroffe, R., Ogue, N.O., Frank, L.G., 2003. Limiting depredation by African carnivores: the role of livestock husbandry. *Conservation Biology*, 17, 1521-1530.
- Packer, C., Ikanda, D., Kissui, B., Kushnir, H., 2005. Lion attacks on humans in Tanzania. Understanding the timing and distribution of attacks on rural communities will help to prevent them, 436, 927-928.
- Quammen, D., 2003. *Monster of God: the man-eating predator in the jungles history and the mind*. W.W. Norton, New York.
- Rowcliffe, M.J., Milner-Gulland, J.E., Cowlishaw, G., 2005. Do bushmeat consumers have other fish to fry? *Trends in Ecology and Evolution*, 20, 274-276.
- Rusch, G.M., Stokke, S., Røskaft, E., Mwakalebe, G., Wiik, H., Arnemo, J.M, Lyamuya R., 2005. Human-wildlife interactions in western Serengeti, Tanzania: Effects of land management on migratory routes and mammal population densities. NINA Report 85.
- Røskaft, E., Bjerke, T., Kaltenborn, B.P, Linnell, J.D.C., Anderson, R., 2003. Pattern of self-reported fear towards large carnivores among the Norwegian public. *Evolution and Human Behaviour*, 24, 184-198.
- Røskaft, E., B. Händel, T. Bjerke, Kaltenborn, P.B., 2007. Attitudes of humans towards large carnivores in Norway. *Wildlife Biology*, 13, 172-185.

- Saberwal, V.K., Gibbs, J.P., Chellam, R., Johnsigh, A.J.T., 1994. Lion – Human Conflict in the Gir Forest, India. *Conservation Biology*, 8, 501-507.
- Sillero-Zubiri, C., Laurenson, M.K., 2001. Interactions between carnivores and local communities: conflict or co-existence? In: *Carnivore Conservation*. (Eds. J.L. Gittleman, S.M. Funk, D.W. Macdonald, R.K. Wayne). Cambridge University Press, Cambridge, U.K., pp. 107-122.
- Sinclair, A.R.E., 1995. Serengeti past and present. In: *Serengeti II - Dynamics, Management, and Conservation of an Ecosystem* (Eds. A.R.E. Sinclair, P. Arcese). The University of Chicago Press. Chicago, pp. 3-30.
- Stuart, C. T., Macdonald, I. A. W., Mills, M. G. L., 1985. History, Current Status and Conservation of Large Mammalian Predators in Cape Province, Republic of South Africa. *Biology Conservation*, 31, 7-19.
- Terborgh, J., Lopez, L., Nunez, P., Rao, M., Shahabudin, G., Orihuela, G., Riveros, M., Ascanio, R., Adler, G.H., Lambert, T.D., Balbas, L., 2002. Ecological meltdown in predator-free forest fragments. *Science*, 294, 1923.
- Thirgood, S.J. Mosser, A., Tham, S., Hopcraft, J.G.C., Mwangomo, E., Ikengeya, T., Kilewo, M., Fryxell, J.M., Sinclair, A.R.E., Borner, M., 2004. Can park protect migratory ungulates? The case of Serengeti wildebeest. *Animal Conservation*, 7, 113-120.
- Thirgood, S., Redpath, S., Rothery, P., Aebischer, N., 2000. Raptor predation and population limitation in red grouse. *Journal of Animal Ecology*, 69, 504-516.

- Treves, A., Naughton-Treves, L., 2005. Evaluating lethal control in the management of human-wildlife conflict. In people and wildlife: conflict or coexistence? (Eds. R. Woodroffe, S. Thirgood, A. Rabinowitz).
- UNDP, 2002. World Population Development Program. www.hdr.undp.org/ (accessed on 16-09-2007).
- URT United Republic of Tanzania, 2002. 2002 population and housing census. <http://www.tanzania.org.tz/census/index.html> (accessed 30.03.2007).
- Woodroffe, R., 2000. Predators and people: Using human densities to interpret declines of large carnivores. *Animal Conservation*, 3, 165-173.
- Woodroffe, R., Lindsay, P., Romanach, S., Stein, A., Ole Ranah, S.M.K., 2005. Livestock predation by endangered African wild dogs (*Lycaon pictus*) in northern Kenya. *Biological Conservation* 124, 225-234.
- Woodroffe, R., Ginsberg, J., Macdonald, D., 1997. The African wild dog: status, survey and conservation action plan. IUCN, Gland.
- World Bank, 2006. World Development indicators database: Tanzania data profile. <http://devdata.worldbank.org/external/CPProfile.asp?PTYPE=CP&CCODE=TZA> (accessed 12-08-2007).

Paper I

Benefits and costs of illegal grazing and hunting in the Serengeti ecosystem

J.W. NYAHONGO¹, M.L. EAST^{2*}, F.A. MTURI³ AND H. HOFER²

¹Tanzania Wildlife Research Institute, PO Box 661, Arusha, Tanzania, ²Leibniz-Institute for Zoo and Wildlife Research, Alfred-Kowalke-Strasse 17, D-10315 Berlin, Germany, and ³Department of Zoology and Marine Biology, University of Dar es Salaam, PO Box 35091, Dar es Salaam, Tanzania

Date submitted: 10 December 2004 Date accepted: 11 January 2006

SUMMARY

Two forms of natural resource use (meat hunting and livestock grazing) were investigated at three sites in the western region of the Serengeti ecosystem, Tanzania. Statutory management of natural resources in this region was designated as National Park, Game Reserve or village council. A quasi-experimental design examined factors likely to alter the cost and benefit of illegal use by ranking areas within sites in relation to these factors. Factors likely to alter costs were the chance of arrest, determined by the presence or absence of guard posts, and the distance travelled to the site of exploitation. As all sites experienced large fluctuations in the density of migratory herbivores, it was assumed that the benefit acquired from hunting increased with wild herbivore density. Marked seasonal changes in precipitation were considered likely to alter the value of forage and water to livestock owners. Hunting effort (density of snares) increased as the density of wild herbivores increased. The distribution of hunting effort across sites was more consistent with the prediction that high travel costs were more likely to curtail hunting than a high potential cost of arrest. Unlike hunters, livestock owners mostly avoided the use of resources in protected areas probably because of the high potential cost of arrest and confiscation of stock. Natural resources within protected areas were exploited when benefits outweighed likely costs.

Keywords: illegal hunting, livestock grazing, natural resources, Serengeti ecosystem

INTRODUCTION

Hunting of wildlife to obtain meat for subsistence or trade is important to local economies and a growing problem for wildlife managers in many countries (Arcese *et al.* 1995; Campbell & Hofer 1995; Fa *et al.* 1995; Barnett 2000; Loibooki *et al.* 2002; Rao & McGowan 2002). The extent to which wildlife populations in Africa are used for meat is high in terms of the number of animals killed and the volume of

meat obtained (Hofer *et al.* 1996; Mduma *et al.* 1998; Noss 1998; Barnett 2000). This offtake is mainly achieved through the use of inexpensive methods of prey capture, such as wire snares, self-made traps and poisoned darts or arrows (Turner 1987; Noss 1998), and the use of non-selective capture methods such as snares has a negative impact on populations of non-target species (Hofer *et al.* 1993). The most ubiquitous hunting method is the wire snare, probably because snares cost little and are relatively simple to make; thus hunters can afford to own and set numerous snares. Once set, snares are inconspicuous and law enforcers in areas where hunting is illegal find them difficult to detect.

Use of forage and water can produce conflict between managers of protected areas and local communities (Fleischner 1994; Arcese *et al.* 1995; Homewood *et al.* 2001; Madhusudan 2004; Mishra *et al.* 2004). In comparison to illegal hunting with snares, livestock ownership requires greater financial expenditure and the illegal presence of livestock in protected areas is more difficult to conceal.

The Serengeti ecosystem straddles the international border between Tanzania and Kenya. The major part of the ecosystem lies within the Serengeti National Park (Serengeti NP) where hunting of wildlife and grazing of livestock are prohibited. Situated along sections of the Serengeti NP boundary are game reserves, where licensed hunting is permitted but livestock and unlicensed hunting are prohibited. These reserves form a buffer zone between the Serengeti NP and surrounding communities.

Given that over one million people live within 45 km of the western boundary of the Serengeti NP and associated reserves (Campbell & Hofer 1995), and that the main occupation in the area is subsistence farming plus the rearing of livestock (Loibooki *et al.* 2002), it is perhaps not surprising that natural resources within the protected areas are used by local communities (Arcese *et al.* 1995; Campbell & Hofer 1995; Hofer *et al.* 1996; Loibooki *et al.* 2002). The level of illegal hunting for meat is considerable and has resulted in the local extinction of resident herbivores in some areas (Arcese *et al.* 1995; Campbell & Hofer 1995; Hofer *et al.* 1996). Livestock ownership is viewed as a symbol of wealth and status, and inhabitants of villages close to the Serengeti NP that either own livestock or have access to alternative means to generate income and acquire protein are less likely to participate in illegal hunting (Loibooki *et al.* 2002). The link between poverty and illegal meat hunting is also reflected by the fact that illegal hunters arrested in the Serengeti NP were

*Correspondence: Dr Marion L. East Tel: +49 30 5168512 Fax: +49 30 5168735 e-mail: east@izw-berlin.de

predominantly poorly educated, young males that owned few or no livestock (Loibooki *et al.* 2002).

This study aims to build on previous research in the Serengeti ecosystem on how costs and benefits of illegal hunting influence the spatial and temporal distribution of this illegal activity (Arcese *et al.* 1995; Campbell & Hofer 1995; Hofer *et al.* 1996, 2000; Loibooki *et al.* 2002). Here we compare illegal hunting and illegal livestock grazing to investigate whether the spatial distribution of these activities is consistent with the expectation that natural resources within protected areas will be exploited when likely benefit exceeds estimated cost and to assess which component of cost is likely to matter the most.

METHODS

Study area

The study was conducted in the western section of the Serengeti (Tanzania). The economy of local communities was mainly based on subsistence agriculture with more prosperous farmers owning herds of livestock (Loibooki *et al.* 2002). An average herd of livestock in 2001 consisted of 17 animals that had a total sales value of US\$ 423–735 (Loibooki *et al.* 2002). Inhabitants of villages close to Lake Victoria practised commercial fishing, and those in villages close to all-weather roads practised commercial trade (Loibooki *et al.* 2002).

Illegal hunters from local communities chiefly used wire snares to capture wild herbivores for meat that was typically dried before being carried on foot from protected areas (Arcese *et al.* 1995; Hofer *et al.* 2000). Dried meat was used for home consumption, sold to generate income or bartered for other commodities (Hofer *et al.* 2000; Loibooki *et al.* 2002). An estimated 53 000 people are involved in illegal hunting, including both hunters and porters that transport meat from hunting camp out of the protected areas (Loibooki *et al.* 2002). Hunters arrested in the Serengeti NP come from villages within 45 km of the boundary of the protected areas (Campbell & Hofer 1995). Although a large proportion of the annual offtake of meat from the ecosystem is obtained from large migratory species such as wildebeest *Connochaetes taurinus* and zebra *Equus burchelli*, considerable volumes of meat are also obtained from other migratory and resident herbivores species (Arcese *et al.* 1995; Hofer *et al.* 1996). It is not known what proportion of illegally hunted meat is sold for cash, used for home consumption, or bartered for other commodities. For this reason it is difficult to estimate the monetary value of this illegally acquired commodity to the local economy, even though it is undoubtedly important economically (Loibooki *et al.* 2002). If only a third of the estimated annual offtake of approximately 11 950 tonnes of useable meat (Hofer *et al.* 1996) from migratory and resident herbivore species is sold (at a value of US\$ 0.3 per kg fresh weight of meat; Loibooki *et al.* 2002), trade in illegal meat would annually generate more than US\$ 1 million.

Density of wild herbivores, livestock and snares

Between May 2001 and March 2002, data were collected along ground transects in three areas, namely Kirawira, Mihale and Ndabaka. The Kirawira transect was entirely within the Western Corridor section of the Serengeti NP, two ranger posts both within 1 km of the transect being staffed by a total of 14 rangers. From each ranger post, six rangers patrolled by vehicle and on foot, and one ranger provided patrols with radio communication. There were frequent tourist vehicles in the area, many of which could communicate by radio with the ranger posts. This site was a greater distance from the boundary of the protected areas than the other two study sites. The Mihale transect was on the northern side of the Western Corridor that traversed an equal distance of the Serengeti NP, the Grumeti Game Reserve (Grumeti GR) and the unprotected area outside this Reserve. The Serengeti NP section of this transect was at a greater distance from the protected area boundary than the section of this transect in the Grumeti GR. The nearest ranger post was approximately 15 km from this transect. Grazing of livestock and unlicensed meat hunting were prohibited in the Grumeti GR. Although licensed trophy hunting was permitted within the Reserve, during the study no trophy hunters operated and tourists rarely visited the area. Natural resources outside the Grumeti GR could be legally exploited. The Ndabaka transect was on the southern side and at the western end of the Western Corridor, within 3 km of a ranger post and entrance gate to the Serengeti NP that was staffed by five rangers (three patrolled, one administered the entrance gate, and one was responsible for radio communications). Two-thirds of this transect was within the Serengeti NP and one third was in unprotected land outside the Park. As there was no reserve to act as a buffer zone between the Park and local communities, the distance from the boundary to the section of this transect inside the Park was small.

Each of the three study sites contained a 45-km transect composed of three parallel 15-km transects situated 4 km apart. Transect lines and the location of the National Park and Game Reserve boundaries along transects were determined by a global positioning system (GPS; Garmin 12 XL). A vehicle with a driver and an observer was slowly driven along each transect. In the three study sites, each 45-km transect was driven three times per month for 11 months. The numbers of livestock (cattle, goats, sheep and donkeys) and wildlife observed within 200 m either side of the transect line were counted and the GPS positions recorded. The herbivorous species counted during transects included wildebeest, zebra, eland *Taurotragus oryx*, Thomson's gazelle *Gazella thomsoni*, Grant's gazelle *Gazella granti*, topi *Damaliscus lunatus*, impala *Aepyceros melampus*, buffalo *Syncerus caffer*, giraffe *Giraffa camelopardalis*, kongoni *Alcelaphus buselaphus*, and warthog *Phacochoerus aethiopicus*. All these species are hunted and can be caught by wire snares.

Snares within 20 m of either side of a transect line were recorded. The GPS position of snares was taken and

snare were inconspicuously marked with a permanent pen to prevent recounting previously logged snares at a later date.

The densities of wild herbivores, snares and livestock were calculated for each study site using the equation (Caughley & Sinclair 1994):

$$D = \Sigma x / \Sigma A, \quad (1)$$

where D is the calculated mean density of livestock and/or hunting equipment counted, Σx is the sum of mean livestock and/or hunting equipment counted per month, and ΣA is the sum of the mean area covered during the count.

All three sites experienced a similar pattern of precipitation, with the majority of the annual precipitation falling between November and May (the 'wet season') and little precipitation between June and October (the 'dry season').

Mihale village was approximately 5 km from the Mihale transect and, in 2001, contained 1036 people that owned 0.53 sheep or goats per person and 0.65 cattle per person. Mwabayanda village was within 5 km of the Ndabaka transect and, in 2001, this village contained 2771 people that owned 0.50 sheep or goats per person and 0.99 cattle per person. These two villages were of roughly similar size and were comparable in the number of livestock owned per head, which is an index of village wealth (Loibooki *et al.* 2002).

We applied a quasi-experimental design to investigate the relative effects of different factors likely to influence the profitability of illegal activities in different areas, and the same activities conducted outside protected areas. We chose hunting of wild herbivores for meat as a form of resource use known to yield considerably greater benefits when practised inside protected areas (Campbell & Hofer 1995; Hofer *et al.* 2000; Loibooki *et al.* 2002), and contrasted this with grazing and watering of cattle, which are activities unlikely to yield larger immediate benefits when conducted inside protected areas rather than outside such areas. We assumed that the likely benefits of illegal hunting would increase with increasing wild herbivore density and used natural fluctuations in wild herbivore density to test this assumption. We assumed that the value of forage and water resources to livestock owners would increase during periods of low precipitation (dry season), and that herds of livestock would be more easily detected by law enforcers than snares.

As a model of economic costs and benefits of illegal hunting in the Serengeti indicated that travel cost (calculated as the time taken to travel to and from a hunting site multiplied by the opportunity cost of travel) is more important in determining the spatial distribution of hunting activities than the cost of arrest based on penalties incurred if arrested (Hofer *et al.* 2000), our analysis is based on the expectation that travel costs increased with distance travelled, and that the chance of arrest was greater close to ranger posts than in areas without such posts (Campbell & Hofer 1995; Hofer *et al.* 2000). We selected study areas that varied with respect to both potential costs and benefits and predicted that use should occur where costs are

perceived to be low, and where returns from exploitation are likely to outweigh cost.

Statistical analysis

Statistical analyses were performed using SYSTAT 10 (Wilkinson 2000). As data were not normally distributed, non-parametric tests were applied. For all tests $p < 0.05$ (two-tailed) was considered significant. Densities are presented as mean \pm standard error (SE).

We used density of wild herbivores as one index of the potential benefit hunters might gain and the density of snares as an index of the effort exerted by hunters in an area. We used a post-hoc Kruskal-Wallis ANOVA test (Conover 1980) to compare predicted levels of hunting effort in different areas in relation to the cost of travel and cost of law enforcement. The areas considered were: Kirawira NP, Mihale NP, Mihale GR, Ndabaka NP and Ndabaka outside protected area. The unprotected area of the Mihale transect was excluded from this analysis, as no wild herbivores were observed in this area.

RESULTS

Livestock densities

In accordance with our predictions, no livestock were recorded along the Kirawira transect. The density of livestock within the protected area section (Serengeti NP and Grumeti GR) of both the Ndabaka and Mihale transects was lower (Ndabaka 2.8 ± 0.9 animals per km^2 ; Mihale 3.4 ± 1.4 animals per km^2) than the high density of livestock legally grazed outside the Serengeti NP on the Ndabaka transect (38.0 ± 4.1 animals per km^2 ; Wilcoxon Signed Rank $Z = -2.934$, $n = 11$, $p = 0.004$; Fig. 1a) and outside the Grumeti GR on the Mihale transect (35.3 ± 7.2 animals per km^2 ; $Z = -2.934$, $n = 11$, $p = 0.004$; Fig. 1b). These results indicate that herders knew the location of the Park and Reserve boundaries and mostly avoided taking their livestock into protected areas.

During the dry season, the density of livestock within the Serengeti NP and Grumeti GR along the Mihale transect was higher (7.52 ± 1.80 animals per km^2) than along the Serengeti NP section of the Ndabaka transect (0.68 ± 0.42 animals per km^2 ; Mann Whitney $U = 3.0$, $p = 0.007$, $n = 6$).

During the wet season, livestock was absent from the protected sections of the Mihale transect, but low densities of livestock were present in the Serengeti NP section of the Ndabaka transect (4.70 ± 1.22 animals per km^2), mostly between the Park boundary and the Mbalageti River.

Wildlife densities and illegal hunting effort

When the possible benefit to illegal hunters was scored in terms of wild herbivore densities, the Serengeti NP section of the Ndabaka transect was likely to yield the highest level of benefit (66.93 ± 17.06 animals per km^2). Moderate levels of benefit were likely from the Serengeti NP sections in the Kirawira and




Figure 1 Mean monthly (May 2001–March 2002) livestock density per km² in (a) the Mihale transect, and (b) the Ndabaka transect. Solid bars = livestock outside the Serengeti NP and Grumeti GR; open bars = livestock inside the Serengeti NP and Grumeti GR.

Mihale transects respectively (31.20 ± 12.01 animals per km²; 25.95 ± 5.99 animals per km²), and the Grumeti GR section of the Mihale transect (22.65 ± 11.39 animals per km²). Low levels of benefit were likely from the unprotected sections of the Ndabaka (2.77 ± 0.01 animals per km²) and Mihale transects (no animals observed).

Combined data from the Serengeti NP sector of the Mihale and Ndabaka transect displayed the expected positive correlation between the mean monthly density of snares (hunting effort; Table 1) and the mean monthly density of herbivores




Figure 2 Plot of density of snares per km² against density of wild herbivores per km². (○) Serengeti NP section of the Mihale transect, (△) Grumeti GR section of the Mihale transect, (●) Serengeti NP section of the Ndabaka transect. Unprotected areas that contained few or no wild herbivores, and the Kirawira transect that contained no snares, not included.

(combined data from the Mihale and Ndabaka transects, Spearman Rank Correlation $r = 0.641$, $n = 22$, $p = 0.002$). The expected positive correlation between the mean monthly density of snares and the mean monthly density of herbivores was not found in the Grumeti GR section of the Mihale transect (Fig. 2; Spearman's $r = 0.20$, $n = 11$, not significant). Despite high densities of wild herbivores in Serengeti NP at Kirawira (Table 1, Fig. 3a), no evidence of illegal hunting (no snares, pitfall traps, or fences) was recorded along this transect. Owing to a very low density of wild herbivores outside protected areas (Fig. 3b,c), hunters could expect very poor returns and thus snares were rarely set in these areas.

Factors influencing illegal hunting effort

When areas were ranked according to their likely travel costs, and the hunting effort in these areas was predicted according to these ranks (Table 1), all pairwise comparisons of observed hunting effort between areas conformed to predictions, except for the comparison between the Serengeti NP section of Mihale and Kirawira, for which medium and low hunting efforts were predicted but equally low levels were observed in both areas (Table 1). These results indicate that travel is an important cost factor for hunters.

In contrast, when hunting effort was predicted on the basis of the likely chance of arrest (Table 1), then all pairwise comparisons of observed hunting effort between areas showed differences, however all differences except one were in the opposite direction to that expected. In particular, observed

Table 1 Observed and predicted hunting effort (snare density per km²) based on travel costs and chance of arrest at the Kirawira, Ndabaka and Mihale transects (NP: National Park, GR: Game Reserve, unprotected: area outside both NP and GR). Results of post-hoc comparisons following Kruskal-Wallis test on observed hunting effort data; different letters indicate significant differences between sites, ? = unknown because hunting not observed.

<i>Factor</i>	<i>Ndabaka-unprotected</i>	<i>Mihale GR</i>	<i>Ndabaka NP</i>	<i>Mihale NP</i>	<i>Kirawira NP</i>
Observed hunting effort	0.00 ± 0.00	5.37 ± 1.70 ^a	11.23 ± 2.97 ^a	0.45 ± 0.23 ^b	0.00 ± 0.00 ^b
Travel cost	none	small	small	medium	high
Hunting effort predicted by travel cost	?	high	high	medium	low
Chance of arrest	none	small	medium	small	high
Hunting effort predicted by chance of arrest	?	high	medium	high	low

hunting effort in the Grumeti GR section of Mihale was significantly higher than in the NP section of this site when they were predicted to be equal, and, despite the presence of a guard post in the Serengeti NP sector of Ndabaka, the observed hunting effort was significantly higher than in the Serengeti NP sector of Mihale. These results indicate that the likely chance of arrest is perceived by hunters to be low and thus the potential costs associated with arrest do not have the expected influence on hunting effort. The only comparison that followed the expected direction was between the Serengeti NP section of Ndabaka and Kirawira (Table 1).

Seasonal changes in the densities of wild herbivores

Large fluctuations in the mean monthly densities of wild herbivores in each transect (Fig. 3) were caused by the migratory movements of wildebeest and zebra. High densities of wild herbivores were observed in Kirawira in June (Fig. 3a), in the Serengeti NP section of the Ndabaka transect between November and March (Fig. 3b), and in the Serengeti NP and Grumeti GR section of the Mihale transect in July, August, October, December and January (Fig. 3c). Neither resident nor migratory wild herbivores were present in the unprotected area outside the Grumeti GR along the Mihale transect (Fig. 3c) and were present only at very low densities in some months outside the Serengeti NP along the Ndabaka transect (Fig. 3b). This suggests that either the protected areas adequately encompassed the migratory routes or that migratory herds mostly avoided unprotected areas. Few resident herbivores persisted outside the protected areas, suggesting that populations of these species had been overharvested.

DISCUSSION

The results of this study are consistent with the idea that levels of illegal use of natural resources in the west of the Serengeti ecosystem were influenced by the likely value of the resources acquired and the probable costs associated with their acquisition.

Evidence of illegal hunting was found during the 11 months of this study along two of the three transects, suggesting that benefits of hunting mostly outweighed costs in these areas. Our results (Table 1) conformed to the expectation that level of illegal hunting decreased as the distance hunters travelled on foot to hunting areas increased. Travel cost is likely to be assessed not only in terms of distance travelled but also in terms of time that could be devoted to other activities (opportunity cost).

Illegal hunters mostly work at night by themselves or in small groups and use inconspicuous hunting methods. For this reason the likelihood of illegal hunting activities being detected is low, particularly in areas with dense vegetation and certain types of topography (Campbell & Hofer 1995). This may explain why high hunting effort occurred in the vicinity of the Ndabaka ranger post (Table 1, Fig. 3b). Our data are insufficient to test whether the absence of hunting effort at Kirawira was caused by the high chance of arrest afforded by two ranger posts and numerous tourist vehicles, a high travel cost to this area, or a combination of these factors. In general, the results of this study support optimality models developed for the Serengeti ecosystem that predicted that hunting would be depressed more by the cost of travel than the cost of arrest (Hofer *et al.* 1996; Hofer *et al.* 2000).

The relatively lower density of snares in the Serengeti NP section of the Mihale transect compared to that in the Grumeti GR section is most likely the result of a greater travel cost without increased returns, as herbivore densities in both areas were similar (Table 1).

The positive relationship between the density of snares and that of wild herbivores in the Serengeti NP sections of the Mihale and Ndabaka transects (Fig. 2) indicates that hunters increased their effort as the likely level of return increased. Our results cannot discern whether this was the consequence of a relatively stable number of hunters increasing their hunting effort as profitability increased, or was caused by an increase in the number of hunters setting snares in areas with high densities of herbivore, or both of these processes. The observed increase in the density of snares in areas containing high densities of herbivores was likely to be detrimental to wildlife, including non-target species (Hofer *et al.* 1993).




Figure 3 The density of wild herbivores in the (a) Kirawira transect, (b) Ndabaka transect, and (c) Mihale transect. Black bars = density in areas of the Serengeti NP; open bars = density in areas governed by a village council; black bars = density in the Grumeti GR. No wild herbivores were observed in the area governed by a village council in the Mihale transect between May 2001 and March 2002.

The highest densities of migratory herbivores recorded during the study occurred along the Ndabaka transect during the wet season (Fig. 3b). However, when herbivore densities along the Ndabaka transect were high, the density of snares in this area was lower than might have been expected, given the comparatively high snare density recorded in Grumeti GR section of the Mihale transect at far lower herbivore densities (Fig. 2). One possible explanation for this might be that travel by foot and the crossing of rivers in spate during the wet season are likely to be more costly than in the dry season, and drying illegally hunted meat for preservation and ease of transport is likely to be problematic during the wet season. Furthermore, during the wet season, villagers cultivate crops and, as Ndabaka was close to Lake Victoria, fishing may be more profitable than illegal hunting.

High densities of herbivores occurred in the Grumeti GR section of the Mihale site for a brief period of less than a month (Fig. 3c). The density of snares during this month was lower than that found in the Serengeti NP section of the Ndabaka site when similar densities of herbivores were present for several consecutive months (Fig. 2 and Fig. 3b). This indicates that hunters did not easily locate and immediately exploit large, transient herds of migratory herbivores that occupied an area for a brief period.

Overharvesting appears to have eliminated the wild herbivore populations in village managed areas outside the Grumeti GR at the Mihale site, and has decreased the wild herbivore population in village areas outside the Serengeti NP at the Ndabaka site.

During the dry season, the density of livestock within the Serengeti NP and Grumeti GR along the Mihale transect was higher than along the Serengeti NP section of the Ndabaka transect. This is probably because during the dry season the large river in the protected section of the Ndabaka transect (Mbalageti River) did not contain permanent water, and livestock owners moved their stock towards the shores of Lake Victoria where adequate forage and water were available during the dry season. Throughout the dry season the Grumeti River close to the Mihale transect did contain permanent water.

Despite high densities of livestock close to the boundary of the protected areas, domestic stock was rarely illegally present in these areas. This may indicate that livestock owners considered the chance of detection and likely financial penalties (fines or confiscation of livestock) too high in relation to the benefit gained from illegally acquired forage and the use of watering areas inside protected areas. As livestock owners are relatively wealthy members of local communities, they are likely to have less need to engage in illegal activities (Loibooki *et al.* 2001).

ACKNOWLEDGEMENTS

We thank the Messerli Foundation (Switzerland) and the Leibniz-Institute for Zoo and Wildlife Research (Berlin, Germany) for financial support, the Tanzanian Commission

of Science and Technology for permission to conduct the study, the Tanzanian Wildlife Research Institute, Tanzanian Wildlife Division and Tanzanian National Parks for cooperation and support, P.D. Mochlman for helpful comments, K.L.I Campbell for information, F.F. Banyikwa, K. Wilhelm, R. Fyumagwa, D. Thierer, H.Wiik and T. Shabani for assistance, and two referees for their helpful comments.

References

- Arcese, P., Hando, J. & Campbell, K.L.I. (1995) Historical and present – day anti-poaching efforts in Serengeti. In: *Serengeti II: Dynamics, Management and Conservation of an Ecosystem*, ed. A.R.E. Sinclair & P. Arcese, pp. 506–533. Chicago, IL, USA: University of Chicago Press.
- Barnett, R. (2000) *Food for Thought: The Utilization of Wild Meat in Eastern and Southern Africa*. Nairobi, Kenya: TRAFFIC East/Southern Africa: 264 pp.
- Campbell, K.L.I. & Hofer, H. (1995) People and wildlife: spatial dynamics and zones of interaction. In: *Serengeti II: Dynamics, Management and Conservation of an Ecosystem*, ed. A.R.E. Sinclair & P. Arcese, pp. 534–570. Chicago, IL, USA: Chicago University Press.
- Caughley, G. & Sinclair, R.E. (1994) *Wildlife Ecology and Management*, pp. 190–216. Massachusetts, USA: Blackwell Scientific Publications.
- Conover, W.J. (1980) *Practical Nonparametric Statistics*, pp. 213–338. New York, USA: John Wiley & Sons.
- Fa, J.E., Juste, J., Perez del Val, J. & Castroviejo, J. (1995) Impact of market hunting on mammal species in Equatorial Guinea. *Conservation Biology* 9: 1107–1115.
- Fleischner, T.L. (1994) Ecological costs of livestock grazing in western North America. *Conservation Biology* 8: 629–644.
- Hofer, H., Campbell, K.L.I., East, M.L. & Huish, S.A. (1996) The impact of game meat hunting on target and non-target species in the Serengeti. In: *The Exploitation of Mammal Populations*, ed. V.J. Taylor & N. Dunstone, pp. 117–146. London, UK: Chapman and Hall.
- Hofer, H., Campbell, K.L.I., East, M.L. & Huish, S.A. (2000) Modeling the spatial distribution of the economic costs and benefits of illegal game meat hunting in the Serengeti. *Natural Resource Modeling* 13: 151–177.
- Hofer, H., East, M.L. & Campbell, K.L.I. (1993) Snares, commuting hyaenas and migratory herbivores: humans as predators in the Serengeti. *Symposium of the Zoological Society, London* 65: 347–366.
- Homewood, K., Lambin, E.F., Coast, E., Kariuki, A., Kikula, I., Kivelia, J., Said, M., Serneels, S. & Thompson, M. (2001) Long-term changes in Serengeti-Mara wildebeest and land cover: pastoralism, population, or policies? *Proceedings of the National Academy of Sciences, USA* 23: 12544–12549.
- Loibooki, M.T., Hofer, H., Campbell, K.L.I. & East, M.L. (2002) Bushmeat hunting by communities adjacent to the Serengeti National Park, Tanzania: the importance of livestock ownership and alternative sources of protein and income. *Environmental Conservation* 29: 391–398.
- Madhusudan, M.D. (2004) Recovery of wild large herbivores following livestock decline in a tropical Indian wildlife reserve. *Journal of Applied Ecology* 41: 858–869.
- Mduma, S.A.R., Hilborn, R. & Sinclair, A.R.E. (1998) Limits to exploitation of Serengeti wildebeest and implications for its management. In: *Dynamics of Tropical Communities*, ed. D.M. Newbury, H.H.T. Prins & N. Brown, pp. 243–265. Oxford, UK: Blackwell Science.
- Mishra, C., Van Wieren, S.E., Ketner, P., Heitkönig, I.M.A. & Prins, H.H.T. (2004) Competition between domestic livestock and wild bharal *Pseudois nayaaur* in the Indian Trans-Himalaya. *Journal of Applied Ecology* 41: 344–354.
- Noss, A.J. (1998) The impact of cable snare hunting on wildlife populations in the forests of the Central African Republic. *Conservation Biology* 12: 390–398.
- Rao, M. & McGowan, P.J.K. (2002) Wild-meat use, food security, livelihoods, and conservation. *Conservation Biology* 16: 580–583.
- Turner, M. (1987) *My Serengeti Years*. London, UK: Elm Tree Books.
- Wilkinson, L. (2000) *Systat 10*. Chicago, IL, USA: SPSS Inc.: 197 pp.

Paper II

available at www.sciencedirect.comjournal homepage: www.elsevier.com/locate/biocon

Livestock loss caused by predators outside the Serengeti National Park, Tanzania

Tomas Holmern^{a,*}, Julius Nyahongo^{a,b}, Eivin Røskoft^a

^aDepartment of Biology, Norwegian University of Science and Technology, Realfagbygget, N-7491 Trondheim, Norway

^bTanzania Wildlife Research Institute, P.O. Box 661, Arusha, Tanzania

ARTICLE INFO

Article history:

Received 21 February 2006

Received in revised form

23 October 2006

Accepted 30 October 2006

Available online 12 December 2006

Keywords:

Carnivores

Lethal control

Livestock depredation

Spotted hyena

Tanzania

ABSTRACT

Human–carnivore conflict is a serious management issue often causing opposition towards conservation efforts. In a survey of 481 households in seven different villages outside the Serengeti National Park in Tanzania, 67.4% of respondents owned livestock and 27.4% of all the households surveyed reported losses of a total of 4.5% of their livestock to wild predators over 12 months. This loss equated to an average annual financial loss of 19.2% (US \$26.8) of their cash income. Livestock depredation was reported to be caused most often by spotted hyena (*Crocuta crocuta*) (97.7%), leopard (*Panthera pardus*) (1.6%), baboon (*Papio cynocephalus*) (0.4%), lion (*Panthera leo*) (0.1%) and lastly black-backed jackal (*Canis mesomelas*) (0.1%). Total reported losses during 2003 amounted to US \$12,846 of which spotted hyena kills were reported to account for 98.2%. The mean annual livestock loss per household (of those that reported loss) was 5.3 head of stock, which represents more than two-thirds of the local average annual cash income. Depredation by large felids occurred only in a narrow zone along the protected area (<3 km), whereas spotted hyenas killed livestock even in households located far away (>30 km). Tolerance of livestock depredation among the respondents was low. Logistic regression models indicated that education improved tolerance, while for livestock owners higher depredation rates was linked to approval of lethal retaliation and effective protection measures was associated with a reduced desire of retaliation. We recommend that further research should identify the precise causes of livestock loss and which protection measures that can reduce depredation.

© 2006 Elsevier Ltd. All rights reserved.

1. Introduction

Human population increase and technological development is rapidly reducing and fragmenting the available habitat for large carnivores. Although protected areas in principal are shielded from most human activities, the majority of African reserves are not large enough to maintain viable populations of these wide ranging species (Newmark, 1996; Woodroffe and Ginsberg, 1998). Non-protected and partially protected areas (i.e. IUCN categories < IV) therefore play a vital role in main-

taining the existence of carnivores, both in order to increase population sizes and to allow greater genetic exchange between populations (Linnell et al., 2001; Treves and Karanth, 2003).

Large carnivores differ in their ability to adapt to anthropogenic landscapes. Behavioural plasticity and traits that give ecological flexibility and allow populations to recover rapidly from depletion have been identified as important factors for persisting close to humans (Cardillo et al., 2004). For example, in the Masai Mara National Reserve in Kenya, spotted hyenas

* Corresponding author. Tel.: +47 73596093; fax: +47 73596100.

E-mail addresses: Tomas.Holmern@bio.ntnu.no (T. Holmern), nyhwjulius@yahoo.co.uk (J. Nyahongo), Eivin.Roskaft@bio.ntnu.no (E. Røskoft).

0006-3207/\$ - see front matter © 2006 Elsevier Ltd. All rights reserved.

doi:10.1016/j.biocon.2006.10.049

(*Crocuta crocuta*) changed their daily activity rhythm, demographic structure, social behaviour and use of space as a response to increased disturbance from livestock grazing (Boydston et al., 2003). Small geographic range size, long gestation period, low species population density and high trophic level are all factors associated with high extinction risk in carnivores (Cardillo et al., 2004), but despite these biological traits, large carnivore survival ultimately depends on their conflict level with human interests and their social acceptability to humans, particularly outside protected areas (Linnell et al., 2001; Kleiven et al., 2004; Lindsey et al., 2005). For instance, in the Koyiaki ranches outside the Masai Mara National Reserve, Ogutu et al. (2005) attributed substantially lower densities of lions (*Panthera leo*) outside the reserve in comparison to spotted hyenas, to less tolerance among Maasai pastoralists to lion depredation on livestock.

Lethal control has traditionally been the most common method for resolving conflicts between carnivores and livestock, leading to the eradication campaigns towards lions, spotted hyenas and African wild dogs (*Lycaon pictus*) in Southern Africa (Mills and Hofer, 1998; Rasmussen, 1999; Woodroffe and Frank, 2005). Some large carnivore species are therefore threatened after having experienced severe declines. For example, the African wild dog has been extirpated from 25 out of 39 former range countries, largely due to human persecution and habitat fragmentation (Fanshawe et al., 1997). According to the IUCN Red list, African wild dogs are listed as endangered, lions and cheetahs (*Acinonyx jubatus*) are listed as vulnerable, whereas spotted hyenas and leopards (*Panthera pardus*) are not categorised as threatened (i.e. lower risk and least concern respectively; IUCN, 2006). Although most large carnivores in Africa are by now legally protected, local people have few incentives to conserve them. Retaliatory killings of carnivores are common, since livestock depredation can have serious economic consequences for livestock keepers, and compensation schemes that may offset some of the costs are often lacking (Ogada et al., 2003; Frank et al., 2005; Graham et al., 2005). However, as examples from Europe and North America illustrate, compensation schemes do not provide an easy solution to the problem (Linnell et al., 1996; Treves and Karanth, 2003).

In Africa, Tanzania is one of the most important countries for large carnivore conservation (Nowell and Jackson, 1996; Mills and Hofer, 1998). Despite having an extensive protected area system, with several very large protected areas (>10,000 km²), carnivore populations are still severely affected by human activity (Hofer et al., 1993, 1996; Packer et al., 2000). Moreover, human encroachment upon protected areas is intensifying the conflict between carnivores and livestock keepers. However, up to now most studies investigating livestock depredation in Africa have been conducted in areas with relatively low human density or immediately adjacent to protected areas (Rudnai, 1979; Mizutani, 1993; Karani et al., 1995; Butler, 2000; Ogada et al., 2003; Patterson et al., 2004; Kolowski and Holekamp, 2006). Few studies have investigated livestock depredation in areas with high human densities and how distance from the protected area influence livestock depredation. In this study, we explored through a questionnaire study the extent and impact of conflict between carnivores and agro-pastoralist outside the Serengeti

National Park. Moreover, we quantify the perceived economic losses to local communities, and examine which factors influenced the approval of retaliatory killing as a carnivore depredation deterrent, since this is a common but illegal practice in Tanzania that has serious implications for carnivore persistence.

2. Methods

2.1. Study area

2.1.1. Climate and large mammals

The study was carried out on the north-western side of the Serengeti National Park (1°15'–3°30' S, 34°–36° E, Fig. 1). The Serengeti National Park (14,763 km²) is a World Heritage Site and the largest National Park in Tanzania. On the northern side it is buffered by several partially protected areas: Ikorongo Game Reserve (ca. 563 km²), Grumeti Game Reserve (ca. 416 km²) and the Ikoma Open Area (ca. 600 km²). The average annual temperature in the study area is 21.7 °C, with an average annual precipitation of 800 mm in the east to 1050 mm in the north-western parts. The protected area network in the western Serengeti harbours large populations of resident ungulates including giraffe (*Giraffa camelopardis*), buffalo (*Syncerus caffer*), topi (*Damilliscus korrigum*), impala (*Aepyceros melampus*) and gazelles (*Gazella thomsoni* and *G. granti*), as well as large carnivores, such as spotted hyena, lion, leopard and cheetah (African wild dogs are currently absent from this area). The western corridor of the Serengeti National Park is characterised by the annual wildebeest (*Connochaetes taurinus*) migration, which in June–July travels through the partially protected areas on their way north (Sinclair, 1995). However, the partially protected areas only contain low numbers of resident wildlife, because of illegal bushmeat hunting, while the village areas contain almost no large wildlife (Rusch et al., 2005). In the partially protected areas all the larger carnivores are included in the trophy hunting quota, except cheetahs and African wild dogs.

2.1.2. People and livestock husbandry

In the agro-pastoral areas in the western Serengeti there is a high human population density (70 people/km²), and a population growth rate of 2.5% in the period from 1988 to 2002 (human population in Mara Region in 2002 was 1.37 million) (URT, 2002). The villages are administrative units consisting of widely dispersed houses with no clear cut border to households belonging to other villages (Fig. 1), where the multiethnic villages consist of subsistence farmers who complement their livelihoods to varying degrees with livestock keeping and illegal bushmeat hunting. Generated income from these activities is partly used to pay taxes, village development contributions and levies, buy food and to purchase clothing (Loibooki et al., 2002; Holmern et al., 2004). The areas immediately adjoining the Serengeti National Park are experiencing a high pressure for scarce resources, and have a particularly high immigration rate (Campbell and Hofer, 1995).

In the western Serengeti, livestock husbandry is commonly practiced with mixed species herds of cattle, goats and sheep. A few farmers also keep donkeys and pigs. Livestock are usually taken out in the early morning (<09:00) and returned to night enclosures before sunset. Grazing




Fig. 1 – Distribution of reported livestock kills according to the predator responsible for the kill. The location of villages is shown as grey pentagrams, and the households from the seven villages included in the survey as open circles.

usually takes place close to the villages, but in the villages directly bordering the Ikorongo and Grumeti Game Reserves some illegal livestock grazing takes place inside the game reserves (especially in Grumeti Game Reserve). Livestock is always herded by people, in most cases by 1–3 adults, but sometimes also by children. At night cattle and donkeys are kept inside night enclosures (i.e. bomas), that are constructed by closely spaced vertical tree trunks. Goats and sheep are usually brought together in a separate small-stock hut that is constructed of poles and clay with grass roofing. Pigs are kept in separate pens constructed by poles and acacia bush (branches facing out). In addition, most households keep guard dogs. Extremely few people have access to firearms.

2.2. Data collection

The data were collected through a questionnaire survey between September and November 2004. Our survey encompassed 481 randomly chosen households from seven villages (based on household lists and including an equal proportion from each sub-village) in the western Serengeti, located at different distances from the closest protected area border; Kunzugu (3 km), Misseke (4 km), Kihumbu (5 km), Makundusi (8 km), Nyichoka (11 km), Kurusanga (20 km) Maburi (29 km) (see Fig. 1). The seven villages had, according to village records, a total of 2708 households, which means the survey canvassed 17.8% of the households. Interviews were conducted in Kiswahili by two Tanzanian scientists trained in interview techniques in the informant's home (the head of household or their wife), and the questionnaire included a mixture of fixed and open ended questions, which covered the respondent's background (age, tribe, education, etc.), livestock losses in the year 2003 and the approval of retaliatory killing of carnivores. Livestock losses were calculated against the size of herds in 2004. During interviews we used colour plates in field guides to help distinguish between carnivore

species. Moreover, the respondents did not differentiate between striped hyena (*Hyaena hyaena*) and spotted hyena, but available data suggest that the much more common spotted hyena was the main predator on livestock in the area (Mills and Hofer, 1998). Likewise, black backed jackal (*Canis mesomelas*) is likely to be the jackal species present in the villages.

2.3. Statistical analysis

During the survey, we collected the GPS location of each household and the distance to the closest protected area border (i.e. game reserve or national park) was calculated by using ArcView 9.0 (Environmental Systems Research Institute, Redlands, CA, USA). We used logistic regression, to investigate which factors affected approval of retaliatory killing of carnivores. This was assessed by the statement: "Carnivores that cause damage to livestock are pests and should be shot". First we analysed the full data set, including both respondents with livestock and those without ($n = 411$), where we used the predictor variables: (1) distance to closest protected area (PA) border; (2) gender (male, female); (3) age (in years); (4) education (no education, primary school and secondary school pooled); and (5) livestock ownership. The interactions that were included were: education \times PA distance, education \times gender, education \times age. Moreover, since the degree of dependency on livestock might influence the attitude against retaliatory killings, we regressed livestock numbers against crop area and saved these residuals (i.e. positive residuals less dependent on livestock). Thereafter, we ran an analysis for a subset of the data, including only livestock keepers ($n = 274$), where the residuals were used as a covariate in the model. In addition, this subset model included two more predictor variables: (1) perception of effectiveness of livestock measures; (2) number of livestock killed. All "don't know" answers on attitude were excluded from both analyses. We selected the most parsimonious models according to AIC_c.

(Akaike Information Criterion corrected for small samples) (Burnham and Anderson, 2002). Moreover, we used Mann-Whitney U, Kruskal-Wallis one-way analysis of variance and χ^2 tests to investigate the occurrence of livestock depredation, where the considered significance value was $p < 0.05$. The analyses were done using SPSS 14.0 (SPSS, 2005) and R 2.3.0 Software (R Development Core Team, 2006).

3. Results

3.1. Livelihood and reported occurrence of large carnivores

Ninety-seven percent of the 481 respondents were agriculturalists. The primary source of income for respondents was subsistence farming (76.7%), followed by cash crop farming (21.0%), and other income generating activities (2.2%, i.e. sale of livestock products, gravel making). In addition to agriculture, respondents supplemented their income through livestock keeping (24.3%), trading (8.3%) and formal employment (4%).

In 2004, 67.4% of households ($n = 481$) kept a total of 13,029 livestock, with an average herd size of 27 head (± 58.7 SD) of stock per household (Table 1). There was a substantial variation among households in the number of livestock owned (range: 0–547). Most livestock keeping households (55.5%) owned 50 or less animals, 11.9% owned more than 50 animals, whereas 32.6% did not own livestock. The majority of the herd was made up of cattle (63.8%) and goats (26%), while the rest were sheep, pigs and donkeys (Table 1). Most respondents reported that they kept their livestock in enclosures during the night (98.1%), while the rest left them tethered outside their house during the night. In addition, a total of 835 dogs were kept by 66.7% of the households in the study villages.

When the respondents were asked about the occurrence (in the past year) of large carnivores in close proximity to their village, all respondents in the survey claimed that spotted hyenas were present. In the villages located furthest away from the protected area (Maburi and Kurusanga) or in the far west (Kunzugu), very few respondents (0–4.2%) stated that large felids (lion and leopard) occurred nearby. In the villages

Table 1 – Mean composition of livestock herds per household in the study villages (2004)

Village	N	Cattle	Goats	Sheep	Donkey	Pigs	Mean ^a
Misseke	68	8.4	5.0	0.8	0	0.3	14.0
Nyichoka	56	17.7	8.1	1.4	0.02	0.5	27.1
Makundusi	68	30.4	12.1	4.9	0.2	0.03	53.3
Maburi	76	17.1	5.8	2.9	0.01	0	25.9
Kihumbu	69	28.5	9.6	5.2	0	0	43.2
Kurusanga	72	7.7	2.9	0.6	0.3	0	11.5
Kunzugu	72	6.8	6.5	2.8	0	0	16.1
Livestock per hh		17.3	7.1	2.7	0.08	0.1	27.1
% of the total herd		63.8	26.0	9.8	0.2	0.2	100

N, number of households (hh) sampled in the study villages.

Sixty-seven percent of households kept livestock; mean values estimated from all households, including those that had none.

^a Mean number of livestock held by a household.

Table 2 – Economic valuation (US \$) of reported livestock kills (n) by wild predators in the study villages in 2003

	Unit value (US\$)	Spotted hyena	Leopard	Baboon	Lion	Jackal	Total (US\$)
Cattle	60	5700 (95)	0	0	60 (1)	0	5760
Goats	11	4158 (378)	121 (11)	33 (3)	0	11 (1)	4323
Sheep	11	2343 (213)	0	0	0	0	2343
Donkey	120	120 (1)	0	0	0	0	120
Pigs	60	300 (5)	0	0	0	0	300
Total loss		12,621 (692)	121 (11)	33 (3)	60 (1)	11 (1)	12,846 (708)
Mean loss (\pm SD)							
Per hh ^a		26.35 (70.63)	0.25 (3.51)	0.07 (1.12)	0.12	0.02	26.82 (81.99)
Per hh ^b		96.03 (107.42)	0.92 (6.66)	0.25 (2.13)	0.45	0.08	97.73 (132.85)
Loss as a % of:							
Herd		97.74	1.55	0.42	0.14	0.14	100
Local per capita income ^a		18.82	0.18	0.05	0.09	0.01	19.15
Local per capita income ^b		68.60	0.66	0.18	0.32	0.06	69.82
Country per capita income ^a		8.23	0.08	0.02	0.04	0.01	8.38
Country per capita income ^b		30.01	0.29	0.08	0.14	0.03	30.55

hh, household.

The conversion rate from Tanzanian shillings was 1 US \$ = 1000 Tz.

^a Considering all the respondents ($n = 481$).

^b Considering only the respondents who reported loss ($n = 132$).

closest to the protected area (Misseke, Nyichoka, Makundusi, Kihumbu), 8.8–19.6% of respondents perceived that lions and leopards occurred, but Kihumbu deviated from this trend for lions where 68.1% of the respondents claimed they occurred close to their village. Only a single respondent reported cheetah to occur nearby (Nyichoka).

3.2. Livestock depredation

A total of 708 livestock were reported killed by predators in 2003 (Table 2). The majority of livestock killed were goats (55.5%), followed by sheep (30.1%), cattle (13.6%), pigs (0.7%) and donkeys (0.1%). Respondents attributed livestock depredation to be caused mainly by spotted hyena (97.7%), leopard (1.6%), baboon (0.4%), lion (0.1%) and lastly black-backed jackal (0.1%). In addition, a total of 171 dogs were reported lost to wild predators in 2003. Predation on dogs was perceived to be caused mainly by spotted hyenas (96.6%), jackal (1.1%) and some by unidentified predators (2.2%).

Most losses (74.8%) of livestock occurred during the night from the enclosures, while 25.2% occurred when the livestock were herded in the field during the day. Livestock losses due to spotted hyena did not differ significantly between wet and dry season ($\chi^2 = 0.004$, $df = 1$, $p = 0.953$), and predation by spotted hyena mainly happened at night ($\chi^2 = 93.2$, $df = 1$, $p < 0.001$). The same pattern was also apparent for dogs, where there was no difference between seasons ($\chi^2 = 1.1$, $df = 1$, $p = 0.312$), and significantly more dogs were killed during the night ($\chi^2 = 66.4$, $df = 1$, $p < 0.001$). Predation on dogs by spotted hyenas happened both when the guarding dogs were loose outside (66.2%), but also when they were kept inside the respondent's house (33.8%) during the night. For the other predators most attacks on livestock occurred during the day, except for one leopard and one lion attack which happened during the night.

There was no significant difference in distance to the closest protected area between households reporting loss and those that did not ($M-W U = 22155$, $z = -0.646$, $p = 0.518$). Depredation events caused by spotted hyena occurred in all the study villages ($11.2 \text{ km} \pm 9.5$, range: 0.6–31.3 km, $n = 132$), whereas for the other four predators depredation occurred only in households relatively close to the protected area ($2.6 \text{ km} \pm 1.9$, range: 0.7–6.3 km, $n = 7$), and this difference was significant ($M-W U = 124$, $z = -3.3$, $p = 0.001$). Percentage of reported livestock losses was significantly different between the villages ($K-W H = 32.2$, $df = 6$, $p < 0.001$). The greatest depredation rates occurred in Misseke (7.7%) and Nyichoka (7.6%), and the lowest in Kunzugu (1.6%). The perceived losses of livestock represented a total of 4.5% ($\pm 13.5\%$) of their livestock (considering all respondents) or 6.8% ($\pm 15.9\%$) when considering only livestock keepers. Mean annual livestock loss per household (of those that reported loss) was 5.3 head of stock (range: 1–33) or 16.6% ($\pm 21.6\%$), which would cost two-thirds of their average annual income to restore.

3.3. Economic valuation of loss

The total economic loss of 708 livestock for the households included in the survey in the seven villages was US \$12,846

for the year 2003 (Table 2). Spotted hyena contributed 98.2% of the economic value of livestock kills, while the economic impact of the other predators was low, although the consequences for the affected households may be serious. Despite being less numerously killed, cattle ($n = 96$) was the most important stock species in terms of economic value (44.8%, US \$5760), because of its high value in comparison to goats and sheep. The annual mean economic loss to each household (all respondents) was estimated to be US \$26.8 (19.2% of the local cash income). Average annual losses for those households that reported depredation ($n = 132$) was calculated to be US \$97.7, which represented 69.8% of local income per household (Table 2).

Table 3 – Summary of logistic binomial regressions models of approval of retaliatory killing

Model	K	AIC _c	Δ_i	w_i
Full data set ($n = 411$)				
Education	2	472.5	0	0.075
Education + PA distance	3	472.7	0.16	0.069
Education + livestock owner	3	473.0	0.47	0.059
Education + PA distance + PA distance \times education	4	473.3	0.81	0.050
Education + PA distance + livestock owner + gender + age + PA distance \times education + gender \times education + age \times education	9	480.5	7.98	0.001
Only livestock keepers ($n = 274$)				
Education + effectiveness of protection measures + number of livestock killed	4	303.9	0	0.050
Effectiveness of protection measures + number of livestock killed	3	304.1	0.13	0.047
Education + effectiveness of protection measures + number of livestock killed + PA distance	5	305.4	1.44	0.024
Education + effectiveness of protection measures + number of livestock killed + PA distance + livestock dependency	6	306.6	1.44	0.013
Education + effectiveness of protection measures + number of livestock killed + PA distance + livestock dependency + gender + age + PA distance \times education + gender \times education + age \times education	11	314.9	6.09	<0.001

Model formulas are shown for the four most parsimonious and the global model, including the number of parameters (K, i.e. number of model terms plus 1 for intercept and error term), Akaike information criterion corrected for small samples (AIC_c), AIC_c differences ($\Delta_i = \text{AIC}_{ci} - \text{AIC}_{c\text{min}}$) and Akaike weights (w_i , the model probabilities, i.e. normalized likelihoods of the models). The models are shown according to AIC_c, with the most parsimonious model at the top of the list.

3.4. Approval of retaliatory killing

Among the respondents a total of 73.4% approved the retaliatory killings of carnivores, 25.4% disagreed, and 1.2% did not know. The majority answered that carnivores should be killed as a response to livestock depredation, because they cause loss to farmers (54.9%), whereas the main reason for disagreeing was that carnivores are beneficial for the country (12.3%) (Table 5). Although for the full data set (including also people who did not own livestock, $n = 411$) the difference in AICc and evidence ratio did not clearly support any of the four top ranked models, the most parsimonious (i.e. with the lowest number of predictors) was the one containing only the variable education (Tables 3 and 4). Similarly, the most parsimonious model for the subset (including only people who owned livestock, $n = 274$) contained the variables, education, effectiveness of protection measures and number of livestock

killed (Tables 3 and 4). Respondents with a formal education (primary or secondary school) were more tolerant of depredation, while both those experiencing a high loss of livestock and the respondents who perceived their livestock husbandry measures as not being effective were more likely to approve of retaliatory killing of carnivores.

4. Discussion

Our results show that livestock depredation can extend relatively deep into non-protected areas depending on the prevalent predators, and can inflict serious economic losses to farmers. In the Serengeti National Park, the spotted hyena is the most numerous large carnivore and therefore it is not surprising that it is perceived to cause most of the livestock loss in our survey. In addition, the nocturnal and opportunistic foraging behaviour, together with the ability of spotted hyenas to take long-distance commuting trips, make them particularly adaptable to anthropogenic environments (Kruuk, 1972; Hofer and East, 1993; Mills and Hofer, 1998).

There are several potential weaknesses by relying solely on questionnaires that might have influenced our livestock loss data. Firstly, in Tanzania government taxes are levied partly on grounds of livestock numbers and although we made sure to identify ourselves as independent researchers during the study, we cannot rule out that the respondents deliberately underestimated their stock level because they were afraid that the results would somehow compromise them. Secondly, as Rasmussen (1999) pointed out, livestock holders may wrongly attribute stock that has died of natural causes to being caused by carnivores – through sheer neglect or prejudices towards specific carnivore species. Thirdly, livestock holders might have an interest in overestimating the rate of loss, because they might believe that it may be beneficial, either through benefits from compensation schemes or being targeted by outreach activities. However, in Tanzania farmers receive no form for compensation, and therefore have little incentive to misrepresent livestock losses. Outreach activities in the study area also do not focus on wildlife damages therefore farmers should have little to gain from overestimating loss. Lastly, respondents often bias their recollection of past events in favour of larger species, especially when sampling from multiple years (see Kruuk, 1980 for an example). We attempted to minimise this problem by only using the most recent year (2003), instead of using a longer time period. Despite these caveats, several studies show that livestock keeper's perception of livestock depredation gives a relatively reliable index of livestock depredation (Kruuk, 1980; Woodroffe et al., 2005). However, incorporating ways of verifying questionnaire data, either through use of wildlife officers that inspect kills or by providing an indirect measure through analysing scats, can be very valuable (Woodroffe et al., 2005; Wang and Macdonald, 2006).

Several studies show that low natural prey densities may be a strong contributor to high depredation rates (Meriggi and Lovari, 1996; Woodroffe et al., 2005; Kolowski and Holekamp, 2006). However, the relationship is not straightforward, since wolf (*Canis lupus*) predation on livestock may also be high where wolves have access to high natural prey densities (Treves et al., 2004). The low natural prey densities and high livestock densities around the Serengeti National Park may

Table 4 – Parameter estimates for the most parsimonious model of approval of retaliatory killing as judged by the AICc.

Coefficients	Estimate	SE	z	p
<i>Full data set (n = 411)</i>				
(Intercept)	-0.83	0.13	-6.46	<0.001
Education	-0.68	0.26	-2.56	0.011
<i>Only livestock keepers (n = 274)</i>				
(Intercept)	-0.83	0.19	-4.17	<0.001
Education	-0.49	0.34	-1.45	0.148
Effectiveness of protection measures	-0.66	0.30	-2.19	0.028
Number of livestock killed	0.05	0.03	1.84	0.065

Table 5 – Comments given by respondents on reason for agreeing or disagreeing with the statement “Carnivores that cause damage to livestock are pests and should be shot” (n = 171)

Reason given for attitude	%
<i>Negative responses (agree)</i>	
Carnivores cause loss to farmers	54.9
Carnivores should be killed since no compensation for damage is paid	9.4
Carnivores are dangerous and may even attack people	4.7
Carnivores are not as important as other wildlife	1.2
<i>Positive responses (disagree)</i>	
Carnivores are beneficial to our nation	12.3
Wildlife has a right to live	7.0
Should just scare the carnivores away from the village area	5.8
Some carnivores are beneficial since they remove dead animals	2.3
Should report carnivore losses to wildlife officer	1.8
To kill wildlife would be against the idea of conservation	0.6

therefore contribute to the reported high depredation rates. On the Kenyan side of the Serengeti ecosystem Kolowski and Holekamp (2006) linked the arrival of the wildebeest migration to lower depredation rates on livestock. In contrast, we find no temporal variation in depredation rates, although the migration to some extent utilise the areas outside the Serengeti National Park. However, the migration travels quickly through the study area and does not venture into the villages far away from the protected area, and therefore seasonal fluctuations in prey availability are not likely to affect depredation rates.

At a regional scale livestock depredation is usually not considered a serious loss factor, and compared to other sources of loss (i.e. mismanagement, diseases, and theft) the impact of livestock depredation is usually relatively small. For example, across studies done in Africa, disease as a loss factor is 3–6 times larger in magnitude than livestock depredation (Mizutani, 1993; Karani et al., 1995; Rasmussen, 1999; Frank et al., 2005). Nevertheless, in some cases large carnivores can be a serious impediment for the economic situation of local livestock keepers (Mishra, 1997; Wang and Macdonald, 2006). Our data also emphasise that livestock depredation mainly by spotted hyenas is a severe economic constraint for households in the western Serengeti, where 27.4% of households ($n = 132$) in our survey of 481 households believed they had lost livestock to predators in 2003. The costs due to livestock loss were on average US \$97.7 per household, which is almost one third of the GNI per capita in Tanzania (US \$320 in 2004) (World Bank, 2006). However, local farmers in the study area have considerably lower income. Borge (2003) reported that in a survey covering 297 households from six villages in the western Serengeti the average annual cash income per household was US \$140, which means that the stock loss constitutes two-thirds of the average annual income. Farmers also reported that carnivores sometimes killed several animals in one attack, which increases the cost to individual owners. However, in some cases farmers might be able to recoup some of the meat value of killed livestock by chasing off carnivores. The value of livestock (especially cattle) in pastoral and agro-pastoral society's has also a very important cultural aspect, which might contribute to their low tolerance of depredation compared to more commercially based enterprises (Patterson et al., 2004).

Large carnivores are also a common problem to human safety in Tanzania, and elsewhere (Løe and Røskoft, 2004; Packer et al., 2005). For example, in March 2004, a rabid spotted hyena was speared to death after attacking and badly mauling a woman in one of the study villages (Holmern, pers obs). Concerns for human safety combined with livestock loss may aggravate the situation and result in retaliatory killings, especially when funding, logistics and manpower constrain the response of wildlife management authorities. In the western Serengeti, there is widespread approval of retaliatory killing when carnivores kill livestock, or are perceived as a threat to human safety. Spotted hyenas are among the least liked large carnivore species in Africa and their dominance in our sample might have influenced the results. However, we cannot rule out that the precise wording of our statement might have contributed somewhat to increasing the approval rate, partly because it is a leading statement and it also contains

two parts which can make interpretation of responses ambiguous. However, widespread support of retaliatory action in the western Serengeti was also reported by Kaltenborn et al. (2006), especially when spotted hyenas killed livestock. Likewise, Ogutu et al. (2005) reported that pastoral tribes in Kenya had a low tolerance of livestock depredation, while Ogada et al. (2003) found that retaliatory killings correlated with livestock loss rates. Our results also suggest that the number of livestock lost is associated with support of retaliatory killing. Considering the economic impact depredation can have on households, this is hardly surprising. Reducing the number of livestock lost to carnivores might contribute to less support of retaliatory killing, but even areas with comparatively low depredation rates can have a strong desire of lethal control (Linnell et al., 1996). Strong support of lethal wildlife management is by no means typical only for rural farmers in Africa, but has also been reported for North America (Kellert, 1985). However, identifying problem individual can be difficult, and lethal control of predators is only likely to cause a short-term respite from losses, because the same or other predator species rapidly re-establish themselves (Linnell et al., 1999; Stahl et al., 2001; Herfindal et al., 2005). But removal of problem carnivores, for example through trophy hunting in village areas, might facilitate public approval of protection for the remainder.

Developing ways of enabling farmers to benefit from the existence of protected areas could be a possible way forward (Wang and Macdonald, 2006). But in the case of the Serengeti National Park, benefits from outreach activities are currently grossly inadequate to offset costs associated with wildlife, and revenues from trophy hunting in the adjacent Game Reserves have a poor track record of reaching local farmers (Holmern et al., 2004). This situation seems also to be typical for other protected areas in Tanzania (Baldus and Cauldwell, 2004). Experience from community-based conservation projects show that distribution of benefits can be problematic and does not necessarily improve conservation (Newmark and Hough, 2000; Johannesen and Skonhoft, 2005). However, implementing incentive schemes aimed at conserving endangered carnivores can work, as encouraging results reported by Mishra et al. (2003) for snow leopard (*Uncia uncia*) show. This is further supported by Johannesen (2006) that demonstrate through modelling that it is crucial for such programs to forge a link between benefit levels and conservation friendly behaviour in order to improve wildlife conservation and human welfare.

Compared to other studies in Africa, the livestock loss reported in this study is among the highest recorded and needs to be addressed, both because it is an economic constraint to households, but also because it increases the likelihood of approving of illegal retaliatory killings, which may be of serious concern for the conservation of endangered carnivores (Rudnai, 1979; Kruuk, 1980; Mizutani, 1993; Karani et al., 1995; Rasmussen, 1999; Butler, 2000; Frank et al., 2005; Kolowski and Holekamp, 2006). Our results point out the need of formal education in order to improve attitudes, which is in accordance with many similar studies (Lindsey et al., 2005; Woodroffe et al., 2005). Prejudice against carnivores and misconceptions of the actual causes of loss are quiet common among farmers (Rasmussen, 1999). The development of better

education in the region, particularly the establishment of more primary and secondary schools which at the moment have a poor coverage, along with education programmes on wildlife conservation might lead to increasing tolerance and decreasing misconceptions. Earlier research in Africa and Asia has also identified the need of improving livestock husbandry to reduce conflict levels (Kruuk, 1980; Mishra, 1997; Rasmussen, 1999; Ogada et al., 2003). It is therefore essential that further research should address the precise role of livestock husbandry practices in explaining depredation events outside the Serengeti National Park. The construction of night time enclosures might therefore be of particular importance, since most depredation occurs after dark.

Acknowledgements

This study was part of the TAWIRI-NINA-NTNU Project "Biodiversity and the Human Wildlife Interface in the western Serengeti". We thank the Tanzanian Commission of Science and Technology for permission to conduct the study, Serengeti Regional Conservation Project for facilitating the study and Tanzania Wildlife Research Institute for logistic support. We would also like to thank Robert Fyumagwa and Julius Key-yu for their help during the study, Ivar Herfindal for advice on analyses, the brood parasitism/conservation group at the Norwegian University of Science and Technology for fruitful discussions, and two anonymous referees for very constructive comments on previous versions of this manuscript. We are especially grateful to Lukas Malugu and Michael Mayengo for assistance in the field. The work was partly supported by a grant to T.H. from the Nordic African Institute and from Arnfinn and Lise Hejes Legat.

REFERENCES

- Baldus, R.D., Cauldwell, A.E., 2004. Tourist hunting and its role in development of wildlife management areas in Tanzania. In: Sixth international Game Ranching Symposium, Paris, 45pp.
- Borge, A., 2003. Essays on the economics of African wildlife and utilization and management. Dr.polit. thesis, Norwegian University of Science and Technology, Trondheim, Norway, 187pp.
- Boydston, E.E., Kapheim, K.M., Watts, H.E., Szykman, M., Holekamp, K.E., 2003. Altered behaviour in spotted hyena associated with increased human activity. *Animal Conservation* 6, 207–219.
- Burnham, K.P., Anderson, D.R., 2002. *Model Selection and Multimodel Inference: A Practical Information-theoretic Approach*. Springer, Berlin.
- Butler, J.R.A., 2000. The economic costs of wildlife predation on livestock in Gokwe communal land, Zimbabwe. *African Journal of Ecology* 38, 23–30.
- Campbell, K.L.I., Hofer, H., 1995. People and wildlife: spatial dynamics and zones of interaction. In: Sinclair, A.R.E., Arcese, P. (Eds.), *Serengeti II – Dynamics, Management, and Conservation of an Ecosystem*. The University of Chicago Press, Chicago, pp. 534–570.
- Cardillo, M., Purvis, A., Secrest, W., Gittleman, J.L., Bielby, J., Mace, G.M., 2004. Human population density and extinction risk in the world's carnivores. *PLOS Biology* 2, 0909–0914.
- Fanshawe, J.H., Ginsberg, J.R., Sillero-Zubiri, C., Woodroffe, R., 1997. The status and distribution of remaining wild dog populations. In: Woodroffe, R., Ginsberg, J.R., Macdonald, D.W. (Eds.), *The African Wild Dog: Status Survey and Conservation Action Plan*, pp. 11–57.
- Frank, L.G., Woodroffe, R., Ogada, M.O., 2005. People and predators in Laikipia District, Kenya. In: Woodroffe, R., Thirgood, S., Rabinowitz, A. (Eds.), *People and Wildlife: Conflict or Coexistence?* Cambridge University Press, Cambridge, pp. 286–304.
- Graham, K., Beckerman, A.P., Thirgood, S., 2005. Human-predator-prey conflicts: ecological correlates, prey losses and patterns of management. *Biological Conservation* 122, 159–171.
- Herfindal, I., Linnell, J.D.C., Moa, P.F., Odden, J., Austmo, L.B., Andersen, R., 2005. Does recreational hunting of lynx reduce depredation losses of domestic sheep? *Journal of Wildlife Management* 69, 1034–1042.
- Hofer, H., East, M.L., 1993. The commuting system of Serengeti spotted hyaenas: how a predator copes with migratory prey. I. Social organization. *Animal Behaviour* 46, 547–557.
- Hofer, H., East, M.L., Campbell, K.L.I., 1993. Snares, commuting hyaenas and migratory herbivores: humans as predators in the Serengeti. *Symposia of the Zoological Society London* 65, 347–366.
- Hofer, H., Campbell, K.L.I., East, M.L., Huish, S.A., 1996. The impact of game meat hunting on target and non-target species in the Serengeti. In: Taylor, J., Dunstone, N. (Eds.), *The Exploitation of Mammal Populations*. Chapman and Hall, London, pp. 117–146.
- Holmern, T., Borge Johannesen, A., Mbaruka, J., Mkama, S., Muya, J., Røskft, E., 2004. Human-wildlife conflicts and hunting in the western Serengeti, Tanzania. Norwegian Institute for Nature Research, Project Report 26, Trondheim, Norway, 26pp.
- IUCN The World Conservation Union, 2006. *IUCN Red List of threatened species: A Global Assessment*. Gland, Switzerland.
- Johannesen, A.B., 2006. Designing integrated conservation and development projects (ICDPs): illegal hunting, wildlife conservation, and the welfare of the local people. *Environment and Development Economics* 11, 247–267.
- Johannesen, A.B., Skonhøft, A., 2005. Tourism, poaching and wildlife conservation: what can integrated conservation and development projects accomplish? *Resource and Energy Economics* 27, 208–226.
- Kaltenborn, B.P., Bjerke, T., Nyahongo, J.W., Williams, D.R., 2006. Animal preferences and acceptability of wildlife management actions around Serengeti National Park, Tanzania. *Biodiversity and Conservation*, doi: 10.1007/s10531-005-6196-9.
- Karani, I.W., Dublin, H.T., Koehler, G.M., 1995. Livestock depredation by predators in pastoral areas adjacent to Masai Mara National Reserve, Kenya. In: Bisonette, J.A., Krausman, P.R. (Eds.), *Integrating People and Wildlife for a Sustainable Future*. Wildlife Society, Maryland, pp. 360–363.
- Kellert, S.R., 1985. Public perceptions of predators, particularly the wolf and coyote. *Biological Conservation* 31, 167–189.
- Kleiven, J., Bjerke, T., Kaltenborn, B., 2004. Factors influencing the social acceptability of large carnivore behaviours. *Biodiversity and Conservation* 13, 1647–1658.
- Kolowski, J.M., Holekamp, K.E., 2006. Spatial, temporal, and physical characteristics of livestock depredations by large carnivores along a Kenyan reserve border. *Biological Conservation* 128, 529–541.
- Kruuk, H., 1972. *The Spotted Hyena: A Study of Predation and Social Behaviour*. The University of Chicago Press, Chicago.
- Kruuk, H., 1980. The effect of large carnivores on livestock and animal husbandry in Marsabit District, Kenya. IPAL Technical Report E – 4 (ITE Project 675), 52pp.

- Lindsey, P.A., du Toit, J.T., Mills, M.G.L., 2005. Attitudes of ranchers towards African wild dogs *Lycaon pictus*: conservation implications on private land. *Biological Conservation* 125, 113–121.
- Linnell, J.D.C., Smith, M.E., Odden, J., Kaczensky, P., Swenson, J.E., 1996. Strategies for the reduction of carnivore–livestock conflicts: a review. Norwegian Institute for Nature Research, Oppdragsmelding 443, Trondheim, Norway, 118pp.
- Linnell, J.D.C., Odden, J., Smith, M.E., Aanes, R., Swenson, J.E., 1999. Large carnivores that kill livestock: do “problem individuals” really exist? *Wildlife Society Bulletin* 27, 698–705.
- Linnell, J.D.C., Swenson, J., Andersen, R., 2001. Predators and people: conservation of large carnivores is possible at high human densities if management policy is favourable. *Animal Conservation* 4, 345–350.
- Løe, J., Røskoft, E., 2004. Large carnivores and human safety: a review. *Ambio* 33, 283–288.
- Loibooki, M., Hofer, H., Campbell, K.L.I., East, M., 2002. Bushmeat hunting by communities adjacent to the Serengeti National Park, Tanzania: the importance of livestock ownership and alternative sources of protein and income. *Environmental Conservation* 29, 298–391.
- Meriggi, A., Lovari, S., 1996. A review of wolf predation in southern Europe: does the wolf prefer wild prey to livestock. *Journal of Applied Ecology* 33, 1561–1571.
- Mills, G., Hofer, H., 1998. Hyenas: Status Survey and Conservation Action Plan. World Conservation Union, Gland, Switzerland.
- Mishra, C., 1997. Livestock depredation by large carnivores in the Indian trans-Himalaya: conflict perceptions and conservation prospects. *Environmental Conservation* 24, 338–343.
- Mishra, C., Allen, P., McCarthy, T., Madhusudan, M.D., Bayarjargal, A., Prins, H.H.T., 2003. The role of incentive programs in conserving the snow leopard. *Conservation Biology* 17, 1512–1520.
- Mizutani, F., 1993. Home range of leopards and their impact on livestock on Kenyan ranches. In: Dunstone, N., Gorman, M.L. (Eds.), *In Mammals as Predators*. Clarendon Press, Oxford, pp. 425–439.
- Newmark, W.D., 1996. Insularization of Tanzanian parks and the local extinction of large mammals. *Conservation Biology* 10, 1549–1556.
- Newmark, W.D., Hough, J.L., 2000. Conserving wildlife in Africa: integrated conservation and development projects and beyond. *BioScience* 50, 585–592.
- Nowell, K., Jackson, P., 1996. Status, Survey and Conservation Action Plan, Wild Cats. World Conservation Union, Gland, Switzerland.
- Ogada, M.O., Woodroffe, R., Ouge, N.O., Frank, L.G., 2003. Limiting depredation by African carnivores: the role of livestock husbandry. *Conservation Biology* 17, 1521–1530.
- Ogutu, J.O., Bholá, N., Reid, R., 2005. The effects of pastoralism and protection on the density and distribution of carnivores and their prey in the Mara ecosystem of Kenya. *Animal Conservation* 265, 281–293.
- Packer, C., Altizer, S., Appell, M., Brown, E., Martenson, J., O’Brien, S.J., Roelke-Parker, M., Hofman-Lehman, R., Lutz, H., 2000. Viruses of the Serengeti: patterns of infection and mortality in African lions. *Journal of Animal Ecology* 68, 1161–1178.
- Packer, C., Ikanda, D., Kissui, B., Kushnir, H., 2005. Lion attacks on humans in Tanzania. *Nature* 436, 927–928.
- Patterson, B.D., Kasiki, S.M., Selempo, E., Kays, R.W., 2004. Livestock predation by lions (*Panthera leo*) and other carnivores on ranches neighboring Tsavo National Parks, Kenya. *Biological Conservation* 119, 507–516.
- Rasmussen, G.S.A., 1999. Livestock predation by the painted hunting dog *Lycaon pictus* in a cattle ranching region of Zimbabwe: a case study. *Biological Conservation* 88, 133–139.
- R Development Core Team, 2006. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna.
- Rudnai, J., 1979. Ecology of lions in Nairobi National Park and the adjoining Kitengela Conservation Unit in Kenya. *African Journal of Ecology* 17, 85–95.
- Rusch, G., Stokke, S., Røskoft, E., Mwakalebe, G., Wiik, H., Arnemo, J., Lyamuya, R., 2005. Human–wildlife interactions in western Serengeti, Tanzania. Effects of land management on migratory routes and mammal population densities. NINA Report 85, 47pp.
- Sinclair, A.R.E., 1995. Serengeti past and present. In: Sinclair, A.R.E., Arcese, P. (Eds.), *Serengeti II – Dynamics, Management, and Conservation of an Ecosystem*. The University of Chicago Press, Chicago, pp. 3–30.
- SPSS, 2005. SPSS for Windows. Release 14.0. SPSS Inc., Chicago, IL.
- Stahl, L.P., Vandel, J.M., Herrenschmidt, V., Migot, P., 2001. The effect of removing lynx in reducing attacks of sheep in the French Jura mountains. *Biological Conservation* 101, 15–22.
- Treves, A., Karanth, K.U., 2003. Human–carnivore conflict and perspectives on carnivore management worldwide. *Conservation Biology* 17, 1491–1499.
- Treves, A., Naughton-Treves, L., Harper, E.K., Mladenoff, D.J., Rose, R.A., Sickle, T.A., Wydeven, A.P., 2004. Predicting human–carnivore conflict: a spatial model derived from 25 years of data on wolf predation on livestock. *Conservation Biology* 18, 114–125.
- URT United Republic of Tanzania, 2002. 2002 Population and housing census. <http://www.tanzania.go.tz/census/index.html> (accessed 15.06.2006).
- Wang, S.W., Macdonald, D.W., 2006. Livestock predation by carnivores in Jigme Singye Wangchuck National Park, Bhutan. *Biological Conservation* 129, 558–565.
- Woodroffe, R., Frank, L.G., 2005. Lethal control of African lions (*Panthera leo*): local and regional population impacts. *Animal Conservation* 8, 91–98.
- Woodroffe, R., Ginsberg, J.R., 1998. Edge effects and the extinction of populations inside protected areas. *Science* 280, 2126–2128.
- Woodroffe, R., Lindsay, P., Romanach, S., Stein, A., Ole Ranah, S.M.K., 2005. Livestock predation by endangered African wild dogs (*Lycaon pictus*) in northern Kenya. *Biological Conservation* 124, 225–234.
- World Bank, 2006. World Development Indicators database: Tanzania data profile. <http://devdata.worldbank.org/external/CPProfile.asp?PTYPE=CP&CCODE=TZA> (accessed 15.06.2006).

Paper III

1 **Disease is a major cause of livestock loss in villages surrounding western**
2 **Serengeti, Tanzania**

3

4 Nyahongo, W. J.^{1,3*}, Holmern, T^{1.}, Stokke, B.G.^{1.}, Keyyu, J.D.^{3.}, Kaltenborn, P.
5 B^{2.}, and Røskaft, E^{1.}

6 (1) Department of Biology, Norwegian University of Science and Technology
7 (NTNU), Realfagbygget, N-7491 Trondheim, Norway.

8 (2) Norwegian Institute for Nature Research (NINA), Fakkeltgården, 2624,
9 Lillehammer, Norway.

10 (3) Tanzania Wildlife Research Institute (TAWIRI), P. O. Box 661, Arusha,
11 Tanzania.

12

13 Running title: Causes of livestock loss in villages

14

15 *Corresponding author: email: nyahongo@bio.ntnu.no or

16 nyhwjulius@yahoo.co.uk

17 Tel: +47-48230094, +225-754886247

18 Word counts: Abstracts 242 words, Main text (including references 4979)

19

20

21 Key words: Diseases, livestock losses, western Serengeti, wild carnivores

22 **Abstract**

23 Diseases have been responsible for high livestock losses in sub-Saharan
24 Africa, delaying the introduction of cattle-based economies for many years. In
25 this study, we quantified and compared the magnitude of livestock losses per
26 household due to diseases, theft, depredation, and poor management in the
27 grazing field from April to December 2006 in villages located in the western
28 Serengeti. Furthermore, we compared such losses in villages situated close to
29 or farther away from Serengeti National Park. Diseases were responsible for
30 higher livestock losses than other loss causes. Overall, diseases were
31 responsible for 3.5 – 7.0% livestock loss per household during the period of
32 nine months, costing them US\$ 83.5. This loss is equivalent to 59.6% of the
33 average annual household income. The cost per household of theft and poor
34 management in the grazing field was US \$ 3.0 and US \$ 11.4, respectively. The
35 cost of depredation recorded was higher in the household located far away from
36 the park boundary. The depredation cost per household in the four villages was
37 US\$ 16.5. Spotted hyena *Crocuta crocuta* was the only reported predator killing
38 livestock and it killed more sheep and goats than cattle. We recommend better
39 education on animal husbandry practices and extension service to help in
40 maintenance of livestock health and to prevent theft and loss while grazing. To
41 reduce livestock depredation, we suggest that night enclosures for livestock
42 should be improved, especially in villages that are situated further away from
43 national parks.

44 **Introduction**

45 In sub-Saharan African, diseases have been documented to be responsible for
46 high losses in livestock production (Gifford-Gonzalez, 2000). Historically,
47 diseases have been an important factor that delayed the introduction of cattle-
48 based economies by as much as a thousand years after the first appearance of
49 small stock in both eastern and southern Africa (Gifford-Gonzalez, 2000).
50 Diseases that are often fatal to livestock production, especially for cattle in sub-
51 Saharan, Africa include wildebeest-derived Malignant Catarrhal Fever (MCF),
52 East Coast Fever (ECF), Foot and Mouth Diseases (FMD), worms (helminthes),
53 Rift Valley Fever (RVF), rinderpest, anthrax as well as trypanosomiasis
54 (Rwambo et al., 1999; Thomson et al., 2003; Kock, 2003). Livestock diseases
55 have economic consequences on livestock husbandry at two levels; 1) at the
56 domestic level, the diseases are responsible for direct loss due to mortality or
57 indirectly through lowered production and/or the cost of treatment and
58 prevention (Perry et al. 2002; Kock, 2003). 2) At the international level, diseases
59 may affect any opportunity for export of livestock and livestock products
60 between regions or continents, jeopardizing the exchange of products for
61 foreign currency (OIE, 2003; Kock, 2003). Because of negative attitudes of
62 livestock keepers towards wild carnivores, they often claim wild carnivores
63 being responsible for higher losses of livestock despite the direct and indirect
64 impacts of livestock diseases (Mwangi, 1997; Rasmussen, 1999). However,
65 several factors may equally cause significant livestock loss, for example theft,
66 drought and poor livestock husbandry (Ogada et al., 2003). The high prices
67 received for livestock in the livestock auctions, make the theft of animals a

68 lucrative business. In Africa theft may increase with the number of animals the
69 household own, because it may be difficult to notice a loss of one or few
70 animals in a group of several hundred individuals. Moreover, livestock theft may
71 vary depending on the season or between years. During the rain season, it may
72 be easy to follow the tracks the animal stolen has left behind to the destination.
73 Thus, thieves would avoid this season. The night with a full moon is not
74 conducive for livestock raiders because it is possible for livestock keepers to
75 see the livestock in the night holding enclosure from within the household living
76 quarters. Elsewhere outside Africa, livestock theft has been considered the
77 most significant rural crime (WASDA, 2007). Drought may affect livestock
78 directly by reducing the available food and water; hence animals may be so
79 weak that they easily succumb to diseases. Indirectly drought normally
80 associates with famine which drive the livestock keepers to sale some
81 individuals to buy food.

82

83 The level of livestock depredation may be intentionally exaggerated to attract
84 public attention and/or to mask effects of poor livestock management (Nabane,
85 1995; 1996; Infield, 1996). Such negative attitudes towards carnivores due to
86 perceived levels of predation have been cited as a challenging issue in both
87 wildlife conservation and rural development (Woodroffe et al., 2000). In different
88 parts of the world, conflicts between human and wild carnivores have been well
89 documented (e.g. Treves and Karath, 2003; Treves et al., 2004; Røskaft et al.,
90 2007). This conflict has resulted in direct persecution of carnivores to get rid of
91 them close to human settlements (e.g. Mill and Hofer, 1998; Woodroffe and

92 Frank, 2005), and resulted a general dislike of such animals. For example,
93 American citizens do not like wolves *Canis lupus* and coyotes *C. latrans*
94 (Kellert, 1985). Likewise, sheep farmers in Norway show negative attitudes
95 towards bears *Ursus arctos*, wolves and lynx *Lynx lynx* (Kaltenborn et al., 1998;
96 Vittersø et al., 1998; Kaltenborn et al., 1999; Røskaft et al., 2007). In some
97 parts of Africa, the same negative attitudes towards carnivores have been
98 reported as well (Lindsey, et al 2005; Kalternborn et al., 2006; Holmern et al.,
99 2007). For example, livestock keepers in Africa have been reported to kill and
100 poison carnivores to reduce the perceived conflict over livestock depredation
101 (Stuart et al., 1985; Berry, 1990; Holekamp and Smale, 1992).

102

103 The aim of this study was to record and discuss factors responsible for livestock
104 loss in households from the villages surrounding the western Serengeti. The
105 livestock loss causes that were recorded in each selected household included
106 diseases, theft, depredation and loss in the grazing field. We recorded number
107 of animals slaughtered for meat, sold and/or bought as well as newborn calves.

108

109 **Methods**

110 **Study area**

111 Serengeti National Park (SNP) is situated west of the Rift Valley and the
112 western border is close to Lake Victoria while the northern edge borders Kenya
113 (Fig. 1). The central part of the current park was designated as a Game
114 Reserve in 1929. In 1940 hunting was banned and in 1951 it was declared a
115 national park. In the time following, the borders have been modified as the park

116 has expanded. In 1981 Serengeti was inscribed as a World Heritage Site. The
117 park covers 14 763 km² and is the core of the Serengeti ecosystem that
118 includes Ngorongoro Conservation Area, Maswa Game Reserve, Ikorongo-
119 Grumeti Game Reserves and Loliondo Game Controlled Area, in Tanzania as
120 well as the Maasai Mara Natural Reserve to the north in Kenya.

121

122 The current study was conducted in the villages surrounding western Serengeti
123 (Fig. 1), one of the areas of the SNP that currently suffers from conflict between
124 conservation priorities of the park and priorities of local communities (Hofer et
125 al., 1996; Loibooki, 1997). This is a section of the Serengeti ecosystem that
126 extends westward to Lake Victoria with a relatively high human population
127 density (i.e. 70 people/km²; growing at a rate of 2.5% between 1988 and 2002,
128 URT 2002). The majority of local communities along the boundaries of the
129 western Serengeti are subsistence farmers who keep livestock and practice
130 crop production. Many of the farmers obtain natural resources inside the
131 protected areas for home consumption. For instance, during the dry season,
132 livestock keepers illegally graze and water their livestock in the protected areas
133 (Nyahongo et al., 2006). In addition, illegal hunting within the protected areas is
134 well documented (Arcese et al., 1995; Campbell and Hofer, 1995; Loibooki et
135 al., 2002; Nyahongo et al., 2006). The illegal bushmeat hunters may sell the
136 illegally obtained meat to generate income (Arcese et al., 1995; Hofer et al.,
137 1996; Loibooki, 1997).

138

139

140 **Data collection**

141 The current study was conducted between April and December 2006.
142 Households were selected in the following villages: Robanda, Nyamakendo,
143 Nattambiso and Kowak. The first three villages were within 10 km from the
144 boundary of the park while Kowak village was located about 80 km from the
145 park. Household were selected randomly according to household lists in the
146 villages. For practical reason (livestock counting time), we omitted household
147 with more than 200 individual cattle, goats or sheep. The first three months
148 (January, February and March) were spent in villages to introduce researchers
149 to livestock keepers and to establish baseline data on livestock numbers per
150 selected household. Livestock owners were informed about the essence of this
151 study and was assured that the data was only collected for research purpose
152 and not for other purposes like baseline data for setting livestock levees by the
153 government. After establishing the baseline data (i.e. initial numbers of livestock
154 per selected household), we appointed enumerators from the respective
155 villages that consists of livestock owners enumerating any livestock that
156 suffered loss (death due to diseases, loss while grazing, theft, predation), own
157 consumption (slaughtering) or gain (new-born, bought or paid as dowry). While
158 enumerators were collecting data on a daily bases, the researcher visited each
159 household after every three months to recount the animals again in order to
160 cross check the data that enumerators collected. Furthermore, livestock owners
161 were asked about the livestock status during the period of the past three
162 months. Livestock were either counted in the morning before being sent out for
163 grazing in the field (normally 2 to 3 km away from the night holding enclosures)

164 or in the evening when they were brought back to the night holding enclosures
165 (the counting rate was 15 to 20 households per day and we spent one week in
166 each village).

167

168 All livestock were prized according to matured livestock because market prices
169 for livestock are only set for mature animals. This allowed us to be able to
170 calculate the mean cost of livestock loss causes per household per year.

171

172 **Statistical analyses**

173 All analyses were performed using SPSS 14 statistical package (SPSS, 2005).

174 Descriptive statistics were used to calculate means and standard deviation

175 while non-parametric tests were applied to test the differenced among the loss

176 factors, household and livestock species. The mean number of livestock was

177 the average of the livestock count each three months. The proportions (%) of

178 livestock loss/gain causes or household expenditure were calculated as the

179 ratio of each variable to the calculated mean livestock numbers. For all tests $p <$

180 0.05 was considered significant.

181 **Results**

182 **Livestock gain and loss causes**

183 Mean household livestock population variation and the subsequent cost or
184 benefits when presented in monetary term for the current values of livestock
185 species in each village are summarized in Table 1 and Table 2.

186

187 Regardless of household locality, various loss causes affected livestock
188 differently (cattle: Friedman test, $\chi^2 = 233.72$, $df = 3$, $n = 182$, $p < 0.001$; goats:
189 Friedman test, $\chi^2 = 134.07$, $df = 3$, $n = 155$, $p < 0.001$; sheep: Friedman test, χ^2
190 $= 81.26$, $df = 3$, $n = 123$, $p < 0.001$). Furthermore, the mean number of cattle
191 and goats sold per household was higher than the number slaughtered (cattle:
192 Wilcoxon sign rank test, $Z = -7.24$, $n = 182$, $p < 0.001$; goats: Wilcoxon sign
193 rank test, $Z = -3.214$, $n = 155$, $p = 0.001$) but this was not the case for sheep
194 (Wilcoxon sign rank test, $Z = -0.70$, $n = 123$, $p = 0.484$). In all households, new
195 born calves, and not animals bought or paid as dowry, was the significant
196 source of replenishment of livestock numbers (cattle: Wilcoxon sign rank test, Z
197 $= -8.54$, $n = 182$, $p < 0.001$; goats: Wilcoxon sign rank test, $Z = -8.38$, $n = 155$, p
198 < 0.001 ; Sheep: Wilcoxon sign rank test, $Z = -7.56$, $n = 123$, $p < 0.001$).

199

200 **Comparison of livestock loss causes among villages**

201 The mean number of goats and sheep that was depredated was higher in
202 Kowak than in villages that were closer to the park boundary although this was
203 not significant statistically (Table 1). In all livestock depredation events spotted

204 hyena *Crocuta crocuta* was the only carnivore reported to be responsible for
205 livestock killing.

206

207 Mean number of cattle that died of diseases differed significantly among the
208 villages (Kruskal-Wallis, $H = 17.072$, $df = 3$, $p = 0.001$). Furthermore, the
209 difference in mean number of cattle that were stolen among villages was almost
210 significant (Kruskal-Wallis, $H = 7.124$, $df = 3$, $p = 0.068$). The remaining cattle
211 loss causes did not differ significantly among villages ($p > 0.09$ for all cases).

212

213 Likewise, the effect of all loss causes in goats did not differ significantly among
214 the four villages ($p > 0.076$ for all cases). However, for sheep, losses due to
215 diseases and poor management differed significantly among the villages
216 (Kruskal-Wallis, $H = 9.10$, $df = 3$, $p = 0.028$ and $H = 8.85$, $df = 3$, $p = 0.031$,
217 respectively), while theft and depredation on livestock had similar effect among
218 the villages ($p > 0.118$ for all cases).

219

220 **Comparison of livestock loss causes among livestock species**

221 Generally, regardless of distance from the park boundary, mean number of
222 livestock species that were sold, slaughtered for food and that were killed by
223 spotted hyenas differed significantly between livestock species (sold: Kruskal-
224 Wallis, $H = 10.82$, $df = 2$, $p = 0.005$; slaughtered: Kruskal-Wallis, $H = 17.09$, $df =$
225 2 , $p < 0.001$; predation: Kruskal-Wallis, $H = 14.01$, $df = 2$, $p = 0.001$).
226 Households sold more cattle (mean rank = 248.5) than goats (mean rank =
227 231.4) or sheep (mean rank = 202.7). However, households slaughtered more

228 goats for food (mean rank = 249.6) than sheep (mean rank = 243.2) or cattle
229 (mean rank = 205.6). In contrast, sheep were more frequently killed by spotted
230 hyenas (mean rank = 246.6) than goats or cattle (goat: mean rank = 241.1;
231 cattle: mean rank = 210.6). The remaining loss causes did not differ significantly
232 among species ($p > 0.151$).

233

234 **Economic significance of livestock loss or gain causes**

235 In total, the mean value of livestock that household from four villages own was
236 US\$ 2121 (sum of mean cattle, goats and sheep per household) and newborn
237 calves per household worth US\$ 202.7. When the effect of livestock loss
238 causes were pooled, disease were responsible for US\$ 83 (sum of mean losses
239 in cattle, goats and sheep) per household, while wild carnivores caused only
240 US\$ 12.6 per household (15.2% of loss due to diseases). On average, the value
241 of livestock sold per household was US\$ 57.8. This was 30.4% less than the
242 value the household lost due to diseases. Livestock losses due to theft and poor
243 management were US\$ 14.4 while animals slaughtered for meat worth US\$
244 16.5 per household. Each village cost-benefit analysis of each loss or gain
245 causes is summarized in Table 2.

246 **Discussion**

247 The results of this study suggest that diseases are responsible for higher
248 livestock loss than any other livestock loss causes within and among villages.
249 However, for sheep losses due to diseases and poor management differed
250 significantly among the villages. Mean number of cattle and goats sold per
251 household was higher than the number slaughtered in all villages. In each

252 household, new born calves were the significant source of replenishment of
253 livestock numbers. Mean number of livestock species that were sold,
254 slaughtered for food and that killed by spotted hyenas differed significantly
255 between livestock species whereby goats and sheep were more slaughtered for
256 food than cattle.

257

258 Disease is the major factor responsible for higher livestock losses in sub-
259 Saharan Africa (Gifford-Gonzalez, 2000). This factor alone, although not
260 realized by farmers in Africa (Mwangi, 1997), was responsible for a loss of US\$
261 83.5 per household during the nine months study period (Table 2). When this
262 figures was compared to average annual cash income per household in the
263 western Serengeti (US\$ 140, Borge, 2003), diseases were responsible for
264 59.6% of average annual household income in the target villages. On average,
265 diseases contributed to 5.1 times higher in livestock losses than depredation
266 cost. This observation is consistent with previous studies in the same area
267 when the farmers were requested to rank the major factors that were
268 responsible for higher livestock losses (Nyahongo, 2004). Livestock keepers
269 may not observe the direct effect of diseases to their livestock production due to
270 the fact that the sick animals may be slaughtered and used as food or sold to
271 neighbors while carnivores often consume all edible parts of a kill; leaving
272 nothing for human consumption. Moreover, diseases often kill larger number of
273 new born calves than adults (personal observation, 2006). Livestock keepers
274 may not observe this as an important loss because the capital investment in
275 terms of veterinary service, feeding or grazing time and/or output in terms of

276 meat or money (when sold) is relatively much lower for new-born calves than for
277 adults. Moreover, due to poor livestock management records, livestock keepers
278 may not be able to know how many livestock they loose to diseases within a
279 specific period of time. Most of household we visited did not have any record
280 showing number of livestock, new born or even the last time the animal was
281 treated and the cost implication. In contrast, when a predator breaks in the
282 livestock enclosures, usually at night (Nyahongo, 2004; Kolowski and
283 Holekamp, 2006; Holmern et al., 2007), it may kill several adult animals and this
284 may result in serious economic consequences for livestock keepers. However,
285 since the compensation scheme that may offset some of the costs are always
286 lacking in Tanzania, negative attitudes towards carnivores may have developed
287 among farmers, which have resulted in retaliatory killing practices of carnivores
288 in or close to village proximities (Holekamp and Smale, 1992; Ogada et al.,
289 2003; Dickman, 2005; Frank et al., 2005; Graham et al., 2005; Holmern et al.,
290 2007).

291

292 Sheep and goat depredation by spotted hyena was higher though not
293 statistically significant, in the village that was located far away from the park
294 boundary. This suggest that even in open areas with high anthropogenic
295 activities, still there are some refuges for some large carnivores like spotted
296 hyenas. This observation suggest a change in wildlife policies that insist
297 management of wildlife only in the established protected areas such as national
298 park, game reserves and game controlled areas. It would be of the conservation
299 interest and for the future conservation if wildlife management policies include

300 all wild animals in the protected areas as well as in anthropogenic dominated
301 areas. Certain carnivore species such as spotted hyenas have the ability to
302 commute up to 80 km (Hofer and East, 1993) allowing them to forage even in
303 villages located farther from the protected areas. The findings of the present
304 study gives an alternative idea that is inconsistent to the idea that high
305 depredation are only highest closest to the reserves boundary (Mwangi, 1997).
306 However, as Woodroffe (2000) puts it, behavioral plasticity of certain carnivore
307 species facilitate their adaptive adjustment to an increasingly precarious lifestyle
308 in proximity to human, a fact that was reported for spotted hyena of Maasai
309 Mara ecosystem (Boydston et al., 2003). However, it is difficult to establish that
310 the spotted hyena reported in the distant villages commuted from Serengeti or
311 were resident to the village areas.

312

313 Analyses of our data suggest that cattle are kept for solving household needs
314 that require relatively huge amount of money while goat and sheep are kept to
315 tackle small household needs and/or slaughtered to provide meat protein to the
316 household. This might be due to the fact that the economic value of one cattle is
317 equivalent to about four goats or sheep. These ideas are supported by
318 comparing the number of cattle, goats and sheep that were slaughtered and
319 those that were sold. The proportions of cattle slaughtered were far less than
320 those sold by households in the study villages (Table 1, Fig 2).

321

322 Variables like available water and grazing land, weather, market prices of meat
323 (that could lead to elevated theft rate), animal population dynamics in the

324 villages and in the protected areas adjacent to village areas, diseases
325 occurrence may, as the variables included in the analyses, show considerable
326 between year variations. These confounding variables, which cannot be
327 controlled for in a snap shot study like the present one, might have influenced
328 the data we collected. For instance, death of livestock due to diseases may
329 increase with drought or with rain intensity and duration, which cannot be
330 precisely compared within a year because intensity of rain and duration of rain
331 seasons may differ in different areas each year in Tanzania affecting pasture
332 quality and available water for animals. Drought may also influence the number
333 of livestock sold to buy food, because crop production in the country largely
334 depends on rain. Weather, on the other hand may influence the survival of new
335 born calves or may influence the level of depredation. Woodroffe and Frank
336 (2005) observed that rate of livestock depredation by large carnivores increase
337 with the increase in rainfall. Exclusion of households with very many animals
338 might have further led to underestimation of livestock loss because more death
339 from disease (due to density dependent danger of infectious diseases),
340 livestock depredation, theft and loss due to poor management in the grazing
341 field may be expected to increase with increase in livestock numbers.

342

343 **Concluding remarks**

344 The results from this study showed that diseases are the major cause of
345 livestock loss in the villages and that the levels of loss do not vary much among
346 the households in the western Serengeti. In contrast, livestock depredation by
347 spotted hyena was relatively low, although it was relatively higher for goats and

348 sheep in the households from the distant village. Likewise, poor management
349 and theft that can be managed at household level causes livestock losses as
350 well. However, at the household level, a single depredation event may cause a
351 serious economic loss.

352

353 Livestock depredation may be higher in the areas with high human activities,
354 which encourage wildlife managers, conservationists and wildlife ecologists to
355 think deeply about livestock depredation along the gradient of distance from the
356 park and the future conservation of the carnivores along the same gradient.

357

358 This study suggest that local people would benefit from better education on
359 animal husbandry practices and extension service to help them maintain the
360 health of their livestock and to prevent theft and loss of livestock while grazing.
361 We recommend that diseases control and management should be integral part
362 of regional and national development programs to limit disease transmission
363 between livestock and wildlife and even among livestock in the villages. Further
364 studies on the types and epidemiology of diseases causing major livestock
365 losses in the area should be conducted in order to design appropriate diseases
366 control measures.

367

368 **Acknowledgements**

369 The current study was funded by NORAD through the Ministry of Natural
370 Resources and Tourism, Tanzania and is a part of the Tanzania Wildlife
371 Research Institute (TAWIRI), Norwegian Institute for Nature Research (NINA)

372 and Norwegian University of Science and Technology (NTNU) project called
373 Biodiversity and Human Wildlife Interface in Western Serengeti. We
374 acknowledge the financial support from the Norwegian Peace corps and the
375 Quota Programme scheme (NORAD). We thank Commission of Science and
376 Technology (COSTECH) and Tanzania National Parks (TANAPA) for granting
377 permissions to the foreign researchers to work in Serengeti National Park.
378 Finally, we thank village leaders and our field assistants who in one way or the
379 other contributed to this important task.

380

381

382 References

- 383 Arcese, P., Hando J., Campbell, K. (1995) Historical and Present – Day Anti-
384 poaching Efforts in Serengeti In: Sinclair, A.R.E., Arcese P., (Eds)
385 Serengeti II – Dynamics, Management and Conservation of an
386 Ecosystem, pp. 506-533. The University of Chicago Press.
- 387 Berry, H. (1990) The Lions of Etch. *Cat News*, 13, 11-12.
- 388 Borge, A. (2003) Essays on the economics of African wildlife and utilization and
389 management. Dr. polit. thesis, Norwegian University of Science and
390 Technology, Trondheim, Norway, 187 pp.
- 391 Boydston, E.E., Kapheim, K.M., Watts, H.E., Szykman, M., Holekamp, K.E.
392 (2003). Altered behaviour in spotted hyenas associated with increased
393 human activity. *Animal Conservation*, 6, 1-10.
- 394 Campbell, K. L. I, Hofer, H. (1995) People and Wildlife: Spatial Dynamics and
395 Zones of Interaction. In: Serengeti II – Dynamics, Management and
396 Conservation of an Ecosystem (Eds. A.R.E., Sinclair and P. Arcese).
397 The University of Chicago Press, 534-570.
- 398 Dickman, A.J. (2005) An assessment of pastoralist attitudes and wildlife conflict
399 in the Rungwa-Ruaha region, Tanzania, with particular reference to
400 large carnivores. Msc. thesis. University of Oxford.
- 401 Frank, L.G., Woodroffe, R., Ogada, M.O. (2005) People and predators in
402 Laikipia District, Kenya. In: Woodroffe, R., Thirgood, S., Rabinowitz, A.
403 (Eds.), *people and wildlife conflicts or coexistence?* Cambridge
404 University Press, Cambridge, pp. 286-304.

- 405 Gifford-Gonzalez, D. (2000) Animal disease challenges to the emergence of
406 pastoralism in sub-Saharan Africa. *African Archaeological Review*, 17,
407 95-139.
- 408 Graham, K., Beckerman, A.P., Thirgood, S. (2005) Human-predator-prey
409 conflicts: ecological correlates, prey losses and patterns of
410 management. *Biological Conservation*, 122, 159-171.
- 411 Hofer, H., Campbell, K. L. I., East L.M., Huish S.A (1996) The impact of game
412 meat hunting on target and non - target species in the Serengeti. In: the
413 exploitation of mammal population. Chapman & Hall London. U.K.
- 414 Hofer, H., East, M.L. (1993) The commuting system of Serengeti spotted
415 hyenas: how a predator copes with migratory prey. I. Social
416 organization. *Animal Behaviour*, 46, 547-557.
- 417 Holekamp, K. E., Smale, L. (1992) Human-hyena relations in and around the
418 Maasai Mara National Reserve, Kenya. *IUCN SSC Hyena Specialist*
419 *Group Newsletter*, 5, 19-20.
- 420 Holmern, T., Nyahongo, J., Røskaft, E. (2007) Livestock loss caused by
421 predators outside the Serengeti National Park, Tanzania. *Biological*
422 *Conservation*, 135, 534-542
- 423 Infield, M. (1996) Livestock production and wildlife conservation: opportunities
424 for compatible management and integrated production. *East African*
425 *Livestock Conference, Small Ruminant Collaborative Research Support*
426 *Program, University of California. Entebbe, Uganda.*

- 427 Kaltenborn, B. P., Bjerke, T., Strumse, E. (1998) Diverging attitudes towards
428 predators: Do Environmental Beliefs Play a Part? *Human Ecology*
429 *Review*, 5, 1-9.
- 430 Kaltenborn, B. P., Bjerke, T., Vittersø, J. (1999) Attitudes towards large
431 carnivores among sheep farmers, wildlife managers and research
432 biologists in Norway. *Human Dimensions of Wildlife*, 4, 57-73.
- 433 Kaltenborn, B. P., Bjerke, T., Nyahongo, J.W., William, D.R., (2006) Animals
434 preferences and acceptability of wildlife management actions around
435 Serengeti National Park, Tanzania. *Biodiversity and Conservation*. 15,
436 4633-4649.
- 437 Kellert, S. R. (1985) Public Perception of Predators, Particularly the Wolf and
438 Coyote. *Biological Conservation*, 31, 167-189.
- 439 Kock, R.A. (2003) What is this infamous “wildlife/livestock diseases interface?”
440 A review of current knowledge for the African continent. PACE
441 Epidemiology Unit, AU/IBAR, Nairobi, Kenya. [http://wcs-](http://wcs-ahead.org/book/chapter01.pdf)
442 [ahead.org/book/chapter01.pdf](http://wcs-ahead.org/book/chapter01.pdf) (accessed 03-09-2007).
- 443 Kolowski, J.M., Holekamp, K.E. (2006) Spatial, temporal and physical
444 characteristics of livestock depredations by large carnivores along a
445 Kenyan reserve border. *Biological Conservation*, 128, 529-541.
- 446 Lindsey, P.A., du Toit, J.T., Mills, M.G.L. (2005) Attitudes of ranchers towards
447 African wild dogs *Lycaon pictus*: Conservation implications on private
448 land. *Biological Conservation* 125, 113-121.
- 449 Loibooki, M.T., Hofer, H., Campbell, K.L.I., East, M.L. (2002) Bush meat
450 Hunting by communities adjacent to the Serengeti National Park,

451 Tanzania: the importance of livestock ownership and alternative
452 sources of protein and income. *Environmental Conservation*, 29, 391-
453 398

454 Mills, G., Hofer, H. (1998) *Hyena: Status Survey and Conservation Action Plan*.
455 IUCN/SSC Hyena Specialist Group. Information Press, Oxford, UK.

456 Mwangi, N.S. (1997) A cost-benefit analysis of livestock predation on the group
457 ranches in the dispersal areas of Maasai Mara National Reserve,
458 Kenya. Mphil. thesis, Moi University.

459 Nabane, N. (1995) 'Lacking confidence? A gender sensitive analysis of
460 CAMPFIRE in Masoka Village.' *Wildlife and Development Series No.1*.
461 London: International Institute for Environment and Development.

462 Nabane, N. (1996) 'Zimbabwe: whose CAMPFIRE? Gender issues in
463 community- based conservation: the case of Masoka village'. *The Rural*
464 *Extension Bulletin. Special Issue on Community Conservation*, 10, 46-
465 49.

466 Nyahongo, J. W. (2004) *Impact of Human Activities on the Carnivore*
467 *Populations in the Western Corridor of Serengeti Ecosystem*. MSc
468 thesis. University of Dar Es Salaam, Tanzania.

469 Nyahongo, J.W., East, M.L., Mturi, F.A., Hofer, H. (2006) Benefits and costs of
470 illegal grazing and hunting in the Serengeti ecosystem. *Environmental*
471 *Conservation*, 32, 326-332.

472 Office International des Epizooties (2003) *Terrestrial animal health code 2003*.
473 Office International des Epizooties, Paris, France.

474 http://www.oie.int/eng/normes/MCode/A_summary.htm (accessed on
475 03-09-2007)

476 Ogada, M.O., Woodroffe, R., Oguge, N.O., Frank, L.G., (2003) Limiting
477 depredation by African carnivores: the role of livestock husbandry.
478 Conservation Biology, 17, 1521-1530.

479 Perry, B.D., Randolph, T.F., McDermott, J.J., Sones, K.R. and Thornton, P.K.
480 (2002) Investing in animal health research to alleviate poverty.
481 International Livestock Research Institute (ILRI), Nairobi, Kenya. 148pp

482 Rwambo, P., Grootenhuis, J. Demartini, J., Mkumbo, S. (1999) Assessment of
483 wildlife and livestock disease interactions in the Ngorongoro
484 Conservation Area Authority of Tanzania. Integrated Management and
485 Assessment System (IMAS).
486 [http://www.nrel.colostate.edu/projects/imas/prods/disease/NCAREP.ht](http://www.nrel.colostate.edu/projects/imas/prods/disease/NCAREP.html)
487 [ml](http://www.nrel.colostate.edu/projects/imas/prods/disease/NCAREP.html) (accessed on 04-09-2007).

488 Rasmussen, G.S.A. (1999) Livestock predation by the painted hunting dogs
489 *Lycaon pictus* in a cattle ranching region of Zimbabwe: a case study.
490 Biological Conservation, 88, 133-139.

491 Røskaft, E., Bjerke, T., Kaltenborn, B., Linnell, J.D.C., Andersen, R. (2003)
492 Patterns of self-reported fear towards large carnivores among the
493 Norwegian public. Evolution and Human Behavior 24:184-198.

494 Røskaft, E., Handel, B., Bjerke, T., Kaltenborn, B.P. (2007) Human attitudes
495 towards large carnivores in Norway. Wildlife Biology, 13, 172-185.

496 SPSS (2005). SPSS for Windows. Release 14.0. Chicago IL.

- 497 Stuart, C. T., Macdonald, I. A. W., Mills, M. G. L. (1985) History, Current Status
498 and Conservation of Large Mammalian Predators in Cape Province,
499 Republic of South Africa. *Biology Conservation*, 31, 7-19.
- 500 Thomson, G.R., Vosloo, W., Bastos, A.D.S. (2003) Foot and mouth disease in
501 wildlife. *Virus Research*, 19, 145-161.
- 502 Treves, A., Karanth, K.U. (2003) Human-carnivore conflict and perspectives on
503 carnivore management worldwide. *Conservation Biology* 17, 1491-
504 1499.
- 505 Treves, A., Naughton-Treves, L., Harper, E.K., Mlandenoff, D.J., Rose, R.A.,
506 Sickley, T.A., Wydeven, A.P. (2004) Predicting human-carnivore
507 conflict: a spatial model derived from 25 years of data on wolf predation
508 on livestock. *Conservation Biology*, 18, 114-125.
- 509 URT United Republic of Tanzania, (2002) 2002 population and housing census.
510 <http://www.tanzania.org.tz/census/index.html> (accessed 04.09.2007).
- 511 Vittersø, J., Kaltenborn, B. P., Bjerke, T. (1998) Attachment to livestock and
512 attitudes towards large carnivores among sheep farmers in Norway.
513 *Anthrozoos*, 11, 210-217.
- 514 Washington State Department of Agriculture (WASDA) (2007) Rural crime
515 prevention on livestock theft.
516 <http://agr.wa.gov/FoodSecurity/docs/livestockTheft.pdf> (accessed on
517 23-09-2007).
- 518 Woodroffe, R. (2000) Predators and people: using human densities to interpret
519 declines of large carnivores. *Animal Conservation*, 3, 165-173

520 Woodroffe, R. and Frank, L.G. (2005) Lethal control of African lions (*Panthera*
521 *leo*): local and regional population impacts. *Animal Conservation*, 81,
522 91-98.

523

524 **Biographical sketches**

525 Julius Nyahongo is a senior research scientist with the Tanzanian Wildlife
526 Research Institute (TAWIRI). He is a wildlife ecologist specializing in human-
527 wildlife interactions. He has worked intensively with human-wildlife interactions
528 in Tanzania.

529

530 Tomas Holmern is a researcher with the Department of Biology at the
531 Norwegian University of Science and Technology. He is interested in human-
532 wildlife conflicts and has recently been working on bushmeat hunting and local
533 law enforcement in the Serengeti.

534

535 Bård G. Stokke is a researcher at the Department of Biology, Norwegian
536 University of Science and Technology. He is an evolutionary biologist and
537 behavioural ecologist working on a wide range of birds and on human-wildlife
538 conflicts over the use of limited land.

539

540 Julius D. Keyyu is director of research in TAWIRI and he is interested in
541 veterinary science and conservation biology.

542

543 Bjørn Kaltenborn is a senior research scientist with the Norwegian Institute for
544 Nature Research (NINA). He is a geographer and social scientist specializing in
545 human-environment interactions. He has worked extensively with human-
546 wildlife conflicts in the Nordic countries as well as in East Africa and South Asia.

547

548 Eivin Røskaft is a behavioural ecologist interested in a wide range of birds and
549 mammals species in Europe, North America and Africa, and in human-wildlife
550 conflicts over the use of limited land.

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569 Table 1. Mean number of livestock per household and proportion of livestock loss or gain causes (livestock loss causes: diseases, loss in
 570 the bush (poor management while grazing), depredation and theft; livestock gain: newborn and bought/paid as dowry; household
 571 expenditure: sold and slaughtered for meat)

Livestock numbers and loss/gain	Nyamakendo					Nattambiso					Kowak					Overall		
	Cattle	Goats	Sheep	Cattle	Goats	Sheep	Cattle	Goats	Sheep	Cattle	Goats	Sheep	Cattle	Goats	Sheep	Cattle	Goats	Sheep
Mean numbers (\pm SD)	23.4 (17.2)	9.4 (6.0)	13.0 (22.9)	15.2 (12.9)	13.9 (12.2)	8.3 (14.0)	21.6 (12.2)	16.8 (15.0)	14.6 (14.7)	22.5 (22.1)	8.5 (11.7)	9.0 (11.8)	20.5 (16.8)	12.1 (11.2)	9.0 (11.8)	20.5 (16.8)	12.1 (11.2)	11.2 (15.8)
<u>Livestock gain (%)</u>																		
Newborn	10.3	21.3	16.2	5.9	15.1	10.8	9.3	16.1	18.5	5.3	11.8	11.1	7.7	16.1	7.7	16.1	14.2	14.2
Bought	1.7	1.1	0.8	3.3	3.6	1.2	2.3	2.4	2.0	0.9	2.3	2.2	2.1	2.4	2.1	2.4	1.6	1.6
<u>Livestock loss (%)</u>																		
Diseases	3.4	4.3	5.4	2.6	6.5	2.4	5.1	10.1	5.5	3.1	7.1	6.7	3.5	7.0	3.5	7.0	5.0	5.0
Loss in the bush	0.4	0	1.5	0.2	1.4	0	0.5	1.2	1.4	0.4	1.2	1.1	0.4	0.9	0.4	0.9	1.0	1.0
Depredation (%)	0.4	0.3	1.5	0.1	0.7	1.2	0.1	1.8	0.7	0.3	4.7	5.6	0.2	1.9	0.2	1.9	2.2	2.2
Theft	0	0	0.1	0.2	0.1	1.2	0.1	0.1	0.1	0.2	0.3	0.3	0.1	0.1	0.1	0.1	0.1	0.4
Household expenditure (%)																		
Sold	2.1	3.2	4.6	4.6	5.8	2.4	2.8	2.4	1.4	1.3	3.5	4.4	2.7	3.7	2.7	3.7	3.2	3.2
Slaughtered	0.4	1.1	0.8	0.5	1.4	1.2	0.5	3.0	2.0	0.1	2.3	2.2	0.4	2.0	0.4	2.0	1.6	1.6
Mean recruitment (%)	5.3	13.5	3.1	1.0	2.8	3.6	2.5	-0.1	9.4	0.8	-5.0	-7.0	2.4	2.8	2.4	2.8	2.3	2.3

572 Note: Sample sizes: Robanda (n = 37 households for cattle, n = 10 for goats and n = 15 for sheep); Nyamakendo (n = 49 households for cattle, n = 49
 573 for goats and n = 26 for sheep); Nattambiso (n = 46 households for cattle, n = 45 for goats and n = 28 for sheep); Kowak (n = 50 households for cattle,
 574 n = 51 for goats and n = 54 for sheep).
 575
 576
 577
 578

579

580

581

582

Table 2: Cost and benefit implications of livestock loss and/or gain causes (US \$)

Livestock numbers and loss/gain	Robanda			Nyamakendo			Nattambiso			Kowak			Overall values (US\$)		
	Cattle	Goats	Sheep	Cattle	Goats	Sheep	Cattle	Goats	Sheep	Cattle	Goats	Sheep	Cattle	Goats	Sheep
Mean value of livestock	1872.0	188.0	260.0	1216.0	278.0	166.0	1728.0	336.0	292.0	1800.0	170.0	180.0	1654.0	243.0	224.0
<u>Livestock gain values (US\$)</u>															
Newborn	192.8	40.0	42.1	71.7	42.0	17.9	160.7	54.1	54.0	95.4	20.1	20.0	130.1	39.1	33.5
Bought	31.8	2.1	2.1	40.1	10.0	2.0	39.7	33.9	5.8	16.2	3.9	4.0	31.9	12.5	3.5
<u>Livestock loss (US\$)</u>															
Disease	63.6	8.1	14.0	31.6	18.1	4.0	88.1	10.1	16.1	55.8	12.1	12.1	59.8	12.1	11.6
Depredation	7.5	0.6	3.9	1.2	1.9	2.0	1.7	6.0	2.0	5.4	8.0	10.1	4.0	4.1	4.5
Loss in the bush	7.5	0	3.9	2.4	3.9	0	8.6	4.0	4.1	7.2	2.0	2.0	6.4	2.5	2.5
Theft	0	0	0.3	2.4	0.3	2.0	1.7	0.3	0.3	3.6	0.5	0.5	1.9	0.3	0.8
<u>Household expenditure (US\$)</u>															
Sold	39.3	6.0	12.0	55.9	16.1	4.0	48.4	8.1	4.1	23.4	5.9	7.9	41.8	9.0	7.0
Slaughtered	7.5	6.0	2.1	6.1	3.9	2.0	8.6	10.1	5.8	1.8	3.9	4.0	6.0	6.0	4.5

583 Note: Mean local market price of one cattle in the study area was US\$ 80, and for goat/sheep was US\$ 20 in 2006, (the prices were for mature animals).

584

585

586

587

588

589 Figure legends:

590 Figure 1. Map of the western Serengeti showing the sampled villages.

591 Figure 2. Overall livestock population dynamics (loss and gain) in four villages
592 recorded from April to December 2006.

593

594

595

596

597

598

599


600

601

602

603

604




605

606 Figure 1.

607

608



609

610

611 Figure 2.

Paper IV

1 **Spatial-temporal variation in meat and fish consumption among**
2 **humans in the western Serengeti, Tanzania: the importance of**
3 **migratory herbivores**

4

5 Running headline: meat and fish consumption in Serengeti

6

7 Authors:

8 J. W. NYAHONGO^{1,3*}, T. HOLMERN¹, B. P. KALTENBORN² AND E. RØSKAFT¹

9 ¹Department of Biology, Norwegian University of Science and Technology,
10 Realfagbygget, N-7491, Trondheim, Norway

11 ²Norwegian Institute for Nature Research, Fakkeltgården, 2624, Lillehammer, Norway

12 ³Tanzania Wildlife Research Institute, P. O. Box 661, Arusha, Tanzania

13

14

15 Word counts: Abstract 219; Main text (including references and biographical sketches)

16 4969

17 Key words: Bushmeat, fish, illegal hunting, migratory herbivores, western Serengeti,

18

19

20

21 *Author for correspondence: email: nyahongo@bio.ntnu.no or

22 nyhwjulius@yahoo.co.uk

23 Tel.: +47-73550348

24

25

26 **Abstract**

27 Illegal bushmeat hunting has become a serious problem for wildlife managers in
28 many African countries. We investigated the spatial and temporal pattern in meat and
29 fish consumption by people surrounding the Serengeti National Park, Tanzania to
30 further understand the links between hunting and consumption. We studied 150
31 households in five villages during March – December 2006, along a gradient from the
32 National Park boundary up to 80 km away. In addition, two parallel 10 km transects
33 were conducted monthly near three villages closest to the national park in order to
34 investigate the relationship between household meat consumption and the influx of
35 migratory herbivores. We found that the number of meat meals was higher in the
36 villages closest to protected areas. The weekly number of meat meals per household
37 in all villages within 30 km from the national park boundary increased with the influx
38 of migratory herbivores. Moreover, meat consumption was unrelated to income,
39 except in the most distant village where there was a positive correlation. The number
40 of fish meals in the closest villages to the national park decreased with the influx of
41 migratory herbivores. We recommend a coordinated management of fish harvesting
42 from Lake Victoria and wildlife conservation around the Serengeti National Park to
43 implement a sustainable management of the two ecologically different natural
44 resources in the future.

45 **Introduction**

46 Since immemorial time, local communities have relied heavily on use of natural
47 resources such as water supply, forest products, grazing land, firewood and
48 bushmeat. Bushmeat, derived from wild animals, is an important source of cheap
49 protein for African and Latin American societies where in some countries (i.e. Liberia)
50 up to 75% of total meat consumed is derived from wild animals (e.g. Anstey, 1991;
51 Barnett, 2000; Rao & McGowan, 2002). An extensive use of bushmeat has been well
52 documented in West and Central Africa (Fa *et al.*, 2003; Wilkie *et al.*, 2005). Although
53 the issue has received much less attention in East and Southern Africa, studies
54 suggest that illegal bushmeat hunting has developed to become a serious problem
55 for wildlife managers, because of a growing demand for bushmeat, burgeoning
56 human populations (e.g. the human populations of Kenya and Tanzania increased
57 from 6 and 8 million in 1950 to 34 and 38 million in 2005, respectively) and increasing
58 commercialisation of the bushmeat trade (Edroma & Kenyi, 1985; Dublin *et al.*, 1990;
59 Campbell & Hofer, 1995; Barnett, 2000; UN, 2005). For instance, in Tanzania
60 partially protected areas (IUCN category \geq IV) appear to be particularly hard-hit by
61 illegal bushmeat hunting (Caro *et al.*, 1998), combined with high rates of habitat
62 degradation (Pelkey *et al.*, 2000).

63

64 Little quantitative research has been conducted on the factors that drive consumer
65 demand for bushmeat in poor tropical countries (Wilkie & Godoy, 2001). Increases in
66 household income appear to drive a shift in preference from bushmeat to the meat of
67 domesticated animals. Albrechtsen *et al.* (2005) found that income was positively
68 correlated with volume of small livestock meats consumed per household, but
69 negatively related with bushmeat eaten. This indicates that it is the poor that rely

70 mostly on bushmeat. However, the picture is not so clear, since Barnett (2000)
71 reported that there is also an increasing trend for a preference for bushmeat by the
72 affluent elite in urban areas.

73

74 In addition to wildlife, fish also is a vital source of animal protein in Sub-Saharan
75 Africa (FAO, 2004). The great lakes of East Africa, Lake Victoria, Tanganyika and
76 Nyasa, plays a key role in this respect. However, several studies report that catches
77 are declining due to overexploitation, pollution and environmental degradation (i.e.
78 exotic species introductions, water hyacinth (*Eichhornia crassipes*)) (Matsuishi *et al.*,
79 2006; Balirwa, 2007). For example, on the Kenyan side of Lake Victoria, the
80 important fishery on Nile perch (*Lates niloticus*) has declined steadily from a peak of
81 115,000 tons in 1999 to 57,000 tons in 2004, mainly due to lack of enforcement of
82 fishing regulations and lack of involvement of local stakeholders in fisheries
83 management that has led to overexploitation (Njiru *et al.*, 2007). Similar declines due
84 to over-harvesting have also been noted for other fish species such as Nile tilapia
85 (*Oreochromis niloticus*) and the small indigenous cyprinid (*Rastrineobola argentea*)
86 (Matsuishi *et al.*, 2006; Njiru *et al.*, 2007). The diminishing fish resource mainly due to
87 commercial fishing operations has also negative consequences for local communities
88 along the shores of Lake Victoria who are increasingly using smaller gill nets and in
89 some instances illegal and destructive fishing practices to meet household needs
90 (Balirwa, 2007).

91

92 Recently in West Africa, Brashares *et al.* (2004) indicated that fish and bushmeat
93 exploitation is linked, where low regional supplies of fish caused an intensification of
94 local bushmeat hunting. However, the direction of the linkage is debated (Rowcliffe *et*

95 *al.*, 2005), where for instance Wilkie *et al.* (2005) showed that bushmeat availability
96 might also affect the consumption of fish. These studies strongly suggest that the
97 unsustainable fishing and deterioration of the resource base in Lake Victoria does not
98 only have severe implication for local communities, but might also entail serious
99 consequences for wildlife within the bordering protected areas. There is therefore an
100 urgent need to assess the relationship between the fish and bushmeat resources in
101 the lake region and their role in human welfare.

102

103 In this study we therefore focused upon the relationship between the consumption of
104 fish and bushmeat among villages bordering Lake Victoria and the Serengeti National
105 Park (SNP), Tanzania. By conducting transects within different parts of the national
106 park, we investigated how the seasonal presence of high densities of chiefly
107 migratory herbivores and other socio-economic factors affected consumption levels
108 by using villages along a gradient from the resource source (i.e. lake or park).

109

110 **Methods**

111 **Study area**

112 **Climate and large mammals**

113 This study was conducted in the north-western part of the Serengeti National Park
114 (SNP) (Fig. 1) between March and December 2006. The SNP (14,763 km²) is the
115 largest park in Tanzania. In the west the park is bordered by Ikorongo Game Reserve
116 (ca. 563 km²), Grumeti Game Reserve (ca. 416 km²) and the Ikoma Open Area (ca.
117 600 km²) that act as buffer zones between the park and the village areas. The
118 common large resident herbivores in the area include giraffe, buffalo, topi
119 (*Damaliscus korrigum*) and impala (*Aepyceros melampus*). The western corridor of

120 the SNP is characterised by the annual wildebeest migration, which in June – August
121 moves through the partially protected and village areas on their way north (Thirgood
122 *et al.*, 2004). In addition to the migratory herbivores there is also a resident
123 population of wildebeest in the western corridor, that move towards Lake Victoria
124 during the wet season (Maddock, 1979). However, the populations of the resident
125 herbivores in the game reserves are relatively low probably due to high levels of
126 illegal bushmeat hunting (Campbell & Hofer, 1995; Rusch *et al.*, 2005).

127

128 **Local people and livelihoods**

129 Human population density is high (70 people / km²) in the north-western Serengeti,
130 with an annual growth rate of 2.5% between 1988 and 2002 (URT, 2002). The
131 economy of local communities is mainly based on subsistence farming and livestock
132 husbandry. However, erratic rainfall, poor soils, tse tse fly infestation makes farming
133 and keeping livestock very difficult, and alternative sources of protein are therefore
134 vital. For the villages adjacent to Lake Victoria fishing is considered important, where
135 fish species like Nile perch, tilapia and the small indigenous cyprinid (locally called
136 dagaa) dominate the catch. Fishing is done mainly with small gill nets from dug out
137 canoes throughout the year. Fishing in Lake Victoria is not legally restricted and there
138 is no limitation on the number of boats or gears used (Cowx *et al.*, 2003). In addition
139 to fish, many of the subsistence farmers and livestock keepers surrounding the SNP
140 also rely on bushmeat hunting in terms of food security and income generation
141 (Loibooki *et al.*, 2002; Holmern *et al.*, 2002; Kaltenborn *et al.*, 2005).

142

143 In the areas outside the park trophy hunting as well as resident legal hunting is
144 carried out. Legal offtake is low because quotas are set conservatively (Holmern *et*

145 *al.*, 2004). On the other hand, illegal bushmeat hunting originating from the local
146 communities in the west is very common. The great majority of arrested hunters in
147 the protected areas come from villages within 45 km from the closest boundary
148 (Campbell & Hofer, 1995; Holmern *et al.*, 2007). The main hunting method is the use
149 of wire snares, but also other methods are being used (Arcese *et al.*, 1995; Holmern
150 *et al.*, 2006). Bushmeat is commonly sun dried before being transported on foot to
151 the villages. The hunters use dried meat for home consumption, to sale in order to
152 generate income and/or to barter for other commodities (Loibooki *et al.*, 2002;
153 Kaltenborn *et al.*, 2005). An estimated 53,000 people are involved in illegal hunting,
154 including both hunters and porters that transport the meat out of the protected areas
155 (Loibooki *et al.*, 2002). The annual offtake of meat from the ecosystem has been
156 estimated at approximately 11,950 tons (Hofer *et al.*, 1996).

157

158 **Data collection**

159 Five villages were randomly selected along a gradient of distance from the SNP
160 boundaries. Three villages, Robanda, Nyamakendo and Nyatwali, were located
161 within 10 kilometres from the park, with Nyatwali also being located near to Lake
162 Victoria (see Fig 1). Thirty households were randomly selected from the list of
163 households from each village office. Household was defined as including all people in
164 the living quarters that permanently lived there (where we used last name to identify
165 temporary visitors in the household). Every household was visited once a month and
166 was requested to produce the data on the number of meals that consisted of meat,
167 fish and vegetables during the last week preceding the visit.

168

169 During the pre-testing of the questionnaire it was noted that most households were
170 unable to remember the amount of meat (in kg) they had bought or consumed and in
171 some cases some butchers did not have the weighing machines (i.e. relied on hand
172 estimates). Moreover, in most local markets, fish were sold as individuals or as bulk
173 and sometimes sold per volume using specified containers. We therefore solely
174 collected information on the number of meals of fish and meat. Several questions
175 were asked before raising the meat consumption issue. This was important due to
176 the sensitivity of this subject to local communities. It was initially attempted to
177 differentiate between legal meat and illegal bushmeat, but it was not possible due to
178 deep reluctance among the respondents to talk about bushmeat hunting. However, a
179 significant proportion of meat consumed in rural Africa is usually bushmeat (Barnett
180 *et al.*, 2000). Each household member was allowed to enter the discussion as it was
181 previously noted that one member may not remember well the number and type of
182 meals they had been eating in the previous seven days and/or might not be at home
183 every day. Children were encouraged to join the discussion because in most cases
184 they are at home and literally consume every meal prepared. Relying partly on
185 responses from children in order to get an unbiased result has also been used by
186 others (Haule *et al.*, 2002). The questionnaire also included general information on
187 the household size (number of people), age (of respondent), sex, occupation status,
188 education level, number of livestock (physically counted in the boma) and the
189 monthly household income (the average estimated from three consecutive month's
190 income).

191

192 In order to investigate the relationship between the number of meat meals the
193 households consumed and influx of migratory herbivores, two parallel transect lines

194 (each spaced one kilometre apart) were established at Robanda, Nyamakendo and
195 Nyatwali sampling sites. These were run once each month. Each transect line was 10
196 km long. All animals sighted within 200 metres on either side of a vehicle were
197 recorded for subsequent density calculation. We counted only herbivores because
198 these are the targeted species of illegal bushmeat hunting, but we excluded
199 elephants (*Loxodonta africana*) because they are not usually killed for bushmeat.
200 Moreover the animals were placed into two groups according to their seasonal
201 movement within the ecosystem. The migratory group included wildebeest, zebra,
202 eland (*Taurotragus oryx*), and Thomson gazelle (*Gazella thomsoni*); the remaining
203 herbivores were grouped as resident animals. All the three transect originated from
204 the boundary and was driven towards the interior of the park. The animals were
205 counted three hours after and before dawn and dusk, respectively.

206

207 **Statistical analyses**

208 Differences between samples were tested using non-parametric tests due to non-
209 normality in distribution. We also used Pearson correlation to explore the correlation
210 between explanatory variables. We used linear-mixed models, to account for the
211 spatial pseudo-replication of households within the villages, in order to explore the
212 variables influencing the number of meat and fish meals in the households during the
213 study period (10 months), with village set as a random effect (see Crawley, 2002).
214 The predictor variables included in the models were: livestock numbers, distance to
215 resource (in km), household size (HH size), monthly income (income) and education
216 (none, primary, secondary/college). The same was done for the analysis on fish
217 meals where the variable distance to resource was calculated from the Lake shore.
218 Since the use of stepwise multiple regression in ecology has been criticised for

219 having several drawbacks (Whittingham *et al.*, 2006), we instead used an information
220 theoretic approach which allows for several competing models to describe the data.
221 We evaluated the strength of evidence for the model based on Akaike Information
222 Criterion corrected for small samples (AIC_c) following Burnham and Anderson (2002),
223 and selected the most parsimonious model with highest Akaike weights (ω_i). The
224 Akaike weight indicates the probability that the model is the best among the whole
225 set of candidate models and was used to compare the relative performance of
226 models rather than only absolute AIC_c. The analyses were performed using SPSS
227 14.0 (SPSS, 2001) and R 2.3.0 Software (R Development Core Team, 2006) with
228 significant levels set at 0.05. We reported \pm 1 standard deviation (SD) throughout,
229 except for the linear mixed effect models where SE is used.

230

231 **Results**

232 **Household characteristics and consumption**

233 Generally, the studied local communities had many similar socio-economic
234 characteristics (Table 1 and Table 2). Across the five villages the number of meat
235 meals were higher in the villages located close to the SNP boundary (Kruskal-Wallis,
236 $H = 85.2$, $df = 4$, $p = 0.0001$). Similarly, there was a significant difference in the
237 number of fish meals between villages (Kruskal-Wallis, $H = 79.9$, $df = 4$, $p = 0.0001$)
238 where the villages close to Lake Victoria had the highest number of fish meals. The
239 household size had a negative effect on fish consumption in Nyatwali village although
240 the same variables positively influenced the consumption of fish in Rwamkoma
241 village ($r_p = -0.372$, $n = 30$, $p = 0.043$; $n = 30$, $r_p = 0.370$, $p = 0.040$, respectively).
242 Furthermore, in Kowak village there was a significant positive correlation between

243 income and meat consumption ($r_p = 0.521$, $n = 30$, $p = 0.003$). Whereas none of the
244 tested variables had any influence on meat consumption in Robanda village (Table 1).
245

246 **The influence of migratory and resident herbivores**

247 Households who were close to the SNP but far from Lake Victoria consumed more
248 meat and less fish during the influx of migratory herbivores. Mean weekly household
249 meat consumption at Robanda, Nyatwali, Nyamakendo and Rwamkoma villages
250 increased when the densities of migratory herbivores increased (Robanda: $r_p = 0.920$,
251 $n = 10$ months, $p < 0.001$; Nyatwali: $r_p = 0.711$, $n = 10$ months, $p = 0.021$;
252 Nyamakendo: $r_p = 0.953$, $n = 10$ months, $p = 0.0001$; Rwamkoma: $r_p = 0.682$, $n = 10$
253 months, $p = 0.03$; Kowak: $r_p = -0.100$, $n = 10$ months, $p = 0.783$) (Fig 2 a-e).

254
255 In contrast fish consumption was negatively correlated to the densities of migratory
256 herbivore close to Robanda, Nyamakendo and Rwamkoma villages (Robanda: $r_p = -$
257 0.684 , $n = 10$ months $p = 0.029$; Nyamakendo: $r_p = -0.684$, $n = 10$ months, $p = 0.030$;
258 Rwamkoma: $r_p = -0.813$, $n = 10$ months, $p = 0.004$). No significant correlation was
259 found between mean household fish consumption and densities of migratory
260 herbivores for Nyatwali and Kowak villages ($r_p = 0.538$, $n = 10$ months, $p = 0.109$; $r_p =$
261 0.247 , $n = 10$ months, $p = 0.492$, respectively). In all villages, the mean household
262 fish consumption were uncorrelated to the densities of resident herbivores (Robanda:
263 $r_p = -0.289$, $n = 10$ month, $p = 0.418$ s; Nyatwali: $r_p = -0.293$, $n = 10$ months, $p = 0.412$;
264 Nyamakendo: $r_p = 0.260$, $n = 10$ months, $p = 0.464$; Rwamkoma: $r_p = -0.613$, $n = 10$
265 months, $p = 0.059$; Kowak: $r_p = 0.329$, $n = 10$ months, $p = 0.354$).

266

267 **Factors affecting the consumption of meat and fish**

268 There were two models that had a $\Delta AIC_c < 2$ in the analysis on meat consumption
269 (Table 3). The most parsimonious model included the variables livestock and
270 distance to the protected area, where increasing livestock numbers led to a higher
271 consumption, whereas increasing distance to the protected area decreased the
272 number of meat meals (Table 3 and 4). The predictor variables present in the most
273 parsimonious model also ranked high in importance (Table 5) when we summed AIC_c
274 weights of models containing them over the whole set of candidate models. In the
275 analysis on fish consumption, there were also two models that had a $\Delta AIC_c < 2$. The
276 most parsimonious model here included both the variable distance to Lake Victoria
277 and distance to the protected area. Here both increasing distance to Lake Victoria
278 and the protected area reduced the number of fish meals in households, these
279 variables also ranked high in importance (Table 5).

280

281 **Discussion**

282 Our results suggest that households that are close to protected areas consume more
283 meat during the period of the migration than those located farther away where the
284 peak meat consumption in the villages that were close to SNP corresponded to the
285 peak influx of migratory herbivores. Moreover, increased supply of fish did not seem
286 to reduce consumption of meat when this was readily available. In most villages
287 income did not influence meat consumption, only in the most distant ones.

288

289 Although our data cannot distinguish between the legal and illegal sources of
290 consumed meat, the positive correlation between influx of migratory herbivores and
291 the mean number of meat meals consumed in the villages that were close to the park
292 boundary suggest an extended utilization of bushmeat during this period. This

293 interpretation is further supported by other studies, which show that rural residents
294 (and increasingly even in urban areas) rely to a large extent on bushmeat for most of
295 their animal protein (Barnett, 2000; Fa *et al.*, 2003; Albrechtsen *et al.*, 2005).
296 However, one of the study villages (i.e. Robanda) is included in the Serengeti Region
297 Conservation Program (SRCP) game cropping scheme, but this operation has very
298 low quotas that are unlikely to impact the results (Holmern *et al.*, 2002; 2004). Since
299 extremely few residents have had a license to hunt, it may be assumed that the
300 excess meat consumed in the villages that were close to SNP during the influx of
301 migration was illegally obtained. This is supported by the observation of meat
302 consumption at Kowak village that was about 80 km from the park boundary where
303 the meat consumption increased with income and not with influx of migratory
304 herbivores.

305

306 The seasonal availability of herbivores also affects bushmeat prices that are almost
307 halved when the migration arrives in the village areas (Holmern *et al.*, 2002).
308 Bushmeat prices are unrelated to wildlife species, but are primarily determined by the
309 weight of the dry meat pieces which is the most common unit of trade (Holmern, 2000;
310 Holmern *et al.*, 2002). Although there is evidence for a preference for certain
311 bushmeat species among the different tribes (Ndibalema & Songorwa, 2007),
312 generally the identification of species by meat taste is poor (Nyahongo *et al.*, in prep.).
313 The limited cash availability among the very poor people in this area, together with
314 the fact that fish and domestic meat is more expensive (in the villages distant from
315 Lake Victoria) than bushmeat, means that these alternatives cannot currently out-
316 compete wildlife as the primary protein source.

317

318 Furthermore, the access to meat or fish sources may depend on several other factors,
319 such as logistical difficulties because of having to circumvent impassable areas,
320 likelihood of being arrested, the distance to travel and time to reach the profitable
321 source area may also reduce the utility to such resources (Nyahongo *et al.*, 2006).
322 For example, the distance between Kowak village and the closest point to Lake
323 Victoria is only 17.8 km by air (Table 2) but the access is denied due to a large area
324 in between covered by swamp, steep hills, gullies and thick acacia bushes. Thus,
325 people are forced to take a longer route (i.e. use the old Sirari-Musoma road, where
326 the distance becomes approximately 60 km) to reach the closest fishing station
327 located at Kinesi along the shore of Lake Victoria.

328

329 Mean household meat and fish consumption per month fluctuated with movement of
330 migratory herbivores in or close to villages that were less than 30 km from the park.
331 In the western Serengeti, it is generally known that during the influx of wildebeest
332 migrations huge herds of animals roam through villages and agricultural lands and
333 large numbers of animals may be slaughtered literally at the doorstep. Therefore, it is
334 not surprising to see the higher household meat consumption during the influx of
335 migratory herbivores. Similar higher household meat consumption was observed
336 during the influx of migratory herbivores in the study area close to Robanda, Nyatwali,
337 Nyamakendo and Rwamkoma villages where in some villages the peak meat
338 consumption corresponded to the peak migration (Fig. 2a-d). There are also strong
339 traditional cultural motives for hunting in this region (Kaltenborn *et al.*, 2005), which
340 may further influence the consumption and resistance to the introduction of
341 alternative animal protein sources. However, the consumption of fish was relatively
342 low in the above villages during the peak migration suggesting that bushmeat and

343 fish may complement each other, especially in the villages that are farther from the
344 lake but located close to the park boundary.

345

346 **Concluding remarks**

347 In order to reduce the dependence on bushmeat, alternative sources of meat protein
348 together with some income generating projects need to be considered in the general
349 local and national development planning. The contribution of fish to household diet as
350 an alternative to bushmeat should be emphasized so that the limitation on processing
351 fresh fish and transportation to local market is solved. Industrial harvesting of fish
352 from the Lake Victoria need to be coordinated as it may reduce availability of fish to
353 the local market in both villages located close to the lake and the distant villages and
354 thus increasing the pressure and reliance on bushmeat. Therefore the policy markers
355 need to understand the link and the need for coordinated management between
356 these two ecologically very different resources.

357

358 **ACKNOWLEDGEMENTS**

359 This study was funded by NORAD through the Ministry of Natural Resources and
360 Tourism, Tanzania and is a part of the TAWIRI, NINA and NTNU project called
361 “Biodiversity and Human Wildlife Interface in Western Serengeti”. We acknowledge
362 the financial support from the Norwegian Fredskorpset and the Quota Programme
363 scheme (NORAD). We also thank Tanzania National Parks (TANAPA) for granting
364 permissions to the foreign research team to work in Serengeti National Park. Finally,
365 we thank village leaders, the respondents and our field assistants for their help. Ms
366 Claudia Melis assisted with graphs.

367

368 **References**

- 369 Albrechtsen, L., Fa, J.E., Barry, B. & Macdonald, D.W. (2005) Contrasts in availability
370 and consumption of animal protein in Bioko Island, West Africa: the role of
371 bushmeat. *Environmental Conservation*, **32**, 340-348.
- 372 Anstey, S. (1991) Wildlife Utilisation in Liberia. WWF/FDA Wildlife Survey Report.
- 373 Arcese, P., Hando J. & Campbell, K. (1995) Historical and Present-Day Anti-
374 poaching Efforts in Serengeti. In *Serengeti II – Dynamics, Management and*
375 *Conservation of an Ecosystem* (eds A.R.E. Sinclair & P. Arcese), pp. 506-
376 533. The University of Chicago Press, Chicago, USA.
- 377 Balirwa, J.S. (2007) Ecological, environmental and socioeconomic aspects of the
378 Lake Victoria's introduced Nile perch fishery in relation to the native fisheries
379 and the species future potential: lessons to learn. *African Journal of Ecology*,
380 **45**, 120-129.
- 381 Barnett, R. (2000) *Food for Thought. The utilization of wild meat in Eastern and*
382 *Southern Africa*. TRAFFIC East/Southern Africa, pp. 264. Nairobi, Kenya.
- 383 Brashares, J.S., Arcese, P. Sam, M.K., Coppolillo, P.B., Sinclair, A.R.E. & Balmford,
384 A. (2004) Bushmeat hunting, wildlife decline and fish supply in West Africa.
385 *Science*, **306**, 1180-1183.
- 386 Burnham, K.P. & Anderson, D.R. (2002) Model Selection and Multimodel Inference:
387 A practical Information-Theoretic Approach. Springer, New York.
- 388 Campbell, K.L.I & Hofer, H. (1995) People and Wildlife: Spatial Dynamics and Zones
389 of Interaction. In *Serengeti II–Dynamics, Management and Conservation of*
390 *an Ecosystem* (eds A. R. Sinclair & P. Arcese), pp. 534-570. The University
391 of Chicago Press, Chicago, USA.

392 Caro, T.M., Pelkey, N., Borner, M., Campbell, K.L.I., Woodworth, B.L., Farm, B.P.,
393 Ole Kuwai, J., Huish, S.A. & Severre, E.L.M. (1998) Consequences of
394 different forms of conservation for large mammals in Tanzania: preliminary
395 analyses. *African Journal of Ecology*, **36**, 303-320.

396 Cowx I.G., van der Knaap, M., Muhoozi, L.I. & Othina, A. (2003) Improving Fishery
397 Catch Statistics for Lake Victoria. *Aquatic Ecosystem Health & Management*.
398 **6**, 299-310.

399 Crawley, M.J. (2002) Statistical Computing: An introduction to data analysis using S-
400 Plus. John Wiley & Sons Ltd, England.

401 Dublin, H.T., Sinclair, A.R.E., Boutin, S., Anderson, E., Jago, M. & Arcese, P. (1990)
402 Does competition regulates ungulate populations? Further evidence from
403 Serengeti, Tanzania. *Oecologia*, **82**, 283-288.

404 Edroma, E.L. & Kenyi, J.M. (1985) Drastic decline in bohor reedbuck (*Redunca*
405 *redunca* Pallas 1977) in Queen Elizabeth National Park, Uganda. *African*
406 *Journal of Ecology*, **23**, 53-55.

407 Fa, J.E., Currie, D., & Meeuwig, J. (2003) Bushmeat and food security in the Congo
408 Basin: linkage between wildlife and people's future. *Environmental*
409 *Conservation*, **30**,71-78.

410 FAO (Food and Agriculture Organization) (2004) Aquaculture Extension in Sub-
411 Saharan Africa. Food and Agricultural Organisation of the United Nations FAO.
412 Fisheries Circular No. 1002. 55pp.

413 Haule, K.S., Johnsen, F.H & Maganga, S.L.S (2002) Striving for sustainable wildlife
414 management: the case of Kilombero Game Controlled Area, Tanzania.
415 *Environmental Management*, **66**, 31-42.

- 416 Hofer, H., Campbell, K.L.I., East, L.M. & Huish, S.A. (1996) The impact of game meat
417 hunting on target and non-target species in the Serengeti. In *The exploitation*
418 *of mammal population* (eds V.J. Taylor & N. Dunstone), pp. 117-146.
419 Chapman & Hall, London, U.K.
- 420 Holmern, T. (2000) Hunting and community-based conservation outside the
421 Serengeti National Park – A sustainable approach by Serengeti Regional
422 Conservation Project (SRCP)? MSc thesis, Norwegian University of Science
423 and Technology, Trondheim, Norway.
- 424 Holmern, T., Røskaft, E., Mbaruka, J.Y., Mkama, S.Y. & Muya, J. (2002)
425 Uneconomical game cropping in a community-based conservation project
426 outside the Serengeti National Park, Tanzania. *Oryx*, **36**, 364-372.
- 427 Holmern, T., Johannesen, A.B., Mbaruka, J., Mkama, S., Muya, J. & Røskaft, E.
428 (2004) *Human-Wildlife Conflicts and Hunting in the Western Serengeti,*
429 *Tanzania*, pp. 26. NINA Project Report, **26**.
- 430 Holmern, T., Mkama, S. Y., Muya, J. & Røskaft, E. (2006) Intraspecific prey choice of
431 bushmeat hunters outside the Serengeti National Park, Tanzania: a
432 preliminary analysis. *African Zoology*, **41**, 81-87.
- 433 Holmern, T., Muya, J. & Røskaft, E. (2007) Local law enforcement and illegal
434 bushmeat hunting outside the Serengeti National Park, Tanzania.
435 *Environmental Conservation*, **34**, 55-63.
- 436 Kaltenborn, B.P., Nyahongo, J.W. & Tingstad K.M. (2005) The nature of hunting
437 around the Western Corridor of Serengeti National Park, Tanzania. *European*
438 *Journal of Wildlife Research*, **51**, 213–222.
- 439 Loibooki, M., Hofer, H., Campbell, K.L.I. & East M.L. (2002) Bushmeat hunting by
440 communities adjacent to the Serengeti National Park, Tanzania: the

441 importance of livestock ownership and alternative sources of protein and
442 income. *Environmental Conservation*, **29**, 391-398.

443 Maddock, L. (1979) The "Migration" and grazing succession. In *Serengeti I –*
444 *Dynamics of an ecosystem* (eds A.R.E. Sinclair & N. Norton-Griffiths), pp.
445 104-129. The University of Chicago Press, Chicago, USA.

446 Matsuishi, T., Muhoozi, L., Mkumbo, O., Budeba, Y., Njiru, M., Asila, A., Othina, A. &
447 Cowx, I.G. (2006) Are the exploitation pressures on the Nile perch fisheries
448 resources of Lake Victoria a cause for concern? *Fisheries Management and*
449 *Ecology* **13**, 52-71.

450 Ndibalema, V.G. & Songorwa, A. (2007) Illegal meat hunting in Serengeti: dynamics
451 in consumption and preferences. *African Journal of Ecology*, DOI:
452 10.1111/j.1365-2028.2007.00836.x

453 Njiru, M., Nzungi, P., Getabu, A., Wakwabi, E., Othina, A., Jembe, T. & Wekesa, S.
454 (2007) Are fisheries management, measures in Lake Victoria successful? The
455 case of Nile perch and Nile tilapia fishery. *African Journal of Ecology*, **45**, 315-
456 323.

457 Nyahongo, J.W., East, M.L., Mturi, F.A. & Hofer, H. (2006) Benefits and costs of
458 illegal grazing and hunting in the Serengeti ecosystem. *Environmental*
459 *Conservation*, **32**, 326 – 332.

460 Nyahongo, J.W., Holmern, T., Stokke, B.G., Kaltenborn, B. & Røskaft, E. (in prep.)
461 Bushmeat preference and species identification based on meat taste by
462 humans in the western Serengeti, Tanzania.

463 Rao, M. & McGowan, P.J.K. (2002) Wild meat use, food security, livelihoods and
464 conservation. *Conservation Biology*, **16**, 580-583.

465 R Development Core Team (2006) R language and environment for statistical
466 computing. R Foundation for Statistical computing, Vienna, Austria.

467 Rowcliffe, M.J., Milner-Gulland, J.E. & Cowlishaw, G. (2005) Do bushmeat
468 consumers have other fish to fry? *Trends in Ecology and Evolution*, **20**, 274-
469 276.

470 Rusch, G.M., Stokke, S., Røskoft, E., Mwakalebe, G., Wiik, H., Arnemo, J.M &
471 Lyamuya R. (2005) *Human-wildlife interactions in western Serengeti, Tanzania:
472 Effects of land management on migratory routes and mammal population
473 densities*. NINA Report 85.

474 Pelkey, N.W., Stoner, C.J. & Caro, T.M. (2000) Vegetation in Tanzania: assessing
475 long term trends and effects of protection using satellite imagery. *Biological
476 Conservation*, **94**, 297-309.

477 SPSS, (2001) SPSS for windows. Release 12.0.1. SPSS Inc., Chicago, IL.

478 UN (United Nations) (2005). World population prospects: The 2004 revision.
479 Population Division, United Nations, New York, USA, pp. 105. Available from:
480 <<http://esa.un.org/unpp/>> (accessed 15 August 2007).

481 URT United Republic of Tanzania, (2002) 2002 population and housing census.
482 <http://www.tanzania.org.tz/census/index.html> (accessed 30.03.2007).

483 Whittingham, M.J., Stephens, P.A., Bradbury, R.B. & Freckleton, R.P. (2006) Why do
484 we still use stepwise modelling in ecology and behaviour? *Journal of Animal
485 Ecology*, **75**, 1182-1189.

486 Wilkie, D.S. & Godoy, R.A. (2001) Income and price elasticities of bushmeat demand
487 in lowland Amerindian societies. *Conservation Biology*, **15**, 761-769.

488 Wilkie, D.S.; Malcolm, S., Kate, A., Ernestine, N.E., Paul, T. & Ricardo, G. (2005)
489 Role of prices and wealth in consumer demand fro bushmeat in Gabon, Central
490 Africa. *Conservation Biology*, **19** (1), 268-274.

491 **Biographical sketches**

492 Julius Nyahongo is a senior research scientist with the Tanzanian Wildlife Research
493 Institute. He is a wildlife ecologist specializing in human-wildlife interactions. He has
494 worked intensively with human-wildlife interactions in Tanzania.

495

496 Tomas Holmern is a researcher with the Department of Biology at the Norwegian
497 University of Science and Technology. He is interested in human-wildlife interactions
498 and has recently been working on bushmeat hunting and local law enforcement in the
499 Serengeti.

500

501 Bjørn Kaltenborn is a senior research scientist with the Norwegian Institute for Nature
502 Research. He is a geographer and social scientist specializing in human-environment
503 interactions. He has worked extensively with human-wildlife conflicts in the Nordic
504 countries as well as in East Africa and South Asia.

505

506 Eivin Røskaft is a behavioural ecologist interested in a wide range of birds and
507 mammals species in Europe, North America and Africa, and in human-wildlife
508 conflicts over the use of limited land.

509

510

511

512

513

514

515

516
517
518

Table 1: Household social characteristics and mean weekly meat and fish consumption in relation to household size, wealth and level of education, distance from the park boundaries and the lake (Pearson correlation, n = 10 months)

Villages	Household social characteristics				Pearson correlation: (Pvalue, n = 10 months)						
	Level of education achieved (%)		Occupation (%)		Household size		Household wealth		Goats/sheep		
	No education	Primary	Sec/college	Peasant	Employed	Meat	Fish	Meat	Fish	Meat	Fish
Robanda	26.7	66.7	6.7	86.7	13.3	ns	Ns	ns	ns	ns	ns
Nyamakendo	46.7	40.0	13.3	86.7	13.3	ns	Ns	0.030	(18)	(18)	(18)
Nyatwali	33.3	66.7	0	100.0	0	ns	0.043	(27)	(27)	(23)	(23)
Rwamkoma	6.7	90.0	3.3	96.7	3.3	ns	0.040	(29)	(29)	(25)	(25)
Kowak	6.7	83.3	10.0	83.3	16.3	ns	Ns	(29)	(29)	(25)	(25)
Pearson Chi-square test	0.001	< 0.0001	ns	Ns	ns	0.003	Ns	(28)	(28)	(24)	(24)

Note: ns is non-significant; numbers in brackets indicate sample size

519
520
521

Table 2: Mean household size and annual income and the village distance from the park and Lake Victoria (Mean (SD))

Category	Robanda	Nyatwali	Nyamakendo	Rwamkoma	Kowak	Statistics
Mean household size (number of individuals)	11.8 (6.8)	10.6 (5.1)	9.9 (4.9)	11.4 (5.7)	11.7 (8.0)	P = 0.864
Mean household annual income (US\$)	339.6 (159.6)	200.4 (356.4)	187.2 (325.2)	154.8 (283.2)	216.0 (344.4)	P = 0.157
Distance in km from the park boundaries	4	2	9	28	78	-
Distance in km from the Lake Victoria	95	0	78	37	17.8	-

524 **Table 3** – The set of 12 candidate models for explaining the number of meat and fish meals (10 months) in the villages adjacent to
 525 the Serengeti. The models are ranked by the Akaike Information Criterion corrected for small samples (AIC_c). (K = number of
 526 parameters; ΔAIC_c = difference in AIC_c between the best model and the actual model; ω_j = Akaike weights). The most
 527 parsimonious model is on the top of the list. Analyses are based on a total of 147 households in 5 villages.

Model	K	AIC _c	ΔAIC_c	ω_j
a) Meat consumption:				
Livestock + distance to protected area	5	957.2	0.000	0.449
Livestock + distance to protected area + hh size	6	959.5	1.799	0.182
Livestock + distance to protected area + income	6	959.8	2.172	0.152
Livestock + distance to protected area + education	7	961.0	3.337	0.085
Livestock + distance to protected area + hh size + income	7	961.7	4.004	0.061
Livestock + distance to protected area + hh size + education	8	962.9	5.232	0.032
Livestock	4	965.1	7.450	0.010
Livestock + distance to protected area + hh size + income + education	9	965.1	7.462	0.010
Distance to protected area	4	965.8	8.111	0.007
Livestock + hh size	5	967.0	9.301	0.004
Livestock + income	5	967.0	9.585	0.004
Livestock + education	6	968.7	11.050	0.002
b) Fish consumption:				
Distance to Lake Victoria + distance to protected area	5	991.5	0.000	0.393
Distance to Lake Victoria + distance to protected area + hh size	6	993.4	1.916	0.150
Distance to Lake Victoria + distance to protected area + livestock	6	993.5	2.045	0.141
Distance to Lake Victoria + distance to protected area + income	6	993.6	2.136	0.135
Distance to Lake Victoria + distance to protected area + education	7	995.0	3.534	0.067
Distance to Lake Victoria + distance to protected area + livestock + hh size	7	995.3	3.857	0.057
Distance to Lake Victoria + distance to protected area + hh size + livestock	7	995.7	4.238	0.047
Distance to Lake Victoria + distance to protected area + hh size + livestock + income + education	10	1001.0	9.669	0.004
Distance to Lake Victoria	4	1001.0	10.000	0.002
Distance to Lake Victoria + hh size	4	1003.0	11.870	0.001
Distance to Lake Victoria + livestock	5	1003.0	12.010	0.001
Distance to Lake Victoria + income	5	1004.0	12.120	0.001

529 **Table 4** – Estimate for the most parsimonious model of the number of meat and fish meals. For more details see Table 3.

Coefficients	Estimate	SE	T	P-value
a) Meat consumption:				
Intercept	27.919	2.029	13.759	<0.001
Livestock	0.052	0.016	3.206	0.002
Distance to protected	-0.218	0.052	-4.168	0.025
b) Fish consumption:				
Intercept	87.03	3.34	26.06	<0.001
Distance to protected	-0.931	0.205	-4.549	0.045
Distance to Lake	-0.742	0.155	-4.787	0.041

530

531

532

533

534

535

536

537

538 **Table 5** – Cumulative AIC_c weights for the candidate set of models with variables that influenced meat and fish meals. The sum of
 539 AIC_c weights was calculated across all models in the confidence set where the variable (n) was present.

Variable	Meat			Fish		
	n	Sum AIC _c	n	Sum AIC _c	n	Sum AIC _c weights
Intercept	12	1.0	12	1.0	12	1.0
Livestock	11	0.99	5	0.25		
Distance from protected area	8	0.98	8	0.99		
Distance from Lake Victoria	-	-	12	1.0		
HHsize	5	0.29	5	0.259		
Income	4	0.23	3	0.14		
Education	4	0.13	2	0.071		

540 **Figure legends:**

541 Figure 1. Map of the protected areas in the western Serengeti showing the villages
542 as grey pentagons (black are the 5 study villages). The 3 transect locations are
543 shown as grey stripes. Arrows give an illustration of some the different migratory
544 pathways for the wildebeest. Arrow size is roughly proportional to the abundance of
545 herds along the respective pathways. On their northward migration the wildebeest
546 herds use parts of the Western Corridor, as well as the partially protected and village
547 areas, depending upon the rainfall pattern.

548

549 Figure 2. Mean monthly migratory and resident herbivore densities and mean meat
550 and fish meals consumed per week in five villages from March to December 2006.

551

552

553

554


555

556

557


558

559



560
 561
 562
 563
 564
 565
 566

Fig. 1



567
568 Fig. 2

Paper V

1 **Bushmeat preference and species recognition based on meat taste by humans**
2 **in the western Serengeti, Tanzania**

3

4 Running headline: Bushmeat preference

5

6 Authors:

7 Julius W. NYAHONGO^{a,c*}, Tomas HOLMERN^a, Bård G. STOKKE^a, Bjørn P.
8 KALTENBORN^b AND Eivin RØSKAFT¹

9 ^aDepartment of Biology, Norwegian University of Science and Technology (NTNU),
10 Realfagbygget, N-7491, Trondheim, Norway

11 ^bNorwegian Institute for Nature Research (NINA), Fakkeltgården, 2624, Lillehammer,
12 Norway

13 ^cTanzania Wildlife Research Institute (TAWIRI), P. O. Box 661, Arusha, Tanzania

14

15 Word count: Abstract is 246; Main text (including abstract and references is 7884)

16

17

18 *Author for correspondence: email: nyahongo@bio.ntnu.no or
19 nyhwjulius@yahoo.co.uk (J. Nyahongo).

20 Tel. (Office): +47-73550348

21 Fax:+47-73596100

22 Mobile: +47-48230094 or +255-754886247

23 Emails of co-authors: Tomas.Holmern@bio.ntnu.no (T. Holmern),

24 Bard.Stokke@bio.ntnu.no (B.G. Stokke), Eivin.Roskaft@bio.ntnu.no (E. Røskaft),

25 bjorn.kaltenborn@nina.no (B.P. Kaltenborn).

26 **Abstract**

27 Wildlife meat has been an important source of meat protein to human kind since pre-
28 historical time. Even now, some African and Latin American societies survive largely
29 on meat obtained from wild animals. We used human taste ability to rank meat
30 preference and species recognition by providing each test-person with boiled pieces
31 of meat from different wild ungulates (topi, impala, zebra and wildebeest) and beef.
32 Nine-hundred test-persons of different age and sex from nine selected villages in the
33 western Serengeti were included in the experiment. Every test-person was provided
34 with a piece of meat from two of the above mentioned species, yielding 10 groups
35 with 90 persons in each. In order of preference, beef was most preferred followed by
36 topi, impala, zebra and wildebeest. In logistic regression analyses, we investigated
37 the possible influence of age, sex and distance from national park boundaries on
38 meat preference and species recognition. Generally, distance explained a significant
39 amount of variation in meat preference and species recognition in most two-species
40 comparisons. In addition, sex explained some of the variation in preference in the
41 impala-wildebeest, zebra-wildebeest and beef-zebra comparisons. Overall,
42 identification of species by meat taste was poor. The most identifiable meat was beef
43 and the least was impala. A combination of reduced beef prices and improved wildlife
44 management including discouragement of local people the bushmeat supply, could
45 encourage more domestic meat utilization and reduced hunting pressure on resident
46 wild herbivores in Serengeti.

47 Key words: Bushmeat, meat taste, preference ranks, Serengeti

48

49

50

51 **Introduction**

52 Before the onset of agriculture (about 10 000 years B.C., Diamond, 2005), the killing
53 of wild animals for food and survival purposes was one of the most important social
54 activities of the pre-historical *Homo sapiens* (Blain, 2005). Even presently, some
55 African and Latin American societies survive mainly on meat obtained from wild
56 animals (bushmeat) (e.g. Bennet, 2006; Rao and McGowan, 2002). Bushmeat
57 derived from wild animals by local communities has, in modern times, become easy
58 due to efficient hunting techniques such as the use of guns and wire snares (Blain,
59 2005). In Africa, the extensive utilization of bushmeat is well documented in some
60 western and central African countries (Barnett, 2000; Fa et al., 2003; Wilkie et al.,
61 2005). However, bushmeat is often obtained illegally inside protected areas (Hofer et
62 al., 1996; Loibooki et al., 2002; Holmern et al., 2002). Such illegal bushmeat
63 utilization has resulted in a serious conflict between conservation priorities and
64 priorities of local communities surrounding such protected areas (Campbell and
65 Hofer, 1995). One important question is whether current offtake will ensure future
66 sustainability of bushmeat species as demand increases due to a rapidly growing
67 human population. This situation is not only causing widespread depletion of wildlife
68 populations but also enhances encroachment on their habitats (Bodmer, 1994;
69 Alvard et al., 1997; Barnett 2000; Loibooki et al., 2002; Rao and McGowan, 2002; Fa
70 et al., 2003).

71

72 Generally, many species are utilized for bushmeat and species selection within
73 particular areas depends largely on location, habitat type and availability (Barnett,
74 2000). In most African countries, targeted species include insects, reptiles, birds and
75 mammals of various species including primates (FitzGibbon et al., 1996; Noss, 1998;

76 Stein, 2001; John et al., 2003). In Serengeti, northern Tanzania, the common large
77 herbivore species usually utilized for bushmeat include wildebeest (*Connochaetes*
78 *taurinus*), Cape buffalo (*Syncerus caffer*), impala (*Aepyceros melampus*), zebra
79 (*Equus burchelli*), eland (*Tragelaphus oryx*), Thomson gazelle (*Gazella thomsonii*),
80 Grant gazelle (*G. granti*) and giraffe (*Giraffa camelopardalis*). Other species include
81 topi (*Damaliscus korrigum*), kongoni (*Alcelaphus buselaphus*), warthog
82 (*Phacochoerus aethiopicus*), waterbuck (*Kobus ellipsiprymnus*), bush buck
83 (*Tragelaphus scriptus*) and ostrich (*Struthio camelus*) (Campbell and Hofer, 1995;
84 Hofer et al., 1996; Mduma et al., 1998; Holmern et al., 2004).

85

86 Generally, studies on bushmeat utilization have mostly been concerned with the
87 identification of utilized species (Barnett, 2000), dietary contribution of bushmeat to
88 the local people, the cost incurred and profit obtained from the sale of bushmeat.
89 Moreover, some studies have related bushmeat utilization directly to the loss of
90 biodiversity while others have related the human population growth and thereby
91 habitat destruction to the loss of bushmeat species. Yet, some studies have
92 suggested that the contribution of bushmeat may be an important factor in poverty
93 reduction in the rural areas (see, Campbell and Hofer, 1995; Hofer et al., 1996;
94 McKinney, 2001; Haule et al., 2002; Loibooki et al., 2002; Rao and McGowan, 2002;
95 Fa et al., 2003; Rowcliffe et al., 2003; Wilkie et al., 2005; Bennett et al., 2006;
96 Nyahongo et al., 2006).

97

98 Currently, two-thirds of African people (615 million) live in small-scale, low
99 productivity farms and are facing food insecurity that affect 194 million people, most
100 of them being children (Conway and Toenniessen, 2003). Factors that contribute to

101 the food insecurity in Africa include poor distribution of farm products, soil fertility,
102 crop losses due to diseases, pests, and international trade policies such as quota
103 systems, tariffs and subsidies as well as abiotic stress (Orr and Mwale, 2001; Pretty
104 et al., 2002; Rao and McGowan, 2002; Conway and Toenniessen, 2003). In addition,
105 the affordable technological innovations such as improved seeds and fertilizers and
106 capital (human capital, natural capital, produced capital, social capital and cultural
107 capital) are lacking due to abject poverty in Africa (Bebbington, 1999; Conway and
108 Toenniessen, 2003). For the local communities surrounding the western Serengeti,
109 like many other poor African communities, relying on bushmeat hunting is considered
110 important for food security and income generation (Loibooki et al., 2002; Holmern et
111 al., 2004; Kaltenborn et al., 2005).

112

113 Nevertheless, a study on species preference as a result of meat quality and species
114 identification based on meat taste has not been carried out though this might be
115 important in order to conserve the most preferred ungulate species in a sustainable
116 way. Moreover, establishment of game ranches in order to provide alternative meat
117 sources to domestic meat may require knowledge on preference, palatability and
118 acceptance based on perceived taste quality (flavour). In addition, studies that may
119 link illegal bushmeat hunting and local market demand for certain species of
120 ungulates based on perceived meat taste quality may also require prior knowledge
121 on meat quality, species preference and acceptability.

122

123 The aim of this study was to answer two basic questions: 1) Do local people have
124 specific species preference patterns based on meat taste? 2. Are local people
125 capable of identifying species of animals by meat taste? Related to the two main

126 questions, the following three hypotheses were tested: i) People from villages close
127 to the park boundary have long experience with bushmeat species and have eaten
128 more meat from different bushmeat species, hence they are more capable of
129 recognizing different species by meat taste than people from distant villages. ii).
130 Women have more experience preparing meat for food and also do most family
131 cooking, hence they have a certain preference pattern for different species and they
132 are able to identify the bushmeat species by meat taste more correctly than men. iii)
133 Adult people have more experience with bushmeat species and are therefore able to
134 rank the meats and can recognize the species by meat taste more correctly than
135 younger people.

136

137 **Methods**

138 **Study area**

139 The current study was carried out in villages surrounding the north-western part of
140 the Serengeti National Park (here after referred to as “national park”) (Fig. 1) in
141 northern Tanzania, between July and December 2006. The SNP (14,763 km²)
142 established in 1951, is a World Heritage Site (1981) and the largest national park in
143 Tanzania. Ikorongo Game Reserve (ca. 563 km²), Grumeti Game Reserve (ca. 416
144 km²) and the Ikoma Open Area (ca. 600 km²) act as buffer zones between the park
145 and the village areas to the north-west. The average annual temperature in the study
146 area is 21.7°C, with an average annual precipitation of 800 mm in the eastern part to
147 1050 mm in the western parts. The most common large resident herbivores in the
148 area are: giraffe, buffalo, topi, impala and the gazelles, *Gazella sp.* The western
149 Serengeti experiences the annual wildebeest and zebra migration in June–July that
150 moves through the partially protected and village areas on their way north (Sinclair,

151 1995; Thirgood et al., 2004). However, the populations of the resident herbivores are
152 relatively low in the buffer zones and in the open areas probably due to illegal
153 bushmeat hunting (Rusch et al., 2005).

154

155 **Local people and livelihoods**

156 In western Serengeti the human population density is relatively high (70 people/km²)
157 experiencing an annual growth rate of 2.5% between 1988 and 2002 (URT, 2002).
158 Village areas are made up of small units consisting of widely dispersed houses with
159 no clear cut border between the household areas within and between villages. The
160 economy of local communities is mainly based on subsistence crop farming and
161 livestock husbandry that consist of cattle, goats and sheep though a few farmers also
162 keep donkeys and pigs (Loibooki et al., 2002). The human population density within a
163 45 km belt from the park is relatively high partly due to immigration from other distant
164 villages with the intention of free access to natural resources (Campbell and Hofer,
165 1995).

166

167 Illegal bushmeat hunters from adjacent local communities essentially use wire snares
168 to capture wild herbivores for bushmeat (Loibooki et al., 2002). Such meat is
169 commonly sun dried before transported to the villages on foot from protected areas
170 (Hofer et al., 2000). The hunters use dried meat for home consumption, sale to
171 generate income, or bartering with other commodities (Hofer et al., 2000; Loibooki et
172 al., 2002; Kaltenborn et al., 2005). The majority of captured herbivores are large
173 migratory species such as wildebeest and zebra. An estimated 53,000 people are
174 involved in illegal hunting, including both hunters and porters that transport the meat
175 out of the protected areas (Loibooki et al., 2002).

176 **Data collection**

177 Nine villages were selected based on the distance from the national park boundary.
178 Robanda, Nyamakendo and Nattambisso (here after referred to as “immediate”)
179 villages were located within 10 kilometres from the park boundaries. Rwamkoma,
180 Busegwe and Butiama (“intermediate”) villages were located about 40 km from the
181 park, while Kowak, Chereche and Omuga (“distant”) villages were situated more than
182 80 km from the park. All distances are given as shortest air distance from villages to
183 the national park boundaries.

184

185 A visit to each village office prior to the meat taste experiment was done in order to
186 discuss the essence of the experiment with local leaders and elaborate how the meat
187 should be obtained and prepared. Moreover, we asked village leaders to help in
188 convincing people to attend to the meat taste experiment.

189

190 Meat from similar parts of the two different species was compared by giving test-
191 persons two pieces of boiled meat; one from each species. Meat used in the
192 experiments were obtained from two sources; beef was bought from local markets
193 while meat from wild animals were obtained by shooting eight mature males (two
194 animals from each species) from the following species: wildebeest, topi, impala and
195 zebra between July and December 2006 in Ikoma and Sibora Game Control Areas.
196 The hunting was carried out by professional hunters from Grumeti Reserve Fund and
197 game rangers from Grumeti - Ikorongo Game Reserves. The hunting operations
198 were done carefully; no untargeted animal was killed or injured. The shot animals
199 died instantly and were immediately skinned, and carcasses were thereafter stored in
200 a cold room (-15°C) for subsequent meat taste experiments.

201 Meat chopped from hindquarters of the carcasses were cut into small pieces and
202 boiled under constant temperature and pressure for 30 minutes. To control for the
203 effect of meat size on boiling time, we counted pieces of meat that were roughly
204 equally cut and in each pot we placed 120 small-cut pieces. Equal volume of water
205 (i.e. 500 ml) was added to boil the meat and equal weight of salt (one tea spoonful)
206 was added to season it. The temperature was maintained by adjusting the gas
207 cooker knob that controls the flame intensity while the pressure was maintained by
208 placing a lid tightly on the cooking pot. Experiments started after two sets of meat
209 from two different species were boiled. We recorded age and sex of the test-persons
210 and village location in relation to the distance from the national park preceding the
211 meat test. Before introducing the first piece of meat to test-persons, we provided
212 them with clean water to rinse their mouths in order to remove any food remnants
213 from previous meals. We repeated the same procedure after swallowing the first
214 piece of meat.

215

216 The test-persons, who voluntarily came to test meat, were asked to line-up in three
217 different lines based on age and gender. This arrangement was important to
218 encourage more women and children who were shy to mix with adult men. Only the
219 researcher had knowledge of which species that was tested at any time. After each
220 test-person declared that he or she had swallowed the second piece of meat (which
221 was done randomly), we asked him or her to tell us, based on the meat taste, which
222 of the two pieces tasted best. We recorded each individual's opinion and thereafter
223 we requested each test-person to identify the species whose pieces of meat were
224 tasted (choice of two species). Similar experiment and approach was repeated for all
225 different combinations of animals under study and in all nine randomly selected

226 villages (30 test-persons for each two-species comparisons in each of the three
227 village categories based on distance to the national park).

228

229 **Statistical analyses**

230 Differences between samples were tested by using non-parametric tests due to non-
231 normality in distribution. Within each two-paired comparison, forward-step-wise
232 binary logistic regression analyses were carried out to investigate the influence of
233 age (children versus adults), gender and distance from the park (immediate,
234 intermediate and distant villages) and their interactions (independent variables) on
235 meat preference and species recognition (dependent variables). For all tests $p < 0.05$
236 was considered significant. All analyses were performed using SPSS 14.0 statistical
237 package (SPSS, 2005).

238

239 **Results**

240 **General characteristics of test-persons**

241 Overall, 900 test-persons participated in the study, 300 from each of the categories
242 “immediate”, “intermediate” and “distant” villages. The socio-economic characteristics
243 of the test-persons are provided in Table 1.

244

245 **Species preference by meat taste**

246 *Overall preference*

247 Regardless of distance from the park, age and gender, the meat that scored the
248 highest overall preference rank in the two-species comparisons was beef, closely
249 followed by topi and impala. Zebra and wildebeest were least preferred taste trials
250 per species, Table 2). Overall, distance from the park reduced the preference of topi

251 (Pearson Chi-square: $p < 0.001$), impala and zebra. Distance, age or gender did not
252 affect the preference for the remaining species (Table 3).

253

254 Regardless of age and gender, topi scored the highest rank followed by beef in the
255 immediate villages while impala and wildebeest were less preferred. The least
256 preferred animal species in the immediate villages was zebra (Table 2). In the
257 immediate villages only gender had significant effect on the preference for zebra
258 (Pearson Chi-square: $p = 0.037$ (Table 3), whereas females preferred zebra than
259 men (mean rank for men = 63.3, $n = 83$; mean rank for women = 54.2, $n = 37$;
260 Wilcoxon Sign Rank: $W = 2007.0$, $p = 0.037$. Age and gender had no significant
261 effect on preference for the other species in the immediate village (Table 3).

262

263 In the intermediate villages, impala meat was highly favoured followed by beef. Topi
264 and zebra were less preferred. The least favoured species was wildebeest (Table 2).
265 Age and gender had no significant effect on meat preference in intermediate villages
266 (Table 3).

267

268 In distant villages, the most favoured meat was beef followed by impala. Topi ranked
269 the third while wildebeest and zebra scored below average (Table 2). Age and
270 gender had no significant effect on meat preference in distant villages (Table 3).

271

272 *Predictors explaining variation in preference in specific two-species comparisons*

273 *Beef-topi*

274 Overall, there was no statistical difference in test-persons preferring beef (46) over
275 topi (44) (Wilcoxon Sign Rank, $W = -1.42$, $n = 90$, $p = 0.833$). In the logistic

276 regression analysis, none of the independent variables could explain the variation in
277 preference.

278 *Beef-impala*

279 Overall, test persons preferred beef (59) over impala (31) (Wilcoxon Sign Rank, $W =$
280 -2.951 , $n = 90$, $p = 0.003$). In the logistic regression analysis, none of the
281 independent variables could explain the variation in preference.

282 *Beef-wildebeest*

283 Overall, test persons preferred beef (73) over wildebeest (17) (Wilcoxon Sign Rank,
284 $W = -5.90$, $n = 90$, $p < 0.001$). In the final regression model, the only variable that was
285 significant predictor of differences in meat preference was distance to the park
286 boundary (Table 4), and 81.1% of the cases were classified correctly by using this
287 model. Removing this variable resulted in a significantly poorer fit of the model
288 (change in $-2 \log$ likelihood = 8.82, $df = 2$, $p = 0.012$). Although test persons in all
289 three villages preferred beef over wildebeest, more inhabitants of immediate villages
290 preferred wildebeest (36.7%, $n = 30$) than those further away (both 10.0%, $n = 30$ in
291 both cases).

292 *Beef-zebra*

293 Overall, test-persons preferred beef (64) over zebra (26) (Wilcoxon Sign Rank, $W =$
294 Wilcoxon Sign Rank, $W = -7.48$, $n = 90$, $p < 0.001$). In the final regression model, the
295 interaction between distance and gender were significant predictors of differences in
296 meat preference (Table 4), and 78.9% of the cases were classified correctly by using
297 this model. Removing this variable resulted in a significantly poorer fit of the model
298 (change in $-2 \log$ likelihood = 16.51, $df = 2$, $p < 0.001$). Test-persons in the immediate
299 and distant villages preferred beef over zebra (80.0% and 86.7%, respectively, $n = 30$
300 in both cases), while inhabitants of intermediate villages generally preferred zebra

301 (53.3%, n = 30). Moreover, in the intermediate villages, females preferred zebra
302 (88.9%, n = 9) while men preferred beef (61.9%, n = 21). Both sexes in the
303 immediate and distant villages preferred beef (immediate: females 70% (n = 10),
304 males 85 (n = 20); distant: females 100% (n = 5), males 84% (n = 25)), but in general
305 (all distances), females preferred zebra more (45.8%, n = 24) than males (22.7%, n =
306 66).

307 *Topi-impala*

308 Regardless of the distance from the park, age or gender, test-persons preferred topi
309 (57) over impala (33) (Wilcoxon Sign Rank, $W = -2.53$, $n = 90$, $p = 0.011$). In the final
310 regression model, distance to park boundary was a significant predictor of
311 differences in meat preference (Table 4), and 76.7% of the cases were classified
312 correctly by using this model. Removing this variable resulted in a significantly poorer
313 fit of the model (change in -2 log likelihood = 23.33, $df = 2$, $p < 0.001$). Topi was
314 preferred in immediate (86.7%, n = 30) and intermediate villages (73.3%, n = 30)
315 while impala was preferred in the distant villages (70.0%, n = 30).

316 *Topi-wildebeest*

317 Regardless of the distance from the park, age or gender, test-persons preferred topi
318 (61) over wildebeest (29) ($W =$ Wilcoxon Sign Rank, $W = -5.66$, $n = 90$, $p < 0.001$). In
319 the final regression model, the only variable that was significant predictor of meat
320 preference was distance to the park boundary (Table 4), and 81.1% of the cases
321 were classified correctly by using this model. Removing this variable resulted in a
322 significantly poorer fit of the model (change in -2 log likelihood = 29.91, $df = 2$, $p <$
323 0.001). Test-persons in the immediate and intermediate villages preferred topi
324 (immediate: 83.3%, n = 30; intermediate: 90.0%, n = 30), while those in distant
325 villages preferred wildebeest (70.0%, n = 30).

326 *Topi-zebra*

327 Overall, test-persons who participated in the meat taste experiment preferred topi (62)
328 over zebra (28) (Wilcoxon Sign Rank, $W = \text{Wilcoxon Sign Rank}, W = -3.24, n = 90, p$
329 < 0.001). Distance from the park boundary was the only independent variable in the
330 final regression model that significantly predicted the meat preference (Table 4),
331 whereas 80.0% of the cases were classified correctly by using this model. Removing
332 this variable (“distance”) resulted in a significantly poorer fit of the model (change in -
333 $2 \log \text{likelihood} = 28.74, df = 2, p < 0.001$). The test-persons in the immediate (93.3%,
334 $n = 30$) and distant (80.0%, $n = 30$) villages preferred topi, while those in intermediate
335 villages preferred zebra (66.7%, $n = 30$).

336 *Impala-wildebeest*

337 Overall, test-persons, regardless of the distance from the park boundary, age and
338 gender, preferred impala (66) over wildebeest (24), ($W = \text{Wilcoxon Sign Rank}, W = -$
339 $4.43, n = 90, p < 0.001$). In the final regression model, the distance from the park and
340 the interaction between distance and gender of test-persons were significant
341 predictor of the meat preference, whereas 78.9% of the cases were classified
342 correctly by using this model (Table 4). Removing these variables (“distance and
343 gender-distance interaction”) resulted in a significantly poorer fit of the model
344 (change in $-2 \log \text{likelihood} = 17.41, df = 4, p = 0.002$). Although impala was
345 preferred in all villages, the analysis suggest that test-persons in villages that were
346 classified as immediate to the park boundary preferred impala less (56.7%, $n = 30$)
347 than those in intermediate (83.3%, $n = 30$) or distant villages (80.0%, $n = 30$).
348 Moreover, in the immediate villages, males preferred wildebeest (63.2%, $n = 19$)
349 while females preferred impala (90.9, $n = 11$).

350

351 *Impala-zebra*

352 Regardless of the distance from the park boundary, test-person preferred impala (72)
353 over zebra (18) (Wilcoxon Sign Rank, $W = -5.62$, $n=90$, $p < 0.001$). Distance from the
354 park boundary was the only independent variable in the final regression model that
355 significantly predicted the meat preference (Table 4), whereas 80.0% of the cases
356 were classified correctly by using this model. Removing this variable (“distance”)
357 resulted in a significantly poorer fit of the model (change in $-2 \log \text{likelihood} = 7.11$, df
358 $= 2$, $p = 0.028$). The test-persons in the immediate (80%, $n = 30$) and intermediate
359 (90.0%, $n = 40$) villages preferred impala, while those in distant villages had the least
360 preference for impala (60.0%, $n = 20$).

361 *Zebra-wildebeest*

362 Overall, there was no statistical difference in test-persons preferring zebra (48) or
363 wildebeest (42), (Wilcoxon Sign Rank, $W = -6.32$, $n= 90$, $p = 0.527$). In the final
364 regression model, the distance from the park and gender of test-persons were
365 significant predictors of the meat preference, whereas 75.6% of the cases were
366 classified correctly by using this model (Table 4). Removing these variables resulted
367 in a significantly poorer fit of the model (change in $-2 \log \text{likelihood} = 37.02$, $df = 3$, p
368 < 0.001). Test-persons in the intermediate villages mostly preferred zebra (90.0%, n
369 $= 30$), while those in the immediate (76.7%, $n = 30$) and distant (53.3%, $n = 30$)
370 villages preferred wildebeest. Furthermore, overall, females preferred wildebeest
371 (60.6%, $n = 33$), while males preferred zebra (61.4%, $n = 57$).

372

373

374

375

376 ***Species recognition by meat taste***

377 *Overall recognition*

378 Generally, a correct recognition based on meat taste was poor among test-persons
379 of different age and gender in all villages (25.9%, n = 1800 trials). For the pooled
380 data, the most recognizable meat was beef while the least recognized species was
381 impala (Table 2). Overall, distance from the park had reduces the recognition of all
382 species while recognition of topi, impala and zebra increases with age (Table 3).
383 Gender did not have effect on recognition of any species.

384

385 In the immediate village, of all test-persons, 55.8% (n = 120) were able to recognize
386 beef (Table 2). The recognition of the remaining species scored less than 44% (n =
387 120 for each species). Neither age nor gender did influence the recognition of any
388 species by meat taste (Table 3).

389

390 In the intermediate villages, 47.5% (n = 120) of all test-persons were able to
391 recognize beef while none of the wild ungulate scored more than 30% (n = 120 for
392 each species, Table 2). Neither age nor gender did influence the recognition of any
393 species by meat taste (Table 3).

394

395 In the distant village, general recognition of species was relatively low (< 17%) and
396 the meat that was highly recognized (16.7%, n = 120) was beef. The remaining wild
397 ungulates scored 10% or less (Table 2). Age and gender had an effect on recognition
398 of all wild ungulates in the distant village where adult male were able to recognize
399 species than children and female. Beef was recognized similarly by all test-person of
400 different age and sex (Table 3).

401 *Predictors explaining variation in species recognition in specific two-species*
402 *comparisons*

403 *Beef-topi*

404 Overall, 67.8%, (n = 90) of the test persons were able to correctly recognize what
405 species of meat was tasted. In the binary regression analysis, distance from the park
406 was the only variable that could explain the variation in recognition (Table 5), and
407 70.0% of the cases were classified correctly by using this model. Removing this
408 variable resulted in a significantly poorer fit of the model (change in -2 log likelihood =
409 11.52, df = 2, p = 0.003). Test-persons from the immediate villages were better able
410 to identify the two species (53.3%, n = 30) than test-persons from the intermediate
411 (30.0%, n = 30) and distant (13.3%, n = 30) villages.

412 *Beef-impala*

413 Overall, 24.4% (n = 90) of the test-persons were able to correctly recognize what
414 species of meat were tasted. In the binary regression analysis, distance from the
415 park was the only variable that could explain the variation in recognition (Table 5),
416 and 70.0% of the cases were classified correctly by using this model. Removing this
417 variable resulted in a significantly poorer fit of the model (change in -2 log likelihood =
418 11.63, df = 2, p < 0.003). Test-persons in the intermediate villages were better in
419 species recognition (63.3%, n = 30) than test-persons from the immediate (23.3%, n
420 =30) or distant (30.0%, n = 30) villages.

421 *Beef-wildebeest*

422 Overall, 24.4% (n = 90) of the test-persons were able to correctly recognize what
423 species of meat were tested. In the final regression model, the only variable that was
424 a significant predictor of species recognition was distance to the park boundary
425 (Table 5) and 77.8% of the cases were classified correctly by using this model.

426 Removing this variable resulted in a significantly poorer fit of the model (change in -2
427 log likelihood = 28.61, df = 2, $p < 0.001$). Test-persons in immediate villages were
428 better in recognizing species than in the intermediate villages (53.3% versus 20.0%,
429 $n = 30$ in both cases), and test-persons in distant villages did not correctly recognize
430 a single piece of meat (0%, $n = 30$).

431 *Beef-zebra*

432 Regardless of the distance from the park, age or gender, only 32.3% ($n = 90$) of the
433 test-persons were able to recognize the correct species by meat taste. In the final
434 regression model, the only variable that was significant predictor of species
435 recognition was distance from the park boundary (Table 5), and 72.2% of the cases
436 were classified correctly by using this model. Removing this variable resulted in a
437 significantly poorer fit of the model (change in -2 log likelihood = 12.54, df = 2, $p =$
438 0.002). Test-persons from the immediate villages were better able to recognize
439 species (56.7%, $n = 30$) by meat taste than the test persons from intermediate
440 (23.3%, $n = 30$) or distant villages (16.7%, $n = 30$).

441 *Topi-impala*

442 Regardless of the distance from the park, age and gender, 21.1% ($n = 90$) of test-
443 persons were able to recognize the correct species by meat taste. None of the
444 independent variables entered into the binary logistic regression analysis could
445 explain the observed variation in species recognition between these two species.

446 *Topi-wildebeest*

447 Regardless of the distance from the park, age and gender, 16.7% ($n = 90$) of the test-
448 persons were able to recognize the correct species by meat taste. None of the
449 independent variables entered into the binary regression analysis could explain the
450 observed variation in species recognition between these two species.

451 *Topi-zebra*

452 Regardless of the distance from the park, age and gender, 21.1% (n = 90) of the test-
453 persons were able to recognize the correct species by meat taste. Distance from the
454 national park boundary was the only independent variable in the final regression
455 model that significantly predicted the success or failure of a test-person to recognize
456 species by meat taste (Table 5), whereas 78.9% of the cases were classified
457 correctly by using this model. Removing this variable resulted in a significantly poorer
458 fit of the model (change in -2 log likelihood = 15.9, df = 2, p < 0.001). Despite the low
459 recognition rate, the test-persons from the immediate villages were able to recognize
460 species by meat taste better (43.3%, n = 30) than test-persons from intermediate
461 (16.7%, n = 30) and distant (3.3% n = 30) villages.

462 *Impala-wildebeest*

463 Regardless of the distance from the park, age and gender, 20.0% (n = 90) of the test-
464 persons were able to recognize the correct species by meat taste. In the final
465 regression model, the distance from the park was the only independent variable that
466 could explain the variation in ability of a test-person to recognize the species of
467 animals whose meat was included in the experiment (Table 5), and 80.0% of the
468 cases were classified correctly by using this model. Removing this variable resulted
469 in a significantly poorer fit of the model (change in -2 log likelihood = 10.52, df = 2, p
470 = 0.005). The analysis suggests that the ability to correctly recognize species decline
471 with distance from the park boundary (immediate: 33.3%; intermediate: 23.3%;
472 distant: 3.3%, n = 30 at each distance).

473 *Impala-zebra*

474 Regardless of the distance from the park, age and gender, 11.1% (n = 90) of the test-
475 persons were able to recognize the correct species by meat taste. In the binary

476 regression analysis, distance from the park was the only variable that could explain
477 the variation in recognition (Table 5), and 85.6% of cases were classified correctly by
478 using this model. Removing this variable resulted in a significantly poorer fit of the
479 model (change in $-2 \log$ likelihood = 8.44, $df=2$, $p < 0.015$). The analysis suggests
480 that the ability to correctly recognize species declined with distance to the park
481 boundary (immediate: 26.7%, $n = 30$; intermediate: 5.0%, $n = 40$; distant: 0.0%, $n =$
482 20).

483 *Zebra-wildebeest*

484 Regardless of the distance from the park, age and gender, 35.6% ($n = 90$) of the test-
485 persons were able to recognize the correct species by meat taste. In the binary
486 regression analysis, distance from the park was the only variable that could explain
487 the variation in recognition (Table 5), and 71.1% of the cases were classified
488 correctly by using this model. Removing this variable resulted in as significantly
489 poorer fit of the model (change in $-2 \log$ likelihood = 11.62, $df= 2$, $p = 0.003$.
490 Inhabitants from the immediate villages were better able to recognize the two species
491 (60.0%, $n = 30$) than inhabitants from intermediate (23.3%, $n = 30$) or distant villages
492 (23.3%, $n = 30$).

493

494 **Discussion**

495 Our overall results show that test-persons favoured beef, followed by topi and impala.
496 The preference patterns and the ranking position of beef, topi and impala alternated
497 along the gradient of distance from the park suggesting high preference and
498 acceptability of the three species by test-persons from different villages along the
499 gradient of distance from the park boundary. Distance from the park boundary and
500 gender-distance interactions had influence on meat preferences and subsequent

501 species recognition by meat taste of different combinations of beef and four wild
502 ungulates meat (Table 4). In contrast, the results indicated that most test-persons
503 were not able to identify the species based on the meat test. Generally, the most
504 correctly identified meat was beef while the least identified species was impala.

505

506 The test-persons included in this study were of different age and gender all having
507 different experience with wild animals which in a way might have influenced the
508 preference and ultimate identification of animal species based on taste, aroma and
509 texture. Studies have shown that the meat preference may be influenced by fatty
510 aroma, texture and taste. The meat with considerable fat marbling and soft texture is
511 considered best (Matsuishi et al., 2001). Other studies have reported that people
512 differ in their eating process that vary with individual breathing and chewing patterns,
513 composition and amount of saliva, and volume of their oral cavities which may affect
514 the taste perception (van Ruth and Roozen, 2000; Pionnier et al., 2004; Geary et al.,
515 2004). In addition, a study suggests that the moment and completeness of the
516 velopharyngeal closure might vary between individuals, which affect the amount of
517 volatiles transferred to the nasal cavity affecting sensation sensitivity towards
518 different volatile aroma components among people (Buettner and Schieberle, 2000).
519 In addition, in some villages that were close to the park boundary, alleged illegal
520 bushmeat hunters did not want to participate in this experiment because they
521 consider this experiment as a trick by the government to identify the illegal bushmeat
522 hunters. Hence, we missed their valuable experience.

523

524 Furthermore, the results we discuss might have been influenced by age and gender
525 because the data set included more adults than children and more men than women.

526

527 The most preferred meat based on the meat taste experiment was beef. The test-
528 persons might have favoured beef due to relatively high intramuscular fat contents in
529 the beef carcass. Some studies reported that a zebu bull has up to 40% fat content in
530 the dressed carcass compared with 2.5% in wild ungulates (FAO, 1992). Meat with
531 relatively high fat content produces suitable aroma; and when this is associated with
532 soft meat texture (tenderness) which is a characteristic of beef; the meat becomes
533 more palatable and is highly acceptable by consumers (Matsuishi et al., 2004).
534 These two combinations; texture and fatty aroma might be the reasons for most test-
535 persons to favour beef. Moreover, beef is locally available and can be obtained
536 throughout the year hence people may be used to the meat and were at least able to
537 distinguish its taste from the wild meats. Furthermore, the different types of wild
538 fodders the wild ungulates consume and crops residues that the livestock feed on
539 may account for the difference observed. For example, in the study area, the
540 harvesting period lasts from July to September each year; the herdsmen take the
541 livestock to feed on straws of the harvested maize, finger millet and/or sorghum
542 which are relatively of high quality compared to dry and over grazed feeds in the
543 protected areas (personal observation, 2006). Feeding on wide variety of fodders of
544 relatively low qualities has been reported to affect the meat flavour (Duckett and
545 Kuber, 2001).

546

547 In addition, constant vigilance and flights the wild herbivores evolved in response to
548 predation by both wild carnivores and humans (Krebs and Davies, 1987; Caro 2005),
549 may affect meat quality due to high muscle activities which reduces the intramuscular
550 fat contents and finally affecting the meat aroma and texture.

551

552 In the immediate villages, topi was preferred to all other species. Because topi is a
553 resident herbivore, the local communities close to the park boundary might have
554 illegally utilized topi meat for many years. In fact, the Serengeti Regional
555 Conservation Programme (SRCP) cropping operation did not include topi initially as
556 part of the species cropped to provide legal game meat to the villages, but was later
557 included because of substantial pressure from the villagers that said they preferred
558 this species to that of wildebeest and zebra (Holmern et al., 2002). Moreover, a
559 recent study in Serengeti suggested that the population of topi is declining compared
560 to other resident herbivores, which raise the concern that illegal bushmeat hunters
561 might have been targeting the species for perceived quality meat (Rusch et al., 2005).
562 However, impala that is also a resident herbivore scored the highest preference rank
563 in the intermediate villages probably due to the fact that it was a common illegally
564 obtained bushmeat in the area or was confused with topi and/or beef. A general
565 index of how test-persons might have confused topi and impala in the meat taste
566 experiments can be gained from a comparison of preference and subsequent
567 recognition of the two species in the pooled data, in immediate and intermediate
568 villages (Table 2). This finding supports the recent observation on the preference and
569 acceptability of different species animals in the similar area of Serengeti (Kaltenborn
570 et al., 2006).

571

572 In contrast, the zebra and wildebeest meats were not highly appreciated (scored less
573 than 34% in all trials for pooled data). Despite of their large populations in the
574 Serengeti, the two species are migratory herbivores that are only in the village
575 proximities for a period lasting for only three months each year. Although it is known

576 that local communities utilize these species highly during the period when the
577 animals are in the village proximities (Sinclair, 1995, Thirgood et al. 2004), the
578 harvesting period may not be long enough to warrant a fair comparison with other
579 resident herbivores and livestock. In addition, due to vast movement of wildebeest
580 and zebra within the ecosystem of varying habitats, landscapes and the seasonal
581 variation may also subject these migratory ungulates in large resource variations,
582 predation pressure and constant flight that may affect the meat quality (i.e. marbling
583 fat and texture).

584

585 Generally, age of test-persons did not influence the meat preference for all species
586 we studied, suggesting that the five different meats from five different species were
587 distinguishable and ranked specifically and that ranking patterns were independent of
588 age but was due to meat texture and fatty aroma. In contrast, the distance from the
589 park boundary reduced the topi preference from all topi meat taste combinations.
590 This may be due to the fact that test-persons from the immediate villages were used
591 to topi meat and had a relatively strong preference compared to test-persons from
592 distant villages who might have never tasted topi meat. To the distant test-persons,
593 the topi meat might have revealed a strange or unusual aroma and texture; hence
594 they preferred beef that they are used to. However, it is not clear as to why the
595 distance from the park favoured the preference of impala in topi-impala combination
596 and impala-wildebeest combination, zebra in topi-zebra combination and in beef-
597 zebra combination. The observed association might have occurred just by chance or
598 impala was confused with topi and unique fatty aroma of zebra attracted test-persons
599 from distant villages who had never tasted zebra meat.

600

601 Poor identification of species based on meat taste for all species suggest that most
602 test-persons had either low or no experience with different types of wild meats. This
603 can be justified by relatively high beef identification (40%, n = 360) because at least
604 every test-person had interacted with beef quite regularly. However, studies show
605 that taste sensitivity decreases with age and health status. For instance, old people
606 and those who are either taking medication and/or consume excessive alcohol and/or
607 smoke have reduced sensory stimuli (Fukunga et al., 2005). Most of people who
608 participated in this experiment were adults whose health status and social
609 characteristics were not established prior to this experiment.

610

611 Generally distance from the park and gender-distance interaction was significant
612 variables in logistic regression model that tested the influence of distance, age and
613 gender on meat preference and subsequent recognition of species by meat taste
614 among the test-persons (Table 4 and table 5). The preference of a combination of
615 beef and zebra and that of impala and wildebeest was possible to predict in the
616 intermediate villages using female as predictor (Table 4). Women are the cooks of
617 most families in Africa; hence they may prefer certain species based on their
618 experience and the taste of the meat they cook. However, women in the immediate
619 villages, preferred impala more compared to men from the same area that preferred
620 wildebeest more. Likewise, the combination of beef and the four wild ungulates was
621 possible to predict along the gradient of distance from the park while only
622 combinations of topi-zebra and that of impala-wildebeest, impala-zebra and
623 wildebeest-zebra was possible to predict along the gradient of distance from the park
624 (Table 4). Distance from the park explains how test-persons had experience with

625 different species of animals whereby the test-person from distant villages had less
626 experience with wild ungulates but had long experience with beef.

627

628 **Concluding remarks**

629 The results obtained from this study in which meat tastes were conducted to rank the
630 preference based on the meat quality and species recognition revealed that meat
631 taste by humans may be useful to rank meat preferences of different ungulates
632 although the test proved as an unsuitable approach for species recognition. The
633 distance from the park may affect the preference rank of different animal species. In
634 contrast, age of test-persons is not good parameters for the meat preference ranking
635 and subsequent species recognition. A substantial number of taste-persons preferred
636 beef and were able to identify the beef by meat taste approach. This suggests that if
637 the price of beef is reduced and wildlife management somehow manages to limit
638 bushmeat supply (preferably by cooperating actively with communities) many people
639 may choose to eat more beef rather than wild ungulates. This will inevitably reduce
640 the hunting pressure on resident herbivores. The preference of beef also highlights
641 that outreach activities could greatly increase their impact by focusing their attention
642 on improving services for livestock, such as cattle dips, water points and veterinary
643 services.

644

645 Future studies on the effect of bushmeat processes before transportation to the
646 market place is recommended. This is important because the bushmeat consumers
647 may be used to sun-dried meat that may influence the fatty aroma and the texture in
648 different levels among the different species. Finally, the findings from this study
649 suggest the need for special conservation attention to resident herbivores population

650 close to village proximities. Otherwise the long term harvest and uncontrolled illegal
651 bushmeat hunting based on current meat preferences and habitat location may
652 seriously deplete the resident herbivore species from their key habitats.

653

654 **Acknowledgements**

655 The current study was funded by NORAD through the Ministry of Natural Resources
656 and Tourism, Tanzania and is a part of the TAWIRI, NINA/NTNU project called
657 Biodiversity and Human Wildlife Interface in Western Serengeti. We acknowledge the
658 financial support from the Norwegian Peace corps and the Quota Programme
659 scheme (NORAD). We also thank Tanzania National Parks (TANAPA) for granting
660 permissions to foreign research team to work in Serengeti National Park. Finally, we
661 thank village leaders and our field assistants who in one way or the other contributed
662 to this important task.

663

664

665

666

667

668

669

670

671

672

673

674

675 **References:**

- 676 Alvard, M., 1994. Conservation by native people: prey choice in a depleted habitat.
677 Human Nature, 5, 127-154.
- 678 Barnett, R., 2000. Food for Thought. The utilization of wild meat in Eastern and
679 Southern Africa. TRAFFIC East/Southern Africa. Nairobi, Kenya.
- 680 Bebbington, A., 1999. Capitals and capabilities: a framework for analyzing peasant
681 viability, rural livelihoods and poverty. World Development, 27, 2021-2044.
- 682 Bennett, E.L., Blencowe, E., Brandon, K., Brown, D., Burn, R.W., Cowlshaw, G.,
683 Davies, G., Dublin, H., Fa, J.E., Milner-Gulland, E.J., Robinson, J.G., Rowcliffe,
684 J.M., Underwood, F.M., Wilkie D.S., 2006. Hunting for consensus: Reconciling
685 bushmeat harvest, Conservation and development policy in Western Africa.
686 Conservation Biology, 1-4
- 687 Blain, R., 2005. The evolution of man – the history of hunting.
688 <http://ezinearticles.com/?The-Evolution-of-man> (accessed on 10.05.2007).
- 689 Bodmer, R.E., 1994. Managing wildlife with local communities in Peruvian Amazon:
690 the case of Reserva Comunal Tamshiyacu-Tahuayo. In Western, D., Wright,
691 R.M. (Eds), Natural Connections: Perspectives in Community Based
692 Conservation. Island Press, Washington, DC, pp 113-134.
- 693 Buettner, A., Schieberle, P., 2000. Influence of mastification on the concentrations of
694 aroma volatiles – some aspects of flavour release and flavour perception.
695 Food Chemistry, 71, 347-354.
- 696 Campbell, K.L.I, Hofer, H., 1995. People and Wildlife: Spatial Dynamics and Zones of
697 Interaction. In: Sinclair, A.R.E, Arcese, P. (Eds), Serengeti II–Dynamics,
698 Management and Conservation of an Ecosystem. The University of Chicago
699 Press, Chicago, USA, pp. 534-570.

700 Caro, T., 2005. Anti-predator defences in birds and mammals. University of Chicago
701 Press, Chicago.

702 Conway, G., Toenniessen, G., 2003. Science for African food security. *Science*, 299,
703 1187-1188.

704 Diamond, J., 2005. *Guns, germs and steel: the fates of human societies*. W.W.
705 Norton and Co. Inc. New York, USA.

706 Duckett, S.K., Kuber, P.S., 2001. Genetic and nutritional effects on lamb flavour.
707 *Journal of Animal Science*, 79, 249-259.

708 Fa, J.E., Currie, D., Meeuwig, J., 2003. Bushmeat and food security in the Congo
709 Basin: linkage between wildlife and people's future. *Environmental*
710 *Conservation*, 30, 71-78.

711 FAO, 1992. Meat and meat products in human nutrition in developing countries.
712 <http://www.fao.org/docrep/T0562E/T0562E01.htm> (Accessed on 24.04.2007).

713 FitzGibbon, C.D., Lazarus, J., 1995. Anti-predator Behaviour of Serengeti ungulates:
714 Individual Differences and Population Consequences. In: Sinclair, A.R.E.,
715 Arcese, P. (Eds), *Serengeti II—Dynamics, Management and Conservation of*
716 *an Ecosystem*. The University of Chicago Press, Chicago, USA, pp. 258-274.

717 FitzGibbon, C.D., Mogaka, H., Fanshawe, J.H., 1996. Subsistence hunting and
718 mammal conservation in Kenyan coastal forest: resolving a conflict. *The*
719 *exploitation of mammal population*. Chapman & Hall London. U.K.

720 Fukunga, A., Uematsu, H., Sugimoto, K., 2005. Influences of aging on taste
721 perception and oral somatic sensation. *Journal of Gerontology*, 60, 109-113.

722 Geary, M.D., van Ruth, S.M., Delahunty, C.M., Lavin, E.H., Acree, T.E., 2004. Effects
723 of oral physiological characteristics on the release of aroma compounds from
724 oil-in-water emulsions using two model mouth systems. In: Deibler, K.D,

725 Delwiche, J.(Eds.), Handbook of flavour characterization. Marcel Dekker, New
726 York, USA. pp. 345-363

727 Haule, K.S., Johnsen, F.H., Maganga, S.L.S., 2002. Striving for sustainable wildlife
728 management: the case of Kilombero Game Controlled Area, Tanzania.
729 Environmental Management, 66, 31-42.

730 Hofer, H., Campbell, K.L.I., East, L.M., Huish, S.A., 1996. The impact of game meat
731 hunting on target and non-target species in the Serengeti. In Taylor, V.J.,
732 Dunstone, N (Eds.), The exploitation of mammal population. Chapman and
733 Hall, London, U.K., pp. 117-146.

734 Hofer, H., Campbell, K.L.I., East, M.L. and Huish, S.A., 2000. Modelling the spatial
735 distribution of the economic costs and benefits of illegal game meat hunting in
736 the Serengeti. Natural Resource Modeling, 13, 151 – 177.

737 Holmern, T., Røskaft, E., Mbaruka, J., Mkama, S.Y. and Muya., J. 2002.
738 Uneconomical game cropping in a community-based conservation project
739 outside the Serengeti National Park, Tanzania. Oryx, 36 (4), 364-372.

740 Holmern, T., Johannesen, A.B., Mbaruka, J., Mkama, S., Muya, J. and Røskaft, E.,
741 2004. Human-Wildlife Conflicts and Hunting in the Western Serengeti,
742 Tanzania. NINA Project Report, No. 26.

743 John, E.F.A., Currie, D. Meeuwig, J., 2003. Bush meat and food security in the
744 Congo Basin: linkages between wildlife and people's future. Environmental
745 Conservation. 30, 71-78.

746 Kaltenborn, B.P., Bjerke, T., Nyahongo, J.W., Williams, D.R., 2006. Animal
747 preferences and acceptability of wildlife management actions around
748 Serengeti National Park, Tanzania. Biodiversity and Conservation, 15, 4633-
749 4649.

750 Kaltenborn, B.P., Nyahongo, J.W., Tingstad K.M. 2005., The nature of hunting
751 around the Western Corridor of Serengeti National Park, Tanzania. *European*
752 *Journal of Wildlife Research*, 51, 213–222.

753 Krebs, J.R., Davies, N.B., 1987. *An introduction to Behavioural Ecology*, 2nd Ed.
754 Blackwell Scientific Publications. Oxford Press, London.

755 Loibooki, M., Hofer, H., Campbell, K.L.I., East M.L., 2002. Bushmeat hunting by
756 communities adjacent to the Serengeti National Park, Tanzania: the
757 importance of livestock ownership and alternative sources of protein and
758 income. *Environmental Conservation*, 29, 391-398.

759 Matsuishi, M., Igeta, M., Takeda, S., Okitani, A., 2004. Sensory factors contributing to
760 the identification of the animal species of meat. *Journal of Food Science*, 69,
761 218-220.

762 McKinney, M.L., 2001. Role of human population size in raising bird and mammal
763 threat among nations. *Animal Conservation*, 4, 45-57.

764 Mduma, S.A.R., Hilborn, R., Sinclair, A.R.E., 1998. Limits to exploitation of Serengeti
765 wildebeest and implications for its management. In: Newbery, D.M.,
766 Underwood, H.B., Prins, H.H.T.(Eds.), *Dynamics of Tropical Communities*.
767 Blackwell Science, Oxford, UK., pp. 243-265.

768 Noss, A.J., 1998. The impact of cable snares hunting on wildlife population in the
769 forest of the Central Africa Republic. *Conservation Biology*, 12, 390-397.

770 Nyahongo, J.W., East, M.L., Mturi, F.A., Hofer, H., 2006. Benefits and costs of illegal
771 grazing and hunting in the Serengeti ecosystem. *Environmental Conservation*,
772 32, 326–332.

773 Orr, A. and Mwale, B., 2001. Adapting to adjustment: smallholder livelihood
774 strategies in Southern Malawi. *World Development*, 29, 1325-1343.

775 Pionnier, E., Chabanet, C., Mioche, L., Le Quéré, J.-L., Salles, C., 2004. In vivo
776 aroma release during eating of a model cheese: Relationships with oral
777 parameters. *Journal of Agriculture and Food Chemistry*, 52, 557-564.

778 Pretty, J.N., Morison, J.I.L., Hine, R.E., 2003. Reducing food poverty by increasing
779 agricultural sustainability in developing countries. *Agriculture, Ecosystems and*
780 *Environment*, 95, 217-234.

781 Rao, M. and McGowan, P.J.K., 2002. Wild meat use, food security, livelihoods and
782 conservation. *Conservation Biology*, 16, 580-583.

783 Rowcliffe, M.J., Miller-Gulland, E.J., Cowlshaw, G., 2005. Do bushmeat consumers
784 have other fish to fry? *Trends in Ecology and Evolution*, 20, 274-276.

785 Rusch, G.M., Stokke, S., Røskoft, E., Mwakalebe, G., Wiik, H., Arnemo, J.M,
786 Lyamuya R., 2005. Human-wildlife interactions in western Serengeti, Tanzania:
787 Effects of land management on migratory routes and mammal population
788 densities. NINA Report No. 85.

789 Stein, J., 2001. Species affected by the bushmeat trade in Africa.
790 <http://www.bushmeat.org/html/SpeciesAffected.htm>

791 Sinclair, A.R.E., 1995. Serengeti past and present, In: Sinclair, A.R.E, Arcese, P.
792 (Eds.), *Serengeti II—Dynamics, Management and Conservation of an*
793 *Ecosystem*. The University of Chicago Press, Chicago, USA, pp. 3-33.

794 Thirgood, S.J. Mosser, A., Tham, S., Hopcraft, J.G.C., Mwangomo, E., Mlengeya, T.,
795 Kilewo, M., Fryxell, J.M., Sinclair, A.R.E., Borner, M. 2004. Can park protect
796 migratory ungulates? The case of Serengeti wildebeest. *Animal Conservation*,
797 7, 113-120.

798 URT United Republic of Tanzania, 2002. 2002 population and housing census.
799 <http://www.tanzania.org.tz/census/index.html> (accessed 30.03.2007).

800 van Ruth, S.M. and Roozen, J.P., 2000. Influence of mastication and saliva on aroma
801 release in a model mouth system. *Food Chemistry*, 71, 339-345.

802 Wilkie, D.S.; Malcolm, S., Kate, A., Ernestine, N.E., Paul, T., Ricardo, G., 2005. Role
803 of prices and wealth in consumer demand for bushmeat in Gabon, Central
804 Africa. *Conservation Biology*, 19, 268-274.

805

806

807 **Biographical sketches**

808 Julius Nyahongo is a senior research scientist with the Tanzanian Wildlife Research
809 Institute (TAWIRI). He is a wildlife ecologist specializing in human-wildlife interactions.
810 He has worked intensively with human-wildlife interactions in Tanzania.

811

812 Tomas Holmern is a researcher with the Department of Biology at the Norwegian
813 University of Science and Technology. He is interested in human-wildlife conflicts
814 and has recently been working on bushmeat hunting and local law enforcement in
815 the Serengeti.

816

817 Bård G. Stokke is a researcher at the Department of Biology, Norwegian University of
818 Science and Technology. He is an evolutionary biologist and behavioural ecologist
819 working on a wide range of birds and on human-wildlife conflicts over the use of
820 limited land.

821

822 Bjørn Kaltenborn is a senior research scientist with the Norwegian Institute for Nature
823 Research (NINA). He is a geographer and social scientist specializing in human-
824 environment interactions. He has worked extensively with human-wildlife conflicts in
825 the Nordic countries as well as in East Africa and South Asia.

826

827 Eivin Røskaft is a behavioural ecologist interested in a wide range of birds and
828 mammals species in Europe, North America and Africa, and in human-wildlife
829 conflicts over the use of limited land.

830

831

Table 1: The socio-demographic characteristics of test persons

<u>Location from the park boundary</u>	<u>Age (in years) categories (%)</u>		<u>Gender (%)</u>	
	5-17 years	18 and above years	Male	Female
Overall (n = 900)	20.5	79.5	65.7	34.3
Immediate (n = 300)	12.4	87.6	61.5	38.5
Intermediate (n = 300)	24.7	75.3	69.3	30.7
Distant (n = 300)	24.6	75.4	66.1	33.9

832
833

834 Table 2: Meat preference ranking and species identification by meat taste (Overall = all villages regardless of the distance from the park)

Category	Meat preference based on meat taste(%)					Species recognition by meat taste (%)				
	Beef	Topi	Impala	Zebra	Wildebeest	Beef	Topi	Impala	Zebra	Wildebeest
Overall: (n = 360)	69.7	63.3	56.1	33.0	29.7	40.0	20.0	12.0	29.7	27.8
Immediate villages (n = 120)	71.7	82.5	44.2	15.8	43.3	55.8	34.2	15.0	59.2	43.3
Intermediate villages (n = 120)	61.7	57.5	63.3	54.2	10.2	47.5	20.8	19.2	23.3	30.0
Distant villages (n = 120)	75.8	50.0	60.8	29.2	35.0	16.7	5.0	4.2	6.7	10.0

835
836
837
838
839
840
841
842
843
844
845
846
847

848 Table 3: The influences of distance, age and gender on meat preference ranking and species recognition (Pearson Chi-square test).
 849 Overall = all villages, age and gender. Overall = all villages regardless of the distance from the park). Number presented are p-value
 850 and significant effects are highlighted.

Category	Meat preference by meat taste				Species recognition by meat taste					
	Beef	Topi	Impala	Zebra	Wildebeest	Beef	Topi	Impala	Zebra	Wildebeest
Overall: (p value)										
Age (n = 360)	0.720	0.314	0.295	0.448	0.212	0.650	0.027	0.003	< 0.001	0.236
Gender (n = 360)	0.221	0.690	0.826	0.420	0.132	0.341	0.870	0.117	0.421	0.552
Distance (n = 360)	0.519	< 0.001	0.012	0.028	0.154	< 0.001	< 0.001	0.012	< 0.001	< 0.001
Immediate village:										
Age (n = 120)	0.442	0.746	0.423	0.436	0.404	0.426	0.767	0.058	0.053	0.924
Gender (n = 120)	0.649	0.597	0.798	0.037	0.766	0.327	0.929	0.322	0.659	0.503
Intermediate villages:										
Age (n = 120)	0.730	0.207	0.293	0.903	0.417	0.369	0.079	< 0.001	0.619	0.033
Gender (n = 120)	0.330	0.450	0.779	0.524	0.581	0.937	0.886	0.469	0.163	0.308
Distant villages:										
Age (n = 120)	0.730	0.207	0.293	0.903	0.417	0.071	< 0.001	0.002	0.006	< 0.001
Gender (n = 120)	0.330	0.450	0.779	0.524	0.581	0.657	0.267	0.122	0.879	0.949

851
 852
 853
 854
 855
 856
 857
 858
 859
 860
 861
 862
 863
 864
 865
 866

867 Table 4: Final logistic regression model of influence of distance from national park boundaries (immediate, intermediate and distant) , gender (male/female) and age
868 (child/adult), on meat preference among villagers close to Serengeti National Park, Tanzania of five species arranged in between-species comparisons. Preference refers to
869 overall preference of 90 test persons in each two-species comparison. Only scores of parameters included in the final model are presented. Parameter estimates (β), are
870 presented with their standard errors (SE). Wald statistic = $(\beta/SE)^2$. Odds-ratio = $\exp(\beta)$. Odds-ratio = $\exp(\beta)$, represents the ratio-change in the odds of the events of a one-unit
871 change in the predictor. * = interaction between main effects

<u>Species comparison</u>	<u>Parameter in the model</u>	<u>Estimate (β)</u>	<u>SE</u>	<u>Wald χ^2</u>	<u>P</u>	<u>Odds-ratio</u>
Beef-zebra	Gender*distance			9.637	0.008	
	Gender*distance (1)	-0.470	0.749	0.394	0.530	0.625
	Gender*distance (2)	-3.397	1.100	9.539	0.002	0.033
	Constant	1.317	0.291	20.530	<0.001	3.733
Beef-wildebeest	Distance			8.289	0.016	
	Distance (1)	-1.651	0.717	5.302	0.021	0.192
	Distance (2)	-8.71x10 ⁻¹⁷	0.861	1.025x10 ⁻³²	1.000	1.000
	Constant	2.197	0.609	13.035	<0.001	9.000
Topi-impala	Distance			19.461	<0.001	
	Distance (1)	2.719	0.669	16.533	<0.001	15.167
	Distance (2)	1.859	0.574	10.497	0.001	6.417
	Constant	-0.847	0.398	4.523	0.033	0.429
Topi-wildebeest	Distance			24.303	<0.001	
	Distance (1)	2.457	0.631	15.137	<0.001	11.667
	Distance (2)	3.045	0.727	17.519	<0.001	21.000
	Constant	-0.847	0.398	4.523	0.033	0.429
Topi-zebra	Distance			21.796	<0.001	
	Distance (1)	1.253	0.863	2.109	0.146	3.500
	Distance (2)	-2.079	0.599	12.067	0.001	0.125
	Constant	1.386	0.456	9.225	0.002	4.000
Impala-wildebeest	Distance			13.059	0.001	
	Distance (1)	-1.925	0.659	8.531	0.003	0.146
	Distance (2)	0.811	0.874	0.861	0.353	2.250
	Gender*distance			7.855	0.020	
Impala-zebra	Gender*distance (1)	2.842	1.152	6.089	0.014	17.143
	Gender*distance (2)	-1.350	1.016	1.766	0.184	0.259
	Constant	1.386	0.456	9.225	0.002	4.000
	Distance			6.743	0.034	
Impala-zebra	Distance (1)	-0.981	0.645	2.309	0.129	0.375
	Distance (2)	-1.792	0.697	6.604	0.010	0.167
	Constant	-0.405	0.456	0.789	0.374	0.667

Wildebeest-zebra	Distance				21.244	<0.001	
	Distance (1)	-1.375	0.611	5.059	0.025	0.253	
	Distance (2)	2.347	0.740	10.064	0.002	10.455	
	Gender	-1.392	0.587	5.622	0.018	0.249	
	Constant	0.496	0.464	1.141	0.285	1.641	

872

873 Table 5: Final logistic regression model of influence of distance from national park boundaries (immediate, intermediate and distant), gender (male/female) and age
874 (children/adult), on species recognition by meat taste among villagers close to Serengeti National Park, Tanzania of five species arranged in between-species comparisons.
875 Species recognition refers to overall recognition of 90 test persons in each two-species comparison. Only scores of parameters included in the final model are presented.
876 Parameter estimates (β), are presented with their standard errors (SE). Wald statistic = $(\beta/SE)^2$. Odds-ratio = $\exp(\beta)$, represents the ratio-change in the odds of the events of
877 the interest for a one-unit change in the predictor.

<u>Species comparison</u>	<u>Parameter in the model</u>	<u>Estimate (β)</u>	<u>SE</u>	<u>Wald χ^2</u>	<u>P</u>	<u>Odds-ratio</u>
Beef-topi	Distance			9.996	0.007	
	Distance (1)	2.005	0.650	9.520	0.002	7.429
	Distance (2)	1.025	0.669	2.347	0.126	2.786
Beef-impala	Constant	-1.872	0.537	12.146	<0.001	0.154
	Distance			10.837	0.004	
	Distance (1)	-0.342	0.587	0.340	0.560	0.710
Beef-wildebeest	Distance (2)	1.394	0.550	6.427	0.011	4.030
	Constant	-0.847	0.398	4.523	0.033	0.429
	Distance			14.799	0.001	
Beef-zebra	Distance (1)	3.636	1.082	11.295	0.001	37.923
	Distance (2)	2.178	1.105	3.885	0.049	8.826
	Constant	-3.367	1.017	10.961	0.001	0.034
Topi-zebra	Distance			11.609	0.003	
	Distance (1)	1.878	0.613	9.383	0.002	6.538
	Distance (2)	0.420	0.653	0.413	0.520	1.522
Impala-wildebeest	Constant	-1.609	0.490	10.793	0.001	0.200
	Distance			10.884	0.004	
	Distance (1)	3.099	1.082	8.207	0.004	22.176
Impala-zebra	Distance (2)	1.758	1.129	2.425	0.119	5.800
	Constant	-3.367	1.017	10.961	0.001	0.034
	Distance			6.116	0.047	
Wildebeest-zebra	Distance (1)	2.674	1.088	6.037	0.014	14.500
	Distance (2)	2.178	1.105	3.885	0.049	8.826
	Constant	-3.367	1.017	10.961	0.001	0.034
Impala-zebra	Distance			6.117	0.047	
	Distance (1)	19.859	8189.042	5.881×10^{-6}	0.998	4.200×10^8
	Distance (2)	17.745	8189.042	4.695×10^{-6}	0.998	5.087×10^7
Wildebeest-zebra	Gender	-19.378	7081.818	7.487×10^{-6}	0.998	0.000
	Constant	-20.419	8189.042	6.217×10^{-6}	0.998	0.000
	Distance			10.964	0.004	
	Distance (1)	1.595	0.570	7.823	0.005	4.929

	Distance (2)	0.000	0.610	0.000	1.000	1.000
	Constant	-1.190	0.432	7.594	0.006	0.304

878

879 Figure legends:

880 Figure 1. Map of the western Serengeti showing some villages where the meat taste
881 were conducted.

882

883

884

885

886

887

888

889

890

891

892

893

894

895

896

897

898

899



900
 901 Figure 1.
 902
 903
 904
 905
 906

Doctoral theses in Biology
Norwegian University of Science and Technology
Department of Biology

Year	Name	Degree	Title
1974	Tor-Henning Iversen	Dr. philos. Botany	The roles of statholiths, auxin transport, and auxin metabolism in root gravitropism
1978	Tore Slagsvold	Dr. philos. Zoology	Breeding events of birds in relation to spring temperature and environmental phenology.
1978	Egil Sakshaug	Dr. philos. Botany	"The influence of environmental factors on the chemical composition of cultivated and natural populations of marine phytoplankton"
1980	Arnfinn Langeland	Dr. philos. Zoology	Interaction between fish and zooplankton populations and their effects on the material utilization in a freshwater lake.
1980	Helge Reinertsen	Dr. philos. Botany	The effect of lake fertilization on the dynamics and stability of a limnetic ecosystem with special reference to the phytoplankton
1982	Gunn Mari Olsen	Dr. scient. Botany	Gravitropism in roots of <i>Pisum sativum</i> and <i>Arabidopsis thaliana</i>
1982	Dag Dolmen	Dr. philos. Zoology	Life aspects of two sympatric species of newts (<i>Triturus</i> , <i>Amphibia</i>) in Norway, with special emphasis on their ecological niche segregation.
1984	Eivin Røskoft	Dr. philos. Zoology	Sociobiological studies of the rook <i>Corvus frugilegus</i> .
1984	Anne Margrethe Cameron	Dr. scient. Botany	Effects of alcohol inhalation on levels of circulating testosterone, follicle stimulating hormone and luteinizing hormone in male mature rats
1984	Asbjørn Magne Nilsen	Dr. scient. Botany	Alveolar macrophages from expectorates – Biological monitoring of workers exposed to occupational air pollution. An evaluation of the AM-test
1985	Jarle Mork	Dr. philos. Zoology	Biochemical genetic studies in fish.
1985	John Solem	Dr. philos. Zoology	Taxonomy, distribution and ecology of caddisflies (<i>Trichoptera</i>) in the Dovrefjell mountains.
1985	Randi E. Reinertsen	Dr. philos. Zoology	Energy strategies in the cold: Metabolic and thermoregulatory adaptations in small northern birds.
1986	Bernt-Erik Sæther	Dr. philos. Zoology	Ecological and evolutionary basis for variation in reproductive traits of some vertebrates: A comparative approach.
1986	Torleif Holthe	Dr. philos. Zoology	Evolution, systematics, nomenclature, and zoogeography in the polychaete orders <i>Oweniimorpha</i> and <i>Terebellomorpha</i> , with special reference to the Arctic and Scandinavian fauna.
1987	Helene Lampe	Dr. scient. Zoology	The function of bird song in mate attraction and territorial defence, and the importance of song repertoires.
1987	Olav Hogstad	Dr. philos. Zoology	Winter survival strategies of the Willow tit <i>Parus montanus</i> .

1987 Jarle Inge Holten	Dr. philos Bothany	Autecological investigations along a coast-inland transect at Nord-Møre, Central Norway
1987 Rita Kumar	Dr. scient Botany	Somaclonal variation in plants regenerated from cell cultures of <i>Nicotiana sanderae</i> and <i>Chrysanthemum morifolium</i>
1987 Bjørn Åge Tømmerås	Dr. scient. Zoology	Olfaction in bark beetle communities: Interspecific interactions in regulation of colonization density, predator - prey relationship and host attraction.
1988 Hans Christian Pedersen	Dr. philos. Zoology	Reproductive behaviour in willow ptarmigan with special emphasis on territoriality and parental care.
1988 Tor G. Heggberget	Dr. philos. Zoology	Reproduction in Atlantic Salmon (<i>Salmo salar</i>): Aspects of spawning, incubation, early life history and population structure.
1988 Marianne V. Nielsen	Dr. scient. Zoology	The effects of selected environmental factors on carbon allocation/growth of larval and juvenile mussels (<i>Mytilus edulis</i>).
1988 Ole Kristian Berg	Dr. scient. Zoology	The formation of landlocked Atlantic salmon (<i>Salmo salar</i> L.).
1989 John W. Jensen	Dr. philos. Zoology	Crustacean plankton and fish during the first decade of the manmade Nesjø reservoir, with special emphasis on the effects of gill nets and salmonid growth.
1989 Helga J. Vivås	Dr. scient. Zoology	Theoretical models of activity pattern and optimal foraging: Predictions for the Moose <i>Alces alces</i> .
1989 Reidar Andersen	Dr. scient. Zoology	Interactions between a generalist herbivore, the moose <i>Alces alces</i> , and its winter food resources: a study of behavioural variation.
1989 Kurt Ingar Draget	Dr. scient Botany	Alginate gel media for plant tissue culture,
1990 Bengt Finstad	Dr. scient. Zoology	Osmotic and ionic regulation in Atlantic salmon, rainbow trout and Arctic charr: Effect of temperature, salinity and season.
1990 Hege Johannesen	Dr. scient. Zoology	Respiration and temperature regulation in birds with special emphasis on the oxygen extraction by the lung.
1990 Åse Krøkjke	Dr. scient Botany	The mutagenic load from air pollution at two work-places with PAH-exposure measured with Ames Salmonella/microsome test
1990 Arne Johan Jensen	Dr. philos. Zoology	Effects of water temperature on early life history, juvenile growth and prespawning migrations of Atlantic salmon (<i>Salmo salar</i>) and brown trout (<i>Salmo trutta</i>): A summary of studies in Norwegian streams.
1990 Tor Jørgen Almaas	Dr. scient. Zoology	Pheromone reception in moths: Response characteristics of olfactory receptor neurons to intra- and interspecific chemical cues.
1990 Magne Husby	Dr. scient. Zoology	Breeding strategies in birds: Experiments with the Magpie <i>Pica pica</i> .
1991 Tor Kvam	Dr. scient. Zoology	Population biology of the European lynx (<i>Lynx lynx</i>) in Norway.
1991 Jan Henning L'Abêe Lund	Dr. philos. Zoology	Reproductive biology in freshwater fish, brown trout <i>Salmo trutta</i> and roach <i>Rutilus rutilus</i> in particular.
1991 Asbjørn Moen	Dr. philos Botany	The plant cover of the boreal uplands of Central Norway. I. Vegetation ecology of Sølendet nature reserve; haymaking fens and birch woodlands
1991 Else Marie Løbersli	Dr. scient Botany	Soil acidification and metal uptake in plants

1991 Trond Nordtug	Dr. scient. Zoology	Reflctometric studies of photomechanical adaptation in superposition eyes of arthropods.
1991 Thyra Solem	Dr. scient Botany	Age, origin and development of blanket mires in Central Norway
1991 Odd Terje Sandlund	Dr. philos. Zoology	The dynamics of habitat use in the salmonid genera <i>Coregonus</i> and <i>Salvelinus</i> : Ontogenic niche shifts and polymorphism.
1991 Nina Jonsson	Dr. philos.	Aspects of migration and spawning in salmonids.
1991 Atle Bones	Dr. scient Botany	Compartmentation and molecular properties of thioglucoside glucohydrolase (myrosinase)
1992 Torggrim Breiehagen	Dr. scient. Zoology	Mating behaviour and evolutionary aspects of the breeding system of two bird species: the Temminck's stint and the Pied flycatcher.
1992 Anne Kjersti Bakken	Dr. scient Botany	The influence of photoperiod on nitrate assimilation and nitrogen status in timothy (<i>Phleum pratense</i> L.)
1992 Tycho Anker-Nilssen	Dr. scient. Zoology	Food supply as a determinant of reproduction and population development in Norwegian Puffins <i>Fratercula arctica</i>
1992 Bjørn Munro Jenssen	Dr. philos. Zoology	Thermoregulation in aquatic birds in air and water: With special emphasis on the effects of crude oil, chemically treated oil and cleaning on the thermal balance of ducks.
1992 Arne Vollan Aarset	Dr. philos. Zoology	The ecophysiology of under-ice fauna: Osmotic regulation, low temperature tolerance and metabolism in polar crustaceans.
1993 Geir Slupphaug	Dr. scient Botany	Regulation and expression of uracil-DNA glycosylase and O ⁶ -methylguanine-DNA methyltransferase in mammalian cells
1993 Tor Fredrik Næsje	Dr. scient. Zoology	Habitat shifts in coregonids.
1993 Yngvar Asbjørn Olsen	Dr. scient. Zoology	Cortisol dynamics in Atlantic salmon, <i>Salmo salar</i> L.: Basal and stressor-induced variations in plasma levels and some secondary effects.
1993 Bård Pedersen	Dr. scient Botany	Theoretical studies of life history evolution in modular and clonal organisms
1993 Ole Petter Thangstad	Dr. scient Botany	Molecular studies of myrosinase in Brassicaceae
1993 Thrine L. M. Heggberget	Dr. scient. Zoology	Reproductive strategy and feeding ecology of the Eurasian otter <i>Lutra lutra</i> .
1993 Kjetil Bevanger	Dr. scient. Zoology	Avian interactions with utility structures, a biological approach.
1993 Kåre Haugan	Dr. scient Bothany	Mutations in the replication control gene trfA of the broad host-range plasmid RK2
1994 Peder Fiske	Dr. scient. Zoology	Sexual selection in the lekking great snipe (<i>Gallinago media</i>): Male mating success and female behaviour at the lek.
1994 Kjell Inge Reitan	Dr. scient Botany	Nutritional effects of algae in first-feeding of marine fish larvae
1994 Nils Røv	Dr. scient. Zoology	Breeding distribution, population status and regulation of breeding numbers in the northeast-Atlantic Great Cormorant <i>Phalacrocorax carbo carbo</i> .
1994 Annette-Susanne Hoepfner	Dr. scient Botany	Tissue culture techniques in propagation and breeding of Red Raspberry (<i>Rubus idaeus</i> L.)
1994 Inga Elise Bruteig	Dr. scient Bothany	Distribution, ecology and biomonitoring studies of epiphytic lichens on conifers

1994 Geir Johnsen	Dr. scient Botany	Light harvesting and utilization in marine phytoplankton: Species-specific and photoadaptive responses
1994 Morten Bakken	Dr. scient. Zoology	Infanticidal behaviour and reproductive performance in relation to competition capacity among farmed silver fox vixens, <i>Vulpes vulpes</i> .
1994 Arne Moksnes	Dr. philos. Zoology	Host adaptations towards brood parasitism by the Cuckoo.
1994 Solveig Bakken	Dr. scient Bothany	Growth and nitrogen status in the moss <i>Dicranum majus</i> Sm. as influenced by nitrogen supply
1995 Olav Vadstein	Dr. philos Botany	The role of heterotrophic planktonic bacteria in the cycling of phosphorus in lakes: Phosphorus requirement, competitive ability and food web interactions.
1995 Hanne Christensen	Dr. scient. Zoology	Determinants of Otter <i>Lutra lutra</i> distribution in Norway: Effects of harvest, polychlorinated biphenyls (PCBs), human population density and competition with mink <i>Mustela vison</i> .
1995 Svein Håkon Lorentsen	Dr. scient. Zoology	Reproductive effort in the Antarctic Petrel <i>Thalassoica antarctica</i> ; the effect of parental body size and condition.
1995 Chris Jørgen Jensen	Dr. scient. Zoology	The surface electromyographic (EMG) amplitude as an estimate of upper trapezius muscle activity
1995 Martha Kold Bakkevig	Dr. scient. Zoology	The impact of clothing textiles and construction in a clothing system on thermoregulatory responses, sweat accumulation and heat transport.
1995 Vidar Moen	Dr. scient. Zoology	Distribution patterns and adaptations to light in newly introduced populations of <i>Mysis relicta</i> and constraints on Cladoceran and Char populations.
1995 Hans Haavardsholm Blom	Dr. philos Bothany	A revision of the <i>Schistidium apocarpum</i> complex in Norway and Sweden.
1996 Jorun Skjærmo	Dr. scient Botany	Microbial ecology of early stages of cultivated marine fish; impact fish-bacterial interactions on growth and survival of larvae.
1996 Ola Ugedal	Dr. scient. Zoology	Radiocesium turnover in freshwater fishes
1996 Ingibjörg Einarsdottir	Dr. scient. Zoology	Production of Atlantic salmon (<i>Salmo salar</i>) and Arctic charr (<i>Salvelinus alpinus</i>): A study of some physiological and immunological responses to rearing routines.
1996 Christina M. S. Pereira	Dr. scient. Zoology	Glucose metabolism in salmonids: Dietary effects and hormonal regulation.
1996 Jan Fredrik Børseth	Dr. scient. Zoology	The sodium energy gradients in muscle cells of <i>Mytilus edulis</i> and the effects of organic xenobiotics.
1996 Gunnar Henriksen	Dr. scient. Zoology	Status of Grey seal <i>Halichoerus grypus</i> and Harbour seal <i>Phoca vitulina</i> in the Barents sea region.
1997 Gunvor Øie	Dr. scient Bothany	Eevaluation of rotifer <i>Brachionus plicatilis</i> quality in early first feeding of turbot <i>Scophthalmus maximus</i> L. larvae.
1997 Håkon Holien	Dr. scient Botany	Studies of lichens in spruce forest of Central Norway. Diversity, old growth species and the relationship to site and stand parameters.
1997 Ole Reitan	Dr. scient. Zoology	Responses of birds to habitat disturbance due to damming.
1997 Jon Arne Grøttum	Dr. scient. Zoology	Physiological effects of reduced water quality on fish in aquaculture.

1997 Per Gustav Thingstad	Dr. scient. Zoology	Birds as indicators for studying natural and human-induced variations in the environment, with special emphasis on the suitability of the Pied Flycatcher.
1997 Torgeir Nygård	Dr. scient. Zoology	Temporal and spatial trends of pollutants in birds in Norway: Birds of prey and Willow Grouse used as Biomonitor.
1997 Signe Nybø	Dr. scient. Zoology	Impacts of long-range transported air pollution on birds with particular reference to the dipper <i>Cinclus cinclus</i> in southern Norway.
1997 Atle Wibe	Dr. scient. Zoology	Identification of conifer volatiles detected by receptor neurons in the pine weevil (<i>Hylobius abietis</i>), analysed by gas chromatography linked to electrophysiology and to mass spectrometry.
1997 Rolv Lundheim	Dr. scient. Zoology	Adaptive and incidental biological ice nucleators.
1997 Arild Magne Landa	Dr. scient. Zoology	Wolverines in Scandinavia: ecology, sheep depredation and conservation.
1997 Kåre Magne Nielsen	Dr. scient. Botany	An evolution of possible horizontal gene transfer from plants to soil bacteria by studies of natural transformation in <i>Acinetobacter calcoaceticus</i> .
1997 Jarle Tufto	Dr. scient. Zoology	Gene flow and genetic drift in geographically structured populations: Ecological, population genetic, and statistical models
1997 Trygve Hesthagen	Dr. philos. Zoology	Population responses of Arctic charr (<i>Salvelinus alpinus</i> L.) and brown trout (<i>Salmo trutta</i> L.) to acidification in Norwegian inland waters
1997 Trygve Sigholt	Dr. philos. Zoology	Control of Parr-smolt transformation and seawater tolerance in farmed Atlantic Salmon (<i>Salmo salar</i>) Effects of photoperiod, temperature, gradual seawater acclimation, NaCl and betaine in the diet
1997 Jan Østnes	Dr. scient. Zoology	Cold sensation in adult and neonate birds
1998 Seethaledsumy Visvalingam	Dr. scient. Botany	Influence of environmental factors on myrosinases and myrosinase-binding proteins.
1998 Thor Harald Ringsby	Dr. scient. Zoology	Variation in space and time: The biology of a House sparrow metapopulation
1998 Erling Johan Solberg	Dr. scient. Zoology	Variation in population dynamics and life history in a Norwegian moose (<i>Alces alces</i>) population: consequences of harvesting in a variable environment
1998 Sigurd Mjøen Saastad	Dr. scient. Botany	Species delimitation and phylogenetic relationships between the Sphagnum recurvum complex (Bryophyta): genetic variation and phenotypic plasticity.
1998 Bjarte Mortensen	Dr. scient. Botany	Metabolism of volatile organic chemicals (VOCs) in a head liver S9 vial equilibration system in vitro.
1998 Gunnar Austrheim	Dr. scient. Botany	Plant biodiversity and land use in subalpine grasslands. – A conservation biological approach.
1998 Bente Gunnveig Berg	Dr. scient. Zoology	Encoding of pheromone information in two related moth species
1999 Kristian Overskaug	Dr. scient. Zoology	Behavioural and morphological characteristics in Northern Tawny Owls <i>Strix aluco</i> : An intra- and interspecific comparative approach
1999 Hans Kristen Stenøien	Dr. scient. Botany	Genetic studies of evolutionary processes in various populations of nonvascular plants (mosses, liverworts and hornworts)

1999 Trond Arnesen	Dr. scient. Botany	Vegetation dynamics following trampling and burning in the outlying haylands at Sølendet, Central Norway.
1999 Ingvar Stenberg	Dr. scient. Zoology	Habitat selection, reproduction and survival in the White-backed Woodpecker <i>Dendrocopos leucotos</i>
1999 Stein Olle Johansen	Dr. scient. Botany	A study of driftwood dispersal to the Nordic Seas by dendrochronology and wood anatomical analysis.
1999 Trina Falck Galloway	Dr. scient. Zoology	Muscle development and growth in early life stages of the Atlantic cod (<i>Gadus morhua</i> L.) and Halibut (<i>Hippoglossus hippoglossus</i> L.)
1999 Torbjørn Forseth	Dr. scient. Zoology	Bioenergetics in ecological and life history studies of fishes.
1999 Marianne Giæver	Dr. scient. Zoology	Population genetic studies in three gadoid species: blue whiting (<i>Micromisistius poutassou</i>), haddock (<i>Melanogrammus aeglefinus</i>) and cod (<i>Gradus morhua</i>) in the North-East Atlantic
1999 Hans Martin Hanslin	Dr. scient. Botany	The impact of environmental conditions of density dependent performance in the boreal forest bryophytes <i>Dicranum majus</i> , <i>Hylocomium splendens</i> , <i>Plagiochila asplenigides</i> , <i>Ptilium crista-castrensis</i> and <i>Rhytidiadelphus lokeus</i> .
1999 Ingrid Bysveen Mjølnørød	Dr. scient. Zoology	Aspects of population genetics, behaviour and performance of wild and farmed Atlantic salmon (<i>Salmo salar</i>) revealed by molecular genetic techniques
1999 Else Berit Skagen	Dr. scient. Botany	The early regeneration process in protoplasts from <i>Brassica napus</i> hypocotyls cultivated under various g-forces
1999 Stein-Are Sæther	Dr. philos. Zoology	Mate choice, competition for mates, and conflicts of interest in the Lekking Great Snipe
1999 Katrine Wangen Rustad	Dr. scient. Zoology	Modulation of glutamatergic neurotransmission related to cognitive dysfunctions and Alzheimer's disease
1999 Per Terje Smiseth	Dr. scient. Zoology	Social evolution in monogamous families: mate choice and conflicts over parental care in the Bluethroat (<i>Luscinia s. svecica</i>)
1999 Gunnbjørn Bremset	Dr. scient. Zoology	Young Atlantic salmon (<i>Salmo salar</i> L.) and Brown trout (<i>Salmo trutta</i> L.) inhabiting the deep pool habitat, with special reference to their habitat use, habitat preferences and competitive interactions
1999 Frode Ødegaard	Dr. scient. Zoology	Host specificity as parameter in estimates of arthropod species richness
1999 Sonja Andersen	Dr. scient. Bothany	Expressional and functional analyses of human, secretory phospholipase A2
2000 Ingrid Salvesen, I	Dr. scient. Botany	Microbial ecology in early stages of marine fish: Development and evaluation of methods for microbial management in intensive larviculture
2000 Ingar Jostein Øien	Dr. scient. Zoology	The Cuckoo (<i>Cuculus canorus</i>) and its host: adaptations and counteradaptations in a coevolutionary arms race
2000 Pavlos Makridis	Dr. scient. Botany	Methods for the microbial econtrol of live food used for the rearing of marine fish larvae
2000 Sigbjørn Stokke	Dr. scient. Zoology	Sexual segregation in the African elephant (<i>Loxodonta africana</i>)
2000 Odd A. Gulseth	Dr. philos. Zoology	Seawater tolerance, migratory behaviour and growth of Charr, (<i>Salvelinus alpinus</i>), with emphasis on the high Arctic Dieset charr on Spitsbergen, Svalbard

2000 Pål A. Olsvik	Dr. scient. Zoology	Biochemical impacts of Cd, Cu and Zn on brown trout (<i>Salmo trutta</i>) in two mining-contaminated rivers in Central Norway
2000 Sigurd Einum	Dr. scient. Zoology	Maternal effects in fish: Implications for the evolution of breeding time and egg size
2001 Jan Ove Evjemo	Dr. scient. Zoology	Production and nutritional adaptation of the brine shrimp <i>Artemia</i> sp. as live food organism for larvae of marine cold water fish species
2001 Olga Hilmo	Dr. scient Botany	Lichen response to environmental changes in the managed boreal forest systems
2001 Ingebrigt Uglem	Dr. scient. Zoology	Male dimorphism and reproductive biology in corkwing wrasse (<i>Symphodus melops</i> L.)
2001 Bård Gunnar Stokke	Dr. scient. Zoology	Coevolutionary adaptations in avian brood parasites and their hosts
2002 Ronny Aanes	Dr. scient	Spatio-temporal dynamics in Svalbard reindeer (<i>Rangifer tarandus platyrhynchus</i>)
2002 Mariann Sandsund	Dr. scient. Zoology	Exercise- and cold-induced asthma. Respiratory and thermoregulatory responses
2002 Dag-Inge Øien	Dr. scient Botany	Dynamics of plant communities and populations in boreal vegetation influenced by scything at Sølendet, Central Norway
2002 Frank Rosell	Dr. scient. Zoology	The function of scent marking in beaver (<i>Castor fiber</i>)
2002 Janne Østvang	Dr. scient Botany	The Role and Regulation of Phospholipase A ₂ in Monocytes During Atherosclerosis Development
2002 Terje Thun	Dr.philos Biology	Dendrochronological constructions of Norwegian conifer chronologies providing dating of historical material
2002 Birgit Hafjeld Borgen	Dr. scient Biology	Functional analysis of plant idioblasts (Myrosin cells) and their role in defense, development and growth
2002 Bård Øyvind Solberg	Dr. scient Biology	Effects of climatic change on the growth of dominating tree species along major environmental gradients
2002 Per Winge	Dr. scient Biology	The evolution of small GTP binding proteins in cellular organisms. Studies of RAC GTPases in <i>Arabidopsis thaliana</i> and
2002 Henrik Jensen	Dr. scient Biology	Causes and consequences of individual variation in fitness-related traits in house sparrows
2003 Jens Rohloff	Dr. philos Biology	Cultivation of herbs and medicinal plants in Norway – Essential oil production and quality control
2003 Åsa Maria O. Espmark Wibe	Dr. scient Biology	Behavioural effects of environmental pollution in threespine stickleback <i>Gasterosteus aculeatus</i> L.
2003 Dagmar Hagen	Dr. scient Biology	Assisted recovery of disturbed arctic and alpine vegetation – an integrated approach
2003 Bjørn Dahle	Dr. scient Biology	Reproductive strategies in Scandinavian brown bears
2003 Cyril Lebogang Taolo	Dr. scient Biology	Population ecology, seasonal movement and habitat use of the African buffalo (<i>Syncerus caffer</i>) in Chobe National Park, Botswana
2003 Marit Stranden	Dr.scient Biology	Olfactory receptor neurones specified for the same odorants in three related Heliothine species (<i>Helicoverpa armigera</i> , <i>Helicoverpa assulta</i> and <i>Heliothis virescens</i>)
2003 Kristian Hassel	Dr.scient Biology	Life history characteristics and genetic variation in an expanding species, <i>Pogonatum dentatum</i>

2003	David Alexander Rae	Dr.scient Biology	Plant- and invertebrate-community responses to species interaction and microclimatic gradients in alpine and Arctic environments
2003	Åsa A Borg	Dr.scient Biology	Sex roles and reproductive behaviour in gobies and guppies: a female perspective
2003	Eldar Åsgard Bendiksen	Dr.scient Biology	Environmental effects on lipid nutrition of farmed Atlantic salmon (<i>Salmo Salar</i> L.) parr and smolt
2004	Torkild Bakken	Dr.scient Biology	A revision of Nereidinae (Polychaeta, Nereididae)
2004	Ingar Pareliussen	Dr.scient Biology	Natural and Experimental Tree Establishment in a Fragmented Forest, Ambohitantely Forest Reserve, Madagascar
2004	Tore Brembu	Dr.scient Biology	Genetic, molecular and functional studies of RAC GTPases and the WAVE-like regulatory protein complex in <i>Arabidopsis thaliana</i>
2004	Liv S. Nilsen	Dr.scient Biology	Coastal heath vegetation on central Norway; recent past, present state and future possibilities
2004	Hanne T. Skiri	Dr.scient Biology	Olfactory coding and olfactory learning of plant odours in heliothine moths. An anatomical, physiological and behavioural study of three related species (<i>Heliothis virescens</i> , <i>Helicoverpa armigera</i> and <i>Helicoverpa assulta</i>).
2004	Lene Østby	Dr.scient Biology	Cytochrome P4501A (CYP1A) induction and DNA adducts as biomarkers for organic pollution in the natural environment
2004	Emmanuel J. Gerreta	Dr. philos Biology	The Importance of Water Quality and Quantity in the Tropical Ecosystems, Tanzania
2004	Linda Dalen	Dr.scient Biology	Dynamics of Mountain Birch Treelines in the Scandes Mountain Chain, and Effects of Climate Warming
2004	Lisbeth Mehli	Dr.scient Biology	Polygalacturonase-inhibiting protein (PGIP) in cultivated strawberry (<i>Fragaria x ananassa</i>): characterisation and induction of the gene following fruit infection by <i>Botrytis cinerea</i>
2004	Børge Moe	Dr.scient Biology	Energy-Allocation in Avian Nestlings Facing Short-Term Food Shortage
2005	Matilde Skogen Chauton	Dr.scient Biology	Metabolic profiling and species discrimination from High-Resolution Magic Angle Spinning NMR analysis of whole-cell samples
2005	Sten Karlsson	Dr.scient Biology	Dynamics of Genetic Polymorphisms
2005	Terje Bongard	Dr.scient Biology	Life History strategies, mate choice, and parental investment among Norwegians over a 300-year period
2005	Tonette Røstelién	PhD Biology	Functional characterisation of olfactory receptor neurone types in heliothine moths
2005	Erlend Kristiansen	Dr.scient Biology	Studies on antifreeze proteins
2005	Eugen G. Sørmo	Dr.scient Biology	Organochlorine pollutants in grey seal (<i>Halichoerus grypus</i>) pups and their impact on plasma thyroid hormone and vitamin A concentrations.
2005	Christian Westad	Dr.scient Biology	Motor control of the upper trapezius

2005 Lasse Mork Olsen	PhD Biology	Interactions between marine osmo- and phagotrophs in different physicochemical environments
2005 Åslaug Viken	PhD Biology	Implications of mate choice for the management of small populations
2005 Ariaya Hymete Sahle Dingle	PhD Biology	Investigation of the biological activities and chemical constituents of selected <i>Echinops</i> spp. growing in Ethiopia
2005 Ander Gravbrøt Finstad	PhD Biology	Salmonid fishes in a changing climate: The winter challenge
2005 Shimane Washington Makabu	PhD Biology	Interactions between woody plants, elephants and other browsers in the Chobe Riverfront, Botswana
2005 Kjartan Østbye	Dr.scient Biology	The European whitefish <i>Coregonus lavaretus</i> (L.) species complex: historical contingency and adaptive radiation
2006 Kari Mette Murvoll	PhD Biology	Levels and effects of persistent organic pollutants (POPs) in seabirds Retinoids and α -tocopherol – potential biomarkers of POPs in birds?
2006 Ivar Herfindal	Dr.scient Biology	Life history consequences of environmental variation along ecological gradients in northern ungulates
2006 Nils Egil Tokle	Phd Biology	Are the ubiquitous marine copepods limited by food or predation? Experimental and field-based studies with main focus on <i>Calanus finmarchicus</i>
2006 Jan Ove Gjershaug	Dr.philos Biology	Taxonomy and conservation status of some booted eagles in south-east Asia
2006 Jon Kristian Skei	Dr.scient Biology	Conservation biology and acidification problems in the breeding habitat of amphibians in Norway
2006 Johanna Järnegen	PhD Biology	Acesta Oophaga and Acesta Excavata – a study of hidden biodiversity
2006 Bjørn Henrik Hansen	PhD Biology	Metal-mediated oxidative stress responses in brown trout (<i>Salmo trutta</i>) from mining contaminated rivers in Central Norway
2006 Vidar Grøtan	phD Biology	Temporal and spatial effects of climate fluctuations on population dynamics of vertebrates
2006 Jafari R Kideghesho	phD Biology	Wildlife conservation and local land use conflicts in western Serengeti, Corridor Tanzania
2006 Anna Maria Billing	phD Biology	Reproductive decisions in the sex role reversed pipefish <i>Syngnathus typhle</i> : when and how to invest in reproduction
2006 Henrik Pärn	phD Biology	Female ornaments and reproductive biology in the bluethroat
2006 Anders J. Fjellheim	phD Biology	Selection and administration of probiotic bacteria to marine fish larvae
2006 P. Andreas Svensson	phD Biology	Female coloration, egg carotenoids and reproductive success: gobies as a model system
2007 Sindre A. Pedersen	phD Biology	Metal binding proteins and antifreeze proteins in the beetle <i>Tenebrio molitor</i> - a study on possible competition for the semi-essential amino acid cysteine
2007 Kasper Hancke	phD Biology	Photosynthetic responses as a function of light and temperature: Field and laboratory studies on marine microalgae
2007 Tomas Holmern	phD Biology	Bushmeat hunting in the western Serengeti: Implications for community-based conservation

2007 Kari Jørgensen	phD Biology	Functional tracing of gustatory receptor neurons in the CNS and chemosensory learning in the moth <i>Heliothis virescens</i>
2007 Stig Ulland	phD Biology	Functional Characterisation of Olfactory Receptor Neurons in the Cabbage Moth, <i>Mamestra Brassicae</i> /L. (Lepidoptera, Noctuidae). Gas Chromatography Linked to Single Cell Recordings and Mass Spectrometry
2007 Snorre Henriksen	phD Biology	Spatial and temporal variation in herbivore resources at northern latitudes
2007 Roelof Frans May	phD Biology	Spatial Ecology of Wolverines in Scandinavia
2007 Vedasto Gabriel Ndibalema	phD Biology	Demographic variation, distribution and habitat use between wildebeest sub-populations in the Serengeti National Park, Tanzania

