

**Associations of obesity and lifestyle with the risk and mortality of bloodstream infection in a general population: a 15-year follow-up of 64,027 individuals in the HUNT Study**

Julie Paulsen<sup>\*1,2,10</sup>, Åsa Askim<sup>3,4,10</sup>, Randi Marie Mohus<sup>3,4,10</sup>, Arne Mehl<sup>2,5,10</sup>, Andrew Dewan<sup>6</sup>, Erik Solligård<sup>3,4,10</sup>, Jan Kristian Damås<sup>1,7,10</sup>, Bjørn O. Åsvold<sup>8,9,10</sup>

1 Centre of Molecular Inflammation Research, Department of Cancer Research and Molecular Medicine, NTNU, Norwegian University of Science and Technology, Trondheim, Norway

2 Department of Medicine, Levanger Hospital, Nord-Trøndelag Hospital Trust, Norway

3 Department of Circulation and Medical Imaging, NTNU, Norwegian University of Science and Technology, Trondheim, Norway

4 Clinic of Anaesthesia and Intensive Care, St. Olavs Hospital, Trondheim University Hospital, Trondheim, Norway

5 Department of Cancer Research and Molecular Medicine, NTNU, Norwegian University of Science and Technology, Trondheim, Norway

6 Department of Chronic Disease Epidemiology, Yale University School of Public Health, New Haven CT, USA

7 Department of Infectious Diseases, St. Olavs Hospital, Trondheim University Hospital, Trondheim, Norway

8 Department of Public Health and Nursing, NTNU, Norwegian University of Science and Technology, Trondheim, Norway

9 Department of Endocrinology, St. Olavs Hospital, Trondheim University Hospital, Trondheim, Norway

10 Mid-Norway Sepsis Research Center, NTNU, Norwegian University of Science and Technology and St. Olavs Hospital, Trondheim University Hospital, Trondheim, Norway

Corresponding author

Julie Paulsen,

Centre of Molecular Inflammation Research, Department of Cancer Research and Molecular Medicine, NTNU, Norwegian University of Science and Technology

Postboks 8905

N-7491 Trondheim

Norway

Phone: +47-97569485, E-mail: [julie.paulsen@ntnu.no](mailto:julie.paulsen@ntnu.no)

Word count text: 3359

Word count abstract: 240

## Abstract

Background: Bloodstream infections (BSIs) cause considerable morbidity and mortality, and primary prevention should be a priority. Lifestyle factors are of particular interest since they represent a modifiable target.

Methods: We conducted a prospective cohort study among participants in the population-based Norwegian HUNT2 Survey, where 64,027 participants were followed from 1995-97 through 2011 by linkage to prospectively recorded information on BSI at local and regional hospitals. The exposures were baseline body mass index (BMI) measurements, and self-reported smoking habits, leisure time physical activity and alcohol intake. The outcomes were hazard ratios (HR) of BSI and BSI mortality.

Results: During 810,453 person-years and median follow-up of 14.8 years, 1844 (2.9%) participants experienced at least one BSI and 396 (0.62%) died from BSI. Compared with normal weight participants (BMI 18.5-24.9 kg/m<sup>2</sup>), the age- and sex-adjusted risk of a first-time BSI was 31% (95% confidence interval (CI) 14-51%) higher at BMI 30.0-34.9 kg/m<sup>2</sup>, 87% (95% CI 50-135%) higher at BMI 35.0-39.9 kg/m<sup>2</sup> and 210% (95% CI 117-341%) higher at BMI ≥40.0 kg/m<sup>2</sup>. The risk of BSI mortality was similarly increased. Compared with never-smokers, current smokers had 51% (95% CI 34-70%) and 75% (95% CI 34-129%) higher risks of BSI and BSI mortality, respectively. Physically inactive participants had 71% (95% CI 42-107%) and 108% (95% CI 37-216%) higher risks of BSI and BSI mortality, respectively, compared with the most physically active.

Conclusions: Obesity, smoking and physical inactivity carry increased risk of BSI and BSI mortality.

Keywords: Bacteremia; Sepsis; Obesity; Smoking; Exercise; Alcohol drinking

### **Key messages**

- Primary prevention of bloodstream infections should be a priority for reducing their morbidity and mortality. Lifestyle factors are modifiable, and may represent an efficient target.
- Few studies have assessed infection-related mortality at the population level, or the role of multiple lifestyle factors.
- In this population-based study, we found strong dose-response relationships between obesity and the risk of BSI and BSI mortality, and the importance of obesity as a risk factor varied by infecting organism.
- Smoking was associated with increased risk of BSI and BSI mortality, even with pathogens that do not primarily infect the airways, such as *Staphylococcus aureus* and *Escherichia coli*.
- The combined effects of obesity, smoking and physical inactivity increased the risk of BSI and BSI mortality 5- to 6-fold at the population level.

## Introduction

Bloodstream infection (BSI), defined as bacteremia associated with infection, is a cause of considerable morbidity and mortality (1). The overall incidence is approximately 180/100,000 person years in high-income countries, and with mortality rates of 20-30/100,000 person-years it ranks among the top seven causes of death in North-America and parts of Europe (2).

In recent years there have been increased initiatives to identify risk factors for BSI, sepsis and other severe bacterial infections (3-6). Lifestyle factors are of particular interest since they are modifiable, and optimization could reduce the burden of BSI at the population level. Yet, the influence of lifestyle factors on the occurrence of severe infections is insufficiently known. Studies assessing the role of lifestyle factors for the risk of infection have usually focused on specific infections or microorganisms (7-10). Few studies have assessed infection-related mortality at the population level, or the role of multiple lifestyle factors. In this Norwegian population-based cohort study of 64,027 subjects, we assessed the associations of obesity, smoking, physical inactivity and alcohol intake with the risk of BSI and BSI mortality.

## Methods

Nord-Trøndelag county in central Norway has a population of approximately 130,000. The county consists of rural areas and small towns. The county is considered generally representative of Norway with regards to sources of income, age distribution, morbidity and mortality, but the average income and prevalence of higher education and current smoking are a little lower than the Norwegian average, and the county has no large cities (11). The HUNT Study (an acronym for the Nord-Trøndelag Health Study, in Norwegian: Helseundersøkelsen i Nord-Trøndelag) is a series of surveys to which all inhabitants in Nord-Trøndelag aged 20 years or older were invited

to participate: HUNT1 (1984-1986), HUNT2 (1995-1997) and HUNT3 (2006-2008) (12). The present study was performed among all participants of HUNT2, where 93,865 persons were invited and 65,236 (69.5%) participated. Among them, 47,316 (72.5%) had also participated in HUNT1 (12). The participants returned questionnaires covering a wide range of health-related topics including lifestyle factors and prior illnesses. They attended a clinical examination that included, among other items, standardized measurements of height, weight and waist circumference (11). The HUNT study database is regularly updated with information on site of residence and vital status from the Norwegian population registry.

Body mass index (BMI) was calculated as weight in kilograms over squared height in  $m^2$  and categorized according to World Health Organization (WHO) recommendations as  $<18.5$  (underweight), 18.5–24.9 (normal weight), 25.0-29.9 (pre-obese), 30.0–34.9 (obese class I), 35.0-39.9 (obese class II) and  $\geq 40.0$   $kg/m^2$  (obese class III) (13). As an alternative measure of obesity we used waist circumference categorized according to WHO as  $<94$ , 94–102 and  $>102$  cm in males and  $<80$ , 80-88 and  $>88$  cm in females (14). Smoking habits were categorized as current, prior or never-smoking according to self-report at HUNT2. Participants were asked about their leisure time weekly amount of light (not sweaty or breathless) and vigorous (sweaty or breathless) physical activity during the last year. We categorized participants as being non- (no vigorous or light activity), slightly ( $<3$  hours of weekly light activity and no vigorous activity), moderately ( $\geq 3$  hours of weekly light activity or  $<1$  hour of vigorous activity) or highly ( $\geq 1$  hour of vigorous activity per week) physically active. We used self-reported sitting time as an additional measure of physical inactivity ( $\leq 4$  hours, 5-7 hours and  $\geq 8$  hours/day).

Alcohol consumption was categorized as  $<1$ , 1-7, 8-14 and  $\geq 15$  glasses/2 weeks. 10.7% of participants only provided information on the frequency of alcohol intake per month. Reported

intake of 0 times/month was grouped with <1 glasses/2 weeks, 1-3 times/month with 1-7 glasses/2 weeks, 4-7 times/month with 8-14 glasses/2 weeks and >7 times/month with  $\geq 15$  glasses/2 weeks. The correlation between these categories of frequency and amount was good with a Spearman's correlation coefficient of 0.80. For all exposure variables, the reference was the category most closely corresponding to health recommendations, i.e. normal weight, never-smokers, and at least one weekly hour of vigorous activity (activity generating sweat or breathlessness). For alcohol consumption, 1-7 glasses/two weeks was set as the reference, because the group of subjects with no alcohol consumption could include subjects that are abstainers for health reasons that may themselves increase the risk of BSI, such as chronic disease and prior alcohol abuse.

Nord-Trøndelag is served by two community hospitals, Levanger Hospital and Namsos Hospital (Nord-Trøndelag Hospital Trust), and St. Olavs Hospital in Trondheim serves as the tertiary referral center. Using the 11-digit unique personal identification number of Norwegian citizens, we linked the HUNT study information to prospectively recorded information on dates and microbes detected in blood cultures at the microbiology laboratories of Levanger and St. Olavs Hospitals from January 1, 1995, and from Namsos Hospital from September 1, 1999. Isolates solely consisting of microorganisms commonly associated with skin contamination such as coagulase negative *Staphylococcus* species, *Corynebacterium* species and *Propionibacterium* species were not considered as BSI (15). BSI mortality was defined as death within 30 days of detection of a BSI, and dates of death were obtained from the Norwegian population register. In participants with multiple positive blood culture specimens, a new episode of BSI was defined as positive blood culture more than 30 days after the previous one.

## Statistical analysis

For each outcome (first-time BSI and BSI mortality), we used Cox regression analysis to estimate hazard ratios (HRs) with 95% confidence intervals (CIs) by categories of obesity and lifestyle variables. The participants were followed from the date of participation in HUNT2 between August 1995 and June 1997 for residents having Levanger Hospital as their primary hospital or from September 1, 1999 for residents having Namsos Hospital as their primary hospital. In the analyses of first-time BSI, participants were followed until their first BSI, migration out of Nord-Trøndelag, death, or end of follow-up at December 31, 2011, whichever occurred first. In the analyses of BSI mortality, participants were followed until migration out of Nord-Trøndelag, death, or December 31, 2011, whichever occurred first. Prior to the start of follow-up, 47 (0.07%) participants experienced a positive blood culture and 1140 (1.78%) were censored due to migration or death, and 22 were excluded because they lacked information on all exposure variables, leaving 64,027 individuals for the statistical analyses. Primary analyses were adjusted for age (by using age as the underlying time scale) and sex (by stratification). In cohort studies, using age as the underlying time scale has been shown to give a more effective control of age compared to using time on study as time scale while adjusting for age (16). In a second model, we additionally adjusted for education as a marker of socioeconomic status (17), and for lifestyle factors that could confound the association between the independent variable and BSI (18, 19). The associations of physical activity and sitting time with BSI risk were additionally adjusted for self-reported chronic mobility impairment, defined as chronic physical illness or injury that impaired daily functioning. Other comorbid conditions, such as cardiovascular diseases, were considered possible mediators of the associations between lifestyle and BSI and were not adjusted for. Additionally, we examined associations separately by age at HUNT2 (< or ≥65

years) and by sex, as well as the combined effect of obesity, smoking and physical inactivity. We examined the combined association of each pair of exposures to explore whether the association of one exposure may differ by levels of other exposures. To assess how changes in smoking habits and weight were associated with BSI, we examined BSI risk among prior smokers categorized by self-reported time since smoking cessation, and we examined the association of weight change from HUNT1 to HUNT2 with BSI risk overall and by categories of BMI at HUNT1. Finally, we examined the associations of obesity and lifestyle factors with the risk of a first-time BSI with each of the most common infecting microorganisms; *Escherichia coli*, *Streptococcus pneumoniae* and *Staphylococcus aureus*. All analyses were performed using Stata version 13.1 (StataCorp).

## Ethics

This study was approved by the Regional Committee for Medical and Health Research Ethics of Central Norway, and all participants signed an informed consent (11).

## Results

During a median follow-up of 14.8 years, amounting to a total of 810,453 person-years of observation for first-time BSI and 816,831 person-years for BSI mortality, 1844 (2.9%) of 64,027 participants experienced at least one episode of BSI, and 396 (0.62%) died from BSI. This corresponds to an incidence rate of 228/100,000 person-years and a mortality rate of 48/100,000 person-years. Baseline characteristics are given in Table 1, and information on infecting microorganisms in Web Table 1. Information on BMI was available for 63,385 (99.0%) participants, on smoking for 63,541 (99.2%), on alcohol intake for 61,637 (96.3%) and on physical activity for 57,250 (89.4%) participants.



Obesity was associated in a dose-dependent manner with increased risk of BSI and BSI mortality. Compared with normal weight participants (BMI 18.5-24.9 kg/m<sup>2</sup>), the age- and sex-adjusted risk of a first-time BSI was 31% (95% CI 14-51%) higher at BMI 30.0-34.9 kg/m<sup>2</sup>, 87% (95% CI 50-135%) higher at BMI 35.0-39.9 kg/m<sup>2</sup>, and 210% (95% CI 117-341%) higher at BMI ≥40.0 kg/m<sup>2</sup> (Table 2). The corresponding increase in risk of BSI mortality was 35% (95% CI -1 to 83%) at BMI 30.0-34.9, 144% (95% CI 55-286%) at BMI 35.0-39.9 kg/m<sup>2</sup>, and 299% (95% CI 93-724%) at BMI ≥40.0 kg/m<sup>2</sup>, compared with normal weight participants (Table 3).

Associations were essentially unchanged after additional adjustment for lifestyle factors and level of education. Using waist circumference as an alternative measure of obesity, we observed associations between wider waist circumference and higher risks of BSI and BSI mortality similar to those observed for BMI (Web Table 2). Low BMI (<18.5 kg/m<sup>2</sup>) was associated with increased risk of BSI in age- and sex-adjusted analysis (75% increased risk, 95% CI 5-193%), but the statistical evidence for this association was weakened in multivariable adjusted analysis (41% increased risk, 95% CI -30 to 185%) (Table 2). For 71% of participants, weight and BMI were also available from the HUNT1 Survey 11 years prior to HUNT2. Most participants remained in their BMI category between the surveys (Web Table 3), but there was a mean weight gain of 4.6 kg (SD 6.7 kg). Weight loss was associated with an increased risk of subsequent BSI, whereas weight gain was not meaningfully associated with increased BSI risk except when weight increased by 10 kg or more. The association of weight change with BSI risk was similar across BMI categories (Web Table 4).

Compared with never-smokers, the age- and sex-adjusted risk of a first time BSI was 51% (95% CI 34-70%) higher among current smokers and 30% (95% CI 16-46%) higher among prior smokers (Table 2). For BSI mortality, there was a 75% (95% CI 34-129%) increase in risk among

current smokers and 30% (95% CI 1-68%) increase in risk among former smokers. After additional adjustment for education, obesity and lifestyle factors, the effect estimates were unchanged for BSI risk and slightly increased for BSI mortality (Table 3). Among prior smokers, the risk of BSI was highest among those who had recently quit smoking (Web Table 5).

Physically inactive participants had a 71% (95% CI 42-107%) increased risk of first-time BSI and 108% (95% CI 37-216%) increased risk BSI mortality compared to those undertaking at least one hour of vigorous exercise per week. In multivariable adjusted analyses, the risk was attenuated to 41% (95% CI 15-74%) and 70% (95% CI 6-174%) higher risks of BSI and BSI mortality, respectively. We found no increased risk of BSI or BSI mortality associated with sitting time (Web Table 2) or alcohol consumption (Table 3). The observed associations did not differ convincingly by age (Web Table 6) or sex (Web Table 7).

The association of each exposure with BSI risk was broadly similar across levels of other exposures (Web Table 8), except that the association of alcohol consumption differed by levels of BMI ( $P_{\text{interaction}}=0.006$ ). Normal-weight participants with high alcohol consumption had increased risk of BSI compared with normal-weight participants with low alcohol consumption (HR 2.26, 95% CI 1.40-3.65), but a similar association was not seen at higher BMI levels.

To assess the combined effect of obesity, smoking and low physical activity, we examined BSI risk among obese ( $\text{BMI} \geq 35.0 \text{ kg/m}^2$ ) current smokers undertaking no or slight physical activity. In these individuals, the age- and sex-adjusted risk of BSI was nearly five-fold increased (HR 4.78, 95% CI 2.44-9.35), and the risk of BSI mortality more than 6-fold increased (HR 6.42, 95% CI 1.83-22.59), compared to normal weight never-smokers with a moderate or high activity level. In crude rates, this corresponded to 414 BSIs and 121 deaths from BSI per 100,000 person-years

in the high-risk group, and 61 BSIs and 13 deaths from BSI per 100,000 person-years in the comparison group.

In analyses of BSI by the most common bacteria (Table 4), obesity was associated in a dose-dependent manner with a strongly increased risk of BSI with *Staphylococcus aureus* (HR 8.24, 95% CI 3.88-17.51 at BMI  $\geq$ 40 kg/m<sup>2</sup>) and also with increased risk of BSI with *Escherichia coli* (HR 2.65, 95% CI 1.48-4.77 at BMI  $\geq$ 40 kg/m<sup>2</sup>) compared with the normal weight participants. In contrast, there was no association between increased BMI and risk of BSI with *Streptococcus pneumoniae*. Smoking was strongly associated with increased risk of *Streptococcus pneumoniae* BSI with a 207% (95% CI 119-330%) and 63% (95% CI 13-135%) risk increase among current and prior smokers respectively, compared with never-smokers. There were similar, but weaker associations between smoking and increased risk of BSI with *Staphylococcus aureus* or *Escherichia coli*. Physical inactivity was associated with a 149% (95% CI 42-335%) increased risk of BSI with *Staphylococcus aureus* and a 53% (95% CI 9-114%) risk of *Escherichia coli* compared with the most active group. There was no association between physical inactivity and BSI with *Streptococcus pneumoniae*.

## Discussion

In this large population-based cohort study, obesity, smoking and physical inactivity were associated with increased risks of BSI and BSI mortality. These results underscore that lifestyle factors may be important for the risk of acquiring invasive infections, a risk which could be attenuated by lifestyle interventions.

The strengths of this study include the population-based design and linkage to prospectively recorded information on BSI from microbiology laboratories at the local and regional hospitals.

Due to the large sample size we could assess both the risk of any BSI, BSI mortality, and the risk of BSI by infecting organism, which expands our understanding of the influence obesity and lifestyle factors exert on the burden of BSI. We did not have clinical information about the course of infection after registration of a positive blood culture; however, review of medical records of patients with *Staphylococcus aureus* and *Streptococcus pneumoniae* BSI in this study population has shown that 98.4% and 98.6% of patients met the 2001 sepsis criteria, respectively (20, 21). Further, we excluded pathogens most commonly associated with contamination, and the majority of positive culture episodes in this material are likely to represent serious infections. Obesity measures were obtained by standardized clinical examination, whereas the other exposure information was self-reported. The self-reported information on vigorous exercise in HUNT has been shown to correlate moderately well with oxygen uptake in healthy adults, whereas the association was less clear for self-report of light exercise (22).

The exposures were measured up to 15 years before the outcome. We would likely have obtained similar results if exposures had been measured closer to the outcome, because our exposure variables represent quite stable traits in most participants. However, the long time span from exposure measurement to outcome may be an advantage if diseases predisposing to BSI may have influenced our exposure variables. Despite the strong association between obesity and increased BSI risk that we observed, weight loss was more strongly than weight gain associated with increased risk of subsequent BSI. Weight loss among middle-aged or elderly individuals is often unintentional and caused by ill health (23), and the association of weight loss with increased BSI risk may represent confounding by ill health.

Although few studies have assessed a comprehensive set of lifestyle factors with a wide range of infections, associations between single lifestyle factors and specific types of infection have been

reported previously. Obesity has been identified as a risk factor for nosocomial infections (24), but the population-level impact of obesity on the risk of severe infections has received less attention (5, 25). Obesity was associated with increased sepsis risk in one cohort study of 30,000 subjects, with a 50% increased risk at BMI 30-39.9 kg/m<sup>2</sup> and two-fold increased risk at BMI  $\geq$ 40 kg/m<sup>2</sup> (8). Abdominal obesity, indicated by increased waist circumference, has been associated with an a 1.7-fold increased risk of sepsis mortality (10). Our results add to that evidence by demonstrating clear dose-response relationships of obesity with BSI and BSI mortality as well as showing that the importance of obesity as a risk factor varied by infecting organism. Possibly, variation in infecting organisms between studies may explain why previous reports of the association between obesity and of pneumonia have reported conflicting results (7, 26).

Smoking is well established as a risk factor for airways infection as well as sepsis (6, 27, 28), whereas the association between smoking and non-airways bacterial infections has received less attention. The 3-fold increased risk of BSI with *Streptococcus pneumoniae* in our study is likely explained by an increased risk of airways infections in smokers. Nonetheless, smoking was also associated with BSI with *Staphylococcus aureus* and *Escherichia coli*, which are less often causes of airway infections. Our finding that smoking is associated with an increased risk of BSI mortality as well as BSI risk is in line with a recent study, where infections were added to conditions that contribute to excess mortality from smoking (29).

Low amounts of exercise was associated with ~ 50% increased risk of sepsis in a population-based cohort (9) and with ~ 50% increased risk of sepsis mortality in a study population of walkers and runners (10); our study extends these findings to a general adult population.

Previous studies have identified an association between excessive alcohol intake and the risk of severe infections such as pneumonia and sepsis (4, 7). The role of a moderate alcohol intake has been less clear (26, 30). We did not detect any association between alcohol intake and risk of BSI in the main analysis, but high alcohol intake was associated with an increased risk of BSI among normal weight participants. We have no explanation why an adverse effect of alcohol on BSI risk should be stronger at normal weight, and this sub-group finding should be interpreted with caution. Few HUNT participants reported a very high alcohol intake, and we had limited possibility to study BSI risk related to excess alcohol intake.

The associations identified in this study could be due to effects of lifestyle factors on the immune system, mechanical factors related to infection susceptibility, or could be mediated through an increased risk of non-communicable diseases. Obesity, smoking and inactivity are associated with increased risk of diseases such as diabetes, cardiovascular disease, cancer and renal disease (31, 32), which all are associated with an increased risk of bacterial infections (4, 30) or mortality from bacterial infections (33). Obesity has been identified as a risk factor for *Staphylococcus aureus* nasal carriage (34), which is associated with a three-fold increased risk of nosocomial *Staphylococcus aureus* BSI (35). Other plausible mechanisms include alterations in immune function. Obesity is associated with alterations in the interaction between metabolic and immune processes (36), smoking exerts an inhibitory effect on the immune system through several pathways (37), and exercise may have immunomodulatory effects (38).

The worldwide obesity prevalence has increased substantially over the last decades (39), and despite reductions in cigarette consumption in some countries, the overall global consumption is still on the rise (40). Coupled with this, infectious diseases are among the most common causes of disease burden globally, particularly in developing countries (41). With combined effects of

obesity, smoking and inactivity increasing the risk of BSI and BSI mortality 5- to 6-fold at the population level, it is likely that facilitating healthy lifestyle choices and halting the obesity epidemic may have a substantial impact on reducing the global risk of invasive bacterial infections.

#### Funding:

This work was supported by a grant from the Liaison Committee between the Central Norway Regional Health Authority and the Norwegian University of Science and Technology.

Conflict of interest: None declared

#### Acknowledgements.

The Nord-Trøndelag Health Study (The HUNT Study) is a collaboration between HUNT Research Centre (Faculty of Medicine, Norwegian University of Science and Technology NTNU), Nord-Trøndelag County Council, Central Norway Health Authority, and the Norwegian Institute of Public Health.

We would like to thank the microbiology departments at Levanger, Namsos and St. Olavs Hospital for providing the data on positive blood cultures, and the Department for Research at Nord-Trøndelag Hospital Trust for assistance with data linkage.

## References

1. Laupland KB, Church DL. Population-based epidemiology and microbiology of community-onset bloodstream infections. *Clin Microbiol Rev* 2014;27:647-664.
2. Goto M, Al-Hasan MN. Overall burden of bloodstream infection and nosocomial bloodstream infection in North America and Europe. *Clin Microbiol Infect* 2013;19:501-509.
3. Cohen J, Vincent JL, Adhikari NK, et al. Sepsis: a roadmap for future research. *Lancet Infect Dis* 2015;15:581-614.
4. Henriksen DP, Pottgard A, Laursen CB, et al. Risk factors for hospitalization due to community-acquired sepsis - a population-based case-control study. *PLoS One* 2015;10:e0124838.
5. Huttunen R, Syrjanen J. Obesity and the risk and outcome of infection. *Int J Obes (Lond)* 2013;37:333-340.
6. Bagaitkar J, Demuth DR, Scott DA. Tobacco use increases susceptibility to bacterial infection. *Tob Induc Dis* 2008;4:12.
7. Almirall J, Bolibar I, Serra-Prat M, et al. New evidence of risk factors for community-acquired pneumonia: a population-based study. *Eur Respir J* 2008;31:1274-1284.
8. Wang HE, Griffin R, Judd S, Shapiro NI, Safford MM. Obesity and risk of sepsis: A population-based cohort study. *Obesity* 2013;21:E762-E769.
9. Wang HE, Baddley J, Griffin RL, et al. Physical inactivity and long-term rates of community-acquired sepsis. *Prev Med* 2014;65:58-64.
10. Williams PT. Inadequate exercise as a risk factor for sepsis mortality. *PloS one* 2013;8:e79344.
11. Holmen J, Midthjell K, Krüger Ø, et al. The Nord-Trøndelag Health Study 1995–97 (HUNT 2): objectives, contents, methods and participation. *Nor Epidemiol* 2003;13:19-32.
12. Krokstad S, Langhammer A, Hveem K, et al. Cohort Profile: the HUNT Study, Norway. *Int J Epidemiol* 2013;42:968-977.
13. World Health Organization. Global Database on Body Mass Index 2006 [cited 2016 May 15]. Available from: [http://apps.who.int/bmi/index.jsp?introPage=intro\\_3.html](http://apps.who.int/bmi/index.jsp?introPage=intro_3.html).
14. World Health Organization. Waist circumference and waist-hip ratio 2011 [cited 2016 May 15]. Available from: [http://apps.who.int/iris/bitstream/10665/44583/1/9789241501491\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/44583/1/9789241501491_eng.pdf).
15. Pien BC, Sundaram P, Raoof N, et al. The clinical and prognostic importance of positive blood cultures in adults. *Am J Med* 2010;123:819-828.
16. Thiebaut AC, Benichou J. Choice of time-scale in Cox's model analysis of epidemiologic cohort data: a simulation study. *Stat Med* 2004;23:3803-3820.
17. Koch K, Sogaard M, Norgaard M, Thomsen RW, Schonheyder HC. Socioeconomic inequalities in risk of hospitalization for community-acquired bacteremia: a Danish population-based case-control study. *Am J Epidemiol* 2014;179:1096-1106.
18. Lahti-Koski M, Pietinen P, Heliovaara M, Vartiainen E. Associations of body mass index and obesity with physical activity, food choices, alcohol intake, and smoking in the 1982-1997 FINRISK Studies. *Am J Clin Nutr* 2002;75:809-817.
19. Stensvold D, Nauman J, Nilsen TI, Wisloff U, Slordahl SA, Vatten L. Even low level of physical activity is associated with reduced mortality among people with metabolic syndrome, a population based study (the HUNT 2 study, Norway). *BMC Med* 2011;9:109.



20. Paulsen J, Mehl A, Askim A, Solligard E, Asvold BO, Damas JK. Epidemiology and outcome of Staphylococcus aureus bloodstream infection and sepsis in a Norwegian county 1996-2011: an observational study. *BMC Infect Dis* 2015;15:116.
21. Askim A, Mehl A, Paulsen J, et al. Epidemiology and outcome of sepsis in adult patients with Streptococcus pneumoniae infection in a Norwegian county 1993-2011: an observational study. *BMC Infect Dis* 2016;16:223.
22. Kurtze N, Rangul V, Hustvedt BE, Flanders WD. Reliability and validity of self-reported physical activity in the Nord-Trondelag Health Study (HUNT 2). *Eur J Epidemiol* 2007;22:379-387.
23. Sahyoun NR, Serdula MK, Galuska DA, Zhang XL, Pamuk ER. The epidemiology of recent involuntary weight loss in the United States population. *J Nutr Health Aging* 2004;8:510-517.
24. Huttunen R, Karpelin M, Syrjanen J. Obesity and nosocomial infections. *J Hosp Infect* 2013;85:8-16.
25. Falagas ME, Kompoti M. Obesity and infection. *Lancet Infect Dis* 2006;6:438-446.
26. Baik I, Curhan GC, Rimm EB, Bendich A, Willett WC, Fawzi WW. A prospective study of age and lifestyle factors in relation to community-acquired pneumonia in US men and women. *Arch Intern Med* 2000;160:3082-3088.
27. Nuorti JP, Butler JC, Farley MM, et al. Cigarette smoking and invasive pneumococcal disease. Active Bacterial Core Surveillance Team. *N Engl J Med* 2000;342:681-689.
28. Arcavi L, Benowitz NL. Cigarette smoking and infection. *Arch Intern Med* 2004;164:2206-2216.
29. Carter BD, Abnet CC, Feskanich D, et al. Smoking and mortality--beyond established causes. *N Engl J Med* 2015;372:631-640.
30. Wang HE, Shapiro NI, Griffin R, Safford MM, Judd S, Howard G. Chronic medical conditions and risk of sepsis. *PLoS One* 2012;7:e48307.
31. Must A, Spadano J, Coakley EH, Field AE, Colditz G, Dietz WH. The disease burden associated with overweight and obesity. *Jama* 1999;282:1523-1529.
32. World Health Organization. Global status report on non-communicable diseases 2010: WHO; 2011 [cited 2016 May 15]. Available from: [http://apps.who.int/iris/bitstream/10665/44579/1/9789240686458\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/44579/1/9789240686458_eng.pdf).
33. Seshasai SR, Kaptoge S, Thompson A, et al. Diabetes mellitus, fasting glucose, and risk of cause-specific death. *N Engl J Med* 2011;364:829-841.
34. Gorwitz RJ, Kruszon-Moran D, McAllister SK, et al. Changes in the prevalence of nasal colonization with Staphylococcus aureus in the United States, 2001-2004. *J Infect Dis* 2008;197:1226-1234.
35. Wertheim HF, Vos MC, Ott A, et al. Risk and outcome of nosocomial Staphylococcus aureus bacteraemia in nasal carriers versus non-carriers. *Lancet* 2004;364:703-705.
36. Chawla A, Nguyen KD, Goh YP. Macrophage-mediated inflammation in metabolic disease. *Nat Rev Immunol* 2011;11:738-749.
37. Mehta H, Nazzal K, Sadikot RT. Cigarette smoking and innate immunity. *Inflamm Res* 2008;57:497-503.
38. Walsh NP, Gleeson M, Shephard RJ, et al. Position statement. Part one: Immune function and exercise. *Exerc Immunol Rev* 2011;17:6-63.
39. NCD Risk Factor Collaboration. Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19· 2 million participants. *The Lancet* 2016;387:1377-1396.

40. The Tobacco Atlas. Cigarette use globally 2015 [cited 2016 May 15, 2016]. Available from: <http://www.tobaccoatlas.org/topic/cigarette-use-globally/>.
41. Murray CJ, Barber RM, Foreman KJ, et al. Global, regional, and national disability-adjusted life years (DALYs) for 306 diseases and injuries and healthy life expectancy (HALE) for 188 countries, 1990-2013: quantifying the epidemiological transition. *Lancet* 2015;386:2145-2191.

**Table 1 Baseline characteristics<sup>a</sup> of 64,027 participants in the HUNT Study, Norway 1995-97**

<b>Variable</b>	<b>Total population (n=64,027)</b>	<b>BSI during follow-up (n=1844)</b>
<b>Age, median (IQR)</b>	48.6 (36.4- 63.6)	67.7 (55.5-74.7)
<b>Female sex, No. (%)</b>	34,079 (53.2)	898 (48.7)
<b>Highest education achieved, No. (%)</b>		
<b>≤ 9 years</b>	22,170 (36.5)	952 (57.7)
<b>10-12 years</b>	26,452 (43.6)	532 (32.2)
<b>&gt;12 years</b>	12,062 (19.9)	167 (10.1)
<b>Chronic mobility impairment, No. (%)</b>		
<b>None</b>	55,084 (86.0)	1414 (76.7)
<b>Slight</b>	3,661 (5.7)	134 (7.3)
<b>Moderate</b>	2,848 (4.5)	145 (7.9)
<b>Severe</b>	2,434 (3.8)	151 (8.2)
<b>BMI kg/m<sup>2</sup>, mean (SD)</b>	26.4 (4.1)	27.7 (4.6)
<b>Waist circumference cm, mean (SD)</b>	86.4 (11.7)	91.7 (12.5)
<b>Smoking, No. (%)</b>		
<b>Never</b>	28,880 (45.5)	703 (38.6)
<b>Prior</b>	16,505 (26.0)	626 (34.3)
<b>Current</b>	18,156 (28.6)	494 (27.1)
<b>Alcohol intake, No. (%)</b>		
<b>&lt;1 glass/two weeks</b>	22,787 (37.0)	906 (52.4)
<b>1-7glasses/two weeks</b>	30,395 (49.3)	665 (38.5)
<b>8-14 glasses/two weeks</b>	6,531 (10.6)	115 (6.7)
<b>≥15 glasses/two weeks</b>	1,924 (3.12)	42 (2.4)
<b>Physical activity<sup>b</sup>, No. (%)</b>		

<b>None</b>	4,642 (8.11)	207 (14.7)
<b>Slight</b>	17,816 (31.1)	497 (35.4)
<b>Moderate</b>	19,344 (33.8)	447 (31.8)
<b>High</b>	15,448 (27.0)	253 (18.0)
<b>Sitting time, No. (%)</b>		
<b>≤ 4 hours/day</b>	15,416 (31.5)	447 (33.5)
<b>5-7 hours/day</b>	15,668 (32.0)	444 (33.3)
<b>≥8 hours/day</b>	17,927 (36.6)	443 (33.2)

---

BSI: Bloodstream infection, IQR: Interquartile range, SD: Standard deviation

- a) The % refers to the proportion within the total population or those with BSI, respectively.
- b) None: No light or vigorous activity; Slight: <3 hours light activity/week and no vigorous activity; Moderate: ≥3 hours light activity or <1 hour vigorous activity/week; High: ≥1 hour vigorous activity/week.

**Table 2 Associations of BMI and lifestyle factors with the risk of bloodstream infection among 64,027 participants in the HUNT Study, Norway 1995-2011**

Lifestyle variable	Age- and sex-adjusted					Age-, sex-, education- and lifestyle-adjusted <sup>a</sup>				
	Person-years at risk	No. BSI	Incidence rate per 100 000 person- years	HR	95% CI	Person-years at risk	No. BSI	Incidence rate per 100 000 person- years	HR	95% CI
<b>BMI, kg/m<sup>2</sup></b>										
<18.5	5,270	15	285	<b>1.75</b>	<b>1.05-2.93</b>	4,402	8	182	<b>1.41</b>	<b>0.70-2.85</b>
18.5-24.9	321,042	523	163	<b>1.00</b>	<b>Reference</b>	288,423	388	135	<b>1.00</b>	<b>Reference</b>
25.0-29.9	348,025	817	235	<b>1.04</b>	<b>0.93-1.17</b>	302,915	568	188	<b>1.02</b>	<b>0.90-1.17</b>
30.0-34.9	103,982	339	326	<b>1.31</b>	<b>1.14-1.51</b>	85,412	234	274	<b>1.38</b>	<b>1.17-1.63</b>
35.0-39.9	21,404	90	420	<b>1.87</b>	<b>1.50-2.35</b>	17,137	58	338	<b>1.77</b>	<b>1.34-2.35</b>
≥40.0	5,196	33	635	<b>3.10</b>	<b>2.17-4.41</b>	4,181	23	550	<b>3.14</b>	<b>2.05-4.81</b>
<b>Smoking</b>										
Never	367,417	703	191	<b>1.00</b>	<b>Reference</b>	317,988	451	142	<b>1.00</b>	<b>Reference</b>
Prior	206,261	626	304	<b>1.30</b>	<b>1.16-1.46</b>	180,356	464	257	<b>1.31</b>	<b>1.14-1.51</b>
Current	231,577	494	213	<b>1.51</b>	<b>1.34-1.70</b>	203,405	364	179	<b>1.53</b>	<b>1.32-1.78</b>

**Physical activity level<sup>b</sup>**

None	53,199	207	389	<b>1.71</b>	<b>1.42-2.07</b>	48,098	170	353	<b>1.41</b>	<b>1.15-1.74</b>
Slight	228,970	497	217	<b>1.18</b>	<b>1.01-1.38</b>	216,129	455	210	<b>1.10</b>	<b>0.93-1.30</b>
Moderate	250,954	447	178	<b>1.00</b>	<b>0.85-1.17</b>	239,738	417	174	<b>0.97</b>	<b>0.83-1.15</b>
High	204,086	253	124	<b>1.00</b>	<b>Reference</b>	197,784	237	120	<b>1.00</b>	<b>Reference</b>

**Alcohol intake**

<1 glass/two weeks	272,422	906	333	<b>1.01</b>	<b>0.91-1.13</b>	219,150	572	261	<b>0.97</b>	<b>0.85-1.10</b>
1-7glasses/two weeks	401,370	665	166	<b>1.00</b>	<b>Reference</b>	377,542	569	151	<b>1.00</b>	<b>Reference</b>
8-14 glasses/two weeks	85,625	115	134	<b>0.89</b>	<b>0.73-1.09</b>	81,741	100	122	<b>0.86</b>	<b>0.69-1.06</b>
≥15 glasses/two weeks	24,312	42	173	<b>1.24</b>	<b>0.91-1.70</b>	23,316	38	163	<b>1.21</b>	<b>0.87-1.69</b>

---

BSI, Bloodstream infection; HR, Hazard ratio; CI, Confidence interval; BMI, Body mass index

a) Adjusted for age, sex, education, smoking, BMI, physical activity and alcohol intake. The association of physical activity with BSI risk was additionally adjusted for chronic mobility impairment

b) None: No light or vigorous activity; Slight: <3 hours light activity/week and no vigorous activity; Moderate: ≥3 hours light activity or <1 hour vigorous activity/week; High: ≥1 hour vigorous activity/week.

**Table 3 Associations of BMI and lifestyle factors with mortality from bloodstream infection<sup>a</sup> among 64,027 participants in the HUNT Study, Norway**

**1995-2011**

Lifestyle variable	Age-and sex- adjusted					Age-, sex-, education-, and lifestyle-adjusted <sup>b</sup>				
	Person-years	No.	Mortality	HR	95% CI	Person-years	No.	Mortality	HR	95% CI
	at risk	BSI	rate per			at risk	BSI	rate per		
		deaths	100 000				deaths	100 000		
			person-years					person-years		
<b>BMI, kg/m<sup>2</sup></b>										
<18.5	5,323	2	38	<b>1.20</b>	<b>0.30-4.88</b>	4,445	1	23	<b>0.96</b>	<b>0.13-6.94</b>
18.5-24.9	323,165	101	31	<b>1.00</b>	<b>Reference</b>	290,075	71	24	<b>1.00</b>	<b>Reference</b>
25.0-29.9	350,823	183	52	<b>1.12</b>	<b>0.88-1.43</b>	304,978	117	38	<b>1.09</b>	<b>0.81-1.47</b>
30.0-34.9	105,218	72	68	<b>1.35</b>	<b>0.99-1.83</b>	86,296	48	56	<b>1.49</b>	<b>1.02-2.17</b>
35.0-39.9	21,806	23	105	<b>2.44</b>	<b>1.55-3.86</b>	17,420	15	86	<b>2.56</b>	<b>1.45-4.53</b>
≥40.0	5,316	8	150	<b>3.99</b>	<b>1.93-8.24</b>	4,272	7	164	<b>5.67</b>	<b>2.56-12.58</b>
<b>Smoking</b>										
Never	370,124	150	41	<b>1.00</b>	<b>Reference</b>	320,130	82	26	<b>1.00</b>	<b>Reference</b>
Prior	208,458	137	66	<b>1.30</b>	<b>1.01-1.68</b>	182,284	98	54	<b>1.47</b>	<b>1.06-2.04</b>
Current	233,420	106	45	<b>1.75</b>	<b>1.34-2.29</b>	205,073	79	39	<b>2.23</b>	<b>1.59-3.14</b>

**Physical activity level<sup>c</sup>**

None	53,918	51	95	<b>2.08</b>	<b>1.37-3.16</b>	48,751	40	82	<b>1.70</b>	<b>1.06-2.74</b>
Slight	230,884	107	46	<b>1.39</b>	<b>0.97-2.00</b>	218,140	98	45	<b>1.42</b>	<b>0.96-2.11</b>
Moderate	252,722	94	37	<b>1.14</b>	<b>0.79-1.64</b>	241,663	85	35	<b>1.19</b>	<b>0.80-1.77</b>
High	205,083	43	21	<b>1.00</b>	<b>Reference</b>	198,931	36	18	<b>1.00</b>	<b>Reference</b>

**Alcohol intake**

<1 glass / two weeks	275,556	206	75	<b>0.96</b>	<b>0.76-1.22</b>	221,482	122	55	<b>0.89</b>	<b>0.67-1.19</b>
1-7 glasses/two weeks	404,120	130	32	<b>1.00</b>	<b>Reference</b>	380,317	108	28	<b>1.00</b>	<b>Reference</b>
8-14 glasses/two weeks	86,088	26	30	<b>1.05</b>	<b>0.69-1.60</b>	82,233	22	27	<b>0.98</b>	<b>0.62-1.56</b>
≥15 glasses/two weeks	24,451	9	37	<b>1.35</b>	<b>0.68-2.66</b>	23,453	7	30	<b>1.13</b>	<b>0.52-2.45</b>

---

BSI, Bloodstream infection; HR, Hazard ratio; CI, Confidence interval; BMI, Body mass index

a) Defined as death within 30 days after detection of a BSI

b) Adjusted for age, sex, education, smoking, BMI, physical activity and alcohol intake. The association of physical activity with risk of BSI mortality was additionally adjusted for chronic mobility impairment.

c) None: No light or vigorous activity; Slight: <3 hours light activity/week and no vigorous activity; Moderate: ≥3 hours light activity or <1 hour vigorous activity/week; High: ≥1 hour vigorous activity/week.



**Table 4 Age- and sex-adjusted hazard ratios of bloodstream infection by the 3 most common microorganisms among 64,027 participants in the HUNT**

**Study, Norway 1995-2011**

Lifestyle variable	<i>Escherichia coli</i>			<i>Staphylococcus aureus</i>			<i>Streptococcus pneumoniae</i>		
	No. BSI /Person-years at risk	HR	95% CI	No. BSI /Person- years at risk	HR	95% CI	No. BSI /Person-years at risk	HR	95% CI
<b>BMI, kg/m<sup>2</sup></b>									
<18.5	6/5,270	<b>1.78</b>	<b>0.79-4.02</b>	1/5,270	<b>1.16</b>	<b>0.16-8.40</b>	4/5,270	<b>2.97</b>	<b>1.08-8.12</b>
18.5-24.9	192/321,042	<b>1.00</b>	<b>Reference</b>	56/321,042	<b>1.00</b>	<b>Reference</b>	82/321,042	<b>1.00</b>	<b>Reference</b>
25.0-29.9	296/348,025	<b>1.04</b>	<b>0.87-1.25</b>	86/348,025	<b>0.99</b>	<b>0.70-1.39</b>	110/348,025	<b>0.94</b>	<b>0.70-1.25</b>
30.0-34.9	132/103,982	<b>1.32</b>	<b>1.06-1.66</b>	47/103,982	<b>1.75</b>	<b>1.18-2.59</b>	28/103,982	<b>0.73</b>	<b>0.47-1.13</b>
35.0-39.9	38/21,404	<b>1.95</b>	<b>1.37-2.77</b>	10/21,404	<b>2.15</b>	<b>1.09-4.25</b>	6/21,404	<b>0.83</b>	<b>0.36-1.91</b>
≥40.0	12/5,196	<b>2.65</b>	<b>1.48-4.77</b>	8/5,196	<b>8.24</b>	<b>3.88-17.51</b>	1/5,196	<b>0.61</b>	<b>0.09-4.42</b>
<b>Smoking</b>									
Never	288/367,417	<b>1.00</b>	<b>Reference</b>	76/367,417	<b>1.00</b>	<b>Reference</b>	63/367,417	<b>1.00</b>	<b>Reference</b>
Prior	230/206,261	<b>1.36</b>	<b>1.12-1.64</b>	73/206,261	<b>1.24</b>	<b>0.88-1.76</b>	70/206,261	<b>1.63</b>	<b>1.13-2.35</b>
Current	160/231,578	<b>1.35</b>	<b>1.10-1.66</b>	60/231,578	<b>1.59</b>	<b>1.12-2.27</b>	96/231,578	<b>3.07</b>	<b>2.19-4.30</b>

**Physical****activity level<sup>a</sup>**

None	69/53,199	<b>1.53</b>	<b>1.09-2.14</b>	27/53,199	<b>2.49</b>	<b>1.42-4.35</b>	19/53,199	<b>1.24</b>	<b>0.70-2.20</b>
Slight	172/228,970	<b>1.13</b>	<b>0.86-1.48</b>	62/228,970	<b>1.64</b>	<b>1.02-2.64</b>	63/228,970	<b>1.10</b>	<b>0.72-1.68</b>
Moderate	170/250,954	<b>1.10</b>	<b>0.84-1.44</b>	49/250,954	<b>1.17</b>	<b>0.72-1.91</b>	68/250,954	<b>1.12</b>	<b>0.74-1.69</b>
High	79/204,086	<b>1.00</b>	<b>Reference</b>	25/204,086	<b>1.00</b>	<b>Reference</b>	36/204,086	<b>1.00</b>	<b>Reference</b>

---

BSI, Bloodstream infection; HR, Hazard ratio; CI, Confidence interval

a) None: No light or vigorous activity; Slight: <3 hours light activity/week and no vigorous activity; Moderate: ≥3 hours light activity or <1 hour vigorous activity/week; High: ≥1 hour vigorous activity/week.