

PROCEEDINGS B

On being the right size: increased body size is associated with reduced telomere length under natural conditons

Journal:	<i>Proceedings B</i>
Manuscript ID	Draft
Article Type:	Research
Date Submitted by the Author:	n/a
Complete List of Authors:	Ringsby, Thor Harald; Norwegian University for Technology and Science, Centre for Biodiversity Dynamics Jensen, Henrik; Norwegian University for Technology and Science, Centre for Biodiversity Dynamics Pärn, Henrik; Norwegian University for Technology and Science, Centre for Biodiversity Dynamics Kvalnes, Thomas; Norwegian University for Technology and Science, Centre for Biodiversity Dynamics Boner, Winnie; University of Glasgow, Institute of Biodiversity, Animal Health & Comparative Medicine, College of Medical, Veterinary and Life Sciences Gillespie, Robert; University of Glasgow, Institute of Biodiversity, Animal Health & Comparative Medicine, College of Medical, Veterinary and Life Sciences Holand, Håkon; Norwegian University for Technology and Science, Centre for Biodiversity Dynamics Hagen, Ingerid; Norwegian University for Technology and Science, Centre for Biodiversity Dynamics Rønning, Bernt; Norwegian University for Technology and Science, Centre for Biodiversity Dynamics Sæther, Bernt-Erik; Norwegian University for Technology and Science, Centre for Biodiversity Dynamics Monaghan, Pat; University of Glasgow, Institute of Biodiversity, Animal Health & Comparative Medicine, College of Medical, Veterinary and Life Sciences
Subject:	Ecology < BIOLOGY, Evolution < BIOLOGY, Genetics < BIOLOGY
Keywords:	body size, telomere length, artificial selection experiment, <i>Passer domesticus</i>
Proceedings B category:	Evolutionary Biology

On being the right size: increased body size is associated with reduced telomere length under natural conditions

Thor H. Ringsby,¹ Henrik Jensen,¹ Henrik Pärn,¹ Thomas Kvalnes,¹ Winnie Boner,² Robert Gillespie,² Håkon Holand,¹ Ingerid Julie Hagen¹, Bernt Rønning,¹ Bernt - E. Sæther,¹ Pat Monaghan²

¹ Centre for Biodiversity Dynamics, Department of Biology, Norwegian University for Science and Technology, NO-7491 Trondheim, Norway. ² Institute of Biodiversity, Animal Health & Comparative Medicine, College of Medical, Veterinary and Life Sciences, Graham Kerr Building, University of Glasgow, Glasgow G12 8QQ UK.

Abstract

Evolution of body size is likely to involve trade-offs between body size, growth rate and longevity. Within species, larger body size is associated with faster growth and ageing, and reduced longevity, but the cellular processes driving these relationships are poorly understood. One mechanism that might play a key role in determining optimal body size is the relationship between body size and telomere dynamics. However, we know little about how telomere length is affected when selection for larger size is imposed in natural populations. We report here on the relationship between structural body size and telomere length in wild house sparrows at the beginning and end of a selection regime for larger parent size that was imposed for four years in an isolated population of house sparrows. A negative

relationship between fledgling size and telomere length was present at the start of the selection; this was extended when fledgling size increased under the selection regime, demonstrating a persistent co-variance between structural size and telomere length. Changes in telomere dynamics, either as a correlated trait or a consequence of larger size, could reduce potential longevity and the consequent trade-offs could thereby play an important role in the evolution of optimal body size.

Key words: body size, telomere length, life history trade-off, longevity, *Passer domesticus*, selection experiment.

1. Introduction

Understanding the mechanisms that shape life history strategies is at the heart of evolutionary ecology. Much of our current theoretical and empirical research is based around the premise that the balance of costs and benefits of different life history components results in natural selection producing the optimal compromise [1, 2]. Hence we predict that trade-offs will occur among growth, self-maintenance, and reproduction. Differences in the optimal resolution of these trade-offs through individual differences in resource acquisition and allocation [3] underlies the individual variation that we see in growth rate, body size, reproductive rate and longevity at the population level.

It is well recognised that in practice such trade-offs are difficult to demonstrate [1, 3-6]. Considerable effort has been devoted to investigating how variation in reproductive investment influences other life history traits [7-9]. The relationships among growth, body

size and longevity have received much less attention [10], but are coming more to the fore as the importance of understanding intra-and inter-specific variation in the rate of ageing is recognised as an important question in modern biology [6, 11]. A paradox that has puzzled evolutionary ecologists and biogerontologists is the opposing patterns that frequently are observed in the relationship between body size and longevity among and within species. Across species, there is a strong positive relationship between body size and maximal lifespan; big species tend to live longer than small ones. Within species however, the opposite relationship has often been found: larger individuals have shorter lives than conspecifics of smaller size, and this has been demonstrated in the wild and in captivity [e.g. 12, 13-18]. The within species variation in lifespan is also exemplified in species subjected to artificial selection to produce variation in body size such as the domestic dog (*Canis lupus familiaris*). Great variation in body size is present in dogs; larger dog breeds have shorter lifespans, and this has recently been shown to occur because they age faster [13]. Similarly, when laboratory species such as mice and rats are selected or genetically engineered for larger or smaller size, lifespan is again reduced or increased [17, 19, 20].

We do not currently understand how the long term costs associated with larger size are incurred. There are a number of potential, non-exclusive routes whereby this could occur. A candidate process is reduced telomere length due to faster cell division rates and/or telomere attrition rate [5, 6, 10, 21]. Given that telomere length is linked to organismal senescence [22-24], and that average telomere length in tissues at the end of growth is predictive of lifespan [25, 26], this may point to a mechanistic process that could explain the lifespan penalty associated with larger size. This could be the case in dogs, where lifespan is negatively correlated with body size among breeds and telomere length in large dogs breeds is reported to be shorter than in the smaller breeds. However, the extreme variation created in this species (100 fold) could have created aberrant patterns that are not evident under natural

conditions [27]. A key question then is whether telomere length covaries with intraspecific variation in body size in natural conditions, in the absence of such extreme size variation.

We used samples that had been collected during an experiment conducted in a population of wild house sparrows (*Passer domesticus*) in northern Norway, in which a selection pressure for larger body size was experimentally imposed on the breeding population over a four year period, which clearly increased the size of both adults and their offspring (see details in methods). We examined the extent to which telomere length and body size were related in fledglings, and, importantly, whether any relationship was maintained when fledgling body size was increased under the selection regime.

2. Material and methods

(a) Study site

The study was conducted in the house sparrow population on the island Leka situated ca 3 km from the mainland in the Nord-Trøndelag county in northern Norway (65°06'N, 11°038'E, see map in Hagen, Billing [28]) and was initiated in 2002. Leka covers 57 km² and is dominated by cultivated areas (mainly silage production for dairy cattle), marsh, deciduous forests and mountain and heath areas.

Artificial selection experiment

As part of a large study on body size variation, each year the whole adult house sparrow population at Leka was captured by mist netting in February and held temporarily in a barn for up to 12 days, which was the time it took to capture all the individuals on the island. The house sparrows were given *ad libitum* water and food, which included concentrate feeds for

cattle, sunflower seeds and bread. The temperature in the room was kept at 10-12°C, which is the normal temperature inside cowsheds in this area during winter. All the sparrows were measured for morphological traits as well as blood sampled before they were released into the barn. We used tarsus length to indicate structural body size in this study as tarsus size has been shown to be a stable indicator of body size and is relatively resistant to local variations in environmental conditions during growth and in adulthood in house sparrows [29]. After practically all (> 95%) individuals at Leka had been captured, we separately estimated the mean and the standard deviation of tarsus lengths within males and females. Based on this information, we determined the cut-off tarsus size where all individuals with shorter tarsus lengths were removed and released in distant populations of house sparrows away from Leka [30]. This artificial selection procedure was conducted annually from 2002 until 2005 (i.e. 4 selection events). Thus, each winter we removed ca 50-60 % of the individuals with the smallest tarsus lengths. As a consequence of this annual artificial selection protocol at Leka, which allowed only the larger individuals to breed, tarsus length of males produced at Leka progressively increased during the 4 years of selection. This is indicated in a generalized linear mixed model (brood identity as random effect, fitted with the glmer package [31] R Core Team [32]) where the average size among fledglings that were later recaptured as juveniles or adults showed that their full-grown tarsus lengths increased from 2002 (19.473 mm, *s.e.* = 0.211) to 2005 (20.360 mm, *s.e.* = 0.369), increasing annually compared to the year 2002, which is represented by the intercept ($\beta_{Intercept} = 19.473, s.e. = 0.211, t = 92.3$, $\beta_{2003} = 0.633, s.e. = 0.308, t = 2.06$, $\beta_{2004} = 1.227, s.e. = 0.459, t = 2.67$, $\beta_{2005} = 0.887, s.e. = 0.425, t = 2.09, N = 28$).

In the present study we examined telomere lengths from red blood cell samples collected from known age fledglings from whom body size measures (tarsus length) were also taken. This was done for a sample of fledglings produced in the first year of selection

(2002) and a sample produced in the last year of selection (2005). We could only use samples from which DNA of sufficient quality and quantity could be extracted, and for which tarsus length data had been collected. The sex of the birds was not known at the time of sample collection. We confined our analysis to males, for which we had the larger sample presumably for stochastic reasons, since the pattern of growth in body size differs between the sexes in house sparrows [33], and the sample size for females was too small to enable investigation of the differences between the sexes.

(b) Sampling procedures

In the study region, the house sparrows lay 1-3 clutches during the breeding season, early May-to mid - August [34]. During this period we searched for nests and recorded all that were accessible. Typically, the nests were located under the ceiling in barns and cowsheds at dairy farms. Each nest was visited about once a week and the number of eggs, the number of nestlings and their age were recorded. Fledglings were measured for tarsus length between 9-15 days of age, when we also collected a small sample of blood ($25\mu\text{l}$) used here for , genetic sexing and analyses of telomere lengths ([for more details on fieldwork procedures see 35, 36]). The age range at which the 44 fledglings in this study were sampled was small, covering a six day period.

(c) Molecular methods

Until DNA extraction, the blood was preserved in 96% ethanol at room temperature. The blood was lysed in 60 μl Lairds buffer (Ausubel et al. 1989), with 90 μg proteinase K (Sigma

Aldrich), and incubated at 50°C for 3 hours. The genomic DNA was extracted from the lysate using the ReliaPrep Large Volume HT gDNA Isolation System (Promega), automated on a Biomek NXp robot (Beckman Coulter) following the manufacturer's recommendations, the only exception being elution of DNA in 25 mM Tris HCl (pH 8). The DNA concentrations were measured using a Flurostar Omega scanner (MBG Labtech). Samples with DNA concentrations above 60 ng/μl were normalized to a concentration of 50 ng/μl with 25 mM Tris HCl (pH 8). For further details see [28].

The sex of fledglings was determined by molecular genetic methods using DNA extracted from the blood samples. A copy of the CHD-gene is located on both the avian Z and W chromosomes. Because the two copies vary in size in house sparrows, CHD-Z is 370 bp and CHD-W is 388 bp, they can be separated by electrophoresis. We amplified the CHD-gene by Polymerase Chain Reaction (PCR) using P2 and P8 primers [37]. PCR amplification was carried out in 10μl reactions containing 5 μl 2x QIAGEN Multiplex PCR Master Mix (QIAGEN), 3μl extracted DNA (ca. 20 ng/μl) and 0.125 μM of each CHD primer. The CHD primers were included in a 2 μl multiplex primer mix which also contained the seven microsatellite loci Ase18 and Pdo10 [38], Pdoμ1 and Pdoμ3 [39], Pdoμ5[40], Pdo33 and Pdo40[41]. We used a touchdown PCR profile: 94°C for 15 minutes; 12 cycles of 94°C for 30 seconds, 62°C for 1 minute 30 seconds, and 72°C for 1 minute, where the annealing temperature was dropped 1°C each cycle; 19 cycles of 94°C for 30 seconds, 50°C for 1 minute 30 seconds, and 72°C for 1 minute; 60°C for 5 minutes. The PCR product was kept at 4°C until 1 μl was taken out and mixed with 0.5 μl GeneScan 600 LIZ size standard (Applied Biosystems) and 10 μl HiDi Formamide (Applied Biosystems) prior to capillary electrophoresis on an ABI 3130xl Genetic Analyzer (Applied Biosystems). The P8 primer was fluorescently labeled using NED (Applied Biosystems) enabling us to score Z- and W-copies of the CHD-gene in the GeneMapper 4.0 software (Applied Biosystems). In birds,

males are the homogametic sex and were thus identified by being homozygous for the CHD-gene (i.e. carrying two short copies).

The concentration and quality of DNA samples were assessed using a Nanodrop-8000 Spectrophotometer. DNA samples were stored at -20°C. Relative telomere lengths were measured using the qPCR method. While this gives a relative rather than an absolute measure, it is widely used in telomere studies and very suitable for the very small blood samples we had available[42]. This method also includes any interstitial repeats of the telomere sequence, the level of which is not known for this species. We therefore further examined a small sample of fledglings for which we had sufficient DNA, using both the standard southern blot TRF method and in-gel TFR method[42] ; telomere length was in the 15-20 kb range, and no substantial levels of interstitial repeats were observed when the gels were examined. Telomere measurements were made using the qPCR method as described by Criscuolo et al. [43] with the following modifications. We verified that Gapdh was a suitable control gene in this species by carrying out a melt curve analysis and checking that the dissociation curve was a single peak. As the melting temperature of the product is sequence dependent, a single peak indicates amplification of a single product. The amplification products, from a random selection of samples, were also run on a gel to confirm that the amplification was of a single product. DNA samples (10ng) were assayed using the Absolute blue qPCR SYBR green Low Rox master mix (Thermo scientific) with telomere primers (Tel1b and Tel2b) at a final concentration of 500nM and Gapdh primers at a final concentration of 200nM. Forward Gapdh primer was 5'- GAG GTG CTG CTC AGA ACA TTA T-3' and reverse Gapdh primer was 5'- ACG GAA AGC CAT TCC AGT AAG-3'. The telomere thermal profile was 15 minutes at 95°C, followed by 27 cycles of 15 seconds at 95°C, 30 seconds at 58°C, 30 seconds at 72°C. The Gapdh thermal profile was 15 minutes at 95°C, followed by 40 cycles of 15 seconds at 95°C, 15 seconds at 60°C. Both assays were

followed by melt curve analysis of (58 – 95°C 1oc/5s ramp). The reference sample was serially diluted (from 40 to 2.5 ng/well) to produce a standard curve for each plate. This was used to calculate plate efficiencies, all of which fell well within the acceptable range (i.e. 100±10%) and only samples that fell within the bounds of the standard curve were included. As described in Criscuolo et al. [43], relative telomere measurements were calculated using the $\Delta\Delta C_t$ method. This provides a ratio of the abundance of the telomeric sequence to the abundance of the reference single copy gene (henceforth, T/S ratio). Since the qPCR efficiencies were tightly controlled, these data were highly correlated with the values obtained when the calculations were done using the Pfaffl method, which corrects for differences in plate efficiencies ($r_{Pearson} = 0.998$).

(d) Statistical analyses

We used linear mixed models assuming Gaussian residual distribution, where all models included the identity of the nest as random factor in order to account for any non independence of fledglings from the the same brood (R version 3.0.2. [32], package lme4, function lmer [31]). The dataset comprised 24 and 20 male fledglings in 2002 and 2005, respectively, distributed among 21 nests, with structural body size being represented by tarsus length. The significance of the explanatory variables were evaluated by likelihood-ratio tests. We tested whether the average fledgling tarsus length changed between 2002 and 2005 and accounted for any effect of the slight variation in fledgling age at sampling by retaining this variable in all models. Likewise we tested whether the average telomere length differed between 2002 and 2005, respectively. We also tested whether a linear relationship between telomere length (response variable) and tarsus length was present when the samples from 2002 and 2005 were pooled and whether a similar linear relationship existed within each year

by testing the significance of the interaction term (year:tarsus length). Also here we included the fledgling age at sampling in the model. Age was not significantly correlated with tarsus length in the small age range covered by this data set ($r_{Pearson} = 0.05$, $p > 0.1$; $n = 44$).

Assumptions of all statistical models were evaluated visually by diagnostic plots.

The growth of the fledglings was not monitored during the selection experiment, since they were measured only once for logistic reasons and to minimize disturbance. However, to give an insight into how tarsus measured as a fledgling related to adult size, we used data for fledglings that were recaptured later as juveniles and/or adults and from which the full-grown tarsus length was then obtained. The sample size for this was small (10 individuals from the 2002 cohort and 5 from the 2005 cohort, distributed over 10 nests). We examined the relationship between tarsus size (standardized to the age of 10 days to account for variation in the age of measurement of fledglings), and tarsus size when fully grown by using mixed generalized linear model (package lme4, function lmer, accounting for nest identity as random factor). To obtain the standardized size at the age of 10 days, we modeled the relationship between age and tarsus length (as response variable) by a quadratic model based on a large sample ($n = 10517$) from our main study area in northern Norway. From this model we extracted the predicted tarsus length at the age of 10 days for each individual.

3. Results

(a) Body size variation

During the period over which selection for larger structural size in the breeding population was imposed, the average tarsus length in the fledglings increased significantly (by 4.3%) in the 2005 cohort as compared to the 2002 cohort ($\beta_{\text{tarsus 2005}} = 0.823$, s.e. = 0.243; $\chi^2 = 7.887$, $p = 0.005$; $n = 44$; figure 1a). This is because (as can be seen in figure 3a) the upper range of

tarsus lengths in fledglings was extended as expected. Our capacity to examine whether the increase in fledgling tarsus length in 2005 also resulted in an increase in full grown tarsus length compared to in 2002 was limited, since we only had data for the 15 birds measured both as fledglings and as full grown adults. Nonetheless, figure 2 shows that tarsus length standardised to 10 days of age was predictive of adult tarsus size ($\beta = 0.593$, $s.e. = 0.076$; $\chi^2 = 27.184$, $p < 0.001$; $n = 15$; figure 2). As can be seen in this figure, the larger tarsus sizes in fledglings produced as a consequence of the selection regime gave rise to larger adults.

(b) Size and telomere length

This increase in fledgling tarsus length was accompanied by a significant reduction in average telomere lengths from 2002 to 2005 ($\beta_{\text{telomere length 2005}} = -0.163$, $s.e. = 0.070$, $\chi^2 = 4.640$, $p = 0.0312$, $n = 44$, figure 1b). We examined the relationship between tarsus length and telomere length within the two cohorts, in order to see if there was a significant negative relationship between size and telomere length, whether this was evident in 2002 at the start of the selection regime, and whether the relationship changed when structural body size had increased significantly at the end of the selection regime. Figure 3 shows these relationships. An overall negative relationship between tarsus length and telomere length was present for fledglings when the samples from 2002 and 2005 were pooled ($\beta_{\text{tarsus length}} = -0.105$, $s.e. = 0.030$; $\chi^2 = 8.710$, $p = 0.003$; $n = 44$) as seen in figure 3. Fledglings with larger structural size had shorter telomeres. The slopes of the negative relationships between telomere length and tarsus length did not differ significantly between 2002 and 2005, as the interaction effect (year:tarsus length) was not significant ($p > 0.1$). Thus, structural size and telomere length were negatively related, and the the extension of the upper range of tarsus length in fledglings

extended, but did not change, this relationship. All the models above controlled for the actual age of the fledglings at measurement which had no significant effect ($p > 0.05$) in all models.

4. Discussion

In an isolated population of house sparrows in northern Norway, we have shown a negative relationship in fledglings between structural size (as indicated by tarsus length) and telomere length. This relationship continued when fledgling size was increased by an experimental protocol that increased the structural body size of the breeding population, demonstrating the persistent phenotypic link between the two traits. To our knowledge this is the first study that demonstrates that large size is accompanied by a reduction in telomere length in a wild vertebrate species. Because the study was carried out in the wild, unmeasured, correlated factors could influence the results, and the sample size was relatively small. However, that the negative relationship between structural size and telomere length was maintained when the size was extended at the upper end by the selection regime suggests that the negative co-variation between the two traits is not co-incidental.

The reduced telomere length could be a consequence of associations between structural size and gene expression [44, 45], or a downstream phenotypic consequence of larger size, for example by influencing oxidative damage incurred during growth [46]. Alternatively, or additionally, it could involve gametic, embryonic or post natal changes in cell division rates to enable a larger size to be achieved [47, 48]. The size of fledglings is predictive of adult size, and the larger fledglings produced under the selection regime were larger both as chicks and as adults (figure 1*b*). It is also possible that the reduction in telomere length occurred as a consequence not simply of larger size, but also as a result of

faster growth needed to reach this adult size within the defined time frame likely to occur in seasonally breeding species. Thus, growth and body size could both be involved. With respect to the effect of extending the upper range of body size, it would obviously have been useful to have replicate of the selection experiment, [49] but this was not possible due to the logistical effort required to undertake this kind of work in the field. Furthermore, the negative relationship is unlikely to have arisen as a consequence of processes such as genetic drift, since the same relationship was present in the first year, and before the selection experiment was really underway.

Directional selection on body size is commonly reported in the literature [50] and it is well established within quantitative genetics that directional selection on one focal trait will often alter other genetically correlated traits indirectly, depending on the strength and the sign of the genetic correlations [44, 51]. Since telomere length at the end of the growth period is related to lifespan [25], it seems likely that the negative fitness consequences of reduced longevity that accompany larger size could constrain evolution of larger body size. A prerequisite for such an evolutionary mechanism is that body size and telomere length are heritable traits [51]. The heritability of body size is well documented in avian literature [52, 53], and significant heritability of telomere lengths has so far been demonstrated non-human species such as sand lizard (*Lacerta agilis*) [54] collared flycatcher (*Ficedula albicollis*), kakapo (*Strigops habroptilus*) [55, 56], king penguins (*Aptenodytes patagonicus*) [57] and the great reed warbler (*Achrocephalus arundinaceus*) [58]. Heritability of telomere length has also been found in humans [59-61]. However, the genetic correlations between telomere length and other phenotypic traits [51] have yet to be investigated in detail. The present study reveals a evidence of a mechanism that could underpin a trade-off between body size and longevity within species. Since lifespan is a key component of lifetime reproductive success

in iteroparous breeders like birds, the link between structural size and telomere length could have important fitness consequences.

Ethics statement: The project was conducted in accordance with permits from the Norwegian Environmental Agency (Permit Numbers; 2001/6427-ARTS/VI/ARE and 2004/1671 ARTS-VI-ID) and from Bird Ringing Centre at Stavanger Museum, Norway. The project was conducted in accordance with Norwegian legal policy for animal welfare approved by the Norwegian Animal Research Authority (Permit Numbers: S-2603-01 and S-2004-8032-1). These permits cover all research activity involving animals described in this study. No animals were killed or sacrificed in this study.

Author Contributions: BES, THR and HJ designed the selection experiment. THR and HJ and BR collected data. HJ was responsible for the extraction of DNA in Trondheim, whereas WB and RG were responsible for the analyses of telomere lengths in Glasgow. THR conducted the statistical analyses with significant contributions from HP, TK and HH. HP also made the figures. THR and PM wrote the manuscript. BR, TK, IJH, HJ, HP, HH, WB, RG and BES contributed during the process with discussions and manuscript revisions.

Acknowledgements: We are grateful to the fieldworkers that collected the data, in particular to Ole Roar Davidsen. We are also grateful for the help by lab technician Randi Røsbak in Trondheim, and Neil Metcalfe for useful comments.

Data accessibility: The dataset on which the statistical analyses in this study are based is available as electronic supplementary material online (dataset 1).

Funding statement: This work was funded by the European Research Council (ERC-2010-AdG 268562) and the Research Council of Norway (2232571F50). The laboratory work in Glasgow was supported by ERC Advanced Grant 268926.

References

1. Stearns S.C. 1992 *The evolution of life histories*. Oxford, UK Oxford University Press.
2. Roff D.A. 2002 *Life History Evolution* Sunderland, MA, Sinauer.
3. van Noordwijk A.J., de Jong G. 1986 Acquisition and allocation of resources - their influence on variation in life-history tactics. *Am Nat* **128**(1), 137-142. (doi:10.1086/284547).
4. Reznick D.N. 1997 Life history evolution in guppies (*Poecilia reticulata*): Guppies as a model for studying the evolutionary biology of aging. *Experimental Gerontology* **32**(3), 245-258. (doi:10.1016/s0531-5565(96)00129-5).
5. Metcalfe N.B., Monaghan P. 2001 Compensation for a bad start: grow now, pay later? *Trends in Ecology & Evolution* **16**(5), 254-260. (doi:10.1016/s0169-5347(01)02124-3).
6. Metcalfe N.B., Monaghan P. 2003 Growth versus lifespan: perspectives from evolutionary ecology. *Exp Gerontol* **38**, 935-940. (doi:doi:10.1016/S0531-5565(03)00159-1).
7. Martin T.E. 1995 Avian life-history evolution in relation to nest sites, nest predation and food. *Ecol Monogr* **65**(1), 101-127. (doi:10.2307/2937160).
8. Reznick D., Nunney L., Tessier A. 2000 Big houses, big cars, superfleas and the costs of reproduction. *Trends in Ecology & Evolution* **15**(10), 421-425. (doi:10.1016/s0169-5347(00)01941-8).
9. Nilsson J.A., Svensson E. 1996 The cost of reproduction: A new link between current reproductive effort and future reproductive success. *Proc R Soc B-Biol Sci* **263**(1371), 711-714. (doi:10.1098/rspb.1996.0106).
10. Blanckenhorn W.U. 2000 The evolution of body size: What keeps organisms small? *Q Rev Biol* **75**(4), 385-407. (doi:10.1086/393620).
11. Austad S.N. 2010 Methusaleh's Zoo: How Nature provides us with Clues for Extending Human Health Span. *Journal of Comparative Pathology* **141**, S10-S21. (doi:10.1016/j.jcpa.2009.10.024).
12. Blanckenhorn W. 2000 The evolution of body size: what keeps organisms small? *Quart Rev Biol* **75**, 385 - 407.
13. Kraus C., Pavard S., Promislow D.E.L. 2013 The Size-Life Span Trade-Off Decomposed: Why Large Dogs Die Young. *Am Nat* **181**(4), 492-505. (doi:10.1086/669665).
14. Bartke A. 2012 Healthy Aging: Is Smaller Better? - A Mini-Review. *Gerontology* **58**(4), 337-343. (doi:10.1159/000335166).
15. Austad S.N. 2010 Animal size, metabolic rate, and survival, among and within species. In *The comparative biology of aging* (ed. Wolf N.S.), pp. 27-41. Netherlands, Springer.
16. Bernstein R.M. 2010 The Big and Small of It: How Body Size Evolves. *American Journal of Physical Anthropology*, 46-62. (doi:10.1002/ajpa.21440).
17. Miller R.A., Harper J.M., Galecki A., Burke D.T. 2002 Big mice die young: early life body weight predicts longevity in genetically heterogeneous mice. *Aging Cell* **1**(1), 22-29. (doi:10.1046/j.1474-9728.2002.00006.x).

18. Samaras T.T. 2009 Should we be concerned over increasing body height and weight? *Experimental Gerontology* **44**(1-2), 83-92. (doi:10.1016/j.exger.2008.02.002).
19. Miller R.A., Chrisp C., Atchley W. 2000 Differential longevity in mouse stocks selected for early life growth trajectory. *J Gerontol Ser A-Biol Sci Med Sci* **55**(9), B455-B461.
20. Patronek G.J., Waters D.J., Glickman L.T. 1997 Comparative longevity of pet dogs and humans: Implications for gerontology research. *J Gerontol Ser A-Biol Sci Med Sci* **52**(3), B171-B178.
21. Arendt J.D. 1997 Adaptive intrinsic growth rates: An integration across taxa. *Q Rev Biol* **72**(2), 149-177. (doi:10.1086/419764).
22. Tarry-Adkins J.L., Chen J.H., Smith N.S., Jones R.H., Cherif H., Ozanne S.E. 2009 Poor maternal nutrition followed by accelerated postnatal growth leads to telomere shortening and increased markers of cell senescence in rat islets. *Faseb J* **23**(5), 1521-1528. (doi:10.1096/fj.08-122796).
23. Monaghan P. 2010 Telomeres and life histories. the long and the short of it. In *Year in Evolutionary Biology* (eds. Schlichting C.D., Mousseau T.A.), pp. 130-142. Malden, Wiley-Blackwell.
24. Aubert G., Lansdorp P.M. 2008 Telomeres and aging. *Physiol Rev* **88**(2), 557-579. (doi:10.1152/physrev.00026.2007).
25. Heidinger B.J., Blount J.D., Boner W., Griffiths K., Metcalfe N.B., Monaghan P. 2012 Telomere length in early life predicts lifespan. *Proceedings of the National Academy of Sciences* **109**(5), 1743-1748. (doi:10.1073/pnas.1113306109).
26. Aviv A., Susser E. 2013 Leukocyte Telomere Length and the Father's Age Enigma: Implications for Population Health and for Life Course. *Int J Epidemiol* **42**(2), 457-462. (doi:10.1093/ije/dys236).
27. Fick L.J., Fick G.H., Li Z.C., Cao E., Bao B., Heffelfinger D., Parker H.G., Ostrander E.A., Riabowol K. 2012 Telomere Length Correlates with Life Span of Dog Breeds. *Cell Reports* **2**(6), 1530-1536. (doi:10.1016/j.celrep.2012.11.021).
28. Hagen I.J., Billing A.M., Ronning B., Pedersen S.A., Parn H., Slate J., Jensen H. 2013 The easy road to genome-wide medium density SNP screening in a non-model species: development and application of a 10K SNP-chip for the house sparrow (*Passer domesticus*). *Mol Ecol Resour* **13**(3), 429-439. (doi:10.1111/1755-0998.12088).
29. Killpack T.L., Karasov W.H. 2012 Growth and development of house sparrows (*Passer domesticus*) in response to chronic food restriction throughout the nestling period. *J Exp Biol* **215**(11), 1806-1815. (doi:10.1242/jeb.066316).
30. Skjelseth S., Ringsby T.H., Tufto J., Jensen H., Saether B.E. 2007 Dispersal of introduced house sparrows *Passer domesticus*: an experiment. *Proc R Soc B-Biol Sci* **274**(1619), 1763-1771. (doi:10.1098/rspb.2007.0338).
31. Bates D., Mächler M., Bolker B., Walker S. 2015 Fitting Linear Mixed-Effects Models using lme4. *Journal of Statistical Software* (ArXiv e-print; <http://arxiv.org/abs/1406.5823>).
32. R Core Team. 2013 R: A Language and Environment for Statistical Computing. (Vienna, Austria, R Foundation for Statistical Computing.
33. Cleasby I.R., Burke T., Schroeder J., Nakagawa S. 2011 Food supplements increase adult tarsus length, but not growth rate, in an island population of house sparrows (*Passer domesticus*). *BMC Research Notes* **4**(431).
34. Ringsby T.H., Saether B.E., Tufto J., Jensen H., Solberg E.J. 2002 Asynchronous spatiotemporal demography of a house sparrow metapopulation in a correlated environment. *Ecology* **83**(2), 561-569.
35. Jensen H., Saether B.E., Ringsby T.H., Tufto J., Griffith S.C., Ellegren H. 2004 Lifetime reproductive success in relation to morphology in the house sparrow *Passer domesticus*. *J Anim Ecol* **73**(4), 599-611.

36. Parn H., Ringsby T.H., Jensen H., Saether B.E. 2012 Spatial heterogeneity in the effects of climate and density-dependence on dispersal in a house sparrow metapopulation. *Proc R Soc B-Biol Sci* **279**(1726), 144-152. (doi:10.1098/rspb.2011.0673).
37. Griffiths R., Double M.C., Orr K., Dawson R.J.G. 1998 A DNA test to sex most birds. *Mol Ecol* **7**(8), 1071-1075. (doi:10.1046/j.1365-294x.1998.00389.x).
38. Griffith S.C., Dawson D.A., Jensen H., Ockendon N., Greig C., Neumann K., Burke T. 2007 Fourteen polymorphic microsatellite loci characterized in the house sparrow *Passer domesticus* (Passeridae, Aves). *Mol Ecol Notes* **7**(2), 333-336. (doi:10.1111/j.1471-8286.2006.01598.x).
39. Neumann K., Wetton J.H. 1996 Highly polymorphic microsatellites in the house sparrow *Passer domesticus*. *Mol Ecol* **5**(2), 307-309.
40. Griffith S.C., Stewart I.R.K., Dawson D.A., Owens I.P.F., Burke T. 1999 Contrasting levels of extra-pair paternity in mainland and island populations of the house sparrow (*Passer domesticus*): is there an 'island effect'? *Biol J Linnean Soc* **68**(1-2), 303-316. (doi:10.1111/j.1095-8312.1999.tb01171.x).
41. Dawson D.A., Horsburgh G.J., Krupa A.P., Stewart I.R.K., Skjelseth S., Jensen H., Ball A.D., Spurgin L.G., Mannarelli M.E., Nakagawa S., et al. 2012 Microsatellite resources for Passeridae species: a predicted microsatellite map of the house sparrow *Passer domesticus*. *Mol Ecol Resour* **12**(3), 501-523. (doi:10.1111/j.1755-0998.2012.03115.x).
42. Nussey D.H., Baird D., Barrett E., Boner W., Fairlie J., Gemmell N., Hartmann N., Horn T., Hausmann M., Olsson M., et al. 2014 Measuring telomere length and telomere dynamics in evolutionary biology and ecology. *Methods in Ecology and Evolution* **5**(4), 299-310. (doi:10.1111/2041-210x.12161).
43. Criscuolo F., Bize P., Nasir L., Metcalfe N.B., Foote C.G., Griffiths K., Gault E.A., Monaghan P. 2009 Real-time quantitative PCR assay for measurement of avian telomeres. *J Avian Biol* **40**(3), 342-347. (doi:10.1111/j.1600-048X.2008.04623.x).
44. Lande R., Arnold S.J. 1983 The measurement of selection on correlated characters. *Evolution* **37**(6), 1210-1226.
45. Roff D.A., Fairbairn D.J. 2007 The evolution of trade-offs: where are we? *J Evol Biol* **20**(2), 433-447. (doi:10.1111/j.1420-9101.2006.01255.x).
46. Richter T., von Zglinicki T. 2007 A continuous correlation between oxidative stress and telomere shortening in fibroblasts. *Experimental Gerontology* **42**(11), 1039-1042. (doi:10.1016/j.exger.2007.08.005).
47. Allsopp R.C., Chang E., Kashefi-Azham M., Rogaev E.I., Piatyszek M.A., Shay J.W., Harley C.B. 1995 Telomere Shortening Is Associated with Cell Division in Vitro and in Vivo. *Experimental Cell Research* **220**(1), 194-200. (doi:<http://dx.doi.org/10.1006/excr.1995.1306>).
48. Herborn K.A., Heidinger B.J., Boner W., Noguera J.C., Adam A., Daunt F., Monaghan P. 2014 Stress exposure in early post-natal life reduces telomere length: an experimental demonstration in a long-lived seabird. *Proc R Soc B-Biol Sci* **281**(1782). (doi:10.1098/rspb.2013.3151).
49. Henderson N. 1997 Spurious Associations in Unreplicated Selected Lines. *Behav Genet* **27**(2), 145-154. (doi:10.1023/A:1025689425738).
50. Kingsolver J.G., Diamond S.E. 2011 Phenotypic Selection in Natural Populations: What Limits Directional Selection? *Am Nat* **177**(3), 346-357. (doi:10.1086/658341).
51. Lynch M., Walsh B. 1998 *Genetics and Analysis of Quantitative traits*. Sunderland, MA, Sinauer.
52. Merila J., Sheldon B.C. 2001 *Avian quantitative genetics* 179-181 p.
53. Potsma E. 2014 Four decades of estimating heritabilities in wild vertebrate populations: improved methods, more data, better estimates? In *Quantitative genetics in the wild* (eds. Charmantier A., Garant D., Kruuk L.). Oxford, UK, Oxford University Press.


54. Olsson M., Pauliny A., Wapstra E., Uller T., Schwartz T., Blomqvist D. 2011 Sex Differences in Sand Lizard Telomere Inheritance: Paternal Epigenetic Effects Increases Telomere Heritability and Offspring Survival. *PLoS One* **6**(4), e17473. (doi:10.1371/journal.pone.0017473).
55. Voillemot M., Hine K., Zahn S., Criscuolo F., Gustafsson L., Doligez B., Bize P. 2012 Effects of brood size manipulation and common origin on phenotype and telomere length in nestling collared flycatchers. *BMC Ecol* **12**. (doi:10.1186/1472-6785-12-17).
56. Horn T., Robertson B.C., Will M., Eason D.K., Elliott G.P., Gemmell N.J. 2011 Inheritance of Telomere Length in a Bird. *PLoS One* **6**(2). (doi:10.1371/journal.pone.0017199).
57. Reichert S., Rojas E.R., Zahn S., Robin J.P., Criscuolo F., Massemin S. 2015 Maternal telomere length inheritance in the king penguin. *Heredity* **114**(1), 10-16. (doi:10.1038/hdy.2014.60).
58. Asghar M., Bensch S., Tarka M., Hansson B., Hasselquist D. 2014 Maternal and genetic factors determine early life telomere length. *Proceedings of the Royal Society of London B: Biological Sciences* **282**(1799).
59. Graakjaer J., Der-Sarkissian H., Schmitz A., Bayer J., Thomas G., Kolvraa S., Londono-Vallejo J.A. 2006 Allele-specific relative telomere lengths are inherited. *Hum Genet* **119**(3), 344-350. (doi:10.1007/s00439-006-0137-x).
60. Broer L., Codd V., Nyholt D.R., Deelen J., Mangino M., Willemsen G., Albrecht E., Amin N., Beekman M., de Geus E.J.C., et al. 2013 Meta-analysis of telomere length in 19 713 subjects reveals high heritability, stronger maternal inheritance and a paternal age effect. *Eur J Hum Genet* **21**(10), 1163-1168. (doi:10.1038/ejhg.2012.303).
61. Njajou O.T., Cawthon R.M., Damcott C.M., Wu S.-H., Ott S., Garant M.J., Blackburn E.H., Mitchell B.D., Shuldiner A.R., Hsueh W.-C. 2007 Telomere length is paternally inherited and is associated with parental lifespan. *Proceedings of the National Academy of Sciences* **104**(29), 12135-12139. (doi:10.1073/pnas.0702703104).

Figure legends:


Figure 1. An artificial selection experiment for increased tarsus size was conducted among breeding adults in a house sparrows population at the island Leka in northern Norway during 2002-2005. (a) The tarsus length for fledglings increased significantly from 2002 to 2005 ($n = 44$). (b) During the same period the telomere lengths of the fledglings decreased significantly ($n = 44$). In both figures the fledgling tarsus length was standardized to 10 days age for illustration (but not in the statistical analyses, see methods). The boxplots represents the 25%-75% quartiles, where the horizontal line represents the median of the data and the notches corresponds to the 95% confidence interval. Individual values are given as grey circles. See Results for further details.

Figure 2. The tarsus length of house sparrows at fledgling stage was positively predictive for the tarsus length measured when individuals were recaptured later at full-grown size ($n = 15$) at the island Leka in northern Norway. Individuals from 2002 and 2005 are indicated with white and dark circles, respectively. In this analysis the fledgling tarsus length was standardized to 10 days age to allow for small differences in the age at measurement. See Results for further details.


Figure 3. The relationship between the tarsus length of fledglings and their telomere lengths. In this figure the fledgling tarsus length was standardized to 10 days age for illustration (but not in the statistical analyses, see methods). Data from the first and the last year of the selection experiment (2002 and 2005, respectively) are indicated by white and dark circles, respectively ($n = 44$).



67x127mm (300 x 300 DPI)



69x69mm (300 x 300 DPI)



69x69mm (300 x 300 DPI)