

Acta Crystallographica Section E

**Structure Reports**

**Online**

ISSN 1600-5368

Editors: **W.T.A. Harrison, J. Simpson and M. Weil**

## 7-Azathieno[2,3-*c*]cinnoline


**Lars Kr. Hansen, Vegar Stockmann and Anne Fiksdahl**

*Acta Cryst.* (2007). **E63**, o3896

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site or institutional repository provided that this cover page is retained. Republication of this article or its storage in electronic databases other than as specified above is not permitted without prior permission in writing from the IUCr.

For further information see <http://journals.iucr.org/services/authorrights.html>



*Acta Crystallographica Section E: Structure Reports Online* is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. In 2007, the journal published over 5000 structures. The average publication time is less than one month.

**Crystallography Journals Online** is available from [journals.iucr.org](http://journals.iucr.org)

## 7-Azathieno[2,3-c]cinnoline

Lars Kr. Hansen,<sup>a\*</sup> Vegar Stockmann<sup>b</sup> and Anne Fiksdahl<sup>b</sup>

<sup>a</sup>Department of Chemistry, University of Tromsø, N-9037 Tromsø, Norway, and

<sup>b</sup>Department of Chemistry, Norwegian University of Science and Technology, N-7491 Trondheim, Norway

Correspondence e-mail: larsk@chem.uit.no


Received 21 June 2007; accepted 21 August 2007

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$ ;  $R$  factor = 0.041;  $wR$  factor = 0.132; data-to-parameter ratio = 17.3.

The title compound (systematic name: pyrido[4,3-e]thieno[2,3-c]pyridazine),  $C_9H_5N_3S$ , consists of three fused heterocyclic rings. The molecule is planar and the  $\text{N}=\text{N}$  bond length of  $1.302(2)\text{ \AA}$  is in good agreement with values observed in similar compounds. The molecules show  $\pi-\pi$  stacking interactions, forming molecular stacks along the  $b$  axis with interplanar distances of  $3.39(2)$  and  $3.49(2)\text{ \AA}$ . The C—H groups of the thiophene ring are involved in C—H···N interactions, joining molecules into two-dimensional sheets parallel to  $(0\bar{1}1)$ .

## Related literature

For related literature, see: Barton *et al.* (1985); Hansen *et al.* (2007); Holt & Fiksdahl (2006); Hökelek *et al.* (1990, 1991a,b); Stockmann & Fiksdahl (2007); van der Meer *et al.* (1972).



## Experimental

### Crystal data

$C_9H_5N_3S$   
 $M_r = 187.22$   
Triclinic,  $P\bar{1}$   
 $a = 6.8340(18)\text{ \AA}$   
 $b = 7.656(2)\text{ \AA}$   
 $c = 8.847(2)\text{ \AA}$   
 $\alpha = 104.172(3)^\circ$   
 $\beta = 101.376(4)^\circ$

$\gamma = 107.042(5)^\circ$   
 $V = 410.63(18)\text{ \AA}^3$   
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.34\text{ mm}^{-1}$   
 $T = 293(2)\text{ K}$   
 $0.40 \times 0.30 \times 0.20\text{ mm}$

### Data collection

Enraf–Nonius CAD-4 diffractometer  
Absorption correction:  $\psi$  scan [ABSCALC in OSCAIL (McArdle & Daly, 1999; North *et al.*, 1968)]  
 $T_{\min} = 0.882$ ,  $T_{\max} = 0.935$   
2646 measured reflections  
2387 independent reflections  
1805 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.008$   
3 standard reflections  
frequency: 120 min  
intensity decay: 1%

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$   
 $wR(F^2) = 0.132$   
 $S = 1.09$   
2387 reflections  
138 parameters  
All H-atom parameters refined  
 $\Delta\rho_{\max} = 0.37\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.28\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}1-\text{H}1\cdots\text{N}3^{\text{i}}$	0.943 (19)	2.441 (18)	3.352 (3)	159.1 (18)
$\text{C}2-\text{H}2\cdots\text{N}1^{\text{ii}}$	0.89 (2)	2.52 (2)	3.360 (2)	158 (2)

Symmetry codes: (i)  $x - 1, y - 1, z - 1$ ; (ii)  $x - 1, y, z$ .

Data collection: CAD-4-PC Software (Enraf–Nonius, 1992); cell refinement: CELDIM in CAD-4-PC Software; data reduction: XCAD (McArdle & Higgins, 1995); program(s) used to solve structure: SHELXS97 (Sheldrick, 1997); program(s) used to refine structure: SHELXL97 (Sheldrick, 1997); molecular graphics: ORTEX (McArdle, 1995) and ORTEPIII (Burnett & Johnson, 1996); software used to prepare material for publication: OSCAIL (McArdle, 1993).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2090).

## References

- Barton, J. W., Lapham, D. J. & Rowe, D. J. (1985). *J. Chem. Soc. Perkin Trans. I*, pp. 131–133.  
Burnett, M. N. & Johnson, C. K. (1996). ORTEPIII. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.  
Enraf–Nonius (1992). CAD-4-PC Software. Enraf–Nonius, Delft, The Netherlands.  
Hansen, L. K., Stockmann, V. & Fiksdahl, A. (2007). *Acta Cryst. E* **63**, o3290.  
Hökelek, T., Kılıç, E. & Tüzün, C. (1991a). *Acta Cryst. C* **47**, 373–376.  
Hökelek, T., Kılıç, E. & Tüzün, C. (1991b). *Acta Cryst. C* **47**, 369–373.  
Hökelek, T., Watkin, D. J., Kılıç, E. & Tüzün, C. (1990). *Acta Cryst. C* **46**, 1027–1029.  
Holt, J. & Fiksdahl, A. (2006). *J. Heterocycl. Chem.* **43**, 417–423.  
McArdle, P. (1993). *J. Appl. Cryst.* **26**, 752.  
McArdle, P. (1995). *J. Appl. Cryst.* **28**, 65.  
McArdle, P. & Daly, P. (1999). ABSCALC. PC version. National University of Ireland, Galway, Ireland.  
McArdle, P. & Higgins, T. (1995). XCAD. National University of Ireland, Galway, Ireland.  
Meer, H. van der (1972). *Acta Cryst. B* **28**, 367–370.  
North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.  
Sheldrick, G. M. (1997). SHELXS97 and SHELXL97. University of Göttingen, Germany.  
Stockmann, V. & Fiksdahl, A. (2007). In preparation.

## **supplementary materials**

*Acta Cryst.* (2007). E63, o3896 [doi:10.1107/S1600536807041335]

## 7-Azathieno[2,3-*c*]cinnoline

L. K. Hansen, V. Stockmann and A. Fiksdahl

### Comment

The crystal structure of the title compound was solved as part of a study of new tris-heterocyclic compounds with potential biological activity (Stockmann & Fiksdahl, 2007). Thieno[*c*]cinnolines (Barton *et al.*, 1985) have been described in the literature and the crystal structure of benzo[*c*]cinnoline (9,10-diazaphenanthrene) has been solved (van der Meer *et al.*, 1972). A view of the title molecule with the atomic numbering scheme is presented in Fig. 1. The bond lengths are within the normal range of such bonds and also in accordance with the regio-isomer 7-azathieno[3,2-*c*]cinnoline (Hansen *et al.*, 2007) and other benzo[*c*]cinnoline derivatives (Hökelek *et al.*, 1990, 1991*a,b*). The mean N=N distance for seven similar structures is 1.293 (10) Å, ranging from 1.283 (4) to 1.306 (2) Å. The C6—C7 bonds are always significantly shorter than the C7—C8 bonds. Mean C6—C7 bond is 1.364 (10) Å while the mean C7—C8 bond is 1.409 (10) Å. Also the angles show the same systematic differences. The mean C8—C9—C3 angle and the mean C4—C8—C9 angle are 117.8(2.) and 116.1(1.9)°, respectively, while the mean N2—C4—C8 angle is 122.3(2.0)° (see van der Meer, 1972 and references cited therein). The C—H groups of the thiophene ring are involved in intermolecular C—H···N interactions.

### Experimental

7-Azathieno[2,3-*c*]cinnoline was prepared by intramolecular diazo coupling of the diazonium ion intermediate, made by  $\text{NOBF}_4$  diazotization (Holt & Fiksdahl, 2006) of the 3-amino-4-(thiophen-3-yl)pyridine precursor. Single crystals were grown by crystallization from ethyl acetate (Stockmann & Fiksdahl, 2007).

### Refinement

All H atoms were found from a difference map and were refined with isotropic displacement parameters.

### Figures




Fig. 1. A view of the title molecule with the atomic numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.

# supplementary materials

---

## pyrido[4,3-e]thieno[2,3-c]pyridazine

### Crystal data

C <sub>9</sub> H <sub>5</sub> N <sub>3</sub> S	Z = 2
M <sub>r</sub> = 187.22	F <sub>000</sub> = 192
Triclinic, PT	D <sub>x</sub> = 1.514 Mg m <sup>-3</sup>
Hall symbol: -P 1	Mo K $\alpha$ radiation
a = 6.8340 (18) Å	$\lambda$ = 0.71069 Å
b = 7.656 (2) Å	Cell parameters from 25 reflections
c = 8.847 (2) Å	$\theta$ = 12–18°
$\alpha$ = 104.172 (3)°	$\mu$ = 0.34 mm <sup>-1</sup>
$\beta$ = 101.376 (4)°	T = 293 (2) K
$\gamma$ = 107.042 (5)°	Prism, colourless
V = 410.63 (18) Å <sup>3</sup>	0.40 × 0.30 × 0.20 mm

### Data collection

Enraf–Nonius CAD-4 diffractometer	R <sub>int</sub> = 0.008
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 29.9^\circ$
Monochromator: graphite	$\theta_{\text{min}} = 2.5^\circ$
T = 298(2) K	$h = 0 \rightarrow 9$
$\omega/20$ scans	$k = -10 \rightarrow 10$
Absorption correction: $\psi$ scan [ABSCALC in OSCAIL (McArdle & Daly, 1999; North <i>et al.</i> , 1968)]	$l = -12 \rightarrow 12$
$T_{\text{min}} = 0.882$ , $T_{\text{max}} = 0.935$	3 standard reflections
2646 measured reflections	every 120 min
2387 independent reflections	intensity decay: 1%
1805 reflections with $I > 2\sigma(I)$	

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.041$	All H-atom parameters refined
$wR(F^2) = 0.132$	$w = 1/[\sigma^2(F_o^2) + (0.0802P)^2 + 0.041P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.09$	$(\Delta/\sigma)_{\text{max}} = 0.016$
2387 reflections	$\Delta\rho_{\text{max}} = 0.37 \text{ e } \text{\AA}^{-3}$
138 parameters	$\Delta\rho_{\text{min}} = -0.28 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Least-squares planes ( $x,y,z$  in crystal coordinates) and deviations from them (\* indicates atom used to define plane)

$$-0.8879 (0.0025) x + 6.8684 (0.0030) y - 5.7312 (0.0029) z = 1.8667 (0.0014)$$

$$\begin{aligned} * 0.0357 (0.0008) S1 * -0.0053 (0.0012) N1 * -0.0173 (0.0012) N2 * 0.0186 (0.0014) N3 * 0.0049 (0.0013) C1 * -0.0149 \\ (0.0013) C2 * 0.0030 (0.0013) C3 * -0.0112 (0.0013) C4 * 0.0111 (0.0015) C5 * 0.0122 (0.0014) C6 * -0.0038 (0.0013) C7 * \\ -0.0169 (0.0012) C8 * -0.0161 (0.0012) C9 \end{aligned}$$

Rms deviation of fitted atoms = 0.0156

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\text{sigma}(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.31663 (8)	0.41015 (6)	0.11053 (5)	0.05756 (17)
N2	0.78580 (19)	0.7761 (2)	0.48565 (19)	0.0527 (3)
N1	0.6832 (2)	0.6459 (2)	0.34347 (19)	0.0516 (3)
N3	0.7020 (3)	1.0644 (2)	0.83784 (19)	0.0640 (4)
H2	0.009 (3)	0.557 (3)	0.340 (3)	0.064 (6)*
H5	0.937 (4)	1.042 (3)	0.766 (3)	0.071 (6)*
H6	0.419 (4)	1.059 (4)	0.871 (3)	0.083 (7)*
H7	0.203 (4)	0.833 (4)	0.648 (3)	0.084 (7)*
H1	-0.045 (3)	0.332 (3)	0.077 (2)	0.057 (5)*
C1	0.0860 (3)	0.4135 (2)	0.1556 (2)	0.0521 (4)
C2	0.1168 (2)	0.5381 (2)	0.30372 (18)	0.0440 (3)
C3	0.4669 (2)	0.5808 (2)	0.29745 (17)	0.0400 (3)
C4	0.6725 (2)	0.8437 (2)	0.58315 (18)	0.0421 (3)
C5	0.7900 (3)	0.9894 (3)	0.7357 (2)	0.0584 (4)
C6	0.4846 (3)	1.0005 (3)	0.7961 (2)	0.0576 (4)
C7	0.3532 (3)	0.8628 (2)	0.65419 (18)	0.0449 (3)
C8	0.44728 (19)	0.77948 (18)	0.54209 (15)	0.0349 (3)
C9	0.33770 (19)	0.63693 (18)	0.38810 (15)	0.0342 (3)

## Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0703 (3)	0.0558 (3)	0.0418 (2)	0.0245 (2)	0.01587 (18)	0.00583 (17)
N2	0.0308 (6)	0.0565 (8)	0.0718 (9)	0.0165 (5)	0.0131 (6)	0.0227 (7)

## supplementary materials

---

N1	0.0395 (6)	0.0590 (8)	0.0648 (8)	0.0245 (6)	0.0224 (6)	0.0205 (7)
N3	0.0738 (10)	0.0452 (7)	0.0484 (8)	0.0114 (7)	-0.0071 (7)	0.0048 (6)
C1	0.0461 (8)	0.0480 (8)	0.0460 (8)	0.0091 (6)	-0.0026 (6)	0.0096 (6)
C2	0.0311 (6)	0.0470 (7)	0.0487 (8)	0.0118 (5)	0.0057 (5)	0.0139 (6)
C3	0.0390 (6)	0.0418 (7)	0.0426 (7)	0.0178 (5)	0.0136 (5)	0.0139 (5)
C4	0.0314 (6)	0.0393 (7)	0.0492 (7)	0.0090 (5)	0.0024 (5)	0.0151 (6)
C5	0.0450 (8)	0.0463 (8)	0.0611 (10)	0.0050 (7)	-0.0090 (7)	0.0116 (7)
C6	0.0773 (12)	0.0485 (8)	0.0421 (8)	0.0242 (8)	0.0131 (8)	0.0076 (6)
C7	0.0481 (8)	0.0444 (7)	0.0426 (7)	0.0177 (6)	0.0157 (6)	0.0115 (6)
C8	0.0315 (6)	0.0347 (6)	0.0373 (6)	0.0115 (5)	0.0077 (5)	0.0119 (5)
C9	0.0291 (5)	0.0368 (6)	0.0367 (6)	0.0126 (5)	0.0083 (4)	0.0121 (5)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

S1—C1	1.7058 (19)	C2—H2	0.89 (2)
S1—C3	1.7267 (17)	C3—C9	1.3929 (18)
N2—N1	1.302 (2)	C4—C8	1.4114 (18)
N2—C4	1.375 (2)	C4—C5	1.421 (2)
N1—C3	1.3534 (19)	C5—H5	0.92 (2)
N3—C5	1.293 (3)	C6—C7	1.359 (2)
N3—C6	1.361 (3)	C6—H6	0.97 (2)
C1—C2	1.354 (2)	C7—C8	1.4089 (19)
C1—H1	0.95 (2)	C7—H7	0.97 (3)
C2—C9	1.4205 (18)	C8—C9	1.4136 (18)
C1—S1—C3	90.68 (7)	N3—C5—C4	123.69 (16)
N1—N2—C4	119.47 (12)	N3—C5—H5	115.1 (15)
N2—N1—C3	118.30 (13)	C4—C5—H5	121.2 (15)
C5—N3—C6	117.91 (15)	C7—C6—N3	124.48 (17)
C2—C1—S1	114.11 (12)	C7—C6—H6	117.7 (15)
C2—C1—H1	127.8 (12)	N3—C6—H6	117.8 (15)
S1—C1—H1	118.1 (12)	C6—C7—C8	118.13 (16)
C1—C2—C9	111.69 (14)	C6—C7—H7	112.6 (15)
C1—C2—H2	122.8 (14)	C8—C7—H7	129.3 (15)
C9—C2—H2	125.4 (14)	C7—C8—C4	118.27 (13)
N1—C3—C9	126.65 (14)	C7—C8—C9	126.32 (12)
N1—C3—S1	121.77 (11)	C4—C8—C9	115.40 (12)
C9—C3—S1	111.58 (11)	C3—C9—C8	115.56 (12)
N2—C4—C8	124.60 (13)	C3—C9—C2	111.93 (13)
N2—C4—C5	117.89 (14)	C8—C9—C2	132.50 (12)
C8—C4—C5	117.51 (15)		
C4—N2—N1—C3	-0.5 (2)	C6—C7—C8—C9	-179.09 (14)
C3—S1—C1—C2	0.64 (13)	N2—C4—C8—C7	-179.74 (13)
S1—C1—C2—C9	-0.48 (18)	C5—C4—C8—C7	-0.2 (2)
N2—N1—C3—C9	-0.8 (2)	N2—C4—C8—C9	-0.6 (2)
N2—N1—C3—S1	179.35 (11)	C5—C4—C8—C9	178.96 (13)
C1—S1—C3—N1	179.21 (14)	N1—C3—C9—C8	1.4 (2)
C1—S1—C3—C9	-0.63 (11)	S1—C3—C9—C8	-178.80 (9)
N1—N2—C4—C8	1.2 (2)	N1—C3—C9—C2	-179.35 (14)
N1—N2—C4—C5	-178.37 (15)	S1—C3—C9—C2	0.49 (15)

C6—N3—C5—C4	−0.4 (3)	C7—C8—C9—C3	178.46 (13)
N2—C4—C5—N3	−179.99 (16)	C4—C8—C9—C3	−0.58 (18)
C8—C4—C5—N3	0.4 (3)	C7—C8—C9—C2	−0.6 (2)
C5—N3—C6—C7	0.1 (3)	C4—C8—C9—C2	−179.69 (14)
N3—C6—C7—C8	0.1 (3)	C1—C2—C9—C3	−0.02 (18)
C6—C7—C8—C4	−0.1 (2)	C1—C2—C9—C8	179.12 (14)

*Hydrogen-bond geometry ( $\text{\AA}$ , °)*

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
C1—H1···N3 <sup>i</sup>	0.943 (19)	2.441 (18)	3.352 (3)	159.1 (18)
C2—H2···N1 <sup>ii</sup>	0.89 (2)	2.52 (2)	3.360 (2)	158 (2)

Symmetry codes: (i)  $x-1, y-1, z-1$ ; (ii)  $x-1, y, z$ .

## supplementary materials

Fig. 1

