



Use of mechanical and pharmacological restraint over an eight-year period and its relation to clinical factors

Solveig Klæbo Reitan, Anne-Sofie Helvik & Valentina Iversen

To cite this article: Solveig Klæbo Reitan, Anne-Sofie Helvik & Valentina Iversen (2017): Use of mechanical and pharmacological restraint over an eight-year period and its relation to clinical factors, Nordic Journal of Psychiatry, DOI: [10.1080/08039488.2017.1373854](https://doi.org/10.1080/08039488.2017.1373854)

To link to this article: <http://dx.doi.org/10.1080/08039488.2017.1373854>



© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 06 Sep 2017.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

Use of mechanical and pharmacological restraint over an eight-year period and its relation to clinical factors

Solveig Klæbo Reitan^{a,b}, Anne-Sofie Helvik^{a,c,d} and Valentina Iversen^{a,b}

^aDepartment of Mental Health, St. Olav University Hospital HF, Trondheim, Norway; ^bFaculty of Medicine and Health Science, Norwegian University of Science and Technology, Trondheim, Norway; ^cDepartment of Public Health and Nursing, Faculty of Medicine and Health Care, Norwegian University of Science and Technology (NTNU), Trondheim, Norway; ^dNorwegian National Advisory Unit on Ageing and Health, Vestfold Hospital Trust, Tønsberg, Norway

ABSTRACT

Background: Use of restraint and finding the balance between security and ethics is a continuous dilemma in clinical psychiatry. In daily clinic and in planning health-care service, knowledge on the characteristics of restraint situations is necessary to optimize its use and avoid abuse.

Methods: We describe characteristics in the use of pharmacological and mechanical restraint in psychiatric acute wards in a hospital in Middle Norway over an eight-year period. Data on all cases of mechanical and pharmacological restraint from 2004 to 2011 were retrospectively collected from handwritten protocols. Complementary information on the patients was obtained from the hospital patient administrative system.

Results: Restraint in acute wards was used on 13 persons per 100,000 inhabitants annually. The percentage of admitted patients exposed to restraint was 1.7%, with a mean of 4.5 cases per exposed patient. Frequency per 100 admitted patients varied from 3.7 (in 2007) to 10 (in 2009). The majority of restraint cases concerned male patients under 50 years and with substance-abuse, psychotic, or affective disorders. Significantly more coercive means were used during daytime compared to night and morning. There was a significant increase in pharmacological coercion during spring and mechanical coercion during summer.

Conclusions: Restraint was used on 1.7% of admitted patients, representing 13 per 100,000 inhabitants per year. Use of restraint was higher during certain periods of the day and was associated with the patient's diagnosis, age, gender, and legal status of hospitalization. There was a marked variation over the years.

ARTICLE HISTORY

Received 14 March 2017
Revised 21 August 2017
Accepted 27 August 2017

KEYWORDS

Restraint; coercion; mechanical; pharmacological; variation

Background and aim

Background

Any use of coercion raises ethical dilemmas and needs continuous surveillance. In clinical psychiatry, the balance between care, security, and abuse is challenging. Thus, there is an overall intention to limit the use of coercion to an absolute minimum, and there is national [1] as well as international [2–4] focus on how this can be achieved. Knowledge on factors influencing the use of coercion is mandatory to optimize the situation. Both clinicians and administrative decision makers need this information. There have been attempts to structure knowledge on factors influencing the occurrence and the duration of restraint [5,6]. However, there are deviations in reports between countries [7]. The need for a consensus on definitions has been discussed. A time frame of one year to avoid variation over the year and report on the number of incidents of restraint per 100,000 inhabitants in the hospital's catchment area are suggested [8,9].

National legislation regulates the use of different forms of coercion. The Norwegian health-care act §4–8 regulates the use of restraint [10]. Means that can be used include (a) mechanical restraint (straps on limbs and/or chest binding the patient to a bed or, in some cases, straps used to minimize movement during walk), (b) short-time isolation (seclusion), (c) pharmacological restraint in the form of short-acting medications such as benzodiazepines and antipsychotics *per os* or as injection (not to be confused with coerced psychopharmacological medication regulated by §4–4), and (d) short-lasting hold of the patient. The means of restraint can only be used in accordance with a decision by a senior clinical psychiatrist or clinical psychologist seeing the patient in the situation.

Restraint is often regarded as necessary in clinical psychiatry [11,12] and is mainly performed with care. Still, there are reports on unwanted effects such as prolonged admittance, pressure sores, psychological traumas, and even death [11,13–17]. Executing coercion and restraint is also stressful for the health-care personal. Patient's characteristics reported

CONTACT Solveig Klæbo Reitan  solveig.reitan@ntnu.no  St. Olav University Hospital HF, Department Mental Health, PO Box 3250 Sluppen, 7006 Trondheim, Norway

© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group
This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

to influence the use of restraint vary but include the patient's diagnosis [11,18–21], younger age [20,22], immigrant background [20], and aggressive or self-harming behavior [23]. Reports on the effect of gender vary. Some studies report male gender to be a risk factor [20,22], others report a higher risk for female patients [19], and others find no effect of gender [20,24,25]. The time of day (peak in afternoon and early evening) and season (peak in autumn) are non-patient factors influencing the use of restraint [22]. Also, ward and staff factors such as education, work satisfaction, patient/staff ratio, systematic identification of patient crises, and action by separating severely ill patients from others may have an effect [5,26,27].

To evaluate the effect of attempts to reduce and optimize the use of restraint, a reliable registration of restraint is necessary. In Norway, electronic local and central registers are insufficient so far. Retrospective questionnaires also have been used [5]. A more direct way of registering the actual use of restraint is the examination of hand-written protocols at individual wards like Høyer and Drange [28,29] and Wynn [22] have done.

Aim of the study

We wanted to investigate factors associated with the use of mechanical and pharmacological restraints in a heterogeneous, stable acute ward population over an eight-year period. Also, we wanted to study the occurrence of mechanical and pharmacological restraints and the trends over a period of several years to reveal potential variations.

Materials and method

Setting

Data on medical and mechanical restraint according to the Norwegian health-care act §4–8 [10] as described in background were collected retrospectively from hand-written protocols for the period from 1st January 2004 to 31st December 2011. The included wards represented the only acute psychiatric inpatient care for adults in South-Trøndelag County in Middle Norway covering both rural and urban areas with a population of 290,000 inhabitants.

Study population

A total of 1468 cases of pharmacological and mechanical restraint were registered. Because of registration errors, ten cases were excluded, leaving 1458 cases of restraint in this study.

According to the tradition in the wards, isolation (seclusion) was not used as restraint, and holding was not registered until 2008. Thus, we have data only for mechanical and pharmacological restraint.

Measures

Information regarding all coercive episodes had been consecutively recorded in hand-written protocols at each ward by trained nurses and psychiatrists or psychologists. The date

and time of day were registered in these protocols. The patients' age, gender, diagnosis, and legal status of hospitalization (voluntarily or involuntarily admitted) at the time of restraint were collected from the hospital's electronic patient administration system (PAS).

Diagnoses were categorized in six groups based on diagnoses from ICD-10: organic mental disorders (F00–F09), use of psychoactive substances (F10–F19), schizophrenic spectrum disorders (F20–F29), mood disorders (F30–F39), personality disorders (F60–F69), and other diagnostic groups, including neurotic, stress-related, and somatoform disorders (F40–F49), behavior syndromes (F50–F59), mental retardation (F70–F79), disorders of psychological development (F80–F89), behavioral and emotional disorders usually occurring in childhood or adolescence (F90–F98), and unspecified mental disorders (F99).

The year was divided in four seasons: winter (21st December to 20th March), spring (21st March to 20th June), summer (21st June to 20th September), and fall (21st September to 20th December). The day was divided into four intervals: 00:01–06:00, 06:01–12:00, 12:01–18:00, and 18:01–00:00.

Analysis

Chi-square tests were used to compare the categorical data of patients being subjected to restraint (gender, legal referral status, and diagnosis) by type of restraint, and a simple comparison of age by type of restraint was done with the Mann–Whitney test. A cross tabulation was also performed to measure the frequency of the different restraints per season and per time interval and by admissions per year.

In the preliminary analysis, binary logistic regressions (OR 95% CI) were used to assess the correlation between the independent variables (gender, age, legal hospitalisation status, season, 24-h distribution, and diagnostic groups) by mechanical restraints versus pharmacological restraints. The diagnostic groups were categorized with schizophrenic spectrum disorders as a reference. Independent variables which correlated with the outcome in the binary analysis with $p < 0.100$ were included in the adjusted logistic regression analysis.

Ethics

In accordance with the regulations of the Health Research Act of 1st July 2008, the research project was presented to the regional committee for medical and health research ethics (REC). According to REC, there was no need for their consent, because the data were collected for quality-assurance reasons for the hospital. The Data Protection Office at Hospital and the Department Head approved the project (number 13/6804-2).


Results

Use of restraint

In the eight-year period under study, 324 persons were exposed to restraint covering a total of 1458 cases (mean 4.5

Table 1. Characteristics of patients being subjected to restraint.

	Total <i>N</i>	Mechanical		Pharmacological		<i>p</i> Value
		<i>n</i>	(%)	<i>n</i>	(%)	
Patients	1458	863	(59.2)	595	(40.8)	
Age (years)	1457	862	(59.2)	595	(40.8)	
18–29	463	323	(69.8)	140	(30.2)	<0.001 ^a
30–39	297	172	(57.9)	125	(42.1)	
40–49	370	253	(68.4)	117	(31.6)	
>50	327	114	(34.9)	213	(65.1)	
Gender	1458	863	(59.2)	595	(40.8)	
Female	614	387	(63.0)	227	(37.0)	0.011 ^b
Male	844	476	(56.4)	368	(43.6)	
Referral status ^c	1433	850	(59.3)	583	(40.7)	
Voluntary	361	251	(69.5)	110	(30.5)	<0.001 ^b
Involuntary	1072	599	(55.9)	473	(44.1)	

^aMann–Whitney, nonparametric test.^bPearson Chi-square.^cMissing information for 25 cases.**Figure 1.** Total number of admittances, total number of restraints, and number of restraints in relation to number of admittances. Year on X-axis and logistic number scale on Y-axis.

cases per exposed patient) among the 19,283 patients admitted from the catchment area of 290,000 persons.

Annually, an average of 40 persons were exposed to restraint, corresponding to 1.7% of admitted patients. More than half (59.2%) of the cases of restraint were cases of mechanical restraint (see Table 1).

The use of restraint (in total as well as per admission) varied over the years (Figure 1) from 110 cases of restraint in 2883 admittances (3.8 cases per 100 admittances) in 2007 to a peak of 211 cases of restraint in 2638 admittances (8.0 cases per 100 admittances) in 2008.

Table 2. Characteristics of patients being subjected to restraint for the first time per individual *N* = 324.

	Total <i>N</i>	Mechanical	Pharmacological	<i>p</i> Value
		<i>n</i>	<i>n</i>	
Patients				
Age (years) ^a				0.008 F
18–29	101	46	55	
30–39	75	30	45	
40–49	66	22	44	
>50	82	18	64	
Gender				0.009 Q
Male	178	75	103	
Female	146	41	105	
Referral status ^b				0.066 D
Voluntary	87	38	49	
Involuntary	230	74	155	

^aMann–Whitney, nonparametric test.^bMissing information for 27 cases.

D = Chi-square 3.372.

F = Chi-square 11.765.

Q = Chi-square 6.891.

Patient characteristics associated with restraint

Gender

Significantly fewer females were subjected to restraint ($p = 0.009$, Table 2), although slightly more females than males were admitted to the acute wards (e.g. 58.5% females in the period 2005–2007).

Once subjected to restraint, males would significantly more often be subjected to mechanical restraints compared to females ($p = 0.028$, Table 3).

Legal status of hospitalization

Significantly more cases of restraint concerned patients who were involuntarily admitted (1072 patients with known hospitalisation status, 74.8%, $p < 0.001$, Table 1). However, for individual persons subjected to restraint (Table 2), the association to their legal status of hospitalization did not reach significance ($p = 0.066$). Actually, 361 (25.2%) cases of restraint were on voluntarily admitted patients. The data show that 29.5% of cases of mechanical restraints and 18% of cases of pharmacological restraints were on voluntarily admitted patients. Also, there was no correlation between legal status of hospitalization and type of restraint (mechanical versus pharmacological, Table 3). We had no information about the use of coercion according to legal status for 25 cases (1.7%).

Table 3. Unadjusted and adjusted logistic regression analysis (OR^a and 95%CI^b) of mechanical restraints versus pharmacological restraint at first registration per patient ($n = 316$).

	Unadjusted			Adjusted		
	OR	95% CI	<i>p</i> Value	OR	95% CI	<i>p</i> Value
Female gender (ref. male)	0.536	(0.336–0.856)	0.009	0.564	(0.338–0.941)	0.028
Age (pr years)	0.975	(0.960–0.991)	0.002	0.971	(0.971–0.989)	0.002
Involuntary admissions (ref. voluntary admission first time)	0.624	(0.376–1.034)	0.067	0.730	(0.413–1.290)	0.278
<i>Diagnostic groups at admission first time</i>						
Schizophrenic spectrum disorder (F20–F29)	Ref	1.00		Ref	1.00	0.006
Organic mental disorders (F00–F09)	1.929	(0.858–4.339)	0.112	4.004	(1.529–10.488)	0.005
Psychoactive abuse (F10–F19)	1.952	(0.957–3.984)	0.066	1.611	(0.749–3.467)	0.222
Mood disorders (F30–F39)	0.671	(0.336–1.339)	0.271	0.815	(0.394–1.689)	0.583
Personality disorder (F60–F69)	2.678	(0.901–7.955)	0.076	2.451	(0.769–7.807)	0.129
Collective diagnostic group ^c	2.836	(1.373–5.856)	0.003	2.470	(1.137–5.368)	0.022
Adjusted R^{2d}					14.5%	

^aOR = odds ratio.^bCI = confidence interval.^cNeurotic, stress-related, and somatoform disorders (F40–49), behavior syndromes (F50–59), mental retardation (F70–79), disorder of psychological development (F80–89), behavioral and emotional disorders usually occurring in childhood or adolescence (F90–98), and unspecified mental disorders (F99).^dAdjusted R^2 = Nagelkerke R^2 .**Table 4.** Pharmacological and mechanical restraint among diagnostic groups for first-time restraint per patient.

	Total (N)	Mechanical (n)	Pharmacological (n)	<i>p</i> Value ^a
Total (n)	316			0.003
Organic mental disorders (F00–F09)	31	14	17	
Psychoactive abuse (F10–F19)	44	20	24	
Schizophrenic spectrum disorder (F20–F29)	117	35	82	
Mood disorders (F30–F39)	67	15	52	
Personality disorder (F60–F69)	15	8	7	
Collective diagnostic group ^b	42	23	19	

Missing info in eight cases.

^aPearson chi-square.^bNeurotic, stress-related, and somatoform disorders (F40–49), behavior syndromes (F50–59), mental retardation (F70–79), disorders of psychological development (F80–89), behavioral and emotional disorders usually occurring in childhood or adolescence (F90–98), and unspecified mental disorders (F99).

Age

There was a significant effect of age on cases of restraint ($p < 0.001$, Table 1) and on individuals subjected to restraint ($p = 0.008$, Table 2). Age also was seen to affect the type of restraint chosen ($p = 0.002$, Table 3). Mechanical restraint was more often used for younger patients, and pharmacological restraint was more often chosen for older patients if restraint was used.

Diagnoses

Diagnostic groups are significantly differently represented in the group exposed to coercion ($p = 0.003$), with schizophrenic spectrum disorders (F20–29) and mood disorders (F30–39) being the most often represented (Table 4). Once subjected to restraint, the risk of experiencing mechanical rather than pharmacological restraint is significantly higher if the diagnoses are within the groups organic mental disorder ($p = 0.005$) and collective diagnostic group ($p = 0.022$), as can be seen in Table 3.

Non-patient factors

Seasonal and diurnal variation

The use and type of restraint varied significantly by seasonal time ($p = 0.043$, Table 5). During summer, mechanical restraint was used significantly more often than

pharmacological restraint, whereas pharmacological restraint was used significantly more often than mechanical restraint during spring.

There was a significant diurnal variation ($p = 0.007$, Table 5). About 60% of all types of restraint were performed between 12:01 and 00:00. Both mechanical and pharmacological restraints were used more often in this period, and the difference was particularly large for mechanical restraints.

Discussion

Use of restraint

The data show that 1.7% of patients admitted to psychiatric acute wards (13 persons per 100,000 inhabitants per year) were subject to restraint in this eight-year period. Because the routines and culture for hospital admission vary a lot, the use of restraint per inhabitant probably is the best measure to use [9]. However, most studies report the percentage of admitted patients. In a Swiss study, restraint and seclusion were found to be used on 5.6% of admitted patients [18]. However, comparing these results is difficult because the number of inhabitants is not given in the Swiss study, and in our hospital, seclusion is not used. Others report the frequency of restraint among patients involuntarily admitted [23] or present studies based on wards not otherwise specified [28,29]. To our knowledge, there are no reliable numbers regarding the actual use of restraint per inhabitant. We found

Table 5. Seasonal and daytime variations in use of pharmacological and mechanical restraint.

	Total (N)	Mechanical		Pharmacological		p Value ^a
		n	(%)	n	(%)	
Season	1458	863	100	585	100	0.043
Winter	328	190	22.0	138	23.2	
Spring	383	208	24.1	175	29.4	
Summer	396	254	29.4	142	23.9	
Fall	351	211	24.4	140	23.5	0.007
Time interval	1441	861	100	580	100	
00:01–06:00	288	163	18.9	125	21.6	
06:01–12:00	262	136	15.8	126	21.7	
12:01–18:00	432	274	31.8	158	27.2	
18:01–00:00	459	288	33.4	171	29.5	

^aPearson Chi-square.

a marked variation over the years (from a minimum of 110 cases in 2007 to a maximum of 211 cases in 2008) in the same population, independently of potential attempts to reduce restraint in certain periods. This is in line with a Danish study for 2011–2013, which did not find any effect of attempts to reduce restraint at the wards [30]. There are reports on changes between separate periods (e.g. 1986 and 1990 [28] or 1990, 1991, 1994, 1998, and 2004 [21]) but not on the variation over several consecutive years. This variation was revealed as we analyzed data for several consecutive years in the same wards. It may indicate that analyzing the use of restraint for one specific period or comparing only two separate periods to analyze the effect of attempts to reduce restraint may be biased by the inherent inter-annual variation. The frequently found variation in use of coercion between different geographical regions or hospitals also should be viewed in this perspective. Probably, more studies on the absolute frequency of restraint should be performed over consecutive periods to map the use of restraint better.

Patient characteristics

As mentioned in the introduction, previous findings regarding the effect of gender show a large variation. We find significantly more males among those subjected to restraint (although slightly more females are admitted). Once subjected to restraint, males are subjected to mechanical restraint significantly more often. A study done on the staff's emotional reaction to violent behavior in psychiatric inpatients found that males and females provoked different reactions [31]. Thus, females may be perceived as less threatening than males and less susceptible to require restraint.

The Norwegian legislation allows the use of restraint independently of legal referral status. We investigated the association between legal referral status and restraint, and we did not find any significant correlation among those subjected to restraint. We have not adjusted for involuntary admission among the total patient population. A Danish study found a significant effect of being involuntarily admitted [30]. However, both studies illustrate that a substantial number of patients voluntarily admitted are subjected to restraint. Our data are very much in line with the findings for a Danish acute-ward population serving a catchment area of 250,000

registering the use of medical coercion. Here, 33% of those subjected to medical restraint were voluntarily admitted [32].

We find that younger patients are more often represented among those subjected to restraint and that younger age also influences the choice of restraint (mechanical *versus* pharmacological). This is in line with previous reports [20,22], although not everyone has observed this correlation [30]. We find that the diagnoses most frequently being represented among those subjected to restraint are schizophrenic spectrum disorders and mood disorders. This is in line with the recent Danish finding that the diagnoses Schizophrenia, transient psychoses, and mania made up 61% of the population subjected to pharmacological restraint [32].

Regarding diagnoses, once subjected to restraint, the use of mechanical restraint is significantly higher if the patients are diagnosed with organic mental disorder or are in the collective group, compared to the other diagnoses. Diagnostic groups have been found to influence the use of restraint, although with varying results. One reason for the variety of reports may be that different patient populations are studied and thus, so far, no clear conclusions on the overall effects of diagnoses can be reached. Our finding makes sense because these diagnostic groups include patients experiencing scary psychoses and thus acting aggressively in response to psychotic experiences. Simultaneously, these patients will potentially be more susceptible and vulnerable to experiencing the surroundings as threatening, unsafe, and dangerous. Thus, the psychological effect of restraint may be dramatic for this group.

Non-patient factors

We observed a seasonal variation with slightly more restraints in spring and summer. There were significantly more cases of pharmacological restraint in spring and more cases of mechanical restraint in summer.

This differs from Wynn et al. [22] who report a peak of restraint in autumn. However, a peak of cases of violence in spring (potentially leading to restraint) in the acute psychiatric wards in South-Trøndelag has been reported [33] as well as a peak of restraint in spring in Italy [34]. For mania, it is also established that there is an increase in spring [35], but we could not find any reports on seasonal variation in activity of non-affective psychoses. The finding of increased mechanical restraint in summer is partly in line with the findings of Paavola et al. [36] from a forensic psychiatric unit in Finland which showed an increase in restraint in summer and autumn. Medical explanations for increased restraint in summer are not obvious. Paavola et al. suggest effects of staff vacation and unexperienced substitute staff. This is a possibility in our study as well. This may be a target for interventions to reduce restraint. Worth noting, Wynn et al. [22] collected data in Tromsø, north of the polar circle, whereas Morken and Linaker [33] collected data in the middle of Norway, as we did, (just south of the polar circle but still far north on the Earth). The latitude would be similar for Finland. If one assumes that day light affects activity in psychiatric disorders as well as aggression and lifestyle (e.g.

vacation and substance use as well as different groups of staff at the wards), other places may have other findings. Still, the seasonal variation is a factor in preventive efforts in the wards at these latitudes.

Most use of coercion took place during daytime (12:01–18:00 and 18:01–00:00) in line with other reports [22]. This is expected because these are the hours during which most patients are awake, and this constitutes important information regarding preventive efforts in wards.

Study limitations and strengths

Our study has several limitations. It only addresses patients subjected to two (out of four possible) specific restraints. Also, we have not studied other kinds of coercive acts (coercive medication and nutrition, restrictions in social contact, coercive admission, etc.). Variations in diagnostic groups, age, gender, and comorbidity in the admitted population as a whole are not adjusted for. We only investigated the use of restraint in acute wards, leaving out information from high-security wards. Thus, we cannot draw conclusions on the use of restraint in different groups, all use of coercion, and so on. We did not register the duration of stay before use of restraint. Restraint has been shown to take place mainly during the first hours after admittance [30].

We only investigated factors available from hospital patient files. It was reported recently that extramural factors like no or poor outpatient psychiatric care before admittance as well as patients with poor compliance regarding psychotropic medications affected the use of restraint [30]. Such factors should be included in future studies. Also, we did not have data on clinical evaluation and motivation for use of restraint. Information on these issues may increase the knowledge in the field further. Because legislation and tradition differ significantly between countries, it is hard to generalize all findings across countries. Because all our data come from the same hospital and from the same catchment area, it might be hard to generalize the results to other patient populations.

One of the strengths of our study is that we have investigated the on-site hand-written clinical protocols continuously updated on the wards where the patients are and the coercion takes place to avoid deficient reports of coercion in electronic systems or recall bias in studies using retrospective questionnaires. Because there is a known variation during the year [9], we have covered the whole year and all seasons. Another important factor is that we have covered a period of several, consecutive years and not only reported 'point prevalence'. This matters because there is a marked inter-annual variation. The four wards in the study represented the only available inpatient acute psychiatric service for the whole catchment area of South-Trøndelag and thus the main use of restraint in the area. This allows us to calculate the frequency of restraint per 100,000 inhabitants in the catchment area, avoiding bias due to differences in the organization of psychiatric services as emphasized by Janssen et al. [9]. Thus, a representative image of the use of restraint in this area is shown. Also, this means that all social groups are represented.

Conclusions

A total of 13 persons per 100,000 inhabitants per year (1.7% of admitted patients) were subjected to restraint in the acute wards in the eight-year period from 2004 to 2011. There was a marked inter-annual variation not otherwise explained that should be taken into account when analyzing use of restraint. The inter-annual variation seems to be independent of campaigns to reduce restraint, suggesting that other and maybe extramural/pre-hospital factors should be analyzed to explore the field further.

Among those subjected to restraint, we find significantly more males and younger persons, and in these cases, mechanical restraint is used more often than pharmacological restraint. The diagnostic groups including psychoses (schizophrenic spectrum disorder and affective disorders) were significantly more often found among those subjected to restraint. More restraint was performed during the time period 12:01–00:00 (afternoon and evening) and in spring and summer.

We suggest that studies on the use of restraint should take into account potential inter-annual variations with consecutive years included, as well as seasonal and diurnal variations. The significantly higher representation of patients with psychotic disorders among those subjected to restraint is explainable though worrisome because psychotic patients are particularly vulnerable and susceptible to experiencing the restraint as trauma. This phenomenon should be addressed. We suggest that the knowledge on the time of year and day with increased use of restraint is taken into account when running wards regarding presence, quantity, and competency of staff.

The frequency of cases of restraint (e.g. 40 patients per year) as well as per inhabitant (e.g. 13 per 100,000 inhabitants per year) is not very high. However, the experienced burden of restraint from patients as well as staff is high, and society is generally interested in this field. We suggest that qualitative studies in this field, examining the vast experiences, are also performed.

Acknowledgements

The authors would like to thank Mari Brendskag, Rita Småvik, and Wenche Langfjord Mangerud (St. Olav University Hospital) for their contribution in collecting data.

Disclosure statement

No potential conflict of interest was reported by the authors.

References

- [1] Leiknes KA, Dahm KT. Re: seclusion in emergency psychiatry. *Tidsskr nor Laegeforen*. 2015;135:414–415.
- [2] Taxis JC. Ethics and praxis: alternative strategies to physical restraint and seclusion in a psychiatric setting. *Issues Ment Health Nurs*. 2002;23:157–170.
- [3] Sullivan AM, Bezmen J, Barron CT, et al. Reducing restraints: alternatives to restraints on an inpatient psychiatric service—utilizing

- safe and effective methods to evaluate and treat the violent patient. *Psychiatr Q*. 2005;76:51–65.
- [4] Borckardt JJ, Madan A, Grubaugh AL, et al. Systematic investigation of initiatives to reduce seclusion and restraint in a state psychiatric hospital. *Psychiatr Serv*. 2011;62:477–483.
- [5] Bak J, Zoffmann V, Sestoft DM, et al. Comparing the effect of non-medical mechanical restraint preventive factors between psychiatric units in Denmark and Norway. *Nord J Psychiatry*. 2015;69:433–443.
- [6] Deichmann Nielsen L, Bech P, Hounsgaard L, et al. "Mechanical restraint-confounders, risk, alliance score": testing the clinical validity of a new risk assessment instrument. *Nord J Psychiatry*. 2017;71:441–447.
- [7] Lepping P, Masood B, Flammer E, et al. Comparison of restraint data from four countries. *Soc Psychiatry Psychiatr Epidemiol*. 2016;51:1301–1309.
- [8] Bowers L, van der Werf B, Vokkolainen A, et al. International variation in containment measures for disturbed psychiatric inpatients: a comparative questionnaire survey. *Int J Nurs Stud*. 2007;44:357–364.
- [9] Janssen WA, van de Sande R, Noorthoorn EO, et al. Methodological issues in monitoring the use of coercive measures. *Int J Law Psychiatry*. 2011;34:429–438.
- [10] The mental Health Care Act, Act number 62, relating to the provision and implementation of mental health care, with later amendments; 1999.
- [11] Fisher WA. Restraint and seclusion: a review of the literature. *Am J Psychiatry*. 1994;151:1584–1591.
- [12] Sharfstein SS. Commentary: reducing restraint and seclusion: a view from the trenches. *Psychiatr Serv*. 2008;59:197.
- [13] Hatta K, Shibata N, Ota T, et al. Association between physical restraint and drug-induced liver injury. *Neuropsychobiology*. 2007;56:180–184.
- [14] Mohr WK, Petti TA, Mohr BD. Adverse effects associated with physical restraint. *Can J Psychiatry*. 2003;48:330–337.
- [15] Hem E, Steen O, Opjordsmoen S. Thrombosis associated with physical restraints. *Acta Psychiatr Scand*. 2001;103:73–75. discussion 5–6.
- [16] Jarrett M, Bowers L, Simpson A. Coerced medication in psychiatric inpatient care: literature review. *J Adv Nurs*. 2008;64:538–548.
- [17] Strout TD. Perspectives on the experience of being physically restrained: an integrative review of the qualitative literature. *Int J Ment Health Nurs*. 2010;19:416–427.
- [18] Lay B, Nordt C, Rossler W. Variation in use of coercive measures in psychiatric hospitals. *Eur Psychiatry*. 2011;26:244–251.
- [19] Beck NC, Durrett C, Stinson J, et al. Trajectories of seclusion and restraint use at a state psychiatric hospital. *Psychiatr Serv*. 2008;59:1027–1032.
- [20] Knutzen M, Sandvik L, Hauff E, et al. Association between patients' gender, age and immigrant background and use of restraint—a 2-year retrospective study at a department of emergency psychiatry. *Nord J Psychiatry*. 2007;61:201–206.
- [21] Keski-Valkama A, Sailas E, Eronen M, et al. A 15-year national follow-up: legislation is not enough to reduce the use of seclusion and restraint. *Soc Psychiatry Psychiatr Epidemiol*. 2007;42:747–752.
- [22] Wynn R. Medicate, restrain or seclude? Strategies for dealing with violent and threatening behaviour in a Norwegian university psychiatric hospital. *Scand J Caring Sci*. 2002;16:287–291.
- [23] Husum TL, Bjorngaard JH, Finset A, et al. A cross-sectional prospective study of seclusion, restraint and involuntary medication in acute psychiatric wards: patient, staff and ward characteristics. *BMC Health Serv Res*. 2010;10:89.
- [24] Riecher-Rossler A, Rossler W. Compulsory admission of psychiatric patients—an international comparison. *Acta Psychiatr Scand*. 1993;87:231–236.
- [25] Tunde-Ayinmode M, Little J. Use of seclusion in a psychiatric acute inpatient unit. *Australas Psychiatry*. 2004;12:347–351.
- [26] Hadi F, Khosravi T, Shariat SV, Jalali Nadoushan AH. Predictors of physical restraint in a psychiatric emergency setting. *Med J Islam Repub Iran*. 2015;29:296.
- [27] Lorenzo RD, Miani F, Formicola V, et al. Clinical and organizational factors related to the reduction of mechanical restraint application in an acute ward: an 8-year retrospective analysis. *CPEMH*. 2014;10:94–102.
- [28] Hoyer G, Drange H. Use of coercive measures in Norwegian psychiatric institutions. *Tidsskr nor Laegeforen*. 1991;111:1709–1713.
- [29] Hoyer G, Drange H. Changes in the use of compulsory measures in Norwegian psychiatric institutions. *Tidsskr nor Laegeforen*. 1994;114:585–588.
- [30] Andersen K, Nielsen B. Coercion in psychiatry: the importance of extramural factors. *Nord J Psychiatry*. 2016;70:606–610.
- [31] Rossberg JI, Friis S. Staff members' emotional reactions to aggressive and suicidal behavior of inpatients. *Psychiatr Serv*. 2003;54:1388–1394.
- [32] Bauer JO, Stenborg D, Lodahl T, et al. Treatment of agitation in the acute psychiatric setting. An observational study of the effectiveness of intramuscular psychotropic medication. *Nord J Psychiatry*. 2016;70:599–605.
- [33] Morken G, Linaker OM. Seasonal variation of violence in Norway. *Am J Psychiatry*. 2000;157:1674–1678.
- [34] Aguglia A, Moncalvo M, Solia F, et al. Involuntary admissions in Italy: the impact of seasonality. *Int J Psychiatry Clin Pract*. 2016;20:232–238.
- [35] Wang B, Chen D. Evidence for seasonal mania: a review. *J Psychiatr Pract*. 2013;19:301–308.
- [36] Paavola P, Tiihonen J. Seasonal variation of seclusion incidents from violent and suicidal acts in forensic psychiatric patients. *Int J Law Psychiatry*. 2010;33:27–34.