

1 Original Article:

2 **Anthropometry-based obesity phenotypes and risk of colorectal**  
3 **adenocarcinoma: a large prospective cohort study in Norway**

4

5 Yunxia Lu, M.D., Ph.D.<sup>1,2</sup>; Eivind Ness-Jensen, M.D., Ph.D.<sup>1,3</sup>; Anna Martling, M.D.,  
6 Ph.D.<sup>1</sup>; Kristian Hveem, M.D., Ph.D.<sup>1,3</sup>

7

8 **Affiliations:**

9 <sup>1</sup> Department of Molecular Medicine and Surgery, Karolinska Institutet, Stockholm, Sweden.

10 <sup>2</sup> Department of Epidemiology and Biostatistics, Imperial College London, London, United  
11 Kingdom.

12 <sup>3</sup>HUNT Research Centre, Department of Public Health and General Practice, Norwegian  
13 University of Science and Technology, Levanger, Norway.

14

15 **Authorship contributions:** Yunxia Lu and Kristian Hveem: conceptualization and design;  
16 Eivind Ness-Jensen and Kristian Hveem: data collection; Yunxia Lu: data analysis; All:  
17 reporting of results and interpretation of data; Yunxia Lu: drafting of manuscript; All:  
18 contribution to critical comments and revisions; All: final approval of the version to be  
19 published.

20

21

22 ***Acknowledgments:***

23 The authors wish to acknowledge the services of CONOR, the contributing research centers  
24 delivering data to CONOR and all the study participants. Following cohorts from CONOR

25 were used in the analysis: The Tromsø Health Study; The Troms and Finnmark Health Study;  
26 The Nord-Trøndelag Health Study; The Oslo Health Study; The Oppland and Hedmark  
27 Health Study; and The Hordaland Health Study.

28

29 The Nord-Trøndelag Health Study (The HUNT Study) is a collaboration between the HUNT  
30 Research Centre (Faculty of Medicine, Norwegian University of Science and Technology  
31 NTNU), Nord-Trøndelag County Council, Central Norway Health Authority, and the  
32 Norwegian Institute of Public Health”.

33

34 The study has used data from the Cancer Registry of Norway. The interpretation and  
35 reporting of these data are the sole responsibility of the authors, and no endorsement by the  
36 Cancer Registry of Norway is intended nor should be inferred.

37

38 **Conflicts of interest:** No conflicts of interest need to be declared.

39 **Corresponding author:**

40 Yunxia Lu

41 Address:

42 Department of Molecular Medicine and Surgery,

43 Karolinska Institutet, SE-171 76 Stockholm, Sweden.

44 E-mail: yunxia.lu@ki.se

45 Tel: +46-8-51772401

46 Fax: +46-8-51776280

47

48 **Running head:**

49 Obesity and colorectal cancer

50

51 **Category: word count:** (main text: 3700); (abstract: 248)

52

1 **Abstract**

2 **Background:** Whether obesity phenotypes measured by different anthropometric indices are  
3 associated with a risk of colorectal adenocarcinoma by anatomical location is unclear.

4 **Patients and Methods:** A collection of harmonized population-based cohort studies (Cohort  
5 of Norway, CONOR) with 143,477 participants was conducted between 1994 and 2010.

6 General, abdominal, gluteofemoral obesity, and other type were assessed by body mass index  
7 (BMI), waist circumference, hip circumference (HC), and body adiposity index (BAI)  
8 adjusted by BMI or/and waist circumference. Cox proportional hazards regression was  
9 performed to estimate hazard ratios (HRs) and 95% confidence intervals (CIs) of obesity  
10 relative to a risk of colorectal adenocarcinoma.

11  
12 **Results:** In total, 2044 incident cases of colorectal adenocarcinoma were identified. We  
13 observed a positive association between WC ( $\geq 86$ (women) or  $\geq 96$ (men) versus  $< 75$ (women)  
14 or  $< 88$  (men)) and adenocarcinoma in the proximal colon (HR 1.92, 95% CI: 1.47-2.50) and  
15 distal colon(HR 1.71, 95% CI: 1.25-2.33) when adjusted for BMI. The association with WC  
16 was especially evident in men. BMI was not associated with adenocarcinoma in the colon or  
17 rectum after adjusting for WC. No associations were found between HC and colorectal  
18 adenocarcinoma. When adjusted by BMI plus WC, BAI was negatively associated with  
19 adenocarcinoma in the proximal or distal colon

20  
21 **Conclusion:** Abdominal, not general or gluteofemoral obesity, was associated with an  
22 increased risk of adenocarcinoma in the proximal and the distal colon, especially in men.  
23 Muscularity may be negatively associated with risk of adenocarcinoma in the proximal colon.

24 **Key words:** Anthropometrics; Waist circumference; Abdominal obesity; Muscularity;  
25 Adenocarcinoma; Colon; Rectum; CONOR; HUNT

## 26 **Introduction**

27 General obesity (measured using body mass index (BMI)) and abdominal or central obesity  
28 (measured by waist circumference) increases the risk of colorectal cancer.<sup>1-3</sup> The risk of  
29 colorectal cancer associated with obesity is also influenced by sex, age, menopausal status,  
30 and ethnicity.<sup>4-7</sup> However, direct measurements of subcutaneous and visceral obesity by  
31 computer tomography (CT) have shown inconsistent results.<sup>8-10</sup> It is conceivable, therefore,  
32 that the causal relation between obesity and colorectal cancer may not be as simple as  
33 assumed. In addition to different environmental conditions and hereditary factors, the  
34 selection of anthropometric indices to substitute phenotypes of obesity may have profound  
35 effects on the prediction of colorectal cancer risk.

36

37 Several anthropometric indices for the measurement of obesity have been developed and  
38 applied in epidemiological studies. As the most commonly used anthropometric parameter,  
39 BMI is a good index for general obesity, but not sensitive for more specific obese  
40 phenotypes, e.g. abdominal obesity.<sup>11</sup> The latter is widely recognized as the key factor related  
41 to diabetes, cardiovascular diseases, and cancer.<sup>12,13</sup> Waist circumference and waist to hip  
42 ratio have been demonstrated as two important indices for abdominal obesity.<sup>11</sup> Waist to  
43 height ratio ( or called waist to stature ratio) has been associated with cardiovascular diseases  
44 as a new parameter of abdominal obesity, but reports on colorectal cancer are rare.<sup>14,15</sup> Hip  
45 circumference has been suggested as a measurement of gluteofemoral obesity, which has  
46 been negatively associated with a risk of chronic diseases, including cancer.<sup>16</sup> As a newly  
47 developed anthropometric parameter, waist to height index (WHI) was associated with an  
48 increased risk of colorectal cancer in female Japanese subjects, but no further study has been  
49 reported.<sup>17</sup> Another newly introduced anthropometric parameter, body adiposity index (BAI),  
50 has been recognized as an index of estimating percentage of body fat, but the largely

51 inconsistent results achieved with body adiposity index warrant more explorations of this  
52 index.

53

54 Collectively, a series of anthropometric parameters, representing specific obesity phenotypes,  
55 have been developed during the past decades, but few studies have compared these  
56 anthropometric indices and how they are associated differently with colorectal cancer risk by  
57 anatomical location.

58

59 In the present study, the association between different anthropometric indices for obesity and  
60 colorectal adenocarcinoma by anatomical location were investigated in a large, prospective,  
61 population-based cohort study in Norway: the CONOR study. Since adenocarcinoma is the  
62 dominating histological type (more than 90%) and different histological types of colorectal  
63 cancer may entail different causality, only the risk of adenocarcinoma has been assessed in  
64 the current study.

65

## 66 ***Materials and Methods***

### 67 ***Study population***

68 Detailed information on the design of, and data collection in, the CONOR study has been  
69 described previously.<sup>18</sup> Briefly, CONOR was performed in collaboration between the  
70 Norwegian Institute of Public Health and the Universities of Bergen, Oslo, Tromsø, and  
71 Trondheim (NTNU). Data from 10 regional epidemiological studies were merged into a  
72 national database to study risk factors for a wide range of diseases. In total, 180,553  
73 participants from 10 epidemiology studies were included in the CONOR study.<sup>19</sup> After  
74 excluding repeated participants (7310 with two follow-ups), prevalent cancer cases (906),  
75 individuals who died or migrated before the baseline survey (6075), missing waist

76 circumference, hip circumference, height or weight data (21234), and missing smoking data  
77 (1551), a total of 143,477 participants remained for the final analysis. Anthropometric data  
78 were harmonized throughout all the studies based on common questionnaires/similar clinical  
79 measurements.

80

### 81 ***Follow-up and identification of colorectal cancer cases***

82 The CONOR cohort was followed-up based on linkage to the Norwegian Cancer Register  
83 (NCR) and Statistics Norway, using the unique 11-digit national identity number of  
84 Norwegian citizens. Colorectal cancer was registered in the NCR according to the  
85 International Classification of Diseases, 7th edition (ICD-7). The ICD-7 codes were used to  
86 identify the colorectal cancer cases by anatomical location, including: the proximal colon  
87 (ICD-7 codes 1530, 1531, and 1536, including the cecum, ascending colon, transverse colon,  
88 hepatic flexure, the splenic flexure and appendix); the distal colon (ICD-7 codes 1532 and  
89 1533, including the descending colon, the sigmoid colon); the rectum(ICD-7 code 1540,  
90 including the rectum and rectosigmoid junction). The participants were enrolled into the  
91 cohort at the baseline until diagnosis of colorectal cancer, death, censored (i.e. lost to follow-  
92 up, emigration or diagnosis of other malignancies), or end of follow-up on December 31,  
93 2010, whichever occurred first.

94

### 95 ***Assessment of anthropometric data***

96 Body weight (in kilograms(kg), to one decimal place) and height (in centimeters(cm), to one  
97 decimal place) were manually recorded until the year 2000 and thereafter an electronic height  
98 and weight scale was used. BMI was calculated as body weight (kg) divided by the square of  
99 height(meters square). Waist circumference was measured at the umbilicus to the nearest  
100 centimeter and with the subject standing and breathing normally. Hip circumference was

101 measured as the maximum circumference around the buttocks. Waist to hip ratio and waist to  
102 height ratio was calculated from measurements of waist circumference, hip circumference or  
103 height. Waist to height index was calculated by the formula of waist circumference  
104 (cm)/height (m)/height (m).<sup>17</sup> The body adiposity index was computed by the formula of (hip  
105 circumference(cm)/height(meter)<sup>1.5</sup>)-18.

106

107 We examined each obesity phenotype with one specific anthropometric index. BMI was used  
108 for general obesity, waist circumference for abdominal obesity, hip circumference for  
109 gluteofemoral obesity, and body adiposity index for one uncertain type. Due to the limited  
110 space of the manuscript and also in order to complement the results using other related  
111 indices, the results of waist to hip ratio, waist to height ratio and waist to height index were  
112 further showed in supplemental tables.

113

114 Other data collected at the baseline survey included: marital status, country of birth, years of  
115 education, smoking, alcohol consumption, physical activity, anti-hypertensive drug use, and  
116 self-reported diabetes.

117

### 118 *Statistical analysis*

119 Hazard ratios (HRs) and 95% confidence intervals (95% CIs) for the association between the  
120 anthropometric indices and colorectal cancer were estimated using Cox proportional hazard  
121 models. BMI was grouped into four categories (<22.5, 22.5-25, 25-30, >30 kg/m<sup>2</sup>). The  
122 categorization of BMI was slightly different from the WHO standardization because of small  
123 size of cohort members in the group of BMI less than 18.5. Waist circumference was divided  
124 into three categories based on sex-specific cut-offs (women: <75, 75-85.9, ≥86; men <88, 88-  
125 95.9, ≥96 cm). Hip circumference was categorized into two groups (<101 cm and ≥101 cm),

126 and waist to height ratio three groups ( $<0.5$ ,  $0.5-$  and  $\geq 0.6$ ). Other anthropometric data were  
127 analyzed based on continuous variables. Waist to hip ratio and waist to height ratio were  
128 multiplied by 10 in the model to decrease the significant fluctuation of the small values, and  
129 are interpreted as 1/10 change. WHI was divided by 10 and was interpreted as a per 10 units  
130 increase, while body adiposity index was divided by 5 and interpreted as a per 5 units  
131 increase.

132

133 Analyses of BMI were conducted with and without adjustment for waist circumference. The  
134 analyses of waist circumference, hip circumference, waist to hip ratio, waist to height ratio,  
135 and waist to height index were performed both with and without inclusion of BMI in the  
136 models.<sup>20</sup> Body adiposity index was analyzed with adjustment for BMI or/and waist  
137 circumference. An interaction between sex and anthropometric indices (BMI, waist  
138 circumference, waist to height ratio, and waist to height index) was found. Therefore, further  
139 sex-stratified analyses of anthropometric measurements were performed. P values for trend  
140 were computed based on continuous variables of median values of categories of BMI, waist  
141 circumference, or waist to height ratio.

142

143 Compared with weight and height, waist circumference and hip circumference had a  
144 significant number of missing values (20,902 in total) because both were not measured in  
145 1994, the first round of the survey. We analyzed the data based on three approaches. First, we  
146 removed all of the participants with missing waist circumference or hip circumference data.  
147 Second, we imputed waist circumference based on a sex-specific model adjusted for age, sex,  
148 smoking, alcohol drinking, education, physical activity, height and weight. Third, we  
149 analyzed the data when using missing waist circumference as a separate category. Since the



150 overall results were not changed materially, we kept results based on the first approach in the  
151 main report.

152 For each anthropometric indicator, we analyzed data based on a crude model adjusted for age  
153 and sex and a multivariable model adjusted for all potential confounders, but we only  
154 reported the results based on multivariable models because the overall results were not  
155 changed. We selected confounders based on previous etiological studies on colorectal cancer  
156 together with stepwise selection approaches. The following co-variables were included in the  
157 multivariable model: age (<50, 50-60, ≥60), education (none/primary school/secondary  
158 school, high school, university), currently daily smoking (yes, no), alcohol drinking  
159 (never/seldom, several times per week, about once a week, 2-3 times per month, about once a  
160 month), physical activity (none, <1, 1-2, ≥3 hours/week). There are approximately 10% to  
161 20% missing values for education, alcohol drinking and physical activity. We treated the  
162 missing values as a separate category or deleted them from the total dataset for analyses.  
163 Since the final results did not materially alter, we only included the results based on missing  
164 values as separate categories in order to keep as many participants and colorectal cancer cases  
165 as possible for the whole study. Furthermore, we excluded the first two years of follow-up in  
166 order to decrease the potential bias of reverse causality; the results were similar and are not  
167 shown.

168

169 The proportional hazards assumption was tested on the basis of Schoenfeld residuals  
170 afterfitting a Cox regression model. None of the variables violated the assumption except for  
171 the age groups. The age groups were, thus, treated as a strata factor in the model. A two-sided  
172 test with a significance level ( $\alpha$ ) of 0.05 was chosen. All analyses were performed using SAS  
173 9.3 for Windows (SAS Institute Inc., Cary, NC, USA).

174

175 ***Ethics***

176 The present study was approved by the Regional Committee for Medical and Health  
177 Research Ethics, Central Norway (ID: 2012/853/REK midt). The individual studies included  
178 in CONOR were all approved by their respective ethics committees. All participants signed  
179 an informed consent form.

180

181 ***Results***

182 ***Basic characteristics***

183 During an average of 11.3 years of follow-up, 2044 incident cases of colorectal  
184 adenocarcinoma(853 in the proximal colon, 606 in the distal colon and 555 in the rectum, 30  
185 cases with specified locations) were identified. Of these cases, 1101 (54 %) were men and  
186 943 (46 %) were women (Table 1). The average age at study entry was 64.5 years for cases  
187 and 50.9 years for non-cases. Cases were less educated (35.3% of cases versus 22.6% of the  
188 total cohort in the lowest education category), had more family history of cancer (32.9%  
189 versus 25.3%), less physical activity (10.9% versus 6.3% for 3 or more hours per week) and  
190 alcohol drinking (13% versus 12% for drinking alcohol several times per week), whereas  
191 daily smoking seemed to be more common in the total cohort members (Table 1).

192

193 ***General obesity (body mass index, BMI) and colorectal adenocarcinoma***

194 The highest BMI category (BMI>30) was associated with colorectal adenocarcinoma when  
195 the multivariable models were not adjusted for waist circumference (HR 1.17, 95% CI: 1.02-  
196 1.34), but the association disappeared when the models were adjusted for waist circumference  
197 (HR 0.90, 95% CI: 0.76-1.06)(Table 2). The risk estimates were similar for each anatomical  
198 location within the colon and rectum (Table 2). Interestingly, a negative association of BMI  
199 with proximal colon adenocarcinoma was observed when adjusted for waist circumference

200 (HR 0.77, 95%CI: 0.59-0.99, Table 2). This association was attenuated in the sex-stratified  
201 analyses but still existed, especially in women (Table 3).

202

### 203 ***Abdominal obesity (waist circumference) and colorectal adenocarcinoma***

204 Waist circumference(cm,  $\geq 86$ (women) or  $\geq 96$ (men) versus  $< 75$ (women) or  $< 88$  (men))was  
205 positively associated with adenocarcinoma of the proximal and the distal colon (HR 1.51,  
206 95%CI: 1.24-1.83 and HR 1.48, 95% CI: 1.18-1.86, respectively), and the association became  
207 stronger when the model was adjusted for BMI (HR 1.92, 95% CI: 1.47-2.50 and HR 1.71,  
208 95% CI: 1.25-2.33, respectively)(Table 2). For the rectum, no association was observed (HR  
209 1.16, 95% CI: 0.93-1.46; HR 1.12, 95% CI: 0.82-1.54, with or without adjustment for BMI,  
210 respectively).The positive association between waist circumference and adenocarcinoma of  
211 the proximal and distal colon was evident in both sexes, especially in men (Table 3).A  
212 positive association was further observed for rectal adenocarcinoma in women (HR 2.07,  
213 95%: 1.17-3.68) (Table 3).

214

### 215 ***Gluteofemoral obesity (hip circumference) and colorectal adenocarcinoma***

216 Positive associations were found between HC( $< 101$  cm versus  $\geq 101$ ) and adenocarcinoma in  
217 the proximal and the distal colon (HR 1.23, 95% CI: 1.07-1.42; HR 1.19, 95%CI: 1.01-1.40,  
218 respectively), but not in the rectum (HR 1.03, 95% CI: 0.87-1.22) (Table 2).These  
219 associations were more evident in men (Table 3), but disappeared with adjustments for BMI  
220 plus waist circumference (Table 2).

221

### 222 ***Body adiposity index (BAI) and colorectal cancer***

223 Body adiposity index was not associated with colorectal adenocarcinoma (HR 0.98, 95%CI:  
224 0.93-1.04)(Table 2). However, when the analyses were further adjusted for BMI or BMI plus

225 waist circumference, negative associations were observed for adenocarcinoma of the  
226 proximal and the distal colon (adjustment for BMI, HR 0.88, 95%CI: 0.78-0.99; HR 0.81,  
227 95%CI: 0.70-0.94) (Table 2).

228

### 229 ***BMI and waist circumference***

230 All of the results in this section were analyzed based on a comparison with the normal BMI  
231 (22.5-25kg/m<sup>2</sup>) and lower waist circumference category (women <80cm, men <94cm). In the  
232 low BMI category (<22.5 kg/m<sup>2</sup>), higher waist circumference indicated an increased risk of  
233 colorectal adenocarcinoma, especially in the proximal colon and rectum. The latter was a  
234 surprisingly increased HR which was not observed in the aforementioned analyses (HR 2.37;  
235 95%CI: 1.09-5.12), however, only seven cases of rectal adenocarcinoma were identified in  
236 this group. In the normal BMI group, HR are 1.31 (95%CI: 0.95-1.80) and 1.44 (95%CI:  
237 0.97-2.14) for proximal colon and distal colon respectively when a higher waist  
238 circumference compared to the lower category. (Table 5). In the overweight group (BMI 25-  
239 30kg/m<sup>2</sup>), a higher waist circumference displayed an elevated risk of colon adenocarcinoma,  
240 especially in the distal colon, but not the rectum. Similar high circumference results can be  
241 found in the obese group (BMI≥30 kg/m<sup>2</sup>) (Table 5), but a lower circumference may still  
242 entail an increased risk of adenocarcinoma in the proximal colon (HR), although the results  
243 were not statistically significant due to too few cases.

244

### 245 ***Other anthropometric indices and colorectal adenocarcinoma***

246 Waist to hip ratio (per 1/10 increase) was positively associated with adenocarcinoma in the  
247 proximal and distal colon (HR 1.28, 95%CI: 1.16-1.42 and HR 1.20, 95%CI:1.06-1.36,  
248 respectively), and the associations remained almost similar when adjusted for BMI

249 (Supplemental Table 1). In the sex-stratified analyses, positive associations remained for the  
250 proximal colon in both sexes(Supplemental Table 1).

251

252 Waist to height ratio (per 1/10 increase) was positively associated with adenocarcinoma in  
253 the proximal colon regardless of adjustment for BMI (HR 1.18, 95%CI: 1.06-1.31; HR 1.26,  
254 95%CI: 1.07-1.49, with or without adjustment for BMI) (Supplemental Table 1). This  
255 association remained for categorical variables of Waist to height ratio (Supplemental Table  
256 1). No association was observed for adenocarcinoma in the rectum, regardless of adjustment  
257 for BMI (Table 2). In the sex-stratified analysis, a persistent association with colon  
258 adenocarcinoma was observed in men, especially in the proximal colon(Supplemental Table  
259 1).

260

261 WHI was associated with adenocarcinoma in the proximal colon (HR1.18, 95%:1.01-1.39)  
262 but this association disappeared when adjusted for BM I (Supplemental Table 1). Similar  
263 results could be found in the sex-stratified analysis (Supplemental Table 2).

264

## 265 ***Discussion***

266 Abdominal obesity, represented by waist circumference, waist to hip ratio, or waist to height  
267 ratio, seemed to be the most important obesity phenotype that had the strongest association  
268 with adenocarcinoma in the proximal and the distal colon, but no association with  
269 adenocarcinoma in the rectum. General obesity, represented by body mass index (BMI),  
270 seemed to not be associated with colorectal adenocarcinoma when adjusted for abdominal  
271 obesity. While gluteofemoral obesity, represented by hip circumference, was not associated  
272 with colorectal adenocarcinoma.

273

274 The strengths of the current study included the large population-based cohort design with a  
275 long follow-up period, where anthropometric measures were objectively assessed by standard  
276 protocols rather than being self-reported. Potential confounders such as smoking, consuming  
277 alcohol, education, and physical activity were adjusted for as well. The Norwegian Cancer  
278 Register and Statistics Norway provided outcomes of cancer and death with a high validity.  
279 Weaknesses of the study included the possibility of residual confounding produced by  
280 missing information of nutrients/diet. For missing values, we performed sensitivity analyses  
281 based on imputation, deletion, or treating as a separate category. The overall results, however,  
282 were consistent and conclusions were not changed. We also realize that the anthropometric  
283 measures of abdominal obesity may not separate visceral obesity from subcutaneous fat.  
284 Each of them probably have different effects on cancer incidence, while visceral fat may be  
285 worse. Nevertheless, the mutual adjustment of waist circumference and BAI may provide  
286 more evidence for this issue.

287

288 General and abdominal obesity have been associated with colorectal cancer in many studies.  
289 In a large European cohort study, obesity was associated with a higher relative risk of cancer  
290 in the colon than cancer of the rectum<sup>20</sup>. This is consistent with our results. However, whether  
291 general or abdominal obesity played the leading role was not clear in the previous studies.<sup>20,21</sup>  
292 In our study, abdominal adiposity (mainly determined by waist circumference, waist to hip  
293 ratio or waist to height ratio) was statistically associated with colon cancer especially in men  
294 irrespective of BMI. On the other hand, BMI was not associated with colon cancer when  
295 adjusted for waist circumference. This suggests that abdominal adiposity is a more important  
296 risk factor for colon cancer than general adiposity. However, as Hu et al. pointed out, in a  
297 disease model with waist circumference and BMI, waist circumference would still reflect  
298 abdominal adiposity, but BMI would probably be more a measure of lean body mass since

299 body fatness is to a large extent accounted for by waist circumference, especially in older  
300 adults<sup>11</sup>. This might well explain the negative association of BMI with the proximal colon  
301 adenocarcinoma when the analysis was adjusted for waist circumference. On the other hand,  
302 for a given BMI, individuals with an elevated waist circumference will likely have more  
303 abdominal fat and, thus, more visceral, liver, and ectopic fat and therefore a higher risk of  
304 obesity-related metabolic disorders. In the sex-stratified analyses, a positive association of  
305 waist circumference with adenocarcinoma in the proximal and distal colon persisted,  
306 especially in men. This is consistent with the study from a Chinese cohort.<sup>22</sup> Surprisingly, a  
307 strongly positive association of waist circumference with rectal adenocarcinoma in women  
308 was observed. A positive, but not strong association, was also observed for waist to hip ratio  
309 or waist to height ratio with female rectal adenocarcinoma. As this has been rarely reported in  
310 previous studies, further evidence is warranted. When we examined the risk of colorectal  
311 adenocarcinoma for a given BMI and waist circumference, we found a consistently increased  
312 risk of abdominal obesity represented by a higher waist circumference regardless of lower or  
313 normal BMI, overweight, or obesity. This further strengthened our conclusion regarding the  
314 pivotal role of abdominal obesity on adenocarcinoma in the proximal and distal colon.

315

316 There is increasing evidence that the anatomical position of adipose tissue determines the  
317 effects on the individual and predicts the associated morbidity from cancer.<sup>16,23</sup> This has led to  
318 the addition of the new anthropometric indices of obesity in addition to BMI and waist  
319 circumference. Gluteofemoral body fat, assessed by hip circumference, is associated with a  
320 protective lipid and glucose profile, as well as a decrease in cardiovascular and metabolic  
321 risk.<sup>24</sup> However, the association between hip circumference and cancer is inconclusive.<sup>25,26</sup>  
322 The present study found that an increasing hip circumference was associated with an  
323 increased risk of adenocarcinoma of the colon, while this association disappeared with

324 additional adjustments for BMI and waist circumference. This indicates that gluteofemoral  
325 obesity is not associated with colorectal cancer. Furthermore, since BMI cannot reflect the  
326 percentage of body fat (e.g., with a high BMI may be due to lean muscular mass rather than  
327 body fat), the body adiposity index has therefore been purposely developed. In our study,  
328 body adiposity index was negatively associated with adenocarcinoma in the proximal and the  
329 distal colon when adjusted for BMI or BMI plus waist circumference. This seemed to be  
330 similar to the association between BMI and colorectal adenocarcinoma. If this reflects the  
331 true association, body adiposity index may indicate the percentage of muscular mass for a  
332 given BMI or BMI plus waist circumference, because BMI or waist circumference has  
333 represented the body fat. Since body adiposity index was still an anthropometric index under  
334 debate, this result may need further validating studies.

335

336 Other anthropometric indices have also been investigated in previous studies. Waist to height  
337 ratio is associated with cancer and cardio-metabolic risks.<sup>27,28</sup> In a study among Taiwanese  
338 adults, a Waist to height ratio >0.5 was an indicator of centralized obesity, even among  
339 'healthy' individuals according to BMI and waist circumference.<sup>29</sup> A recent systemic review  
340 and meta-analysis also demonstrated that waist to height ratio is a better screening tool for  
341 adult cardiometabolic risk factors than waist circumference and BMI.<sup>27</sup> As far as we know,  
342 no data has been reported about the association between waist to height ratio and colorectal  
343 cancer. In our study, we found a consistent association between waist to height ratio and an  
344 increased risk of adenocarcinoma of the proximal and distal colon, but not the rectum.  
345 Moreover, waist to height ratio is a simple and easily understood anthropometric index,  
346 which may carry much valuable public health implications, e.g., waist to height ratio may  
347 allow the same boundary values (0.5) for children and adults, women and men.<sup>30</sup> As a newly  
348 developed index, waist to height index (WHI is a composite index proposed by a Japanese



349 study.<sup>17</sup> In our study, however, waist to height index (per 10 units of increase) was not  
350 associated with an increased risk of colorectal adenocarcinoma.

351

352 Abdominal fat is comprised of fat stored subcutaneously (e.g., subcutaneous adipose tissue)  
353 and the adipose tissue located in the abdominal cavity. The latter has been commonly  
354 described as intra-abdominal or visceral adipose tissue. Visceral adiposity is the best  
355 adiposity predictor of liver fat content which is closely related to features of metabolic  
356 syndrome. Abdominal obesity, metabolic syndrome, insulin resistance and modifications in  
357 levels of adipocytokines seem to be of great importance for the underlying mechanisms  
358 linking obesity to colorectal cancer, which is certainly a multifactorial process.<sup>3</sup> Adipose  
359 tissue is a highly active tissue that secretes various cytokines, chemokines, and hormones.<sup>31</sup>  
360 Some of these cytokines can act directly in the promotion of cancer.<sup>13,32</sup> Circulating insulin  
361 levels increase with obesity and many obese patients are insulin resistant. Chronic  
362 hyperinsulinemia decreases insulin growth factor binding proteins 1 and 2, resulting in an  
363 increase in circulating insulin and, more importantly, insulin growth factor. This in turn  
364 results in decreased apoptosis and increased cell proliferation in the target tissues.<sup>23</sup> Studies  
365 have shown a correlation of an elevated C-peptide (a surrogate of circulating insulin) with  
366 colorectal cancer.<sup>32</sup> Furthermore, adipokines (leptin and adiponectin) secreted by adipose  
367 tissue have been associated with carcinogenesis.<sup>33,34</sup> Leptin is a pro-inflammatory hormone  
368 and has also been shown to be directly tumorigenic. Adiponectin levels, which are decreased  
369 in obese individuals, are associated with a lower risk of colorectal cancer.<sup>33,34</sup> Further  
370 carcinogenesis has been proposed for obesity-driven low-grade inflammation. Inflammatory  
371 cells are present in abundance in visceral adipose tissue and the secretion of inflammatory  
372 mediators into the body creates a chronic inflammatory state that is thought to generate a pro-

373 tumorigenic environment.<sup>31</sup> Systemic pro-inflammatory markers, such as C-reactive protein  
374 and interleukin-6, are elevated in the obese due to obesity-driven low-grade inflammation.<sup>13</sup>

375

376 In conclusion, obesity, especially abdominal obesity, was positively associated with  
377 adenocarcinoma of the proximal and distal colon, but not certainly with the rectum. Obesity  
378 control, with a focus on abdominal obesity, will be an important factor in the prevention of  
379 malignancy of the colon.

380

## 381 **References**

382

- 383 1. Moghaddam AA, Woodward M, Huxley R. Obesity and risk of colorectal cancer: a  
384 meta-analysis of 31 studies with 70,000 events. *Cancer epidemiology, biomarkers &*  
385 *prevention : a publication of the American Association for Cancer Research,*  
386 *cosponsored by the American Society of Preventive Oncology.* Dec 2007;16(12):2533-  
387 2547.
- 388 2. Matsuo K, Mizoue T, Tanaka K, et al. Association between body mass index and the  
389 colorectal cancer risk in Japan: pooled analysis of population-based cohort studies in  
390 Japan. *Annals of oncology : official journal of the European Society for Medical*  
391 *Oncology / ESMO.* Feb 2012;23(2):479-490.
- 392 3. Bardou M, Barkun AN, Martel M. Obesity and colorectal cancer. *Gut.* Jun  
393 2013;62(6):933-947.
- 394 4. MacInnis RJ, English DR, Hopper JL, Gertig DM, Haydon AM, Giles GG. Body size  
395 and composition and colon cancer risk in women. *International journal of cancer.*  
396 *Journal international du cancer.* Mar 15 2006;118(6):1496-1500.
- 397 5. Giovannucci E. Obesity, gender, and colon cancer. *Gut.* Aug 2002;51(2):147.
- 398 6. Wise LA, Rosenberg L, Palmer JR, Adams-Campbell LL. Anthropometric risk factors  
399 for colorectal polyps in African-American women. *Obesity.* Apr 2008;16(4):859-868.
- 400 7. Thompson CL, Berger NA, Chak A, Li L. Racial differences in measures of obesity  
401 and risk of colon adenoma. *Obesity.* Mar 2012;20(3):673-677.
- 402 8. Nagata N, Sakamoto K, Arai T, et al. Visceral abdominal fat measured by computed  
403 tomography is associated with an increased risk of colorectal adenoma. *International*  
404 *journal of cancer. Journal international du cancer.* Nov 15 2014;135(10):2273-2281.
- 405 9. Choe EK, Kim D, Kim HJ, Park KJ. Association of visceral obesity and early  
406 colorectal neoplasia. *World journal of gastroenterology : WJG.* Dec 7  
407 2013;19(45):8349-8356.
- 408 10. Summers RM, Liu J, Sussman DL, et al. Association between visceral adiposity and  
409 colorectal polyps on CT colonography. *AJR. American journal of roentgenology.* Jul  
410 2012;199(1):48-57.
- 411 11. Hu F. Obesity Epidemiology: Methods and Applications. *Oxford University Press*  
412 *Inc.* 2008.

- 413 12. Zhang C, Rexrode KM, van Dam RM, Li TY, Hu FB. Abdominal obesity and the risk  
414 of all-cause, cardiovascular, and cancer mortality: sixteen years of follow-up in US  
415 women. *Circulation*. Apr 1 2008;117(13):1658-1667.
- 416 13. Vazzana N, Riondino S, Toto V, et al. Obesity-driven inflammation and colorectal  
417 cancer. *Current medicinal chemistry*. 2012;19(34):5837-5853.
- 418 14. Kahn HS, El Ghormli L, Jago R, et al. Cardiometabolic Risk Assessments by Body  
419 Mass Index z-Score or Waist-to-Height Ratio in a Multiethnic Sample of Sixth-  
420 Graders. *Journal of obesity*. 2014;2014:421658.
- 421 15. Savva SC, Lamnisos D, Kafatos AG. Predicting cardiometabolic risk: waist-to-height  
422 ratio or BMI. A meta-analysis. *Diabetes, metabolic syndrome and obesity : targets  
423 and therapy*. 2013;6:403-419.
- 424 16. Lee MJ, Wu Y, Fried SK. Adipose tissue heterogeneity: implication of depot  
425 differences in adipose tissue for obesity complications. *Molecular aspects of  
426 medicine*. Feb 2013;34(1):1-11.
- 427 17. Kaneko R, Nakazaki N, Tagawa T, et al. A new index of abdominal obesity which  
428 effectively predicts risk of colon tumor development in female Japanese. *Asian  
429 Pacific journal of cancer prevention : APJCP*. 2014;15(2):1005-1010.
- 430 18. Naess O, Sogaard AJ, Arnesen E, et al. Cohort profile: cohort of Norway (CONOR).  
431 *International journal of epidemiology*. Jun 2008;37(3):481-485.
- 432 19. Hammer H. [The central population registry in medical research]. *Tidsskrift for den  
433 Norske laegeforening : tidsskrift for praktisk medicin, ny raekke*. Oct 30  
434 2002;122(26):2550.
- 435 20. Pischon T, Lahmann PH, Boeing H, et al. Body size and risk of colon and rectal  
436 cancer in the European Prospective Investigation Into Cancer and Nutrition (EPIC).  
437 *Journal of the National Cancer Institute*. Jul 5 2006;98(13):920-931.
- 438 21. Keimling M, Renehan AG, Behrens G, et al. Comparison of associations of body  
439 mass index, abdominal adiposity, and risk of colorectal cancer in a large prospective  
440 cohort study. *Cancer epidemiology, biomarkers & prevention : a publication of the  
441 American Association for Cancer Research, cosponsored by the American Society of  
442 Preventive Oncology*. Aug 2013;22(8):1383-1394.
- 443 22. Li H, Yang G, Xiang YB, et al. Body weight, fat distribution and colorectal cancer  
444 risk: a report from cohort studies of 134 255 Chinese men and women. *International  
445 journal of obesity*. Sep 18 2012.
- 446 23. Westley RL, May FE. A twenty-first century cancer epidemic caused by obesity: the  
447 involvement of insulin, diabetes, and insulin-like growth factors. *International journal  
448 of endocrinology*. 2013;2013:632461.
- 449 24. Cameron AJ, Magliano DJ, Soderberg S. A systematic review of the impact of  
450 including both waist and hip circumference in risk models for cardiovascular diseases,  
451 diabetes and mortality. *Obesity reviews : an official journal of the International  
452 Association for the Study of Obesity*. Jan 2013;14(1):86-94.
- 453 25. Fagherazzi G, Chabbert-Buffet N, Fabre A, et al. Hip circumference is associated with  
454 the risk of premenopausal ER-/PR- breast cancer. *International journal of obesity*.  
455 Mar 2012;36(3):431-439.
- 456 26. Rubenstein JH, Morgenstern H, Chey WD, et al. Protective role of gluteofemoral  
457 obesity in erosive oesophagitis and Barrett's oesophagus. *Gut*. Feb 2014;63(2):230-  
458 235.
- 459 27. Ashwell M, Gunn P, Gibson S. Waist-to-height ratio is a better screening tool than  
460 waist circumference and BMI for adult cardiometabolic risk factors: systematic  
461 review and meta-analysis. *Obesity reviews : an official journal of the International  
462 Association for the Study of Obesity*. Mar 2012;13(3):275-286.

- 463 28. Hsieh SD, Yoshinaga H, Muto T. Waist-to-height ratio, a simple and practical index  
464 for assessing central fat distribution and metabolic risk in Japanese men and women.  
465 *International journal of obesity and related metabolic disorders : journal of the*  
466 *International Association for the Study of Obesity*. May 2003;27(5):610-616.
- 467 29. Li WC, Chen IC, Chang YC, Loke SS, Wang SH, Hsiao KY. Waist-to-height ratio,  
468 waist circumference, and body mass index as indices of cardiometabolic risk among  
469 36,642 Taiwanese adults. *European journal of nutrition*. Feb 2013;52(1):57-65.
- 470 30. Ashwell M, Hsieh SD. Six reasons why the waist-to-height ratio is a rapid and  
471 effective global indicator for health risks of obesity and how its use could simplify the  
472 international public health message on obesity. *International journal of food sciences*  
473 *and nutrition*. Aug 2005;56(5):303-307.
- 474 31. Doyle SL, Donohoe CL, Lysaght J, Reynolds JV. Visceral obesity, metabolic  
475 syndrome, insulin resistance and cancer. *The Proceedings of the Nutrition Society*.  
476 Feb 2012;71(1):181-189.
- 477 32. Kaaks R, Toniolo P, Akhmedkhanov A, et al. Serum C-peptide, insulin-like growth  
478 factor (IGF)-I, IGF-binding proteins, and colorectal cancer risk in women. *Journal of*  
479 *the National Cancer Institute*. Oct 4 2000;92(19):1592-1600.
- 480 33. Roberts DL, Dive C, Renehan AG. Biological mechanisms linking obesity and cancer  
481 risk: new perspectives. *Annual review of medicine*. 2010;61:301-316.
- 482 34. Tilg H, Moschen AR. Mechanisms behind the link between obesity and  
483 gastrointestinal cancers. *Best practice & research. Clinical gastroenterology*. Aug  
484 2014;28(4):599-610.
- 485  
486

**Table 1. Characteristics of colorectal adenocarcinoma cases and cohort members in CONOR**

Variables	Cohort participants	Colorectal Adenocarcinoma	Colon		Rectum
			Proximal colon	Distal colon	
<b>Total</b>	143477	2044	853	606	555
Sex, n(%)					
Men	70033(49)	1101(54)	410(48)	332(55)	343(62)
Women	73444(51)	943(46)	443(52)	274(45)	212(38)
<b>Age at examination</b>					
Mean (SD <sup>a</sup> )	50.9(16)	64.5(12)	65.8(11)	63.4(12)	63.5(12)
<b>Age by groups (%)</b>					
<50	81232(57)	341( 17)	119(14)	117(19)	103(19)
50-59	17559(12)	269(13)	96(11)	91(15)	80(14)
≥60	44686(31)	1434(70)	638(75)	398(66)	372(67)
<b>Education, n(%)</b>					
None/primary school/secondary school	32423(23)	724(35)	320(38)	198(33)	198(36)
High school	44964(31)	468(23)	193(22)	136(22)	134(24)
University	29227(20)	237(12)	88(10)	94(16)	51(9)
Missing	36863(26)	615(30)	252(30)	178(29)	172(31)
<b>Smoking status, n(%)</b>					
Not daily smoker	101341(71)	1542(75)	653(77)	461(76)	402(72)

Daily smoker	42136(29)	502(25)	200(23)	145(24)	153(28)
<b>Alcohol consumption last year, n(%)</b>					
Never/seldom	41694(29)	690(34)	316(37)	182(30)	185(33)
About 1-3 times per month	45233(32)	465(23)	174(20)	149(25)	135(24)
About once a week	26106(18)	331(16)	123(14)	116(19)	89(16)
Several times per week	17187(12)	265(13)	104(13)	77(13)	80(14)
Missing	13257(9)	293(14)	136(16)	82(13)	66(13)
<b>Physical activity, n(%)</b>					
None	43492(30)	720(35)	318(37)	225(37)	167(30)
Less than once a week	30222(21)	342(17)	117(14)	108(18)	113(20)
1-2 hours per week	28226(20)	284(14)	113(13)	86(14)	84(15)
3 or more hours per week	15581(11)	129(6)	49(6)	41(7)	36(7)
Missing	25956(18)	569(28)	256(30)	146(24)	155(28)
<sup>b</sup> Family history of cancer, n(%)	36309(25)	672(33)	385(33)	211(35)	169(31)
Diabetes, n(%)	4463(4)	122(6)	57(7)	31(5)	33(6)
<sup>c</sup> Cardiovascular diseases, n(%)	11373(8)	301(15)	137(16)	78(13)	81(15)
Asthma, n(%)	12087(8)	210(10)	99(12)	62(10)	46(8)
Body mass index (BMI), mean (SD)	26.2(4.1)	27.0(4.1)	27.1(4.1)	27.0(4.0)	27.0(4.1)
Waist circumference (cm), mean (SD)	86.9(12.1)	90.7(11.8)	90.6(12.2)	90.6(11.5)	91.0(11.5)
Hip circumference (cm), mean (SD)	101.6(7.8)	102.7(7.6)	102.9(7.8)	102.8(7.4)	102.5(7.5)
Waist to hip ratio*10 (WHR), mean (SD)	8.5(0.9)	8.8(0.9)	8.8(0.9)	8.8(0.9)	8.9(0.8)

Waist to height ratio*10 (WHtR), mean (SD)	5.1(0.7)	5.3(0.7)	5.4(0.7)	5.3(0.6)	5.3(0.6)
Waist to height index, (WHI) mean (SD)	30.1(4.5)	31.6(4.4)	31.9(4.5)	31.3(4.2)	31.4(4.3)
Body adiposity index (BAI), mean (SD)	27.9(4.9)	28.4(4.7)	29.2(5.1)	28.4(4.7)	28.3(5.0)

---

a: SD, standard deviation

b. Family history of cancer: self-reported cancer among parents, siblings and children

c. Cardiovascular diseases: including angina pectoris, myocardial infarction and stroke.

**Table 2. HR and 95% CI for risk of colorectal adenocarcinoma in relation to anthropometric characteristics**

Anthropometric indices	Colorectal adenocarcinoma by anatomical location <sup>a</sup>							
	Total	Proximal colon	Distal colon	Rectum				
	HR(95%CI)	HR(95%CI)	HR(95%CI)	HR(95%CI)				
<b>Total</b>	2044	853	606	555				
<b>Overall obesity: Body mass index (BMI)</b>								
<b>Not adjusted for waist circumference</b>								
<22.5	234	0.97(0.82-1.14)	97	0.95(0.74-1.21)	72	1.06(0.79-1.42)	60	0.87(0.64-1.19)
22.5-25	435	Reference	181	Reference	122	Reference	127	Reference
25-30	942	1.03(0.92-1.15)	389	1.01(0.85-1.21)	286	1.13(0.91-1.39)	251	0.94(0.76-1.16)
>30	433	1.17(1.02-1.34)	186	1.15(0.93-1.41)	126	1.26(0.98-1.62)	117	1.14(0.88-1.47)
<b>P value for trend</b>		0.008		0.11		0.10		0.13
<b>Adjusted for waist circumference</b>								
<22.5	234	1.08(0.91-1.28)	97	1.10(0.84-1.43)	72	1.23(0.90-1.68)	60	0.91(0.66-1.26)
22.5-25	435	Reference	181	Reference	122	Reference	127	Reference
25-30	942	0.88(0.77-1.00)	389	0.80(0.65-0.98)	286	0.94(0.74-1.19)	251	0.90(0.71-1.15)
>30	433	0.90(0.76-1.06)	186	0.77(0.59-0.99)	126	0.94(0.69-1.29)	117	1.07(0.78-1.49)
<b>P value for trend</b>		0.17		0.04		0.37		0.41
<b>Abdominal obesity: Waist Circumference (cm)</b>								
<b>Not adjusted for BMI</b>								
<75(women) or <88 (men)	388	Reference	148	Reference	111	Reference	120	Reference
75-85.9(women) or 88-95.9(men)	699	1.16(1.02-1.31)	281	1.17(0.96-1.43)	214	1.27(1.01-1.60)	195	1.08(0.86-1.36)
≥86(women) or ≥96(men)	957	1.37(1.22-1.55)	424	1.51(1.24-1.83)	281	1.48(1.18-1.86)	240	1.16(0.93-1.46)
<b>P value for trend</b>		<.0001		<.0001		0.0006		0.19
<b>Adjusted for BMI</b>								
<75(women) or <88 (men)	388	Reference	148	Reference	111	Reference	120	Reference
75-85.9(women) or 88-95.9(men)	699	1.26(1.09-1.46)	281	1.34(1.07-1.69)	214	1.42(1.08-1.85)	195	1.09(0.84-1.43)



<b>≥86(women) or ≥96(men)</b>	957	1.56(1.32-1.84)	424	1.92(1.47-2.50)	281	1.71(1.25-2.33)	240	1.12(0.82-1.54)
<b>P value for trend</b>		<.0001		<.0001		0.001		0.49

**Gluteofemoral obesity**  
**Hip circumference (cm)**

<b>Not adjusted for BMI</b>								
<101	816	Reference	322	Reference	237	Reference	238	Reference
≥101	1227	1.14(1.04-1.25)	531	1.23(1.07-1.42)	368	1.19(1.01-1.40)	317	1.03(0.87-1.22)
<b>Adjusted for BMI</b>								
<101	816	Reference	322	Reference	237	Reference	238	Reference
≥101	1227	1.12(1.00-1.25)	531	1.28(1.07-1.53)	368	1.16(0.94-1.43)	317	0.95(0.77-1.18)
<b>Adjusted for BM and waist circumference</b>								
<101	816	Reference	322	Reference	237	Reference	238	Reference
≥101	1227	1.02(0.90-1.15)	531	1.12(0.93-1.35)	368	1.05(0.84-1.30)	317	0.92(0.73-1.15)

**Muscularity**  
**Body adiposity index (BAI, continuous, per 5 units)**

<b>Not adjusted for BMI or waist circumference</b>	2043	0.98(0.93-1.04)	853	0.98(0.91-1.07)	605	0.95(0.86-1.06)	555	1.03(0.92-1.15)
<b>Adjusted for BMI</b>	2043	0.87(0.81-0.94)	853	0.88(0.78-0.99)	605	0.81(0.70-0.94)	555	0.95(0.81-1.10)
<b>Adjusted for waist circumference</b>	2043	0.88(0.83,0.94)	853	0.86(0.78,0.95)	605	0.84(0.74,0.95)	555	0.99(0.88,1.13)
<b>Adjusted for BMI and waist circumference</b>	2043	0.86(0.80-0.94)	853	0.87(0.77-0.98)	605	0.81(0.70-0.93)	555	0.94(0.81-1.10)

a. adjusted for education, smoking status, alcohol drinking, physical activity, family history of cancer, study center, and/or anthropometrics when appropriate, stratified by age groups.

**Table 3. HR and 95% CI for risk of colorectal adenocarcinoma in relation to anthropometric characteristics in men**

Anthropometric indices	Total		Proximal colon		Distal colon		Rectum	
	N	HR(95%CI)	N	HR(95%CI)	N	HR(95%CI)	N	HR(95%CI)
<b>Body mass index (BMI)</b>								
<b>Not adjusted for waist circumference</b>								
<22.5	86	0.89(0.69-1.14)	31	0.86(0.57-1.30)	23	0.86(0.54-1.39)	30	0.91(0.60-1.38)
22.5-25	238	Reference	88	Reference	66	Reference	81	Reference
25-30	561	1.02(0.87-1.19)	207	1.01(0.79-1.30)	176	1.16(0.87-1.54)	169	0.90(0.69-1.18)
30	216	1.30(1.08-1.57)	84	1.37(1.01-1.85)	67	1.48(1.05-2.09)	63	1.11(0.79-1.54)
<b>P value for trend</b>		0.002		0.03		0.009		0.59
<b>Adjusted for waist circumference</b>								
<22.5	103	0.96(0.74-1.24)	31	1.02(0.66-1.57)	23	1.03(0.63-1.70)	36	0.87(0.56-1.34)
22.5-25	279	Reference	88	Reference	66	Reference	94	Reference
25-30	644	0.91(0.76-1.08)	207	0.77(0.58-1.02)	176	0.98(0.72-1.35)	205	0.97(0.72-1.31)
30	246	1.08(0.86-1.36)	84	0.86(0.59-1.25)	67	1.20(0.79-1.83)	75	1.26(0.82-1.92)
<b>P value for trend</b>		0.59		0.39		0.52		0.25
<b>Waist Circumference (cm)</b>								
<b>Not adjusted for BMI</b>								
<75(women) or <88 (men)	216	Reference	70	Reference	54	Reference	86	Reference
75-85.9(women) or 88-95.9(men)	364	1.16(0.98-1.38)	125	1.23(0.92-1.65)	120	1.55(1.12-2.14)	116	0.93(0.70-1.23)
≥86(women) or ≥96(men)	521	1.37(1.17-1.61)	215	1.73(1.31-2.28)	158	1.71(1.24-2.34)	141	0.93(0.71-1.23)
<b>P value for trend</b>		<.0001		<.0001		0.002		0.65
<b>Adjusted for BMI</b>								
<75(women) or <88 (men)	216	Reference	71	Reference	54	Reference	86	Reference
75-85.9(women) or 88-95.9(men)	364	1.20(1.00-1.46)	129	1.39(1.00-1.95)	120	1.57(1.09-2.27)	116	0.90(0.65-1.24)
≥86(women) or ≥96(men)	521	1.37(1.10-1.70)	217	2.05(1.41-2.97)	158	1.62(1.06-2.45)	141	0.82(0.57-1.20)
<b>P value for trend</b>		0.01		<.0001		0.05		0.31
<b>Hip circumference (cm)</b>								
<b>Not adjusted for BMI</b>								
<101	419	Reference	141	Reference	124	Reference	145	Reference
≥101	682	1.21(1.07-1.37)	269	1.41(1.15-1.73)	208	1.24(0.99-1.56)	198	1.02(0.82-1.26)
<b>Adjusted for BMI</b>								
<101	419	Reference	141	Reference	124	Reference	145	Reference
≥101	682	1.14(0.99-1.32)	269	1.39(1.09-1.78)	208	1.09(0.83-1.42)	198	0.99(0.76-1.28)
<b>Adjusted for BMI and waist circumference</b>								
<101	419	Reference	141	Reference	124	Reference	145	Reference
≥101	682	1.07(0.92-1.25)	269	1.20(0.92-1.56)	208	0.99(0.75-1.31)	198	1.04(0.79-1.38)

**Body adiposity index (BAI, continuous, per 5 units)**

<b>Not adjusted for BMI or waist circumference</b>	1101	1.07(0.97-1.18)	410	1.12(0.95-1.31)	332	1.08(0.90-1.30)	343	1.02(0.85-1.22)
<b>Adjusted for BMI</b>	1101	0.94(0.83-1.07)	410	0.97(0.79-1.20)	332	0.88(0.70-1.11)	343	0.96(0.76-1.21)
<b>Adjusted for BMI and waist circumference</b>	1101	0.94(0.83-1.07)	410	0.97(0.79-1.19)	332	0.89(0.70-1.12)	343	0.96(0.76-1.22)

---

**Table 4. HR and 95% CI for risk of colorectal adenocarcinoma in relation to anthropometric characteristics in women**

Anthropometric indices	N		HR(95% CI)		N		HR(95% CI)	
<b>Body mass index (BMI)</b>								
<b>Not adjusted for waist circumference</b>								
<22.5	148	1.00(0.81-1.24)	66	0.98(0.71-1.34)	49	1.14(0.77-1.67)	30	0.83(0.53-1.32)
22.5-25	197	Reference	93	Reference	56	Reference	46	Reference
25-30	381	1.05(0.89-1.25)	182	1.02(0.79-1.31)	110	1.11(0.80-1.54)	82	1.02(0.71-1.47)
30	217	1.09(0.89-1.32)	102	1.01(0.76-1.35)	59	1.11(0.76-1.61)	54	1.24(0.83-1.86)
<b>P value for trend</b>		0.36		0.8		0.91		0.104
<b>Adjusted for waist circumference</b>								
<22.5	180	1.15(0.91-1.46)	66	1.11(0.79-1.57)	49	1.27(0.83-1.93)	30	1.04(0.63-1.73)
22.5-25	229	Reference	93	Reference	56	Reference	46	Reference
25-30	422	0.87(0.71-1.05)	182	0.84(0.64-1.12)	110	0.91(0.63-1.30)	82	0.82(0.55-1.23)
30	245	0.77(0.60-0.99)	102	0.72(0.50-1.04)	59	0.76(0.47-1.21)	54	0.88(0.53-1.48)
<b>P value for trend</b>		0.007		0.04		0.08		0.54
<b>Waist Circumference (cm)</b>								
<b>Not adjusted for BMI</b>								
<75(women) or <88 (men)	172	Reference	78	Reference	57	Reference	34	Reference
75-85.9(women) or 88-95.9(men)	335	1.16(0.96-1.40)	156	1.12(0.85-1.47)	94	1.03(0.74-1.44)	79	1.47(0.98-2.21)
≥86(women) or ≥96(men)	436	1.40(1.16-1.68)	209	1.33(1.02-1.74)	123	1.31(0.94-1.82)	99	1.78(1.19-2.67)
<b>P value for trend</b>		0.0002		0.03		0.07		0.01
<b>Adjusted for BMI</b>								
<75(women) or <88 (men)	172	Reference	78	Reference	57	Reference	34	Reference
75-85.9(women) or 88-95.9(men)	335	1.33(1.06-1.65)	156	1.27(0.92-1.76)	94	1.21(0.82-1.81)	79	1.63(1.01-2.63)
≥86(women) or ≥96(men)	436	1.81(1.39-2.36)	209	1.75(1.19-2.58)	123	1.78(1.10-2.88)	99	2.07(1.17-3.68)
<b>P value for trend</b>		<.0001		0.003		0.01		0.02
<b>Hip circumference (cm)</b>								
<b>Not adjusted for BMI</b>								
<101	397	Reference	181	Reference	113	Reference	93	Reference
≥101	545	1.09(0.95-1.24)	262	1.10(0.91-1.34)	160	1.16(0.91-1.49)	119	1.07(0.81-1.41)
<b>Adjusted for BMI</b>								
<101	397	Reference	181	Reference	113	Reference	93	Reference
≥101	545	1.08(0.90-1.29)	262	1.16(0.89-1.50)	160	1.29(0.92-1.80)	119	0.86(0.59-1.26)
<b>Adjusted for BMI and waist circumference</b>								
<101	397	Reference	181	Reference	113	Reference	93	Reference
≥101	545	0.96(0.80-1.16)	262	1.04(0.79-1.37)	160	1.16(0.81-1.65)	119	0.75(0.51-1.11)
<b>Body adiposity index (BAI, continuous, per 5 units)</b>								
<b>Not adjusted for BMI or waist circumference</b>		0.96(0.90-1.03)		0.95(0.86-1.05)	273	0.92(0.81-1.05)		1.07(0.93-1.23)
<b>Adjusted for BMI</b>		0.86(0.77-0.95)		0.87(0.75-1.01)	273	0.82(0.67-1.00)		0.94(0.76-1.17)
<b>Adjusted for BMI and waist circumference</b>		0.84(0.75-0.93)		0.85(0.73-0.99)	273	0.79(0.65-0.97)		0.92(0.74-1.14)

Table 5. HR and 95% CI for risk of colorectal adenocarcinoma in relation to normal weight obesity

Body Mass index (BMI, kg/m <sup>2</sup> )	Waist circumference (cm)	Non-cases participants	Colorectum		Proximal colon		Distal colon		Rectum	
			Cases	HR(95%CI)	Cases	HR(95%CI)	Cases	HR(95%CI)	Cases	HR(95%CI)
Lower weight (<22.5)	Women<80, men<94	23559	218	1.00(0.84-1.19)	90	1.00(0.76-1.31)	70	1.19(0.87-1.64)	53	0.80(0.58-1.12)
	Women≥80, men≥94	937	16	1.47(0.89-2.43)	7	1.42(0.66-3.06)	2	0.70(0.17-2.87)	7	2.37(1.09-5.12)
Normal weight (≥22.5 and <25)	Women<80, men<94	28671	320	Reference	128	Reference	87	Reference	101	Reference
	Women≥80, men≥94	6920	115	1.23(0.99-1.53)	53	1.31(0.95-1.80)	35	1.44(0.97-2.14)	26	0.97(0.63-1.50)
Over weight(≥25 and <30)	Women<80, men<94	23004	273	0.97(0.82-1.14)	88	0.80(0.61-1.06)	85	1.10(0.82-1.49)	94	1.02(0.77-1.35)
	Women≥80, men≥94	37440	669	1.13(0.99-1.30)	301	1.21(0.98-1.49)	201	1.30(1.01-1.68)	157	0.88(0.69-1.14)
Obese (≥30)	Women<80, men<94	443	5	1.07(0.44-2.58)	4	2.22(0.82-6.02)	1	0.77(0.11-5.55)	0	0
	Women≥80, men≥94	22503	428	1.25(1.08-1.44)	182	1.23(0.98-1.55)	125	1.41(1.06-1.86)	117	1.16(0.88-1.52)

a: normal weight without abdominal obesity; b: overweight with abdominal obesity; c: overweight without abdominal obesity; d: overweight with abdominal obesity

**Supplemental Table 1. HR and 95% CI for risk of colorectal adenocarcinoma in relation to anthropometric characteristics**

Anthropometric indices	Colorectal adenocarcinoma by anatomical location <sup>a</sup>								
		Total	Proximal colon		Distal colon		Rectum		
		HR(95%CI)	HR(95%CI)	HR(95%CI)	HR(95%CI)	HR(95%CI)	HR(95%CI)	HR(95%CI)	HR(95%CI)
<b>Total</b>	2044		853		606		555		
<b>Waist to hip ratio (WHR, continuous, per 1/10)</b>									
<b>Not adjusted for BMI</b>	2043	1.14(1.06-1.22)	853	1.28(1.16-1.42)	605	1.20(1.06-1.36)	555	1.15(1.01-1.31)	
<b>Adjusted for BMI</b>	2043	1.22(1.13-1.31)	853	1.30 (1.16-1.46)	605	1.19(1.03-1.36)	555	1.13(0.97-1.31)	
<b>Waist to height ratio (WHtR, continuous, per 1/10)</b>									
<b>Not adjusted for BMI</b>	2044	1.14(1.06-1.22)	853	1.18(1.06-1.31)	606	1.12(0.99-1.27)	555	1.13(0.98-1.29)	
<b>Adjusted for BMI</b>	2044	1.15(1.03-1.28)	853	1.26(1.07-1.49)	606	1.08(0.88-1.32)	555	1.10(0.89-1.36)	
<b>Not adjusted for BMI</b>									
<0.5	596	Reference	237	Reference	177	Reference	170	Reference	
0.5-	1138	1.19(1.07-1.32)	479	1.26(1.07-1.49)	343	1.27(1.05-1.54)	303	1.06(0.87-1.30)	
≥0.6	310	1.29(1.12-1.49)	137	1.37(1.10-1.71)	86	1.30(0.99-1.70)	82	1.20(0.91-1.58)	
<b>P value for trend</b>		0.0002		0.002		0.02			
<b>Adjusted for BMI</b>									
<0.5	596	Reference	237	Reference	177	Reference	170	Reference	
0.5-	1138	1.23(1.08-1.41)	479	1.37(1.12-1.69)	343	1.30(1.02-1.66)	303	1.04(0.81-1.34)	
≥0.6	310	1.28(1.05-1.57)	137	1.49(1.09-2.02)	86	1.27(0.87-1.84)	82	1.06(0.72-1.56)	
<b>P value for trend</b>		0.007		0.006		0.13		0.74	
<b>Waist to height index (WHI, continuous, per 10 units)</b>									
<b>Not adjusted for BMI</b>	2044	1.14(1.02-1.26)	853	1.18(1.01-1.39)	606	1.07(0.88-1.30)	555	1.18(0.96-1.45)	
<b>Adjusted for BMI</b>	2044	1.06(0.91-1.23)	853	1.17(0.93-1.46)	606	0.90(0.68-1.20)	555	1.10(0.82-1.48)	

a. adjusted for education, smoking status, alcohol drinking, physical activity, family history of cancer, study center, and/or anthropometrics when appropriate, stratified by age groups.

**Supplemental Table 2. HR and 95% CI for risk of colorectal adenocarcinoma in relation to anthropometric characteristics by genders**

Anthropometric indices	Total		Proximal colon				Distal colon				Rectum					
	Male		Female		Male		Female		Male		Female		Male		Female	
	N	HR(95%CI)	N	HR(95%CI)	N	HR(95%CI)	N	HR(95%CI)	N	HR(95%CI)	N	HR(95%CI)	N	HR(95%CI)	N	HR(95%CI)
<b>Waist to hip ratio (continuous, per 1/10)</b>																
Not adjusted for BMI	1101	1.22(1.11-1.34)	942	1.21(1.10-1.33)	410	1.37(1.18-1.59)	443	1.21(1.06-1.40)	332	1.25(1.06-1.48)	289	1.16(0.97-1.39)	343	1.07(0.90-1.26)	212	1.26(1.03-1.54)
Adjusted for BMI	1101	1.18(1.06-1.31)	942	1.24(1.11-1.37)	410	1.34(1.13-1.60)	443	1.26(1.08-1.47)	332	1.15(0.95-1.40)	289	1.20(0.98-1.46)	343	1.05(0.86-1.27)	212	1.22(0.97-1.52)
<b>Waist to height ratio (continuous, per 1/10)</b>																
Not adjusted for BMI	1101	1.24(1.12-1.38)	943	1.09(0.99-1.19)	410	1.40(1.18-1.66)	443	1.08(0.94-1.23)	332	1.28(1.05-1.55)	289	1.05(0.88-1.24)	343	1.07(0.88-1.30)	212	1.20(1.00-1.45)
Adjusted for BMI	1101	1.20(1.02-1.42)	943	1.14(0.98-1.32)	410	1.50(1.16-1.95)	443	1.17(0.95-1.45)	332	1.10(0.82-1.49)	289	1.10(0.83-1.45)	343	1.02(0.76-1.38)	212	1.16(0.85-1.57)
<b>Not adjusted for BMI</b>																
<0.5	229	Reference	367	Reference	74	Reference	163	Reference	61	Reference	116	Reference	89	Reference	81	Reference
0.5-	700	1.20(1.03-1.40)	438	1.23(1.06-1.42)	265	1.38(1.06-1.80)	214	1.24(1.00-1.53)	224	1.50(1.12-2.01)	119	1.16(0.88-1.51)	203	0.89(0.69-1.16)	100	1.38(1.01-1.88)
≥0.6	172	1.43(1.17-1.76)	138	1.18(0.96-1.45)	71	1.79(1.27-2.51)	66	1.14(0.84-1.53)	47	1.55(1.05-2.31)	39	1.19(0.82-1.75)	51	1.09(0.76-1.56)	31	1.33(0.86-2.05)
P value for trend		0.0006		0.03		0.0007		0.2		0.014		0.27		0.85		0.08
<b>Adjusted for BMI</b>																
<0.5	229	Reference	367	Reference	74	Reference	163	Reference	61	Reference	116	Reference	89	Reference	81	Reference
0.5-	700	1.19(0.98-1.44)	438	1.32(1.09-1.59)	265	1.48(1.07-2.04)	214	1.37(1.04-1.81)	224	1.41(0.98-2.01)	119	1.26(0.89-1.79)	203	0.86(0.63-1.19)	100	1.36(0.91-2.02)
≥0.6	172	1.26(0.95-1.67)	138	1.30(0.97-1.73)	71	1.76(1.11-2.79)	66	1.33(0.87-2.02)	47	1.20(0.71-2.04)	39	1.40(0.81-2.41)	51	0.96(0.58-1.58)	31	1.18(0.64-2.17)
P value for trend		0.1		0.03		0.02		0.2		0.4		0.18				
<b>Waist to height index (WHI, continuous, per 10 units)</b>																
Not adjusted for BMI	1119	1.30(1.10-1.53)	964	1.08(0.94-1.24)	410	1.50(1.14-1.96)	447	1.07(0.88-1.31)	332	1.32(0.97-1.79)	274	0.99(0.76-1.28)	343	1.10(0.81-1.49)	213	1.30(0.98-1.71)
Adjusted for BMI	1119	1.12(0.89-1.42)	964	1.07(0.87-1.31)	410	1.38(0.95-2.01)	447	1.13(0.84-1.50)	332	0.98(0.64-1.49)	274	0.93(0.64-1.37)	343	1.02(0.67-1.56)	213	1.20(0.79-1.82)