

**FOLLOW-THE-LEADER MODELS CAN BE VIEWED AS A
NUMERICAL APPROXIMATION TO THE
LIGHTHILL–WHITHAM–RICHARDS MODEL FOR TRAFFIC
FLOW**

HELGE HOLDEN AND NILS HENRIK RISEBRO

ABSTRACT. We show how to view the standard Follow-the-Leader (FtL) model as a numerical method to compute numerically the solution of the Lighthill–Whitham–Richards (LWR) model for traffic flow. As a result we offer a simple proof that FtL models converge to the LWR model for traffic flow when traffic becomes dense. The proof is based on techniques used in the analysis of numerical schemes for conservation laws, and the equivalence of weak entropy solutions of conservation laws in the Lagrangian and Eulerian formulation.

1. INTRODUCTION

There are two paradigms in the mathematical modeling of traffic flow. One is based on an individual modeling of each vehicle with the dynamics governed by the distance between adjacent vehicles. The other is based on the assumption of dense traffic where the vehicles are represented by a density function, and individual vehicles cannot be identified. The dynamics is governed by a local velocity function depending solely on the density. The first model is denoted the Follow-the-Leader (FtL) model, and the second is called the Lighthill–Whitham–Richards (LWR) model [13, 14] for traffic flow. Further refinements and extensions of these models are available. Intuitively, it is clear that the the FtL model should approach or approximate the LWR model in the case of heavy traffic, and that is what is proved here. This problem has been extensively studied, see [1, 2, 3, 5, 6, 7, 8, 9, 10, 12, 15]. Using numerical methods for scalar conservation laws we show that FtL models appear naturally as a numerical approximation of the LWR model. Thus we offer a short and direct proof that the FtL model converges to the LWR model, and our analysis is based on a careful study of the relationship between weak solutions in Lagrangian and Eulerian variables.

In the LWR model vehicles are described by a density $\rho = \rho(t, x)$ where x is the position along the road, and t as usual denotes time. Locally, one assumes that the velocity is given by a function v that depends on the density only, that is, $v = v(\rho)$. If we consider unidirectional traffic on a homogenous road without exits or entries, conservation of vehicles requires that the dynamics is governed by the scalar conservation law

$$\rho_t + (\rho v(\rho))_x = 0,$$

which constitutes the LWR model. It is often denoted as “traffic hydrodynamics” due to its resemblance with fluid dynamics.

Date: September 22, 2017.

2010 Mathematics Subject Classification. Primary: 35L02; Secondary: 35Q35, 82B21.

Key words and phrases. Follow-the-Leader model, Lighthill–Whitham–Richards model, traffic flow, continuum limit.

Research was supported by the grant *Waves and Nonlinear Phenomena (WaNP)* from the Research Council of Norway. The research was done while the authors were at Institut Mittag-Leffler, Stockholm.

The FtL model can be described as follows. Consider N vehicles with length ℓ and position $z_1(t) < \dots < z_N(t)$ on the real axis with dynamics given by

$$\dot{z}_i = v\left(\frac{\ell}{z_{i+1} - z_i}\right) \quad \text{for } i = 1, \dots, N-1,$$

$$\dot{z}_N = v_{\max}$$

Here v denotes a given velocity function with maximum v_{\max} , perhaps the speed limit. Our proofs are considerably simpler when we have a uniform bound on $z_{i+1}(t) - z_i(t)$. Having empty road ahead of the first car would mean that “ $z_{N+1} - z_N = \infty$ ”. This is the same as imposing $\dot{z}_N = v_{\max}$, and in this case $z_{i+1}(t) - z_i(t)$ would not be bounded by a constant independent of time. Therefore we will in this paper assume that we model one of two alternatives:

Periodic case: We are in the periodic case in which $z_i \in [a, b]$ for some interval $[a, b]$, and

$$\dot{z}_N(t) = v\left(\frac{\ell}{b - z_N(t) - a + z_1(t)}\right).$$

Non-periodic case: We imagine that there are infinitely many vehicles to the right of z_N , the distance between each of these vehicles is $M\ell$, for a finite, but arbitrary, constant $M > 1$. In this case

$$\dot{z}_N(t) = v\left(\frac{1}{M}\right).$$

Introduce $y_i = (z_{i+1} - z_i)/\ell$ for $i = 1, \dots, N-1$, to obtain

$$\dot{y}_i(t) = \frac{1}{\ell}(v(1/y_{i+1}(t)) - v(1/y_i(t))).$$

In this paper we analyze the limit of this system of ordinary differential equations when $N \rightarrow \infty$. There are two ways to proceed.

We may analyze this system directly, in what we call the semi-discrete case, see Section 2.1. By using methods from the theory of numerical methods for scalar conservation laws we show that the sequence $\{y_i(t)\}_{i=1}^{N-1}$ converges, as $\ell \rightarrow 0$ and $N \rightarrow \infty$, to a function $y(t, x)$ that satisfies the equation

$$(1.1) \quad \begin{cases} y_t - V(y)_x = 0 & t > 0, \quad x \in [0, 1], \\ y(0, x) = y_0(x) & x \in [0, 1], \end{cases}$$

where $V(y) = v(1/y)$, and with boundary condition

$$\begin{cases} y(t, 1) = y(t, 0) & \text{in the periodic case,} \\ y(t, 1) = M & \text{else.} \end{cases}$$

Note that x is the Lagrangian mass coordinate, so that the integer part of x/ℓ measures how many cars there are to the left of x .

Equation (1.1) is an example of a hyperbolic conservation law. It is well-known that solutions develop singularities, denoted shocks, in finite time independent of the smoothness of the initial data. Thus one needs to study weak solutions, and design so-called entropy conditions to identify the unique weak physical solution. For a scalar conservation law $u_t + f(u)_x = 0$ with initial data $u|_{t=0} = u_0$, the unique weak entropy solution $u = u(t, x)$, which is an integrable function of bounded variation, satisfies the Kruřkov entropy condition

$$(1.2) \quad \int_0^\infty \int_0^\infty (|u - k| \phi_t + \text{sign}(u - k)(f(u) - f(k))\phi_x) dt dx + \int |u_0 - k| \phi|_{t=0} dx \geq 0$$

for all real constants $k \in \mathbb{R}$, and all non-negative test functions $\phi \in C_0^\infty(\mathbb{R} \times [0, \infty))$. See [11].

As an alternative approach, see Section 2.2, we may discretize the time derivative by a small positive Δt and write $z_j^n \approx z_j(n\Delta t)$, $y_j^n \approx y_j(n\Delta t)$, we have that

$$z_j^{n+1} = z_j^n + \Delta t V_j^n, \quad \text{and} \quad y_j^{n+1} = y_j^n + \frac{\Delta t}{\ell} (V_{j+1}^n - V_j^n),$$

where $V_j^n = V(y_j^n)$. The key observation is that this is an approximation of the hyperbolic conservation law $y_t - V(y)_x = 0$ by a monotone scheme, and from the classical result of Crandall–Majda [4], see also [11, Thm. 3.9], we know that this scheme converges, as $\ell \rightarrow 0$, $N \rightarrow \infty$, and $\Delta t \rightarrow 0$, to the entropy solution of equation (1.1), namely $y_t - V(y)_x = 0$. Thus in both cases we obtain convergence to the same hyperbolic conservation law in Lagrangian coordinates.

Next we have to transform the result of the two approaches, both in Lagrangian coordinates, to Eulerian coordinates. For smooth solutions this is nothing but a simple exercise in calculus, but for weak entropy solutions this is a deep result due to Wagner [16]. To be specific, we introduce the Eulerian space coordinate $z = z(t, x)$, with $z_x = y$ and $z_t = V(y)$. A straightforward (but formal) calculation reveals that the Eulerian functions satisfy

$$y_t = -\frac{1}{\rho^2}(\rho_t + \rho_z v), \quad V(y)_x = \frac{1}{\rho} v'(\rho) \rho_z,$$

and hence

$$\rho_t + (\rho v(\rho))_z = 0,$$

which is nothing but the LWR model. These formal transformations are not valid in general for weak entropy solutions. However, thanks to the fundamental result of Wagner [16], weak entropy solutions in Lagrangian coordinates transform into weak entropy solutions in Eulerian variables. The approach here bears some resemblance to the approach in [12], where the proof is obtained in a grid-less manner, and it does not depend on the use of Crandall–Majda and Wagner.

2. THE MODEL

Let us first introduce the FtL model. Consider N vehicles moving on a one-dimensional road. Their position is given as a function of time t as $z_1(t), \dots, z_N(t)$. For the moment (we shall actually show that this is so below) we assume that $z_1(t) < z_2(t) < \dots < z_N(t)$. We introduce the “local inverse density” by

$$y_i = \frac{1}{\ell} (z_{i+1} - z_i), \quad i = 1, \dots, N-1,$$

where ℓ is the length of each vehicle. The velocity of the vehicle at z_i is assumed to be a function of the distance to the vehicle in front, at z_{i+1} . This means that

$$(2.1) \quad \dot{z}_i(t) = v\left(\frac{\ell}{z_{i+1}(t) - z_i(t)}\right), \quad i = 1, \dots, N-1.$$

Regarding the first vehicle, located at z_N , we either assume that there are infinitely many equally spaced vehicles in front of it, i.e., $y_N = M$, or that we are in the periodic setting in an interval $[a, b]$, so that the distance from the vehicle at z_N to the vehicle at z_1 is $b - z_N + z_1 - a$, i.e., $y_N = (b - z_N + z_1 - a)/\ell$. We have

$$(2.2) \quad \dot{z}_N(t) = v\left(\frac{1}{y_N(t)}\right).$$

Regarding the velocity function v , we assume it to be a decreasing Lipschitz continuous function such that

$$(2.3) \quad v(0) = v_{\max} \quad \text{and} \quad v(\rho) = 0 \quad \text{for} \quad \rho \geq 1.$$

The prototypical example is $v(\rho) = v_{\max} \max\{0, 1 - \rho\}$. We define the velocity in Lagrangian variables by $V(y) = v(1/y)$. Observe that V is globally bounded,

Lipschitz continuous and increasing for $y \geq 1$, with a bounded Lipschitz constant L_v .

Rewriting (2.1) in terms of $\{y_i\}$ we get

$$(2.4) \quad \dot{y}_i = \frac{1}{\ell} (V(y_{i+1}) - V(y_i)), \quad i = 1, \dots, N-1,$$

and

$$(2.5) \quad y_N = \begin{cases} M, & \text{non-periodic case,} \\ \frac{1}{\ell}(b - z_N + z_1 - a), & \text{periodic case.} \end{cases}$$

Let us also define the *Lagrangian grid* $\{x_{i-1/2}\}_{i=1}^N$ by $x_{i-1/2} = (i-1)\ell$. We shall also assume throughout that there is a constant $1 \leq K < \infty$, K independent of N and ℓ , such that

$$(2.6) \quad 1 \leq y_j(0) \leq K, \quad \sum_{j=1}^{N-1} |y_{j+1}(0) - y_j(0)| \leq K.$$

2.1. The semi-discrete case. In this section we show that the solution of the system (2.4) of ordinary differential equations converges to an entropy solution of (2.20) as $\ell \rightarrow 0$, and that “ $1/y$ ” converges to an entropy solution of (2.12).

Concretely, we define the piecewise constant function

$$(2.7) \quad y_\ell(t, x) = y_j(t), \quad x \in (x_{j-1/2}, x_{j+1/2}].$$

We shall also use the notation

$$D_+ h_j = \frac{1}{\ell} (h_{j+1} - h_j)$$

for the forward difference. Let

$$y^+ = \max\{y, 0\} \quad \text{and} \quad y^- = -\min\{y, 0\},$$

and let H denote the Heaviside function

$$H(y) = \begin{cases} 0 & y \leq 0, \\ 1 & y > 0. \end{cases}$$

Lemma 2.1. *Let $y_j(t)$ solve the system (2.4). Then*

$$(2.8a) \quad \frac{d}{dt} (y_j - k)^+ \leq D_+ [H(y_j - k) (V(y_j) - V(k))],$$

$$(2.8b) \quad \frac{d}{dt} (y_j - k)^- \leq D_+ [-H(k - y_j) (V(y_j) - V(k))],$$

for any constant k .

Proof. Throughout we use the notation $V_j = V(y_j)$. We have that

$$\begin{aligned} \frac{d}{dt} (y_j - k)^+ &= \frac{1}{\ell} H(y_j - k) (V_{j+1} - V_j) \\ &= \frac{1}{\ell} [H(y_{j+1} - k) (V_{j+1} - V(k)) - H(y_j - k) (V_j - V(k))] \\ &\quad - \frac{1}{\ell} (H(y_{j+1} - k) - H(y_j - k)) (V_{j+1} - V(k)) \\ &= D_+ [H(y_j - k) (V_j - V(k))] \\ &\quad - \frac{1}{\ell} (H(y_{j+1} - k) - H(y_j - k)) (V_{j+1} - V(k)). \end{aligned}$$

Now

$$(H(y_{j+1} - k) - H(y_j - k)) (V_{j+1} - V(k))$$

$$\begin{aligned}
&= \begin{cases} 0 & y_j, y_{j+1} \geq k \text{ or } y_j, y_{j+1} \leq k, \\ V_{j+1} - V(k) & y_j < k < y_{j+1}, \\ V(k) - V_{j+1} & y_{j+1} < k < y_j, \end{cases} \\
&\geq 0,
\end{aligned}$$

since $y \mapsto V(y)$ is increasing. This proves (2.8a); estimate (2.8b) is proved similarly. \square

Now define $y_j(t) = y_1(t)$ for $j < 1$ and $y_j(t) = y_{N-1}(t)$ for $j > N - 1$ in the non-periodic case. In the periodic case we define $y_j(t)$ by periodic extension. To save space, we also use the convention that in the non-periodic case, sums over j range over all $j \in \mathbb{Z}$, while in the periodic case, sums range over $j = 1, \dots, N - 1$.

Lemma 2.2. *If $1 \leq y_j(0) \leq K$, then $1 \leq y_j(t) \leq K$ for $t > 0$.*

Proof. From (2.8a) and (2.8b) we have

$$\frac{d}{dt} \sum_j (y_j(t) - k)^\pm \leq 0.$$

Thus if $y_j(0) \leq K$ for all j , then $y_j(t) < k$ for any constant $k > K$. Similarly $y_j(t) > k$ for any constant $k < 1$ if $y_j(0) \geq 1$ for all j . \square

Lemma 2.3. *If $\{\tilde{y}_j(t)\}_{j=1}^{N-1}$ is another solution of (2.4) and (2.5) with initial data $\tilde{y}_j(0)$, then*

$$(2.9) \quad \sum_j |y_j(T) - \tilde{y}_j(T)| \leq \sum_j |y_j(0) - \tilde{y}_j(0)|,$$

for $T > 0$.

Proof. Adding (2.8a) and (2.8b), and observing that

$$(y - k)^+ + (y - k)^- = |y - k| \quad \text{and} \quad H(y - k) - H(k - y) = \text{sign}(y - k),$$

we find that

$$(2.10) \quad \frac{d}{dt} |y_j - k| \leq D_+ [\text{sign}(y_j - k) (V(y_j) - V(k))].$$

Set $q(y, k) = \text{sign}(y - k) (V(y) - V(k))$. Choosing $k = \tilde{y}_j(\tau)$ in the inequality for $y_j(t)$ and $k = y_j(t)$ in the inequality for $\tilde{y}_j(\tau)$, and adding the two inequalities, give

$$\left(\frac{d}{dt} + \frac{d}{d\tau} \right) |y_j(t) - \tilde{y}_j(\tau)| \leq (D_{+,1} + D_{+,2}) q(y_j(t), \tilde{y}_j(\tau)),$$

where $D_{+,1(2)}$ denotes the difference with respect to the first (second) argument. Summing over j , multiplying with a non-negative test function $\varphi(t, \tau)$, where $\varphi \in C_0^\infty((0, \infty)^2)$, and integrating by parts yield

$$\int_0^\infty \int_0^\infty (\varphi_t + \varphi_\tau) \sum_j |y_j(t) - \tilde{y}_j(\tau)| \, d\tau dt \geq 0,$$

since taking the sum makes the right-hand side “telescope”. Now we can use Kruřkov’s trick, see [11, Sec. 2.4], and choose

$$\varphi(t, \tau) = \psi \left(\frac{t + \tau}{2} \right) \omega_\varepsilon(t - \tau),$$

where $\psi \in C_0^\infty((0, \infty))$, $\psi \geq 0$ and ω_ε is a standard mollifier, to obtain, as $\varepsilon \rightarrow 0$, that

$$\int_0^\infty \psi'(t) \sum_j |y_j(t) - \tilde{y}_j(t)| \, dt \geq 0.$$

Choose ψ to be a smooth approximation to the characteristic function of the interval $(t_1, t_2) \subset (0, T)$, to get

$$\sum_j |y_j(t_2) - \tilde{y}_j(t_2)| \leq \sum_j |y_j(t_1) - \tilde{y}_j(t_1)|.$$

The lemma follows by letting $t_1 \downarrow 0$ and $t_2 \uparrow T$. For details, see [11, Sec. 2.4]. \square

Lemma 2.4. *Assume that $1 \leq y_j(0) \leq K$ and that $\sum_j |y_{j+1}(0) - y_j(0)| \leq K$ for some constant K independent of ℓ . Then there is a sequence $\{\ell_i\}$, where $\ell_i \rightarrow 0$ as $i \rightarrow \infty$, and there exists a function $y \in C([0, T]; L^1([0, 1]))$ such that y_{ℓ_i} converges to y in $C([0, T]; L^1([0, 1]))$.*

Proof. Lemma 2.2 shows that $\{y_\ell\}_\ell$ is bounded independently of ℓ ; choosing $\tilde{y}_j = y_{j+1}$ and using Lemma 2.3 yields the BV bound on $\{y_\ell(t)\}_\ell$ uniformly in ℓ and t . Choosing $\tilde{y}_j(t) = y_j(t - \sigma)$ in Lemma 2.3 for some $0 < \sigma < t$ gives

$$\begin{aligned} \|y_\ell(t, \cdot) - y_\ell(t - \sigma, \cdot)\|_{L^1} &= \ell \sum_j |y_j(t) - y_j(t - \sigma)| \\ &\leq \ell \sum_j |y_j(\sigma) - y_j(0)| \\ &\leq \sum_j \int_0^\sigma |V(y_{j+1}(\xi)) - V(y_j(\xi))| d\xi \\ &\leq \|V\|_{\text{Lip}} \sum_j \int_0^\sigma |y_{j+1}(\xi) - y_j(\xi)| d\xi \\ &\leq \|V\|_{\text{Lip}} \sigma \sum_j |y_{j+1}(0) - y_j(0)| \\ &\leq \|V\|_{\text{Lip}} \sigma K. \end{aligned}$$

Hence the map $t \mapsto y_\ell(t, \cdot)$ is L^1 Lipschitz continuous, with a Lipschitz constant independent of ℓ . Thus by [11, Thm. A.11], the family $\{y_\ell\}_{\ell > 0}$ is compact in $C([0, \infty); L^1([0, 1]))$. \square

Furthermore we assume that as N increases, the initial position of the vehicles are such that there is a function $y_0(x)$ such that

$$(2.11) \quad \lim_{\ell \rightarrow 0} y_\ell(0, \cdot) = y_0(\cdot),$$

and that this convergence is in $L^1([0, 1])$. We also assume that $\|y_0\|_{L^\infty([0, 1])} \leq K$, without loss of generality we can then also assume that $\|y_\ell(0, \cdot)\|_{L^\infty([0, 1])} \leq K$.

It is now straightforward, starting from the discrete entropy inequality (2.10), to show that any limit of $\{y_\ell\}_{\ell > 0}$ is the unique entropy solution to (2.20) by following a standard Lax–Wendroff argument, see [11, Thm. 3.4]. Thus the whole sequence $\{y_\ell\}$ converges, and the unique entropy solution to (1.1) is the limit

$$y = \lim_{\ell \rightarrow 0} y_\ell.$$

Introduce the Eulerian spatial coordinate z , given by the equations

$$\frac{\partial z}{\partial x} = y, \quad \frac{\partial z}{\partial t} = V(y),$$

and the variable $\rho = 1/y$. We can now proceed following the argument of Wagner [16] to obtain that ρ is the unique weak entropy solution to the LWR model

$$(2.12) \quad \begin{cases} \rho_t + (\rho v(\rho))_z = 0, & t > 0, \\ \rho(0, z) = \rho_0(z). \end{cases}$$

We can also study the convergence in Eulerian coordinates directly by defining a discrete version of the transformation from Lagrangian to Eulerian coordinates. To define the discrete version of ρ , we need the approximate Eulerian coordinate; $z_\ell(t, x)$. Define

$$z_\ell(t, x) = \frac{1}{\ell}(x_{j+1/2} - x)z_j(t) + \frac{1}{\ell}(x - x_{j-1/2})z_{j+1}(t), \text{ for } x \in [x_{j-1/2}, x_{j+1/2}],$$

where $\{z_j(t)\}$ solves (2.1). Then

$$\begin{aligned} \frac{\partial z_\ell}{\partial t} &= \frac{1}{\ell}(x_{j+1/2} - x)V_j + \frac{1}{\ell}(x - x_{j-1/2})V_{j+1}, \\ \frac{\partial z_\ell}{\partial x} &= y_j, \end{aligned}$$

for $x \in (x_{j-1/2}, x_{j+1/2})$. The sequence $\{z_\ell\}_{\ell>0}$ is uniformly Lipschitz continuous. Hence by the Arzelà–Ascoli theorem, it converges uniformly to a Lipschitz continuous limit $z(t, x)$ satisfying $z_t = V(y)$ and $z_x = y$ almost everywhere. Furthermore the map $x \mapsto z_\ell(t, x)$ is invertible, with inverse $x_\ell(t, z)$. In the periodic case we set

$$(2.13a) \quad z_{l,\ell}(t) = a, \quad z_r(t) = b,$$

otherwise we define

$$(2.13b) \quad z_{l,\ell}(t) = z_\ell(t, 0), \quad z_r(t) = b + tV(M) = z(t, 1) = z_\ell(t, 1).$$

Observe that $z_{l,\ell}(t) = z_\ell(t, 0) \rightarrow z(t, 0) = z_l(t)$ as $\ell \rightarrow 0$. Define

$$(2.14) \quad \rho_\ell(t, z) = \frac{1}{y_\ell(t, x_\ell(t, z))} \text{ for } z \in [z_{l,\ell}(t), z_r(t)].$$

In the periodic case, we define ρ_ℓ by periodic continuation, while in the non-periodic case we define

$$\rho_\ell(t, z) = \begin{cases} 0 & z < z_{l,\ell}, \\ 1/M & z > z_r. \end{cases}$$

Next we claim that

$$(2.15) \quad \rho_\ell(t, z) \rightarrow \rho(t, z) = \tilde{\rho}(t, x(t, z))$$

in $L^1([z_l, z_r])$ as $\ell \rightarrow 0$. To see this, define $\tilde{\rho}_\ell(t, x) = 1/y_\ell(t, x)$, and compute

$$\begin{aligned} \|\rho(t, \cdot) - \rho_\ell(t, \cdot)\|_{L^1} &= \|\tilde{\rho}(t, x(t, \cdot)) - \tilde{\rho}_\ell(t, x_\ell(t, \cdot))\|_{L^1} \\ &\leq \underbrace{\|\tilde{\rho}(t, x(t, \cdot)) - \tilde{\rho}_\ell(t, x_\ell(t, \cdot))\|_{L^1}}_A \\ &\quad + \underbrace{\|\tilde{\rho}_\ell(t, x_\ell(t, \cdot)) - \tilde{\rho}_\ell(t, x_\ell(t, \cdot))\|_{L^1}}_B. \end{aligned}$$

We have that

$$\begin{aligned} A &= \int_{z_l}^{z_r} |\tilde{\rho}(t, x(t, z)) - \tilde{\rho}_\ell(t, x_\ell(t, z))| dz \\ &\leq \left(\int_{\min\{z_l, z_{l,\ell}\}}^{\max\{z_l, z_{l,\ell}\}} + \int_{\max\{z_l, z_{l,\ell}\}}^{z_r} \right) |\tilde{\rho}(t, x(t, z)) - \tilde{\rho}_\ell(t, x_\ell(t, z))| dz. \end{aligned}$$

Since $\tilde{\rho}_\ell$ and $\tilde{\rho}$ are both bounded by 1, and $z_{l,\ell} \rightarrow z_l$ as $\ell \rightarrow 0$, the first of these integrals tend to zero. Since $x_\ell \rightarrow x$ uniformly, the integrand tends to zero almost everywhere, and is bounded by 2. Hence by the dominated convergence theorem, the last integral tends to zero. The same argument applies to B . Thus the claim (2.15) is justified.

Summing up, we have shown the following result.

Theorem 2.5. *Assume that the function v satisfies (2.3). Let $\{y_j\}_{j=1}^{N-1}$ satisfy (2.4), with either periodic boundary conditions; $y_N(t) = y_1(t)$, or $y_N(t) = M$ for some fixed constant $M > 1$. Assume that the initial positions of the vehicles $\{z_i(0)\}_{i=1}^N$ are such that we can define a bounded function y_0 by (2.11), and that (2.6) holds, namely that the initial data are bounded with finite total variation.*

(i) *The piecewise constant (in space) function $y_\ell(t, x)$ defined by (2.7) converges in $C([0, T]; L^1([0, 1]))$ as $\ell \rightarrow 0$ to the unique weak entropy solution y of (1.1). The function $\rho = 1/y$ satisfies the LWR model (2.12) in Eulerian variables.*

(ii) *The function ρ_ℓ defined by (2.14) converges in $C([0, T]; L^1([0, 1]))$ as $\ell \rightarrow 0$ to the unique weak entropy solution ρ of (2.12).*

2.2. Analysis of the Euler scheme for (2.1). The simplest numerical method to approximate solutions of (2.1) is the forward Euler scheme, viz.,

$$(2.16) \quad z_i((n+1)\Delta t) = z_i(n\Delta t) + \Delta t v\left(\frac{\ell}{z_{i+1}(n\Delta t) - z_i(n\Delta t)}\right),$$

where Δt is a (small) positive number.

If we write the Euler scheme (2.16) in the y variable, we get

$$(2.17) \quad y_i^{n+1} = y_i^n + \lambda (V_{i+1}^n - V_i^n), \quad i = 1, \dots, N-1,$$

where $y_i^n = y_i(t^n)$, $t^n = n\Delta t$, $\lambda = \Delta t/\ell$ and $V_i^n = V(y_i^n)$. As a (right) boundary condition we use

$$(2.18) \quad V_N^n = \begin{cases} V(M) & \text{non-periodic,} \\ V_1^n & \text{periodic.} \end{cases}$$

For $t \geq 0$ and $x \in [0, (N-1)\ell]$ define the function

$$y_\ell(t, x) = y_i^n \quad (t, x) \in [t^n, t^{n+1}) \times (x_{i-1/2}, x_{i+1/2}].$$

Observe that we can rewrite (2.17) as

$$y_i^{n+1} = \left(1 - \lambda \theta_{i+1/2}^n\right) y_i^n + \lambda \theta_{i+1/2}^n y_{i+1}^n,$$

where

$$\theta_{i+1/2}^n = -\frac{V_{i+1}^n - V_i^n}{y_{i+1}^n - y_i^n} \geq 0,$$

and since V is Lipschitz continuous, $\theta_{i+1/2}^n \leq L_v$. Hence if the CFL-condition

$$(2.19) \quad \lambda L_v \leq 1,$$

holds, then y_i^{n+1} is a convex combination of y_i^n and y_{i+1}^n . Thus the scheme (2.17) is monotone. In passing, we note that a consequence is that if $1 \leq y_i^0 \leq K$ for all i , then $1 \leq y_i^n \leq K$ for all i . Regarding the position of vehicles, this means that if $z_i(0) \leq z_{i+1}(0) - \ell$, then $z_i(t^n) \leq z_{i+1}(t^n) - \ell$. So from a road safety perspective, the model is rather optimistic.

We are now interested in taking the limit as $\ell \rightarrow 0$. We do this by increasing the number of vehicles such that $(N-1)\ell = 1$; furthermore we assume that (2.11) holds. Now the conditions are such that fundamental results of Crandall and Majda [4], see also [11, Thm. 3.9], can be applied. Thus we know that there is a function $y: \mathbb{R}_0^+ \times [0, 1] \rightarrow \mathbb{R}$, with $y \in C(\mathbb{R}^+; L^1([0, 1]))$, such that

$$y_\ell(t, x) \rightarrow y(t, x),$$

with the limit being in $C(\mathbb{R}^+; L^1([0, 1]))$, and that y is the unique entropy solution to the Cauchy problem

$$(2.20) \quad \begin{cases} y_t - V(y)_x = 0, & t > 0, x \in [0, 1], \\ y(0, x) = y_0(x). \end{cases}$$

If we do not have periodic conditions, this is supplemented with the boundary condition

$$y(t, 1) = M, \quad t > 0.$$

We remark that since the characteristic speeds of (2.20) are strictly negative, this boundary condition can be enforced strictly.

Note that the convergence of y_ℓ and the bounds $1 \leq y_\ell \leq M$, imply the convergence of $\tilde{\rho}_\ell = 1/y_\ell$ to some function $\tilde{\rho}$. We now proceed to show how $\tilde{\rho}$ is related to the solution of the LWR model.

We also define the discrete ‘‘Lagrange to Euler’’ map \tilde{z}_ℓ as follows. Let

$$\tilde{z}_{i+1/2}^n = \tilde{z}_{i-1/2}^n + \ell y_i^n, \quad \text{i.e., } \tilde{z}_{i+1/2}^n = z_{i+1}^n.$$

Since z_i^n solves (2.16), we also have that

$$\tilde{z}_{i+1/2}^{n+1} = \tilde{z}_{i+1/2}^n + \Delta t v_{i+1}^n.$$

Define $\tilde{z}_\ell(t^n, x_{i+1/2}) = \tilde{z}_{i+1/2}^n$, and by bilinear interpolation between these points. For later use we employ the notation for the value of \tilde{z}_ℓ at the edges of the ‘‘Lagrangian grid’’,

$$\begin{aligned} \tilde{z}_{i+1/2}(t) &= \frac{1}{\Delta t} \left((t^{n+1} - t) \tilde{z}_{i+1/2}^n + (t - t^n) \tilde{z}_{i+1/2}^{n+1} \right), \quad \text{for } t \in [t^n, t^{n+1}], \\ \tilde{z}^n(x) &= \frac{1}{\ell} \left((x_{i+1/2} - x) \tilde{z}_{i-1/2}^n + (x - x_{i-1/2}) \tilde{z}_{i+1/2}^n \right), \quad \text{for } x \in [x_{i-1/2}, x_{i+1/2}]. \end{aligned}$$

Observe that $\tilde{z}_{i-1/2}(t)$ coincides with the approximate trajectory of the vehicle starting at $z_i(0)$ calculated by the Euler method (2.16). Since y_ℓ is bounded, we can invoke the Arzelà–Ascoli theorem to establish the convergence

$$\lim_{\ell \rightarrow 0} \tilde{z}_\ell(t, x) = z(t, x),$$

with the limit being in $C([0, T] \times [0, 1])$ and $z \in C([0, T] \times [0, 1])$, and that

$$\frac{\partial z}{\partial x} = y, \quad \frac{\partial z}{\partial t} = V(y),$$

weakly. We have that the map $x \mapsto \tilde{z}_\ell(t, x)$ is invertible for each t , we denote the inverse map by x_ℓ , so that $x_\ell(t, z_\ell(t, x)) = x$. Define $z_{l,\ell}$ and z_r as in (2.13) and ρ_ℓ as in (2.14).

Note that if $z \in (z_{i-1/2}(t), z_{i+1/2}(t))$ and $t \in [t^n, t^{n+1})$, then

$$x_\ell(t, z) \in (x_{i-1/2}, x_{i+1/2}], \quad \rho_\ell(t, z) = \tilde{\rho}_i^n := \frac{1}{y_i^n}.$$

As before we have that

$$\rho_\ell(t, z) \rightarrow \rho(t, z) = \tilde{\rho}(t, x(t, z))$$

in $L^1([z_l, z_r])$ as $\ell \rightarrow 0$.

By Wagner’s result [16], we have proved the following theorem.

Theorem 2.6. *Assume that the function v satisfies (2.3). Let $\ell > 0$ and $N \in \mathbb{N}$, let $\{z_j\}_{j=1}^N$ satisfy (2.16), and assume that either we are in the periodic case $z_j \in [0, 1]$, or that z_N satisfies the boundary condition (2.2), with $y_N = M$. Assume that the initial positions of the vehicles $\{z_i(0)\}_{i=1}^N$ are such that we can define a bounded function y_0 by (2.11), and that (2.6) holds.*

Define the function $\rho_\ell(t, z)$ by (2.14). Let N and ℓ satisfy $(N - 1)\ell = 1$ and assume that the CFL-condition (2.19) holds.

As $\ell \rightarrow 0$, ρ_ℓ converges in $C([0, \infty); L^1)$ to the unique entropy solution ρ of the conservation law (2.12).

To illustrate the ideas in this paper we show how the method works in a concrete example. We have a periodic road in the interval $z \in [-1, 1]$, and choose to position N vehicles in this interval so that

$$\rho_\ell(0, z) \approx \frac{1}{2} (\cos(\pi z) + 1).$$

In Figure 1 we show the Lagrangian grid and the corresponding mapping to Eulerian coordinates for $N = 20$, and $t \in [0, 2]$. The vertical lines in the Eulerian coordinates




FIGURE 1. Left: the Lagrangian grid $\{(t^n, x_{i-1/2})\}_{i=1}^N$. Right: the Eulerian grid $\{(t^n, z_{i-1/2}^n)\}_{i=1}^N$. In both cases $n = 0, \dots, 40$.

are also the paths followed by the vehicles, and the grid in Eulerian coordinates is the result of applying the map z_ℓ to the rectangular grid depicted in Lagrangian coordinates on the left. In Figure 2, we show the approximate density ρ_ℓ at $t = 0$ and $t = 2$ in Eulerian coordinates. We see that the solution at $t = 2$ approximates




FIGURE 2. The approximate density ρ_ℓ for $t = 0$ and $t = 2$ in Eulerian coordinates.

the ubiquitous “ N -wave”.

REFERENCES

- [1] B. Argall, E. Cheleshkin, J. M. Greenberg, C. Hinde, and P.-J. Lin. A rigorous treatment of a follow-the-leader traffic model with traffic lights present. *SIAM J. Appl. Math.* 63(19): 149–168, 2002.
- [2] A. Aw, A. Klar, T. Materne, and M. Rascle. Derivation of continuum traffic flow models from microscopic follow-the-leader models. *SIAM J. Appl. Math.*, 63(1): 259–278, 2002.
- [3] R. M. Colombo and E. Rossi. On the micro-macro limit in traffic flow. *Rend. Sem. Math. Univ. Padova* 131:217–235, 2014.

- [4] M. G. Crandall and A. Majda. Monotone difference approximations for scalar conservation laws. *Math. Comp.*, 34:1–21, 1980.
- [5] E. Cristiani and S. Sahu. On the micro-to-macro limit for first-order traffic flow models on networks. *Networks and Heterogeneous Media* 11(3):395–413, 2016. doi:10.3934/nhm.2016002.
- [6] M. Di Francesco, S. Fagioli, and M. D. Rosini. Deterministic particle approximation of scalar conservation laws. Preprint, [arXiv:1605.05883v1](https://arxiv.org/abs/1605.05883v1), 2016.
- [7] M. Di Francesco, S. Fagioli, M. D. Rosini, and G. Russo. A deterministic particle approximation for non-linear conservation laws. In N. Bellomo, P. Degond, E. Tadmor (eds.) *Active Particles, Volume 1*, Birkhäuser, 2017, pp. 333–378.
- [8] M. Di Francesco and M. D. Rosini. Rigorous derivation of nonlinear scalar conservation laws from follow-the-leader type models via many particle limit. *Arch. Ration. Mech. Anal.*, 217(3):831–871, 2015.
- [9] P. Goatin and F. Rossi. A traffic flow model with non-smooth metric interaction: well-posedness and micro-macro limit. *Comm. Math. Sci.* 15(1):261–287, 2017.
- [10] Ke Han, Tao Yaob, and T. L. Friesz. Lagrangian-based hydrodynamic model: Freeway traffic estimation. Preprint [arXiv:1211.4619v1](https://arxiv.org/abs/1211.4619v1), 2012.
- [11] H. Holden and N. H. Risebro. *Front Tracking for Hyperbolic Conservation Laws*. Springer-Verlag, New York, 2015, Second edition.
- [12] H. Holden and N. H. Risebro. The continuum limit of Follow-the-Leader models — a short proof. *Discrete Cont. Dyn. Syst. A*, to appear.
- [13] M. J. Lighthill and G. B. Whitham. Kinematic waves. II. A theory of traffic flow on long crowded roads. *Proc. Roy. Soc. (London), Series A*, 229(1178):317–345, 1955.
- [14] P. I. Richards. Shockwaves on the highway. *Operations Research*, 4(1): 42–51, 1956.
- [15] E. Rossi. A justification of a LWR model based on a follow the leader description. *Discrete Cont. Dyn. Syst. Series S* 7(3): 579–591, 2014.
- [16] D. H. Wagner. Equivalence of the Euler and Lagrangian equations of gas dynamics for weak solutions. *J. Diff. Eqn.*, 68(1): 118–136, 1987.

(Holden)

DEPARTMENT OF MATHEMATICAL SCIENCES, NTNU NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, NO-7491 TRONDHEIM, NORWAY

E-mail address: helge.holden@ntnu.no

URL: <https://www.ntnu.edu/employees/holden>

(Risebro)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OSLO, P.O. BOX 1053, BLINDERN, NO-0316 OSLO, NORWAY

E-mail address: nilshr@math.uio.no