

RESEARCH ARTICLE

Open Access



# The association between dynamic lung volume and peak oxygen uptake in a healthy general population: the HUNT study

Øystein Rasch-Halvorsen<sup>1,3,5\*</sup> , Erlend Hassel<sup>1,3</sup>, Arnulf Langhammer<sup>2</sup>, Ben M. Brumpton<sup>3,4</sup> and Sigurd Steinshamn<sup>1,3</sup>

## Abstract

**Background:** Although dynamic lung volume is not considered a limiting factor of peak oxygen uptake ( $VO_{2peak}$ ) in healthy subjects, an association between forced expiratory lung volume in one second ( $FEV_1$ ) and  $VO_{2peak}$  has been reported in a healthy population aged 69 – 77 years. We hypothesized that a corresponding association could be found in a healthy general population including young and middle-aged subjects.

**Methods:** In a population-based study in Norway, we investigated the association between  $FEV_1$  above the lower limit of normal (LLN) and  $VO_{2peak}$  using linear regression and assessed the ventilatory reserve (VR) in healthy subjects aged 20 – 79 years ( $n = 741$ ).

**Results:** On average, one standard deviation (SD) increase in  $FEV_1$  was associated with 1.2 ml/kg/min (95% CI 0.7 – 1.6) higher  $VO_{2peak}$ . The association did not differ statistically by sex ( $p$ -value for interaction = 0.16) and was similar (0.9 ml/kg/min, 95% CI 0.2 – 1.5) in a sensitivity analysis including only never-smokers ( $n = 376$ ). In subjects below and above 45 years of age, corresponding estimates were 1.2 ml/kg/min (95% CI 0.5 – 1.8) and 1.2 ml/kg/min (95% CI 0.5 – 1.9), respectively. Preserved VR ( $\geq 20\%$ ) was observed in 66.6% of men and 86.4% of women.

**Conclusions:** Normal dynamic lung volume, defined as  $FEV_1$  above LLN, was positively associated with  $VO_{2peak}$  in both men and women, in never-smokers and in subjects below and above 45 years of age. The majority of subjects had preserved VR, and the results suggest that  $FEV_1$  within normal limits may influence  $VO_{2peak}$  in healthy subjects even when no ventilatory limitation to exercise is evident.

**Keywords:** Exercise test, Oxygen consumption, Respiratory function tests, Forced expiratory volume

## Background

Peak oxygen uptake ( $VO_{2peak}$ ) is an indicator of cardiorespiratory fitness (CRF) and a primary measure in cardiopulmonary exercise testing (CPET). The  $VO_{2peak}$  achieved at CPET is determined by the efficiency of the integrated oxygen transport and utilization system during maximal exercise [1].

The upper functional limit of the cardiovascular system is generally accepted as the primary limiting factor of  $VO_{2peak}$  in healthy subjects [2–4]. More specifically, the maximal cardiac stroke volume is a major factor determining the maximal capacity for oxygen transport during exercise [3]. In contrast, the functional reserves of the healthy pulmonary system for ventilation and gas exchange are considered well preserved in non-athletes exercising at sea level [5, 6]. The impact of dynamic lung volume on  $VO_{2peak}$  is commonly assessed indirectly by the ventilatory reserve (VR), reflecting the difference between estimated ventilatory capacity and measured minute ventilation at peak exercise ( $VE_{peak}$ ). Reduced VR is considered indicative of ventilatory limitation [7].

\* Correspondence: [oystein.rasch-halvorsen@ntnu.no](mailto:oystein.rasch-halvorsen@ntnu.no)

<sup>1</sup>K.G. Jebsen Center for Exercise in Medicine, Department of Circulation and Medical Imaging, Faculty of Medicine and Health Sciences, NTNU, Norwegian University of Science and Technology, Trondheim, Norway

<sup>3</sup>Clinic of Thoracic and Occupational Medicine, St. Olavs Hospital, Trondheim University Hospital, Trondheim, Norway

Full list of author information is available at the end of the article



Although normal dynamic lung volume is not the primary limiting factor of  $VO_{2peak}$ , forced expiratory lung volume in one second ( $FEV_1$ ) above the lower limit of normal (LLN) was recently found to be positively associated with  $VO_{2peak}$  in healthy elderly subjects [8]. However, this has not been studied in young and middle-aged subjects, and the potential influence of  $FEV_1$  above LLN on  $VO_{2peak}$  is largely unknown. In healthy subjects, CRF and  $VO_{2peak}$  are associated with risk of all-cause mortality [9]. Therefore, any influence of  $FEV_1$  on the overall efficiency of the oxygen transport chain could be important.

We aimed to investigate the association between normal dynamic lung volume and CRF in a general population and hypothesized that  $FEV_1$  above LLN is associated with  $VO_{2peak}$  in healthy subjects of a wide age-span. Additionally, we evaluated the VR and explored whether an association between  $FEV_1$  above LLN and  $VO_{2peak}$  differs by sex. Provided the finding of an association between  $FEV_1$  above LLN and  $VO_{2peak}$ , we also hypothesized an association between  $FEV_1$  above LLN and peak oxygen pulse ( $O_{2pulse}$ ), a proposed non-invasive estimator of cardiac stroke volume [10].

## Methods

### Study population

The Nord-Trøndelag Health Study (HUNT) is a population-based cohort study in Nord-Trøndelag County in Norway. The adult part of the third survey (HUNT3), conducted between October 2006 and June 2008, invited all residents aged 20 years or more ( $n = 93,860$ ) and 54.1% participated. The HUNT3 cohort profile has been published previously [11].

Two sub-studies in HUNT3, the Lung Study and the Fitness Study, provided measurements from spirometry and CPET, respectively. Selection into the Lung Study included a 10% random sample in addition to subjects with self-reported asthma, chronic obstructive pulmonary disease (COPD), respiratory symptoms and use of medication for asthma/COPD. In total, 18,244 subjects were invited and 66.5% responded. Selection into the Fitness Study included subjects without self-reported respiratory symptoms, cardiorespiratory disease and medication for hypertension or asthma/COPD. From four selected municipalities, 12,609 subjects were eligible for CPET. At the end of the study 36.7% had completed the test [12].

The present study represents an overlap between the Lung Study and the Fitness Study including only subjects without self-reported respiratory symptoms or cardiopulmonary disease. Subjects aged 20 – 79 years ( $n = 918$ ) were included. We excluded 28 subjects with  $FEV_1 < LLN$  and 53 subjects with peak respiratory exchange ratio ( $RER_{peak}$ )  $\leq 1.00$ , indicating potential submaximal effort in healthy subjects. Furthermore, 96 subjects had missing

values regarding covariates, heart rate reserve (HRR) or  $VE_{peak}$ , leaving 741 subjects for the statistical analyses.

### Dynamic lung volume and ventilatory reserve

Spirometry (MasterScope Jaeger version 5.1; JAEGER, Wuerzburg, Germany) was conducted in accordance with the American Thoracic Society/European Respiratory Society recommendations [13].  $FEV_1$  was chosen as a variable of dynamic lung volume because of high reproducibility and common use in clinical practice, and the LLN was defined as the fifth percentile ( $z$ -score =  $-1.645$ ). Predicted values and  $Z$ -scores were calculated using the Global Lung Function Initiative 2012 (GLI-2012) software [14].

Ventilatory reserve was calculated as  $VR = 1 - VE_{peak}/$  maximal voluntary ventilation (MVV). MVV was estimated by  $FEV_1 \times 40$  [15]. VR was dichotomized into preserved ( $\geq 20\%$ ) and reduced ( $< 20\%$ ) levels [16].

### Oxygen uptake, oxygen pulse and heart rate reserve

CPET was performed on a treadmill (DK7830; DK City, Taichung City, Taiwan) with a facemask (Hans Rudolph, Shawnee, KS, USA) and a heart rate (HR) monitor (Polar S610/RS400; Polar Electro Oy, Kempele, Finland). A portable system (MetaMax II; CORTEX Biophysik GmbH, Leipzig, Germany) measured  $VO_2$ , carbon dioxide output ( $VCO_2$ ) and VE. Incremental exercise was performed by an individualized protocol of increasing speed and/or incline until exhaustion as previously published [17].

Maximal oxygen uptake ( $VO_{2max}$ ), defined as  $RER_{peak} > 1.05$  and  $VO_2$  increase  $< 2$  ml/kg/min despite increased workload, was attained by 87.4%. As these predefined criteria were not met in all subjects, the term  $VO_{2peak}$  was used to designate the highest  $VO_2$  achieved. In the 12.6% (93 subjects) not meeting the  $VO_{2max}$  criteria,  $RER_{peak}$  and HRR were used to assess subject effort. No subjects had  $RER_{peak} \leq 1.00$  (exclusion criterion) and all but 8.6% (8 subjects) had  $HRR \leq 5\%$ .

$VO_{2peak}$  was calculated as the mean of the three highest 10-s values and indexed by bodyweight (ml/kg/min). Peak oxygen pulse was calculated as peak  $O_{2pulse}$  (ml/beat) =  $VO_{2peak}$  (ml/min)/peak HR (beat/min). Heart rate reserve was calculated as  $HRR = 1 - \text{peak HR}/(220 - \text{age})$ .

### Covariates

Covariates were chosen based on a priori knowledge of associations with both the explanatory variable and the outcome variable. Age (years) was recorded at time of CPET. Body mass index (BMI) was calculated as the ratio of bodyweight (kg) and height squared ( $m^2$ ). Physical activity index (PAI) was calculated as the product of scores given to answers on questions of frequency ( $< \text{once a week} = 0$ ,  $\text{once a week} = 1$ ,  $\text{two} - \text{three times a week} = 2$  and  $\text{approximately every day} = 3$ ), duration

(< 30 min = 1, ≥ 30 min = 1.5) and intensity of leisure time physical activity (Borg Rating of Perceived Exertion (RPE) 6 – 11 = 0, 12 – 13 = 5 and 14 – 20 = 10). Apart from redefining low, moderate and vigorous intensity as Borg RPE 6 – 11, 12 – 13 and 14 – 20, respectively [17], scores were assigned concordant with a non-exercise  $VO_{2peak}$  prediction model from HUNT3 [18]. Smoking status (never, former, daily and occasional) was obtained through questionnaire.

### Statistical analyses

Descriptive statistics of the total sample and the sample stratified by sex and VR were calculated as mean and standard deviation (SD) or number of observations and percentages unless otherwise specified. Continuous variables were compared using independent samples t tests.

The association between  $FEV_{1Z-score}$  and  $VO_{2peak}$  was modeled using linear regression. Regression coefficients with 95% confidence intervals (CI) were estimated in the total sample and in subgroups stratified by sex in three models. Model 1 estimated the crude association between  $FEV_{1Z-score}$  and  $VO_{2peak}$ . Model 2 adjusted for BMI and smoking status. Model 3 additionally adjusted for age and PAI. Sex was included as a covariate along with an interaction term ( $FEV_{1Z-score} \times sex$ ) when men and women were combined in model 3. BMI and PAI were included as continuous variables and smoking status and age (categorized into 10-year age-groups) as categorical. To assess potential residual confounding by smoking, the association between  $FEV_{1Z-score}$  and  $VO_{2peak}$  was investigated in a subgroup including only never-smokers ( $n = 376$ ) with adjustment for BMI, age, PAI and sex.

The association between  $FEV_{1Z-score}$  and  $VO_{2peak}$  was also assessed in subgroups stratified by dichotomized age (below and above 45 years) and adjusted for BMI, smoking status, PAI and sex.

The association between  $FEV_{1Z-score}$  and peak  $O_{2pulse}$  was assessed in the total sample and in subgroups stratified by sex. The estimates were adjusted for covariates equivalent to model 3 in the primary analyses. The interaction term ( $FEV_{1Z-score} \times sex$ ) was included when men and women were combined.

Model assumptions were assessed by residual plots and no violations were discovered. Statistical analyses were performed with IBM SPSS statistics 22.0 (IBM Corp., Armonk, NY, USA).

### Ethics

All subjects gave written informed consent. The study was approved by the Regional Committee for Medical and Health Research Ethics (REC Central 2015/1758).

### Results

Compared to women, men were of similar age, had similar BMI, reported similar level of physical activity, were less likely to report daily smoking, had similar HRR, lower VR, higher  $FEV_{1Z-score}$ , higher  $VO_{2peak}$ , higher peak  $O_{2pulse}$  and similar  $RER_{peak}$  (Table 1).

The proportions of subjects with reduced VR were 33.4% (men) and 13.6% (women). Age and BMI were similar in all subgroups stratified by sex and VR. In both sexes, subjects with preserved VR were less active than subjects with reduced VR. The HRR was negative and  $RER_{peak}$  similar in all subgroups (Table 2).

Compared to subjects with preserved VR, subjects with reduced VR had lower  $FEV_{1Z-score}$  in both men (difference 0.57, 95% CI 0.39 – 0.74) and women (difference 0.59, 95% CI 0.35 – 0.84), higher  $VO_{2peak}$  in both men (difference 5.0 ml/kg/min, 95% CI 3.1 – 6.9) and women (difference 1.2 ml/kg/min, 95% CI -1.0 – 3.5) and higher peak  $O_{2pulse}$  in both men (difference 2.4 ml/beat, 95% CI 1.6 – 3.2) and women (difference 1.6 ml/beat, 95% CI 0.7 – 2.5).

In all three models,  $FEV_{1Z-score}$  was positively associated with  $VO_{2peak}$  (Table 3). In women the point estimates ( $\beta$ ) were similar in model 2 ( $\beta = 1.6$ ) and model 3 ( $\beta = 1.6$ ). In men the point estimate in model 2 ( $\beta = 0.3$ ) was lower than in model 3 ( $\beta = 0.9$ ). In model 3, the association between  $FEV_{1Z-score}$  and  $VO_{2peak}$  did not differ statistically by sex ( $p$ -value for interaction = 0.16).

In the sensitivity analysis including only never-smokers,  $FEV_{1Z-score}$  was positively associated with  $VO_{2peak}$  ( $\beta = 0.9$ , 95% CI 0.2 – 1.5), as in the main model.

Stratified by dichotomized age,  $FEV_{1Z-score}$  was positively associated with  $VO_{2peak}$  in both categories (Table 4).

$FEV_{1Z-score}$  was positively associated with peak  $O_{2pulse}$  in both women and men (Table 5). The association did not differ statistically by sex ( $p$ -value for interaction = 0.67).

### Discussion

The main finding in this study was a positive association between  $FEV_1$  above LLN and  $VO_{2peak}$  in healthy subjects, aged 20 – 79 years, where the majority had preserved VR. The association was preserved after performing sub-analyses in both men and women, in never-smokers and in subjects below and above 45 years of age.

This is one of the first studies to investigate the association between normal dynamic lung volume and  $VO_{2peak}$ . Hassel et al. [8] reported positive associations between resting measures of lung function within the normal range and  $VO_{2peak}$  in healthy subjects aged 69 – 77 years. The present study expands on this work by assessing a more general population including young and middle-aged subjects. The results show that age-induced reduction in dynamic lung volume is not likely to fully explain the positive association between  $FEV_{1Z-score}$  and  $VO_{2peak}$ .

**Table 1** Descriptive statistics stratified by sex

	Men (n = 359)	Women (n = 382)
Age median (years) (Interquartile range)	46.5 (36.5 – 55.5)	45.0 (32.6 – 55.2)
Age group (years)		
20 – 29	65 (18.1%)	77 (20.2%)
30 – 39	45 (12.5%)	71 (18.6%)
40 – 49	99 (27.6%)	86 (22.5%)
50 – 59	91 (25.3%)	97 (25.4%)
60 – 69	48 (13.4%)	44 (11.5%)
70 – 79	11 (3.1%)	7 (1.8%)
BMI (kg/m <sup>2</sup> )	26.5 ± 3.0	25.4 ± 3.8
PAI median (Interquartile range)	15.0 (0.0 – 30.0)	15.0 (7.5 – 30.0)
Smoking status		
Never	186 (51.8%)	190 (49.7%)
Former	109 (30.4%)	122 (31.9%)
Daily	24 (6.7%)	40 (10.5%)
Occasional	40 (11.1%)	30 (7.9%)
HRR	-0.05 ± 0.08	-0.04 ± 0.07
VR	0.23 ± 0.12	0.30 ± 0.11
FEV <sub>1Z-score</sub>	0.28 ± 0.83	0.19 ± 0.87
VO <sub>2peak</sub> (ml/kg/min)	45.8 ± 9.1	36.8 ± 7.6
Peak O <sub>2pulse</sub> (ml/beat)	21.3 ± 3.8	14.0 ± 2.4
RER <sub>peak</sub>	1.14 ± 0.05	1.13 ± 0.06

BMI body mass index, PAI physical activity index, HRR heart rate reserve, VR ventilatory reserve, FEV<sub>1Z-score</sub> forced expiratory volume in one second Z-score, VO<sub>2peak</sub> peak oxygen uptake, O<sub>2pulse</sub> oxygen pulse, RER<sub>peak</sub> peak respiratory exchange ratio

Although the results show an association between FEV<sub>1Z-score</sub> and VO<sub>2peak</sub>, potential mechanisms are unclear. The majority of men (66.6%) and women (86.4%) had preserved VR, indicating residual capacity to increase ventilation at peak exercise. Preserved VR is commonly observed in healthy non-athletes performing

maximal exercise at sea level, reflecting large ventilatory capacity relative to metabolic demand. Furthermore, the normal physiological response at peak exercise is characterized by low HRR, indicating maximal utilization of cardiovascular capacity for oxygen transport. Combined, preserved VR and low HRR characterize cardiovascular limitation [5]. As this pattern was observed in the majority of subjects, the positive association between FEV<sub>1Z-score</sub> and VO<sub>2peak</sub> does not seem to be conditional on ventilatory limitation. Additionally, we found a positive association between FEV<sub>1Z-score</sub> and peak O<sub>2pulse</sub> in both sexes. As O<sub>2pulse</sub> is a proposed non-invasive estimator of cardiac stroke volume, we cannot rule out an influence of normal dynamic lung volume on cardiac output in healthy non-athletes. Although such a physiological interaction between the pulmonary and the cardiac system remains hypothetical, it may be appropriate to consider measures of dynamic lung volumes as covariates in future studies of exercise performance in healthy subjects.

Reduced VR is due to reduced ventilatory capacity and/or increased ventilatory demand, reflecting two different mechanisms for ventilatory limitation to exercise. Reduced VR secondary to reduced ventilatory capacity is commonly observed in COPD, where reduced dynamic lung volume with expiratory flow limitation and dynamic hyperinflation are important mechanisms of exercise intolerance [19]. Although FEV<sub>1Z-score</sub> was lower in subjects with reduced VR, all subjects had FEV<sub>1</sub> above LLN. Furthermore, exercise intolerance due to ventilatory limitation in COPD is reflected by low VO<sub>2peak</sub> [5] and high HRR indicating residual capacity of the cardiac system at peak exercise. In contrast, we found higher VO<sub>2peak</sub> in men and women with reduced VR, suggesting higher CRF in these subjects. Additionally, all subgroups had negative HRR, reflecting a physiological response with cardiovascular limitation independent on VR. Accordingly we do not consider mechanical ventilatory limitation likely to explain the

**Table 2** Descriptive statistics stratified by sex and ventilatory reserve (VR)

	Men		Women	
	VR ≥ 20% (n = 239)	VR < 20% (n = 120)	VR ≥ 20% (n = 330)	VR < 20% (n = 52)
Age median (years) (interquartile range)	46.0 (35.7 – 55.7)	46.7 (38.1 – 54.8)	44.1 (31.6 – 54.6)	47.4 (35.0 – 57.8)
BMI (kg/m <sup>2</sup> )	26.4 ± 3.1	26.8 ± 2.8	25.2 ± 3.7	26.8 ± 3.9
PAI median (interquartile range)	15.0 (0.0 – 30.0)	30.0 (8.1 – 30.0)	15.0 (7.5 – 30.0)	22.5 (10.0 – 30.0)
HRR	-0.05 ± 0.08	-0.05 ± 0.08	-0.04 ± 0.07	-0.04 ± 0.07
FEV <sub>1Z-score</sub>	0.47 ± 0.82	-0.09 ± 0.72	0.27 ± 0.85	-0.32 ± 0.77
VO <sub>2peak</sub> (ml/kg/min)	44.1 ± 8.5	49.1 ± 9.3	36.7 ± 7.5	37.9 ± 8.2
Peak O <sub>2pulse</sub> (ml/beat)	20.5 ± 3.5	22.9 ± 3.9	13.7 ± 2.3	15.4 ± 3.0
RER <sub>peak</sub>	1.14 ± 0.05	1.15 ± 0.05	1.13 ± 0.06	1.14 ± 0.06

BMI body mass index, PAI physical activity index, HRR heart rate reserve, FEV<sub>1Z-score</sub> forced expiratory volume in one second Z-score, VO<sub>2peak</sub> peak oxygen uptake, O<sub>2pulse</sub> oxygen pulse, RER<sub>peak</sub> peak respiratory exchange ratio

**Table 3** The association between FEV<sub>1Z-score</sub> and VO<sub>2peak</sub> (ml/kg/min) in total sample and stratified by sex

	n	Model 1 <sup>a</sup>		Model 2 <sup>b</sup>		Model 3 <sup>c</sup>	
		β	95% CI	β	95% CI	β	95% CI
Total	741	1.8	1.0 – 2.6	1.3	0.6 – 2.0	1.2	0.7 – 1.6
Men	359	0.8	-0.4 – 1.9	0.3	-0.7 – 1.3	0.9	0.2 – 1.6
Women	382	2.2	1.3 – 3.0	1.6	0.9 – 2.3	1.6	1.0 – 2.1

FEV<sub>1Z-score</sub> forced expiratory lung volume in one second Z-score, VO<sub>2peak</sub> peak oxygen uptake, β regression coefficient for FEV<sub>1Z-score</sub>  
<sup>a</sup>Crude model. <sup>b</sup>Adjusted for body mass index (BMI) and smoking status.  
<sup>c</sup>Adjusted for BMI, smoking status, age, physical activity index (PAI) with addition of sex in total sample model

association between FEV<sub>1Z-score</sub> and VO<sub>2peak</sub> in healthy subjects with FEV<sub>1</sub> above LLN.

In contrast to reduced VR from reduced ventilatory capacity, reduced VR from increased ventilatory demand is observed in healthy endurance trained subjects with high CRF [5]. High CRF with high VO<sub>2peak</sub> may be associated with respiratory muscle fatigue [20], and affect gas exchange if the upper limits for oxygen transport by the pulmonary system are exceeded [21]. Exercise induced arterial hypoxemia (EIAH) reduces VO<sub>2peak</sub> in significant proportions of endurance trained athletes [22], but is not usually observed in healthy subjects from a general population with lower VO<sub>2peak</sub>. Therefore, EIAH is unlikely to explain the positive association between FEV<sub>1Z-score</sub> and VO<sub>2peak</sub> found in this study.

Although mechanisms associated with mechanical ventilatory limitation and EIAH are unlikely to explain the association between FEV<sub>1Z-score</sub> and VO<sub>2peak</sub>, we cannot exclude this possibility. Stratification on dichotomized VR was considered but not performed, as VR is a likely collider on the association between FEV<sub>1Z-score</sub> and VO<sub>2peak</sub> potentially inducing biased estimates in stratified analyses. Furthermore, FEV<sub>1</sub> may be a marker of non-uniform ventilation due to small airway and/or parenchymal disease/properties causing increased heterogeneity in ventilation-perfusion relationships and variability in

**Table 4** The association between FEV<sub>1Z-score</sub> and VO<sub>2peak</sub> (ml/kg/min) in total sample and stratified by dichotomized age

	n	β	95% CI
Total	741	1.2	0.7 – 1.6
Age < 45 years	357	1.2	0.5 – 1.8
Age ≥ 45 years	384	1.2	0.5 – 1.9

FEV<sub>1Z-score</sub> forced expiratory lung volume in one second Z-score, VO<sub>2peak</sub> peak oxygen uptake, β regression coefficient for FEV<sub>1Z-score</sub> adjusted for body mass index (BMI), smoking status, physical activity index (PAI) and sex with addition of age in total sample model

**Table 5** The association between FEV<sub>1Z-score</sub> and peak O<sub>2pulse</sub> (ml/beat) in total sample and stratified by sex

	n	β	95% CI
Total	741	0.33	0.10 – 0.56
Men	359	0.32	-0.08 – 0.73
Women	382	0.39	0.14 – 0.63

FEV<sub>1Z-score</sub> forced expiratory lung volume in one second Z-score, O<sub>2pulse</sub> oxygen pulse, β regression coefficient for FEV<sub>1Z-score</sub> adjusted for body for mass index (BMI), smoking status, age, physical activity index (PAI) with addition of sex in total sample model

gas exchange, which is not excluded by preserved VR. Accordingly, our study raises further questions that potentially impact current understanding of respiratory and exercise physiology. Future studies should focus on uncovering the underlying mechanisms of the associations we report, and are likely to benefit from including invasive measures from arterial blood gases.

The strengths of this study include the high number of subjects from a general population. Direct measurement of VO<sub>2</sub> is the gold-standard for evaluating CRF and was used in this study. Furthermore, we used the GLI-2012 reference equations, found valid for the Norwegian population [23], taking into account age, sex and height related variance in FEV<sub>1</sub>. Although we did not a priori consider age and PAI as confounders of the association between FEV<sub>1Z-score</sub> and VO<sub>2peak</sub>, these variables are known predictors of VO<sub>2peak</sub> [18], and were included as covariates in the final models. Confounding was detected in men and accordingly adjusted for. Although we can only test for associations, we consider an influence of both FEV<sub>1Z-score</sub> on VO<sub>2peak</sub> and FEV<sub>1Z-score</sub> on peak O<sub>2pulse</sub> more likely than vice versa, thereby reducing the limitations from the cross-sectional study design.

There are some limitations. The external validity may suffer, as subjects recruited to a fitness study may be more active and fit than the general population. All subjects were considered to be healthy, but eligibility for the HUNT Fitness study was evaluated only through self-report. Although secondary criteria of maximal subject effort were met, measured VO<sub>2peak</sub> may have underestimated true VO<sub>2max</sub> in the 12.6% not meeting the VO<sub>2max</sub> criteria [24], potentially influencing the precision of the estimates. Additionally, lack of measurements from arterial blood gases limits further discussion on potential physiological mechanisms related to variability in gas exchange. Finally, O<sub>2pulse</sub> is an indirect approximate of cardiac stroke volume, and the association between FEV<sub>1Z-score</sub> and peak O<sub>2pulse</sub> permits only speculation on a potential influence of normal dynamic lung volume on cardiac performance during exercise.

## Conclusions

Normal dynamic lung volume, defined as FEV<sub>1</sub> above LLN, was positively associated with VO<sub>2peak</sub> in both men and women, in never-smokers and in subjects below and above 45 years of age. The majority of subjects had preserved VR, and the results suggest that FEV<sub>1</sub> within normal limits may influence VO<sub>2peak</sub> in healthy subjects even when no ventilatory limitation to exercise is evident. Future studies should focus on understanding the underlying mechanisms of this association. If causality can be established, it would further emphasize the importance of respiratory health.

## Abbreviations

BMI: Body mass index; CI: Confidence interval; COPD: Chronic obstructive pulmonary disease; CPET: Cardiopulmonary exercise test; CRF: Cardiorespiratory fitness; EIAH: Exercise induced arterial hypoxemia; FEV<sub>1</sub>: Forced expiratory lung volume in one second; GLI-2012: Global Lung Function Initiative 2012; HR: Heart rate; HRR: Heart rate reserve; HUNT: The Nord-Trøndelag Health Study; LLN: Lower limit of normal; MVV: Maximal voluntary ventilation; O<sub>2pulse</sub>: Oxygen pulse; PAL: Physical activity index; REC: Regional Committee for Medical and Health Research Ethics; RER<sub>peak</sub>: Peak respiratory exchange ratio; RPE: Rating of perceived exertion; SD: Standard deviation; VCO<sub>2</sub>: Carbon dioxide output; VE<sub>peak</sub>: Peak minute ventilation; VO<sub>2max</sub>: Maximal oxygen uptake; VO<sub>2peak</sub>: Peak oxygen uptake; VR: Ventilatory reserve

## Acknowledgements

The Nord-Trøndelag Health Study (The HUNT Study) is a collaboration between HUNT Research Centre (Faculty of Medicine and Health Sciences, NTNU, Norwegian University of Science and Technology), Nord-Trøndelag County Council, Central Norway Health Authority, and the Norwegian Institute of Public Health.

## Funding

This project has been made possible by funding from the Norwegian ExtraFoundation for Health and Rehabilitation. The funding body has not been involved in any aspect of the study regarding design of the study or collection, analysis and interpretation of data or in writing of the manuscript.

## Availability of data and materials

The data that support the findings of this study are available from HUNT Research Centre but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available. Data are however available for export from the HUNT databank given approval of application to HUNT Research Centre (<http://www.ntnu.edu/hunt/data>).

## Authors' contributions

ØR-H analyzed the patient data and was a major contributor to the design of the study, interpretation of the data and in writing the manuscript. EH, AL, BMB and SS were all major contributors to the design of the study, interpretation of the data and in writing the manuscript. All authors read and approved the final manuscript.

## Ethics approval and consent to participate

All subjects gave written informed consent. The study was approved by the Regional Committee for Medical and Health Research Ethics (REC Central 2015/1758).

## Consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Author details

<sup>1</sup>K.G. Jebsen Center for Exercise in Medicine, Department of Circulation and Medical Imaging, Faculty of Medicine and Health Sciences, NTNU, Norwegian University of Science and Technology, Trondheim, Norway. <sup>2</sup>Department of Public Health and Nursing, Faculty of Medicine and Health Sciences, NTNU, Norwegian University of Science and Technology, Trondheim, Norway. <sup>3</sup>Clinic of Thoracic and Occupational Medicine, St. Olavs Hospital, Trondheim University Hospital, Trondheim, Norway. <sup>4</sup>K.G. Jebsen Center for Genetic Epidemiology, Department of Public Health and Nursing, Faculty of Medicine and Health Sciences, NTNU, Norwegian University of Science and Technology, Trondheim, Norway. <sup>5</sup>NTNU, Faculty of Medicine and Health Sciences, Department of Circulation and Medical Imaging, 8905, 7491 Trondheim, Norway.

Received: 19 July 2018 Accepted: 10 December 2018

Published online: 06 January 2019

## References

- Wagner PD. Determinants of maximal oxygen transport and utilization. *Annu Rev Physiol.* 1996;58:21–50.
- Blomqvist CG, Saltin B. Cardiovascular adaptations to physical training. *Annu Rev Physiol.* 1983;45:169–89.
- Bassett DR Jr, Howley ET. Limiting factors for maximum oxygen uptake and determinants of endurance performance. *Med Sci Sports Exerc.* 2000;32(1):70–84.
- Stickland MK, Butcher SJ, Marciniuk DD, Bhutani M. Assessing exercise limitation using cardiopulmonary exercise testing. *Pulm Med.* 2012;2012:824091.
- American Thoracic S. American College of Chest P. ATS/ACCP statement on cardiopulmonary exercise testing. *Am J Respir Crit Care Med.* 2003;167(2):211–77.
- Force ERST, Palange P, Ward SA, Carlsen KH, Casaburi R, Gallagher CG, et al. Recommendations on the use of exercise testing in clinical practice. *Eur Respir J.* 2007;29(1):185–209.
- Johnson BD, Weisman IM, Zeballos RJ, Beck KC. Emerging concepts in the evaluation of ventilatory limitation during exercise: the exercise tidal flow-volume loop. *Chest.* 1999;116(2):488–503.
- Hassel E, Stensvold D, Halvorsen T, Wisloff U, Langhammer A, Steinshamn S. Association between pulmonary function and peak oxygen uptake in elderly: the generation 100 study. *Respir Res.* 2015;16:156.
- Kodama S, Saito K, Tanaka S, Maki M, Yachi Y, Asumi M, et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA.* 2009;301(19):2024–35.
- Stringer WW, Hansen JE, Wasserman K. Cardiac output estimated noninvasively from oxygen uptake during exercise. *J Appl Physiol* (1985). 1997;82(3):908–12.
- Krokstad S, Langhammer A, Hveem K, Holmen TL, Midthjell K, Stene TR, et al. Cohort profile: the HUNT study, Norway. *Int J Epidemiol.* 2013;42(4):968–77.
- Aspenes ST, Nilsen TI, Skaug EA, Bertheussen GF, Ellingsen O, Vatten L, et al. Peak oxygen uptake and cardiovascular risk factors in 4631 healthy women and men. *Med Sci Sports Exerc.* 2011;43(8):1465–73.
- Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, et al. Standardisation of spirometry. *Eur Respir J.* 2005;26(2):319–38.
- Quanjer PH, Stanojevic S, Cole TJ, Baur X, Hall GL, Culver BH, et al. Multi-ethnic reference values for spirometry for the 3–95-yr age range: the global lung function 2012 equations. *Eur Respir J.* 2012;40(6):1324–43.
- Campbell SC. A comparison of the maximum voluntary ventilation with the forced expiratory volume in one second: an assessment of subject cooperation. *J Occup Med.* 1982;24(7):531–3.
- Guazzi M, Adams V, Conraads V, Halle M, Mezzani A, Vanhees L, et al. EACPR/AHA joint scientific statement. Clinical recommendations for cardiopulmonary exercise testing data assessment in specific patient populations. *Eur Heart J.* 2012;33(23):2917–27.
- Nes BM, Janszky I, Aspenes ST, Bertheussen GF, Vatten LJ, Wisloff U. Exercise patterns and peak oxygen uptake in a healthy population: the HUNT study. *Med Sci Sports Exerc.* 2012;44(10):1881–9.

18. Nes BM, Janszky I, Vatten LJ, Nilsen TI, Aspenes ST, Wisloff U. Estimating V.O<sub>2</sub> peak from a nonexercise prediction model: the HUNT study, Norway. *Med Sci Sports Exerc.* 2011;43(11):2024–30.
19. O'Donnell DE, Reville SM, Webb KA. Dynamic hyperinflation and exercise intolerance in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 2001;164(5):770–7.
20. Queslati F, Berriri A, Boone J, Ahmaidi S. Respiratory muscle strength is decreased after maximal incremental exercise in trained runners and cyclists. *Respir Physiol Neurobiol.* 2018;248:25–30.
21. Dempsey JA, Hanson PG, Henderson KS. Exercise-induced arterial hypoxaemia in healthy human subjects at sea level. *J Physiol.* 1984;355:161–75.
22. Dempsey JA, Wagner PD. Exercise-induced arterial hypoxemia. *J Appl Physiol* (1985). 1999;87(6):1997–2006.
23. Langhammer A, Johannessen A, Holmen TL, Melbye H, Stanojevic S, Lund MB, et al. Global lung function initiative 2012 reference equations for spirometry in the Norwegian population. *Eur Respir J.* 2016;48(6):1602–11.
24. Poole DC, Jones AM. Measurement of the maximum oxygen uptake Vo<sub>2</sub>max: Vo<sub>2</sub>peak is no longer acceptable. *J Appl Physiol* (1985). 2017; 122(4):997–1002.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

