cmgh RESEARCH LETTER

Subtle Protective Roles of Clusterin in Gastric Metaplasia After Acute Oxyntic Atrophy

Gastric cancer is preceded by mucosal remodeling like spasmolytic polypeptideexpressing metaplasia (SPEM) and intestinal metaplasia. The cytoprotective protein clusterin (CLU) is highly expressed in oxyntic mucosa during the emergence of metaplasia, but it is unknown whether CLU promotes or counteracts the process. $\hat{1}^{-4}$ Here, we use clusterin knockout (CLU-KO) mice to examine the role of CLU in the emergence, recovery, and repair of gastric metaplasia after tamoxifeninduced acute oxyntic atrophy.5 Detailed methodology is described in Supplementary Materials and Methods.

The stomachs of CLU-KO mice had normal gastric gland morphology and normal number of mature cell types, although with a tendency toward pit cell/foveolar and enterochromaffinlike (ECL) cell hyperplasia (Figure 1A and *B*). Interestingly, this could be owing to the hypergastrinemia in CLU-KO mice (Figure 1C).⁶ We performed RNA sequencing to examine whether the absence of CLU affected the gene expression signature of the oxyntic mucosa, showing 62 differentially expressed genes (DEGs) compared with wild-type mice (Supplementary Table 1). Among them were the ECL cell marker histidine decarboxylase and the putative intestinal metaplasia marker deleted in malignant brain tumors 1. However, although several of the genes were involved in CLU-relevant processes,¹ we could not define a gene profile characterizing expression untreated CLU-KO mice compared with wild-type mice. Overall, our results indicate that CLU partly affects gastric epithelial homeostasis because the absence of CLU leads to hypergastrinemia and slightly increased numbers of pit and ECL cells.

In tamoxifen-treated wild-type mice. CLU was increased and highly expressed in proliferative and metaplastic glands with a temporal profile that correlated with emergence and repair (Figure 1D), suggesting that CLU participates in the metaplastic process. Unexpectedly, although they lacked CLU, tamoxifen induced metaplasia with a similar degree of parietal cell atrophy (CLU-KO, 83%; vs wild-type, 87%) (Figure 1E and F), number of apoptotic cells (Figure 1F), transdifferentiating chief cells, proliferating and SPEM-cells (Figure 1G and H) as in wild-type mice. Also, CLU-KO mice were not more susceptible to a lower dose of tamoxifen (Supplementary Figure 1). Hence, CLU is not a necessary promoting factor in the emergence of gastric metaplasia.

Gene expression analysis was performed to further address the role of clusterin on a molecular level (Supplementary Tables 1-11). There was an increased and prominent transcriptional difference (1026 vs 62 DEGs in untreated mice) between the emerging metaplasia (day 3) of CLU-KO and wild-type mice (Supplementary Tables 2 and 3), indicating distinct underlying molecular events. Apparently, the metaplastic mucosa of CLU-KO mice showed more mitochondrial dysfunction, which fits with the known functions of CLU as both a sensor and protector against oxidative stress.7 Intriguingly, oxidative stress resulting from chronic inflammation might play a central role in gastric tumorigenesis,⁸ suggesting that CLU deficiency will be a disadvantage. Indeed, the metaplastic mucosa of CLU-KO mice expressed a higher number of genes associated with gastric metaplasia and (cancer) stem cells,^{3,8,9} and lower levels of mucosal protective factors and suppressors tumor (Figure 2Aand Supplementary Tables 4-6).¹⁰ Thus, although it is not clearly evident histologically, transcriptional

profiling indicates that the emergence of metaplasia is more advanced in CLU-KO than wild-type mice.

During early recovery (day 7 + 10), the metaplastic mucosa of both CLU-KO and wild-type mice were still substantially changed from normal. However, when directly comparing CLU-KO with wild-type mice at this stage, there were only subtle histologic and transcriptional differences (Figure 2*A*, Supplementary Figure 2, and Supplementary Tables 6–8), possibly suggesting more similar mechanisms operating during early recovery.

In contrast, after repair (day 21), histologic analysis showed normal gland morphology and a restored number of mature cell lineages in tamoxifen-treated wild-type mice. However, the oxyntic mucosa of CLU-KO mice tended to be more hyperplastic, with slightly more parietal and neck cell-tochief cell-transition cells per gland (Figure 2B-D). In addition, the gene expression profile of tamoxifen-treated CLU-KO mice was altered markedly from vehicle-treated controls (409 DEGs) (Supplementary Tables 9 and 10), which was not found in wild-type mice (44 DEGs). CLU-KO mice had continuously down-regulated mature cell lineage marker genes, including the chief cell marker Mist1 and the pit cell gene Muc5ac, up-regulation of the tuft cell marker Dclk1, and still several dysregulated gastric metaplasiaassociated and mucosal protective genes, such as Mmp12, Sox9, and Gkn2 (Figure 2A, E, and F).

Thus, in the absence of CLU, restoration of normal mucosa, with re-expression of all cell lineage marker genes and disappearance of metaplastic gene signatures, seemed to be delayed, again suggesting that CLU-KO mice showed a more advanced form of metaplasia and a hampered repair compared with wild-type mice. Hence, instead of being a necessary promoting factor, CLU might protect against emergence and premalignant progression of gastric metaplasia.



Figure 1. Clusterin is not a necessary factor for gastric epithelial homeostasis or tamoxifen-induced gastric metaplasia. (A) Representative H&E images showing normal gland morphology of oxyntic and antral mucosa in an untreated CLU-KO mouse. (B) Quantification of different mature gastric cell lineages per gland in wild-type (WT) (n = 3-4) and CLU-KO (n = 6) mice. (C) Serum gastrin levels (pmol/L) in untreated WT (n = 11) and CLU-KO (n = 11) mice. (D) In situ hybridization showing expression of clusterin messenger RNA (mRNA) (Clu; brown) and triple immunofluorescence staining showing CLU (red) expression in proliferating (Ki67; light blue) metaplastic cells (lectin griffonia simplicifolia II [GSII]; green) of oxyntic mucosa from vehicle- and tamoxifen-treated WT mice on days 3, 7, and 21. (E) Representative H&E images of oxyntic mucosa and immunohistochemistry staining showing expression of H⁺/K⁺adenosine triphosphatase [ATPase] β -subunit (parietal cells; brown) in oxyntic mucosa from vehicle- and tamoxifentreated CLU-KO mice on day 3. (F) Quantification of parietal cells (H+/K+-ATPase β-subunit) and apoptotic cells (cleaved caspase 3-positive [CASP3]) per gland in oxyntic mucosa from vehicle- (n = 4 in each strain) and tamoxifentreated CLU-KO (n = 6) and WT (n = 7) mice on day 3. (G) Double immunofluorescence staining showing cells coexpressing (yellow) pepsinogen II (PEPII) (red) and GSII (green) and cells co-expressing Ki67 (red) and GSII (green) in oxyntic mucosa from vehicle- and tamoxifen-treated CLU-KO mice on day 3. (H) Quantification of cells co-expressing PEPII and GSII (vellow) or cells co-expressing Ki67 and GSII per gland in oxyntic mucosa from vehicle- (n = 4 in each strain) and tamoxifen-treated CLU-KO (n = 6) and WT (n = 7) mice on day 3. Nuclei were counterstained with hematoxylin (blue) or DAPI (blue). Black squares, WT; blue circles, CLU-KO. Each dot represents an individual animal and the black lines mark the means. Scale bars: 100 μ m. Statistical significance: *P < .05, ***P < .0001 by Student t test (B and C) or 1-way analysis of variance with the Sidak multiple comparison test (F and H). D, day; DAPI, 4',6-diamidino-2phenylindole; NE, neuroendocrine; TAM, tamoxifen; Veh, vehicle.



Figure 1. (continued).



Figure 2. Tamoxifen induces transcriptionally more advanced gastric metaplasia with delayed repair in CLU-KO mice compared with wild-type (WT) mice. (A) Gene expression levels (log₂ fold-change [FC]) of metaplasia-associated genes, (cancer) stem cell-associated, and mucosal protective factors expressed at different levels (statistically significant for at least 1 time point) in tamoxifen-treated CLU-KO (n = 4-6) (blue) and WT (n = 3-7) (black) mice compared with vehicle-treated controls (n = 9 in each strain) at emergence (day 3), recovery (day 7 + 10), and repair (day 21) (Supplementary Table 6). #Metaplasia-associated genes that were dysregulated only in CLU-KO mice on day 3. (B) Representative H&E images of oxyntic mucosa from tamoxifen-treated WT and CLU-KO mice on day 21. (C) Quantification of parietal cells H⁺/K⁺ -ATPase β subunit and cells co-expressing Ki67 and lectin griffonia simplicifolia II [GSII] per gland in oxyntic mucosa from tamoxifen-treated WT (n = 5) and CLU-KO (n = 4) mice on day 21. (D) Quantification of cells expressing pepsinogen II (PEPII; red), GSII (green), or co-expressing PEPII and GSII (yellow) per gland in oxyntic mucosa from tamoxifen-treated WT (n = 5) and CLU-KO (n = 4) mice on day 21. Data are presented as means - SD. (E) Gene expression levels ($\log_2 FC$) of mature gastric cell lineage marker genes in tamoxifen-treated CLU-KO (n = 4) (blue) and WT (n = 5) (black) mice compared with vehicle-treated controls (n = 9 in each strain) at repair (day 21) (Supplementary Tables 9 and 10). (F) Immunohistochemistry staining showing expression of DCLK1 (brown) and quantification of cells expressing DCLK1 per gland in oxyntic mucosa from vehicle- and tamoxifen-treated WT and CLU-KO mice (n = 4 in each group) on day 21. Nuclei were counterstained with hematoxylin (blue). Black squares, WT; blue circles, CLU-KO. Each dot represents an individual animal and the black lines mark the means. Statistical significance: *P < .05, ***P < .0001 by Student t test (C and D) or 1-way analysis of variance with the Sidak multiple comparison test (F). Scale bars: 100 µm (B) and 50 µm (F). D, day; TAM, tamoxifen; Veh, vehicle.



PÅL VANGE,¹ TORUNN BRULAND,^{1,2} BJØRN MUNKVOLD,¹ ELIN SYNNØVE RØYSET,^{1,3} MARTIN GLEAVE,⁴ INGUNN BAKKE,^{1,2}

¹Department of Clinical and Molecular Medicine, Faculty of Medicine and Health Sciences, NTNU–Norwegian University of Science and Technology, Trondheim, Norway

²Clinic of Medicine, St. Olav's University Hospital, Trondheim, Norway

³Department of Pathology, St. Olav's University Hospital, Trondheim, Norway

⁴Vancouver Prostate Centre, Vancouver, Canada Figure 2. (continued).

Correspondence author e-mail: ingunn. bakke@ntnu.no.

References

- 1. Rohne P, et al. Biomol Concepts 2016;17:1–15.
- Vange P, et al. PLoS One 2017; 12:e0184514.
- 3. Weis VG, et al. Gut 2013; 62:1270–1279.
- 4. Shimizu T, et al. J Pathol 2016; 239:399–410.
- 5. Huh WJ, et al. Gastroenterology 2012;142:21–24 e7.
- 6. Feng R, et al. Gastroenterology 2014;147:655–666 e9.
- 7. Trougakos IP. Gerontology 2013; 59:514–523.

- 8. Seishima R, et al. Cancer Prev Res (Phila) 2015;8:492–501.
- Engevik AC, et al. Cell Mol Gastroenterol Hepatol 2016;2:605–624.
- 10. Menheniott TR, et al. J Clin Invest 2016;126:1383–1400.

Abbreviations used in this letter: CLU, clusterin; CLU-KO, clusterin knockout; DEG, differentially expressed gene; ECL, enterochromaffin-like cell; SPEM, spasmolytic polypeptide-expressing metaplasia.

Most current article

© 2019 The Authors. Published by Elsevier Inc. on behalf of the AGA Institute. This is an open access article under the CC BY-NC-ND license (http://creativecommons. org/licenses/by-nc-nd/4.0/).

org/licenses/by-nc-nd/4.0/). 2352-345X https://doi.org/10.1016/j.jcmgh.2018.09.013

Received February 6, 2018. Accepted September 7, 2018.

Acknowledgments

The authors thank Anne Kristensen for skillful technical assistance and Knut Grøn and the animal unit of the Comparative Medicine Core Facility, NTNU, for assistance with the animal models. The imaging analyses were performed in collaboration with the Cellular and Molecular Imaging Core Facility, NTNU. The RNA sequencing and bioinformatic analyses were provided by the Genomics Core Facility, NTNU. The Comparative Medicine Core Facility, Cellular and Molecular Imaging Core Facility are funded by the Faculty of Medicine and Health Sciences at

NTNU and the Central Norway Regional Health Authority.

Author contributions

Pål Vange, Torunn Bruland, and Ingunn Bakke contributed to the conception and design of the study; Martin Gleave provided transgenic mice; Pål Vange, Ingunn Bakke, and Bjørn Munkvold performed the experiments; Pål Vange, Torunn Bruland, Bjørn Munkvold, Elin Synnøve Røyset, and Ingunn Bakke contributed to the assembly, analysis, and interpretation of data; Pål Vange, Torunn Bruland, and Ingunn Bakke wrote the manuscript; and Pål Vange, Torunn Bruland, Bjørn Munkvold, Elin Synnøve Røyset, Martin Gleave, and Ingunn Bakke were involved in the revision and final approval of the submitted and published version of the manuscript.

Conflicts of interest

The authors disclose no conflicts.

Funding

This work was supported by The Liaison Committee between the Central Norway Regional Health Authority and NTNU–Norwegian University of Science and Technology, Trondheim, Norway; The Faculty of Medicine and Health Sciences at NTNU; The Cancer Fund at St. Olav's University Hospital, Trondheim, Norway; and Familien Blix fond til fremme av medisinsk forskning. No study sponsor had any role in the study design, collection, analysis, or interpretation of data.



Lack of **clusterin** in homeostasis caused subtle changes with hypergastrinemia, development of slightly more advanced acute metaplasia and hampered repair



Supplemental Graphical Summary.