

VERTICAL-HORIZONTAL DECOMPOSITION OF LAPLACIANS AND COHOMOLOGIES OF MANIFOLDS WITH TRIVIAL TANGENT BUNDLES

SÖNKE ROLLENSKE, ADRIANO TOMASSINI AND XU WANG

ABSTRACT. In this paper we obtain a vertical-horizontal decomposition formula of Laplacians on manifolds with a special foliation structure. Two Nomizu type theorems for cohomologies of nilmanifolds follow as applications.

CONTENTS

1. Introduction	2
2. Motivations	3
2.1. First Motivation: Künneth formula	3
2.2. Second motivation: Nomizu-type theorems	4
3. Foliations of nilpotent type	5
3.1. Nilpotent foliation	5
3.2. Vertical-Horizontal decomposition of d	6
3.3. Vertical-Horizontal decomposition of $\bar{\partial}$	7
4. Vertical–Horizontal decomposition of Laplacians	8
4.1. Fundamental theorem	8
4.2. Proof of the real case	9
4.3. Proof of the complex case	11
5. An example: the Kodaira–Thurston manifold	12
5.1. Nilpotent fibrations	15
6. Nilpotent frame	16
6.1. Real case	16
6.2. Complex case	18
7. Proof of the main theorem	19
7.1. Cohomology description of Theorem 4.1	19
7.2. Spectral sequence for double complex	20
7.3. The final proof	22
7.4. Proof of the strong version of Cordero-Fernández-Gray-Ugarte’s theorem	22
8. An explicit version of our main theorem	22
9. Further examples	24
9.1. A nilmanifold not satisfying Theorem 8.2	24

Date: August 29, 2019.

2010 Mathematics Subject Classification. 53C25, 53C55.

Key words and phrases. Künneth formula; horizontal lift; vertical form; Laplacian; nilpotent foliation; torus fibration; Nomizu type theorem; nilpotent group.

This work was partially supported by the Project PRIN “Varietà reali e complesse: geometria, topologia e analisi armonica” and by GNSAGA of INdAM.

9.2. Nilpotent frame in a non-nilpotent Lie group	25
References	26

1. INTRODUCTION

Let X be an n -dimensional compact smooth manifold. Assume that the tangent bundle T_X of X is trivial (thus the cotangent bundle T_X^* is also trivial). Let

$$\Phi := \{\sigma^1, \dots, \sigma^n\}$$

be a global smooth frame of T_X^* . Inspired by [20] and [9], we introduce the following

Definition 1.1. *We call Φ a nilpotent frame if*

$$(1.1) \quad d\sigma^j = \sum_{k,l>j} A_{kl}^j \sigma^k \wedge \sigma^l, \quad \forall 1 \leq j \leq n,$$

where A_{kl}^j are “real constants”.

In the complex case, assume that the holomorphic tangent bundle $\wedge^{1,0}T_X$ of a compact complex manifold X is *smoothly* trivial (may not be trivial as a holomorphic vector bundle). Now assume that the complex dimension of X is n . Let

$$\Psi := \{\xi^1, \dots, \xi^n\}$$

be a global smooth frame of $\wedge^{1,0}T_X^*$. We shall use the following

Definition 1.2. *We call Ψ a complex nilpotent frame if*

$$(1.2) \quad d\xi^j = \sum_{k,l>j} B_{kl}^j \xi^k \wedge \xi^l + \sum_{k,l>j} B_{k\bar{l}}^j \xi^k \wedge \bar{\xi}^l, \quad \forall 1 \leq j \leq n,$$

where B_{kl}^j and $B_{k\bar{l}}^j$ are “complex constants”.

We have the following generalization of the main results in [20], [9] and [13].

Main Theorem: *Let X be a compact smooth (resp. complex) manifold. Assume that T_X^* (resp. $\wedge^{1,0}T_X^*$) possesses a nilpotent (resp. complex nilpotent) frame Φ (resp. Ψ). Then every de Rham (resp. Dolbeault) cohomology class of X can be represented by \mathbb{R} (resp. \mathbb{C}) linear combination of finite wedge products of forms in Φ (resp. $\Psi \cup \bar{\Psi}$).*

Remark 1: Denote by A^* (resp. $A^{*,*}$) the finite dimension \mathbb{R} (resp. \mathbb{C}) linear space spanned by wedge products of Φ (resp. $\Psi \cup \bar{\Psi}$). Then we know that the d -cohomology is well defined on A^* , the $\bar{\partial}$ -cohomology is well defined on $A^{*,*}$ and they are also called *the Lie algebra cohomologies*. Let us denote them by $H_{d,\Phi}^*$ and $H_{\bar{\partial},\Psi}^{*,*}$ respectively. Then our main theorem is equivalent to say that

$$H_d^* \simeq H_{d,\Phi}^*, \quad H_{\bar{\partial}}^{*,*} \simeq H_{\bar{\partial},\Psi}^{*,*},$$

where H_d^* (resp. $H_{\bar{\partial}}^{*,*}$) denotes the usual *de Rham* (resp. *Dolbeault*) cohomology group. See section 8 for a more explicit description of $H_{d,\Phi}^*$ and $H_{\bar{\partial},\Psi}^{*,*}$ in certain cases and section 9 for related results.

Remark 2: The main ingredient in the proof of our main theorem is the following vertical-horizontal decomposition of Laplacians (see Theorem 4.1)

$$\square_d = \square_{d^v} + \square_{d^h + R_d},$$

associated to the following decomposition

$$d = d^h + d^v + R_d,$$

of d on a smooth manifold with a special foliation structure, where d^h only increases the horizontal degree, d^v only increases the vertical degree and the remaining term R_d is a tensor (see [1], [2], [3], [21], [18], [19], [26] for the background and related results).

Remark 3: The proof of our main theorem in section 7 also gives the following result: Let X be a compact smooth manifold. Assume that $\mathbb{C} \otimes T_X^*$ possesses a nilpotent frame Φ . Then every complex de Rham cohomology class of X can be represented by \mathbb{C} linear combination of finite wedge products of forms in Φ .

Our main theorem suggests to study the following problem:

(\star): *Let G be a Lie group, let Γ be a discrete subgroup of G . Assume that with respect to the left action of Γ , $X := \Gamma \backslash G$ is a compact manifold. When does T_X^* possess a nilpotent frame ?*

If G is nilpotent then of course T_X^* possesses a nilpotent frame. But it is also interesting to study the general case, e.g. $\mathrm{SL}(2, \mathbb{Z}) \backslash \mathrm{SL}(2, \mathbb{R})$ (non-compact!). In section 9, we shall give an example where T_X^* possesses a nilpotent frame but G is not nilpotent. For related results, see [6, 7].

Acknowledgments: Xu Wang would like to thank B. Berndtsson, J.P. Demailly and N. Mok for several useful discussions about the topics of this paper. Sönke Rollenske is grateful to the other authors for the invitation to join the project at a relatively late stage. He is also grateful to A. Fino and J. Ruppenthal for many discussions about the Dolbeault cohomology of nilmanifolds that culminated in the paper [13]. We are also pleased to thank the anonymous referee for valuable remarks and suggestions for a better presentation of our results.

2. MOTIVATIONS

2.1. First Motivation: Künneth formula. Our first motivation comes from the following well known Künneth formula:

Theorem 2.1 (Künneth formula). *Let (X, g_X) and (Y, g_Y) be two compact Riemannian manifolds. Let $(E, h_E), (F, h_F)$ be Hermitian complex vector bundles over X and Y respectively.*

- *If E and F are flat then we have the following formula for de Rham cohomologies:*

$$H_d^k(X \times Y, E \otimes F) = \bigoplus_{p+q=k} H_d^p(X, E) \otimes H_d^q(Y, F);$$

- *If X, Y are complex manifolds and E, F are holomorphic vector bundles then*

$$H_{\bar{\partial}}^{p,q}(X \times Y, E \otimes F) = \bigoplus_{j+k=p, l+m=q} H_{\bar{\partial}}^{j,l}(X, E) \otimes H_{\bar{\partial}}^{k,m}(Y, F).$$

One way to prove the above formulas is to use the Leray spectral sequence for fibrations, see [9]. Our motivation comes from the proof of using the following decomposition formulas of Laplacians:

$$(2.1) \quad \square_d^{X \times Y} = \square_d^X + \square_d^Y,$$

and

$$(2.2) \quad \square_{\bar{\partial}}^{X \times Y} = \square_{\bar{\partial}}^X + \square_{\bar{\partial}}^Y.$$

More precisely, we will study the following problem:

Problem: How to generalize (2.1) and (2.2) to non-product fibrations ?

Remark: One way to study the above problem is to develop the L^2 -theory of the Leray spectral sequence for fibrations (see [11] and [5] for related results). We know that for the spectral sequence of the double complex $(\partial, \bar{\partial})$, $d = \partial + \bar{\partial}$, the associated L^2 -theory is based on the classical Bochner-Kodaira-Nakano formula.

2.2. Second motivation: Nomizu-type theorems. Our second motivation is based on the following celebrated Nomizu's theorem [20] proved in 1954:

Nomizu's theorem (weak version): *Let G be a simply connected nilpotent Lie group with a discrete subgroup Γ . Assume that $X := \Gamma \backslash G$ is compact and the ascending central series of the Lie algebra of G (see section 6.2 for the definition) defines a torus fibration resolution of X . Then the de Rham cohomology of X can be represented by G -invariant forms.*

In 1976, Sakane [24] proved that the Nomizu theorem is also true for compact complex parallelisable solvmanifolds. The following theorem of Cordero-Fernández-Gray-Ugarte [9] is a generalization of Sakane's theorem:

Cordero-Fernández-Gray-Ugarte's theorem (weak version): *Let G be a simply connected nilpotent Lie group with a discrete subgroup Γ . Assume that $X := \Gamma \backslash G$ is a compact manifold with a left invariant integrable almost complex structure J . Assume that the J -compatible ascending series of the Lie algebra of G (see section 6.3 for the definition) defines a holomorphic torus fibration resolution of X . Then the Dolbeault cohomology of X can be represented by G -invariant forms.*

Remark: The assumption that the J -compatible ascending series of the Lie algebra of G defines a holomorphic torus fibration resolution is contained in the proof of the main theorem in [9].

In real case the ascending central series will always give a torus fibration resolution (see page 208 in [10]). Thus the following result is still true:

Nomizu's theorem (original version): *Let G be a simply connected nilpotent Lie group with a discrete subgroup Γ . Assume that $X := \Gamma \backslash G$ is compact. Then the de Rham cohomology of X can be represented by G -invariant forms.*

In complex case, the J -compatible ascending series may not give a torus fibration resolution (see Example 3.6 in [23] or [13]). But our main theorem implies the following result.

Cordero-Fernández-Gray-Ugarte's theorem (strong version): *Let G be a simply connected nilpotent Lie group with a discrete subgroup Γ . Assume that $X := \Gamma \backslash G$ is a compact manifold with a nilpotent complex structure (see [9], page 2, for the definition). Then the Dolbeault cohomology of X can be represented by G -invariant forms. In particular, it is independent of Γ .*

The above result applies in a number of important cases.

Corollary 2.2. *Let G be a simply connected nilpotent Lie group with a discrete cocompact subgroup Γ and left-invariant complex structure J . If G is 2-step nilpotent, then the Dolbeault cohomology of $X = (\Gamma \backslash G, J)$ can be computed by left-invariant forms.*

Proof. We only have to observe that if G is 2-step nilpotent then every left-invariant complex structure on G is nilpotent in the above sense by [22, Prop. 3.3] so the strong version of Cordero-Fernández-Gray-Ugarte's theorem applies. \square

A different way to generalise the weak version of Cordero-Fernández-Gray-Ugarte's theorem was considered in [13] and like in loc. cit. we are able to settle all cases of low dimension.

Corollary 2.3. *Let X be a nilmanifold of real dimension at most 6 with left-invariant complex structure. Then the Dolbeault cohomology of X is computed by left-invariant forms.*

Proof. In dimension 2 and 4 there are only tori and the Kodaira-Thurston manifold to consider, for which the result is well known.

In real dimension 6 there are only finitely many nilpotent Lie algebras and the ones admitting complex structures are classified by Salamon [25]. In [22, Proof of Thm.B] the statement was shown to hold for all complex structures on all such nilmanifolds except possibly for those with Lie algebra \mathfrak{h}_7 , in the notation of Salamon (see also [8] for the original definition). Since \mathfrak{h}_7 is 2-step nilpotent, indeed the free 2-step nilpotent Lie-algebra on 3 generators, the previous corollary applies to this remaining case. \square

3. FOLIATIONS OF NILPOTENT TYPE

3.1. Nilpotent foliation. Let us recall the definition of distribution first.

Definition 3.1 (Distribution). *Let X be a smooth manifold. We call*

$$\mathcal{V} := \{\mathcal{V}_x\}_{x \in X},$$

a rank- r distribution on X if for every $x \in X$, \mathcal{V}_x is an r -dimensional real linear subspace of $T_x X$ (space of vectors at x) and there exist smooth vector fields V_1, \dots, V_r on an open neighborhood, say U_x , of x such that

$$\mathcal{V}_y = \text{Span}_{\mathbb{R}}\{V_1(y), \dots, V_r(y)\},$$

for every $y \in U_x$. We call $\{V_1, \dots, V_r\}$ a local basis of \mathcal{V} .

Remark: If V is a smooth vector field on X such that $V(x) \in \mathcal{V}_x$ for every $x \in X$ then we say that V lies in \mathcal{V} and write $V \in \mathcal{V}$. Denote by $C^\infty(T_X)$ the space of smooth vector fields on X . Then one may look at a rank- r distribution as a subspace of $C^\infty(T_X)$ that is locally generated by r linearly independent smooth vector fields.

Definition 3.2 (Integrable Distribution). *A distribution \mathcal{V} is said to be integrable if $[V, W] \in \mathcal{V}$, for every $V, W \in \mathcal{V}$ (see the remark above). We call an integrable distribution a foliation on X .*

Remark: It is enough to check integrability for local basis of \mathcal{V} . The classical Frobenius theorem tells us that a rank- r distribution \mathcal{V} is integrable if and only if for every $x \in X$ there exists a smooth local coordinate system, say $\{x^1, \dots, x^n\}$, near x such that \mathcal{V} is generated by $\{\partial/\partial x^1, \dots, \partial/\partial x^r\}$ near x (i.e., \mathcal{V} is tangent to the fibers of the map $(x^1, \dots, x^n) \mapsto (x^{r+1}, \dots, x^n)$). Thus a rank- r integrable distribution is equivalent to a foliation of r -dimensional local smooth manifolds.

We shall use the following lemma to define the notion of nilpotent foliation.

Lemma 3.3. *Let \mathcal{V} be a distribution on X . Let g_X be a smooth Riemannian metric on X . Then*

$$\mathcal{V}^\perp := \{\mathcal{V}_x^\perp\}_{x \in X},$$

is also a distribution on X , where each \mathcal{V}_x^\perp denote the orthogonal complement of \mathcal{V}_x in $T_x X$ with respect to g_X .

Proof. Let $\{V_1, \dots, V_r\}$ be a local basis of \mathcal{V} . Then we can extend it to a local frame, say $\{V_1, \dots, V_n\}$, of T_X . Denote by V_j^\perp , $j > r$, the orthogonal projection of V_j to \mathcal{V}^\perp . Then we know that $\{V_j^\perp\}_{j>r}$ generates \mathcal{V}^\perp locally. \square

Definition 3.4 (Nilpotent Foliation). *Let \mathcal{V} be a distribution on a Riemannian manifold (X, g_X) . We call (\mathcal{V}, g_X) a nilpotent foliation structure on X if locally there exists an orthonormal frame $\{V_1, \dots, V_n\}$ of (T_X, g_X) such that*

- 1) $\{V_j\}_{j \leq r}$ is a local basis of \mathcal{V} and $\{V_j\}_{j > r}$ is a local basis of \mathcal{V}^\perp ;
- 2) $[V_j, V_k] = 0$ for every $1 \leq j \leq r, 1 \leq k \leq n$.

Remark: Notice that condition 2) in the above definition implies that a nilpotent foliation is always integrable.

We shall also study nilpotent foliations on complex manifold.

Definition 3.5 (Complex Nilpotent Foliation). *Let \mathcal{V} be a distribution on a complex manifold (X, J) . Let g_X be a J -Hermitian metric on X . We call (\mathcal{V}, J, g_X) a complex nilpotent foliation structure on X if locally there exists an orthonormal frame $\{V_1, \dots, V_n\}$ of $(T_X^{1,0}, g_X)$ such that*

- 1) $\{V_j, \bar{V}_j\}_{j \leq r}$ is a local basis of \mathcal{V} and $\{V_j, \bar{V}_j\}_{j > r}$ is a local basis of \mathcal{V}^\perp ;
- 2) $[V_j, V_k] = [V_j, \bar{V}_k] = 0$ for every $1 \leq j \leq r, 1 \leq k \leq n$.

Remark: Since g_X is J -Hermitian, a complex nilpotent foliation also satisfies $J(\mathcal{V}^\perp) = \mathcal{V}^\perp$.

3.2. Vertical-Horizontal decomposition of d .

Definition 3.6 (Vertical-Horizontal Vector Field). *Let \mathcal{V} be a distribution on a Riemannian manifold (X, g_X) . We call $V \in \mathcal{V}$ a vertical vector field and $W \in \mathcal{V}^\perp$ a horizontal vector field.*

We also need the dual of the notion of vertical-horizontal vector field (motivated by [4], see also formula (1.3) in [1]).

Definition 3.7 (Vertical-Horizontal One-Form). *A differential one-form u on X is said to be horizontal (resp. vertical) if $V \lrcorner u = 0$ for every vertical (resp. horizontal) vector field V on X .*

Definition 3.8 (Vertical-Horizontal Degree). *Denote by T_h^* and T_v^* the subbundles of T^*X generated by horizontal one-forms and vertical one-forms respectively. Then we have*

$$\wedge^p T^*X = \bigoplus_{k+l=p} (\wedge^k T_h^*) \wedge (\wedge^l T_v^*).$$

We call a section of $(\wedge^k T_h^*) \wedge (\wedge^l T_v^*)$ a degree $(k|l)$ -form and say that it has horizontal degree k and vertical degree l .

The following lemma suggests to study vertical-horizontal decomposition of the exterior derivative.

Lemma 3.9. *Let \mathcal{V} be a distribution on a Riemannian manifold (X, g_X) . Let u be a smooth degree $(k|l)$ -form on X . Assume that \mathcal{V} is integrable. Then we can write*

$$du = d^v u + d^h u + R_d u,$$

where $d^v u$ is degree $(k|l+1)$, $d^h u$ is degree $(k+1|l)$ and $R_d u$ is degree $(k+2|l-1)$.

Proof. Let us locally write

$$u = \sum u_h^j \wedge u_v^j,$$

where u_v^j are $(0|l)$ -forms and u_h^j are $(k|0)$ -forms. Since \mathcal{V} is integrable, we know that $d(u_h^j)$ has no degree $(k-1|2)$ components. Thus du has no degree $(k-1|l+2)$ components. \square

Definition 3.10 (Atiyah Tensor). *Let \mathcal{V} be an integrable distribution on a Riemannian manifold (X, g_X) . Then we define d^h as the $(1|0)$ -part of d and d^v as the $(0|1)$ -part of d . We call the following degree $(2|-1)$ tensor*

$$R_d := d - d^h - d^v,$$

the Atiyah tensor.

Remark 1: From the proof of the above Lemma, we know that the Atiyah tensor is zero if and only if \mathcal{V}^\perp is integrable. In case \mathcal{V} is associated to the Lie algebra \mathfrak{g} of a G -bundle, then cohomology class of each Lie-algebra component of the Atiyah tensor is also called the *Atiyah class*.

Remark 2: d^h, d^v are also well defined on the space of all smooth forms on X . The reason is we can always write a smooth form u as

$$u = \sum u^{(k|l)},$$

where each $u^{(k|l)}$ denotes the degree $(k|l)$ -component of u . Then we can define

$$d^h u = \sum d^h u^{(k|l)}, \quad d^v u = \sum d^v u^{(k|l)}.$$

3.3. Vertical-Horizontal decomposition of $\bar{\partial}$. Now let us consider the case that \mathcal{V} is a distribution on a complex manifold (X, J) with a J -Hermitian Riemannian metric g_X (we call (X, J, g_X) a *Hermitian complex manifold*) such that $J(\mathcal{V}) = \mathcal{V}$. Then we have

$$\wedge^{p,q} T^* X = \oplus_{k+j=p, l+s=q} (\wedge^{k,l} T_h^*) \wedge (\wedge^{j,s} T_v^*).$$

We call smooth section of $(\wedge^{k,l} T_h^*) \wedge (\wedge^{j,s} T_v^*)$ a *degree $(k, l|j, s)$ -form* and say that it has *horizontal degree (k, l)* and *vertical degree (j, s)* . Similar as the real case, we have

Lemma 3.11. *Let \mathcal{V} be an integrable distribution on a hermitian complex manifold (X, J, g_X) . Let u be a smooth degree $(k, l|j, s)$ -form on X . Assume that $J(\mathcal{V}) = \mathcal{V}$. Then we can write*

$$\bar{\partial} u = \bar{\partial}^v u + \bar{\partial}^h u + R_{A_1} u + R_{A_2} u + R_{KS} u,$$

where $\bar{\partial}^v u$ is degree $(k, l|j, s+1)$, $\bar{\partial}^h u$ is degree $(k, l+1|j, s)$ and $R_{A_1} u$ is degree $(k+1, l+1|j-1, s)$, $R_{A_2} u$ is degree $(k, l+2|j, s-1)$ and $R_{KS} u$ is degree $(k+1, l|j-1, s+1)$.

Definition 3.12 (Complex Atiyah Tensor and Kodaira-Spencer Tensor). *Let \mathcal{V} be a J -invariant integrable distribution on a Hermitian complex manifold (X, J, g_X) . We define $\bar{\partial}^h$ as the $(0, 1|0, 0)$ -part of $\bar{\partial}$ and $\bar{\partial}^v$ as the $(0, 0|0, 1)$ -part of $\bar{\partial}$. We call*

$$R_A := R_{A_1} + R_{A_2},$$

the complex Atiyah tensor and R_{KS} the Kodaira-Spencer tensor.

Remark: In case \mathcal{V} is given by the fiber-tangent distribution of a proper holomorphic submersion, then cohomology class of each component of R_{KS} is just the well known Kodaira–Spencer class. In general, put

$$R_{\bar{\partial}} = R_A + R_{KS},$$

If $R_{KS} \neq 0$ then $R_d \neq R_{\bar{\partial}} + R_{\partial}$. In fact, we have

$$R_d = R_A + \overline{R_A}, \quad d^v = \bar{\partial}^v + \partial^v,$$

and

$$d^h = R_{KS} + \overline{R_{KS}} + \partial^h + \bar{\partial}^h.$$

In case \mathcal{V} is a complex nilpotent foliation, we can prove that

Lemma 3.13. *Assume that (\mathcal{V}, J, g_X) is a complex nilpotent foliation structure. Then $R_{KS} \equiv 0$.*

Proof. It suffices to show that if u is an vertical $(1, 0)$ -form then $\bar{\partial}u$ has no degree $(1, 0|0, 1)$ -component. Since

$$(3.1) \quad du(W, V) = L_W(u(V)) - L_V(u(W)) - u([W, V]),$$

it is enough to show that for every vertical $(0, 1)$ -vector field V and horizontal $(1, 0)$ -vector field W , the vertical $(1, 0)$ -component of $[V, W]$ is zero, which follows from 2) in Definition 3.5. \square

4. VERTICAL–HORIZONTAL DECOMPOSITION OF LAPLACIANS

4.1. Fundamental theorem. The fundamental theorem in this paper is the following:

Theorem 4.1. *(Real case): Let (X, g_X) be an oriented Riemannian manifold with a nilpotent foliation structure (see Definition 3.4). Then on the space of smooth forms on X , we have*

$$(4.1) \quad \square_d = \square_{d^v} + \square_{d^h + R_d}.$$

(Complex case): Let (X, J, g_X) be a hermitian complex manifold with a complex nilpotent foliation structure (see Definition 3.5). Then on the space of smooth forms on X , we have

$$(4.2) \quad \square_{\bar{\partial}} = \square_{\bar{\partial}^v} + \square_{\bar{\partial}^h + R_{\bar{\partial}}}, \quad \square_{\bar{\partial}^v} = \square_{\partial^v}.$$

Remark 1: In our proof, we shall use the following notation: if P is a differential operator on the space of smooth forms on X then we shall write P^* as the adjoint of P and write

$$\square_P := PP^* + P^*P.$$

Recall that P^* satisfies

$$(Pu, v) = (u, P^*v),$$

if u is a smooth form and v is a smooth form with compact support. Thus P^* and \square_P are well defined on the space of smooth forms. If P maps a degree k form to a degree $(k + p)$ form then we say that P has degree p . If P is a degree p operator and Q is a degree q operator then we write

$$[P, Q] := PQ - (-1)^{pq}QP.$$

Since $d, d^v, d^h + R_d$ are degree one operators, we have

$$\square_d = [d, d^*], \quad \square_{d^v} = [d^v, (d^v)^*],$$

and

$$\square_{d^h+R_d} = [d^h + R_d, (d^h + R_d)^*].$$

Thus (4.1) is equivalent to

$$(4.3) \quad [d^h, (d^v)^*] = 0, \quad [d^v, R_d^*] = 0.$$

Remark 2 : Notice that if X is compact then

$$((\square_P + \square_Q)u, u) = \|Pu\|^2 + \|P^*u\|^2 + \|Qu\|^2 + \|Q^*u\|^2,$$

for every smooth form u on X . Thus $(\square_P + \square_Q)u = 0$ is equivalent to

$$Pu = P^*u = Qu = Q^*u = 0,$$

which gives the following corollary:

Corollary 4.2. (Real case): Let (X, g_X) be an oriented compact Riemannian manifold with a nilpotent foliation structure (see Definition 3.4). Then a smooth form u lies in $\ker \square_d$ if and only if

$$(4.4) \quad d^v u = (d^v)^* u = (d^h + R_d)u = (d^h + R_d)^* u = 0 \quad \text{on } X.$$

(Complex case): Let (X, J, g_X) be a hermitian compact complex manifold with a complex nilpotent foliation structure (see Definition 3.5). Then a smooth form u lies in $\ker \square_{\bar{d}}$ if and only if

$$(4.5) \quad \bar{\partial}^v u = (\bar{\partial}^v)^* u = (\bar{\partial}^h + R_{\bar{d}})u = (\bar{\partial}^h + R_{\bar{d}})^* u = \partial^v u = (\partial^v)^* u = 0 \quad \text{on } X.$$

4.2. Proof of the real case. Let (\mathcal{V}, g_X) be a nilpotent foliation structure on X (see Definition 3.4). Let $\{V_j\}$ be the local frame of (T_X, g_X) in Definition 3.4. Let us write

$$X_j^h = V_{r+j}, \quad X_k^v = V_k, \quad 1 \leq k \leq r, \quad 1 \leq j \leq n-r.$$

We know each X_k^v is vertical and each X_j^h is horizontal. Denote by

$$\{\varphi_h^j, \varphi_v^k\}_{1 \leq k \leq r, 1 \leq j \leq n-r},$$

the dual frame of $\{X_j^h, X_k^v\}$. By 2) in Definition 3.4 and (3.1), we have

$$d\varphi_h^j = \sum_{k,l=1}^{n-r} C_{kl}^j \varphi_h^k \wedge \varphi_h^l, \quad 1 \leq j \leq n-r,$$

and

$$d\varphi_v^k = \sum_{j,l=1}^{n-r} D_{jl}^k \varphi_h^j \wedge \varphi_h^l, \quad 1 \leq k \leq r,$$

where C_{kl}^j and D_{jl}^k are smooth functions. Thus we have

Lemma 4.3. The components R_d, d^v, d^h of d can be written as

$$R_d = \sum_{k=1}^r \sum_{j,l=1}^{n-r} \left(D_{jl}^k \varphi_h^j \wedge \varphi_h^l \right) \wedge (X_k^v \lrcorner),$$

$$d^v = \sum_{p=1}^r \varphi_v^p \wedge (X_p^v),$$

$$d^h = \sum_{j=1}^{n-r} \varphi_h^j \wedge (X_j^h) + \sum_{j,k,l=1}^{n-r} C_{kl}^j \left(\varphi_h^k \wedge \varphi_h^l \right) \wedge (X_j^h \rfloor).$$

Now let us finish the proof of the first identity in (4.3).

Lemma 4.4. $[d^v, R_d^*] = 0$.

Proof. Since R_d is a tensor, we have

$$(4.6) \quad R_d^* = \sum_{k=1}^r \sum_{j,l=1}^{n-r} D_{jl}^k \varphi_v^k \wedge (X_l^h \rfloor)(X_j^h \rfloor).$$

Thus R_d^* commutes with $\varphi_v^p \wedge$. Moreover, $d^2 \varphi_v^k = 0$ gives that $d^v D_{jl}^k \equiv 0$. Thus $[d^v, R_d^*] = 0$. \square

We need the following proposition to prove $[d^h, (d^v)^*] = 0$.

Proposition 4.5. Denote by $*$ the Hodge star operator on our oriented manifold (X, g_X) . Assume that the orientation of X is given by $\Omega_h \wedge \Omega_v$, where

$$\Omega_h := \varphi_h^1 \wedge \cdots \wedge \varphi_h^{n-r}, \quad \Omega_v := \varphi_v^1 \wedge \cdots \wedge \varphi_v^r.$$

Denote by $*_h$ (resp. $*_v$) the Hodge star operator with respect to Ω_h (resp. Ω_v) on the space of horizontal (resp. vertical) forms respectively. Then

$$(4.7) \quad *(u_h \wedge u_v) = (-1)^{(n-r-p)q} *_h u_h \wedge *_v u_v,$$

where u_h is a degree p horizontal form, u_v is a degree q vertical form.

Proof. Notice that

$$(u_h \wedge u_v) \wedge (-1)^{(n-r-p)q} (*_h u_h \wedge *_v u_v) = (u_h \wedge *_h u_h) \wedge (u_v \wedge *_v u_v) = (u_h \wedge u_v) \wedge *(u_h \wedge u_v).$$

Thus (4.7) follows. \square

Lemma 4.6. Let $u = f u_h \wedge u_v$ be a smooth degree $(a|b)$ form, where u_h (resp. u_v) is a finite wedge product of φ_h^k (resp. φ_v^l) and f is a smooth function. Then

$$(4.8) \quad (d^v)^* u = (-1)^a u_h \wedge \left((-1)^{r(b+1)+1} *_v d^v *_v (f u_v) \right),$$

and

$$(4.9) \quad (d^h)^* u = \left((-1)^{(n-r)(a+1)+1} *_h d^h *_h (f u_h) \right) \wedge u_v.$$

Proof. The main idea is to use the fact that $(d^v)^* u$ (resp. $(d^h)^* u$) is the degree $(a|b-1)$ (resp. $(a-1|b)$) part of $d^* u$ and $d^* u = (-1)^{n(a+b+1)+1} * d * u$. Thus the above proposition applies. We shall only prove the first formula. By (4.7), we have

$$*u = (-1)^{(n-r-a)b} f *_h u_h \wedge *_v u_v.$$

Thus

$$d * u = (-1)^{(n-r-a)b} (df \wedge (*_h u_h \wedge *_v u_v) + f d(*_h u_h \wedge *_v u_v)).$$

Using (4.7) again, we know that the degree $(a|b-1)$ -part of $(-1)^{n(a+b+1)+1} * d * u$ is equal to the right hand side of (4.8). \square

Remark: Since

$$(-1)^{r(b+1)} *_v (\varphi_v^j \wedge) *_v u_v = X_j^v \rfloor u_v,$$

(4.8) gives the following formula:

Lemma 4.7. $(d^v)^* = -\sum_{j=1}^r (X_j^v \lrcorner) \wedge (X_j^v)$.

Now we can prove the second identity in (4.3).

Lemma 4.8. $[d^h, (d^v)^*] = 0$.

Proof. Notice that $d^2\varphi_h^j = 0$ gives $d^v C_{kl}^j \equiv 0$. Thus Lemma 4.3 and the above lemma give

$$(4.10) \quad [d^h, (d^v)^*] = \sum ([X_j^h, X_k^v])(X_k^v \lrcorner) \wedge (\varphi_h^j \wedge).$$

But by 2) in Definition 3.4, we have

$$(4.11) \quad [X_j^h, X_k^v] = 0.$$

Thus the lemma follows. \square

The proof of the real case is complete.

4.3. Proof of the complex case. Let (\mathcal{V}, J, g_X) be a complex nilpotent foliation structure on X . Let $\{V_j\}_{1 \leq j \leq n}$ be the local frame of $(T_X^{1,0}, g_X)$ in Definition 3.5). Put

$$X_j^h := V_{r+j}, \quad X_k^v = V_k, \quad 1 \leq k \leq r, \quad 1 \leq j \leq n-s.$$

We know each X_k^v is vertical and each X_j^h is horizontal. Denote by

$$\{\varphi_h^j, \varphi_v^k\}_{1 \leq k \leq r, 1 \leq j \leq n-r},$$

the dual frame of $\{X_j^h, X_k^v\}$. By 2) in Definition 3.5 and (3.1), we have

$$d\varphi_h^j = \sum_{k,l=1}^{n-r} C_{kl}^j \varphi_h^k \wedge \varphi_h^l + \sum_{k,l=1}^{n-r} C_{k\bar{l}}^j \varphi_h^k \wedge \overline{\varphi_h^l}, \quad 1 \leq j \leq n-r,$$

and

$$d\varphi_v^k = \sum_{j,l=1}^{n-r} D_{jl}^k \varphi_h^j \wedge \varphi_h^l + \sum_{j,l=1}^{n-r} D_{j\bar{l}}^k \varphi_h^j \wedge \overline{\varphi_h^l}, \quad 1 \leq k \leq r,$$

where $C_{kl}^j, C_{k\bar{l}}^j, D_{jl}^k$ and $D_{j\bar{l}}^k$ are smooth functions, which gives

Lemma 4.9. *The components $\bar{\partial}^h, \bar{\partial}^v, R_{\bar{\partial}}$ can be written as*

$$\bar{\partial}^h = \sum_{j=1}^{n-r} \overline{\varphi_h^j} \wedge (\overline{X_j^h}) + \sum_{j,k,l=1}^{n-r} \overline{C_{kl}^j (\varphi_h^k \wedge \varphi_h^l) \wedge (X_j^h \lrcorner)} + \sum_{j,k,l=1}^{n-r} C_{k\bar{l}}^j (\varphi_h^k \wedge \overline{\varphi_h^l}) \wedge (X_j^h \lrcorner),$$

$$\bar{\partial}^v = \sum_{k=1}^r \overline{\varphi_v^k} \wedge (\overline{X_k^v}),$$

and $R_{\bar{\partial}} = R_{KS} + R_{A_1} + R_{A_2}$ satisfies

$$R_{KS} = 0, \quad R_{A_1} = \sum_{k=1}^r \sum_{j,l=1}^{n-r} \left(D_{j\bar{l}}^k \varphi_h^j \wedge \overline{\varphi_h^l} \right) \wedge (X_k^v \lrcorner), \quad R_{A_2} = \sum_{k=1}^r \sum_{j,l=1}^{n-r} \overline{\left(D_{jl}^k \varphi_h^j \wedge \varphi_h^l \right) \wedge (X_k^v \lrcorner)}.$$

By a similar proof as the real case, we have

$$(4.12) \quad [\bar{\partial}^v, R_{\bar{\partial}}^*] = 0,$$

and the following analogy of Lemma 4.7.

Lemma 4.10. $(\bar{\partial}^v)^* = -\sum_{j=1}^r (\overline{X_j^v} \lrcorner) \wedge (X_j^v)$.

Similar as the real case, the above lemma gives

$$(4.13) \quad [\bar{\partial}^h, (\bar{\partial}^v)^*] = 0$$

We know that (4.12) and (4.13) together give

$$\square_{\bar{\partial}} = \square_{\bar{\partial}^v} + \square_{\bar{\partial}^h + R_{\bar{\partial}}}.$$

Now it suffices to prove

$$\square_{\bar{\partial}^v} = \square_{\partial^v}.$$

By Lemma 4.10 and Lemma 4.9, we have

$$(4.14) \quad \square_{\bar{\partial}^v} = - \sum_{j=1}^r (X_j^v) (\overline{X_j^v}) = - \sum_{j=1}^r (\overline{X_j^v}) (X_j^v),$$

which gives

$$\square_{\bar{\partial}^v} = \square_{\bar{\partial}^v} = \square_{\partial^v}.$$

The proof of Theorem 4.1 is complete.

Remark: One may also prove the complex case of Theorem 4.1 by using vertical–horizontal decomposition of the following Demailly–Griffiths–Kähler identity (see page 306 in [12], [15] or [27] for a pure algebraic proof)

$$(4.15) \quad \bar{\partial}^* = i[\partial, \Lambda] + [L, \theta^*], \quad Lu := \omega \wedge u, \quad \theta := [\bar{\partial}, L], \quad \Lambda := L^*,$$

where ω denotes the real Hermitian (1, 1)-form associated to (g_X, J) .

5. AN EXAMPLE: THE KODAIRA–THURSTON MANIFOLD

The Kodaira–Thurston surface was first found by Kodaira in [17]. It is the first example [28] of complex symplectic manifold without Kähler structure. Let us recall its definition in [27]. Consider the following group structure

$$a * b := (a^1 + b^1, a^2 + b^2, a^3 + a^1 b^2 + b^3, a^4 + b^4),$$

on \mathbb{R}^4 . The Kodaira–Thurston surface X is defined as the quotient manifold with respect to the *left* action of \mathbb{Z}^4 on (notice that \mathbb{Z}^4 is a discrete subgroup of G)

$$G := (\mathbb{R}^4, *).$$

It is easy to see that X is a compact manifold with respect to the quotient topology. Let (x^1, x^2, x^3, x^4) be the canonical coordinate system on \mathbb{R}^4 . We know

$$(5.1) \quad \begin{cases} \varphi_h = dx^1 + idx^2, \\ \varphi_v = dx^3 - x^1 dx^2 + idx^4, \\ \overline{\varphi}_h = dx^1 - idx^2, \\ \overline{\varphi}_v = dx^3 - x^1 dx^2 - idx^4, \end{cases}$$

is a frame of the space of G -invariant (with respect to the left action of G) 1-forms on X . Let J be the almost complex structure on X such that the associated $\wedge^{1,0} T^* X$ is spanned by $\{\varphi_h, \varphi_v\}$. Notice that

$$(5.2) \quad \begin{cases} d\varphi_h = 0, \\ d\varphi_v = -\frac{i}{2}\varphi_h \wedge \overline{\varphi}_h. \end{cases}$$

implies that J is integrable on X . One may check that

$$\begin{cases} \bar{\partial}(x^1 + ix^2) = 0, \\ \bar{\partial}(x^3 + ix^4 - \frac{i}{2}(x^1)^2) = 0. \end{cases}$$

Thus

$$\begin{cases} z := x^1 + ix^2, \\ w := x^3 + ix^4 - \frac{i}{2}(x^1)^2. \end{cases}$$

are local holomorphic coordinates on X . Now we know that the following holomorphic map from (\mathbb{R}^4, J) to \mathbb{C}

$$(z, w) \mapsto z,$$

defines a holomorphic submersion, say π , from X to the torus $\mathbb{T} := \mathbb{C}/\mathbb{Z}^2$. Let $\{X^h, X^v\}$ be the global frame of $T^{1,0}(X)$ that is dual to $\{\varphi_h, \varphi_v\}$. Then we know that

$$\begin{cases} \partial/\partial z := X^h + ix^1 X^v, \\ \partial/\partial w := X^v. \end{cases}$$

is a holomorphic π -local (i.e., well defined on the π -inverse of a sufficiently small open set in \mathbb{T}) frame for $T^{1,0}X$. Thus we have

Proposition 5.1. $\pi : X \rightarrow \mathbb{T}$ is locally trivial.

Remark: Notice that (5.2) implies that the fibers of π defines a complex nilpotent foliation structure on (X, J, g_X) , where g_X is J -hermitian such that the fundamental form of (g_X, J) is

$$\omega = i\varphi_h \wedge \bar{\varphi}_h + i\varphi_v \wedge \bar{\varphi}_v.$$

We know that $\varphi_h, \bar{\varphi}_h$ are horizontal forms and $\varphi_v, \bar{\varphi}_v$ are vertical forms. By (5.2), we know that

$$R_{\bar{\partial}} = -\frac{i}{2}\varphi_h \wedge \bar{\varphi}_h \wedge (X^v \lrcorner \cdot),$$

is of degree $(1, 1 | -1, 0)$. We shall use Theorem 4.1 and Corollary 4.2 to give another proof of the following well known result (see section 5 in [9]).

Theorem 5.2. Denote by $\mathcal{H}^{p,q}(\bar{\partial})$ the space of $\bar{\partial}$ -harmonic (p, q) -forms on the Kodaira-Thurston surface (X, J, g_X) , we have

$$\begin{cases} \mathcal{H}^{0,0}(\bar{\partial}) = \text{Span}_{\mathbb{C}} \langle 1 \rangle, \\ \mathcal{H}^{1,0}(\bar{\partial}) = \text{Span}_{\mathbb{C}} \langle \varphi_h \rangle, \\ \mathcal{H}^{0,1}(\bar{\partial}) = \text{Span}_{\mathbb{C}} \langle \bar{\varphi}_h, \bar{\varphi}_v \rangle, \\ \mathcal{H}^{2,0}(\bar{\partial}) = \text{Span}_{\mathbb{C}} \langle \varphi_h \wedge \varphi_v \rangle, \\ \mathcal{H}^{1,1}(\bar{\partial}) = \text{Span}_{\mathbb{C}} \langle \bar{\varphi}_h \wedge \varphi_v, \varphi_h \wedge \bar{\varphi}_v \rangle, \\ \mathcal{H}^{0,2}(\bar{\partial}) = \text{Span}_{\mathbb{C}} \langle \bar{\varphi}_h \wedge \bar{\varphi}_v \rangle, \\ \mathcal{H}^{2,1}(\bar{\partial}) = \text{Span}_{\mathbb{C}} \langle \varphi_h \wedge \varphi_v \wedge \bar{\varphi}_h, \varphi_h \wedge \varphi_v \wedge \bar{\varphi}_v \rangle, \\ \mathcal{H}^{1,2}(\bar{\partial}) = \text{Span}_{\mathbb{C}} \langle \varphi_v \wedge \bar{\varphi}_h \wedge \bar{\varphi}_v \rangle, \\ \mathcal{H}^{2,2}(\bar{\partial}) = \text{Span}_{\mathbb{C}} \langle \varphi_h \wedge \varphi_v \wedge \bar{\varphi}_h \wedge \bar{\varphi}_v \rangle. \end{cases}$$

Proof. $\mathcal{H}^{0,0}(\bar{\partial}) = \text{Span}_{\mathbb{C}} \langle 1 \rangle$ is trivial. By Corollary 4.2, we know that all harmonic forms in $\mathcal{H}^{p,q}(\bar{\partial})$ lie in the kernel of $\bar{\partial}^v$, $(\bar{\partial}^v)^*$, $\bar{\partial}^h + R_{\bar{\partial}}$ and $(\bar{\partial}^h + R_{\bar{\partial}})^*$.

Degree (1,0) case: Let

$$u = a\varphi_h + b\varphi_v,$$

be in $\mathcal{H}^{1,0}(\bar{\partial})$. Notice that $\bar{\partial}^v u = 0$ is equivalent to

$$\bar{\partial}^v a = \bar{\partial}^v b = 0,$$

and $(\bar{\partial}^h + R_{\bar{\partial}})u = 0$ is equivalent to

$$\bar{\partial}^h b = 0, \quad \bar{\partial}^h a + \frac{i}{2}b\bar{\varphi}_h = 0.$$

Thus $\bar{\partial}b = 0$ and b is a constant. Notice that $\bar{\partial}^v a = 0$ and $\bar{\partial}^h a + \frac{i}{2}b\bar{\varphi}_h = 0$ together imply $\frac{i}{2}b\bar{\varphi}_h = -\bar{\partial}a$ is $\bar{\partial}$ -exact. Since $\bar{\varphi}_h$ is not $\bar{\partial}$ -exact, we know that $b = 0$. Thus

$$\bar{\partial}^h a = \bar{\partial}^v a = 0,$$

which gives $\bar{\partial}a = 0$ and a is a constant. Thus $\mathcal{H}^{1,0}(\bar{\partial}) = \text{Span}_{\mathbb{C}} \langle \varphi_h \rangle$.

Degree (0,1) case: Let

$$u = a\bar{\varphi}_h + b\bar{\varphi}_v,$$

be in $\mathcal{H}^{0,1}(\bar{\partial})$. $\bar{\partial}^v u = (\bar{\partial}^h + R_{\bar{\partial}})u = 0$ is equivalent to

$$\bar{\partial}^v a = \bar{\partial}^h b = 0.$$

$(\bar{\partial}^v)^*u = 0$ is equivalent to

$$\partial^v b = 0.$$

Since $\square_{\bar{\partial}^v} = \square_{\partial^v}$, we know that $\partial^v b = 0$ implies $\bar{\partial}^v b = 0$. Thus $\bar{\partial}b = 0$ and b is a constant. $(\bar{\partial}^h + R_{\bar{\partial}})^*u = 0$ is equivalent to $\partial^h a = 0$, thus $\partial a = 0$ and a is a constant.

Degree (2,0) case: Let

$$u = a\varphi_h \wedge \varphi_v,$$

be in $\mathcal{H}^{2,0}(\bar{\partial})$. $\bar{\partial}u = 0$ is equivalent to $\bar{\partial}a = 0$, which is equivalent to that a is a constant.

Degree (1,1) case: Let

$$u = a\varphi_h \wedge \bar{\varphi}_h + b\varphi_h \wedge \bar{\varphi}_v + c\varphi_v \wedge \bar{\varphi}_h + f\varphi_v \wedge \bar{\varphi}_v,$$

be a harmonic $(1,1)$ -form. We have

$$\square_{\bar{\partial}^v} u = (\square_{\bar{\partial}^v} a) \varphi_h \wedge \bar{\varphi}_h + (\square_{\bar{\partial}^v} b) \varphi_h \wedge \bar{\varphi}_v + (\square_{\bar{\partial}^v} c) \varphi_v \wedge \bar{\varphi}_h + (\square_{\bar{\partial}^v} f) \varphi_v \wedge \bar{\varphi}_v,$$

Then $\square_{\bar{\partial}^v} u = 0$ is equivalent to

$$\square_{\bar{\partial}^v} a = \square_{\bar{\partial}^v} b = \square_{\bar{\partial}^v} c = \square_{\bar{\partial}^v} f = 0.$$

Together with $\square_{\partial^v} = \square_{\bar{\partial}^v}$, the above identities give

$$d^v a = d^v b = d^v c = d^v f = 0.$$

$(\bar{\partial}^h + R_{\bar{\partial}})u = 0$ is equivalent to

$$\bar{\partial}^h f = 0, \quad \bar{\partial}^h b + \frac{i}{2}f\bar{\varphi}_h = 0,$$

thus $\bar{\partial}f = 0$ and f is a constant. Since $\bar{\partial}^v b = 0$, we have

$$\bar{\partial}b + \frac{i}{2}f\bar{\varphi}_h = 0.$$

By the above computation of $\mathcal{H}^{0,1}(\bar{\partial})$, we know that $\bar{\varphi}_h$ is not $\bar{\partial}$ -exact. Thus $f = 0$ and $\bar{\partial}b = 0$. Now we know that b is a constant. $(\bar{\partial}^h + R_{\bar{\partial}})^*u = 0$ is equivalent to

$$\partial^h a = 0, \quad \frac{i}{2}a\varphi_h + \partial^h c = 0.$$

thus $\partial a = 0$ and a is a constant. Again

$$-\frac{i}{2}\bar{a}\bar{\varphi}_h + \bar{\partial}c = 0,$$

gives $a = 0$ and c is a constant.

For the remaining cases, by the following well known formula

$$\dim \mathcal{H}^{p,q}(\bar{\partial}) = \dim \mathcal{H}^{n-p,n-q}(\bar{\partial}),$$

it is enough to check that the listed forms lie in the $\bar{\partial}$ -harmonic spaces, which follows by a direct computation. \square

5.1. Nilpotent fibrations. For the Kodaira–Thurston manifold, the complex nilpotent foliation structure comes from a holomorphic fibration. The general definition is as follows:

Definition 5.3 (Nilpotent Fibration). *We call a proper smooth submersion $\pi : (X, g_X) \rightarrow (B, g_B)$ between two Riemannian manifolds a nilpotent fibration if the associated foliation \mathcal{V} of the fibers defines a nilpotent foliation structure on (X, g_X) and*

$$(5.3) \quad g_X(V, W) = g_B(\pi_*V, \pi_*W),$$

for every horizontal vector fields V, W on X .

Remark: Let V_B be a vector field on B , we call a vector field V_X on X a lift of V_B if

$$\pi_*V_X = V_B.$$

It is clear that V_B has a unique lift V_X such that V_X is horizontal. (5.3) says that the norm of a vector field on B is equal to the norm of its horizontal lift.

Definition 5.4 (Complex Nilpotent Fibration). *We call a proper holomorphic submersion $\pi : (X, \omega_X) \rightarrow (B, \omega_B)$ between two Hermitian complex manifolds a complex nilpotent fibration if the associated foliation \mathcal{V} of the fibers defines a complex nilpotent foliation structure on (X, ω_X) and*

$$(5.4) \quad \omega_X(V, W) = \omega_B(\pi_*V, \pi_*W),$$

for every horizontal $(1, 0)$ -vector fields V, W on X .

Theorem 4.1 gives

Theorem 5.5. *On the total space of a nilpotent fibration, we have*

$$\square_d = \square_{d^v} + \square_{d^h + R_d}.$$

On the total space of a complex nilpotent fibration, we have

$$\square_{\bar{\partial}} = \square_{\bar{\partial}^v} + \square_{\bar{\partial}^h + R_{\bar{\partial}}}, \quad \square_{\bar{\partial}^v} = \square_{\partial^v}.$$

Remark: Associated to a fibration there is a natural Leray–Serre spectral sequence, which plays a crucial role in the proof of Nomizu-type theorems. But in general a foliation does not give a good fibration structure, thus one has to use other methods. Our main idea is: with the help of Theorem 4.1, one may use the spectral sequence for double complex to continue the reduction process (as in the fibration case), in which the natural setup is a manifold with a nilpotent frame.

6. NILPOTENT FRAME

6.1. **Real case.** Let X be a compact smooth manifold with trivial T_X^* . Let

$$\Phi := \{\sigma^1, \dots, \sigma^n\}$$

be a global smooth frame of T_X^* . Then we have the following Maurer–Cartan equations

$$(6.1) \quad d\sigma^j = \sum_{k,l=1}^n A_{kl}^j \sigma^k \wedge \sigma^l,$$

where A_{kl}^j are globally defined smooth functions on X . Recall that Φ is a *nilpotent frame* if A_{kl}^j are real *constants* and the above equations reduce to

$$(6.2) \quad d\sigma^j = \sum_{k,l>j} A_{kl}^j \sigma^k \wedge \sigma^l.$$

Definition 6.1. Let Φ be a nilpotent frame. Put $r_0 = 0$ and define r_j ($j \geq 1$) inductively by

$$r_j + 1 = \min U_j, \quad U_j := \cup_{j \geq r_{j-1} + 1} \{k, l : A_{kl}^j \neq 0\},$$

where $r_j := n$ if U_j is empty. Fix k such that

$$0 = r_0 < r_1 < \dots < r_{k-1} < r_k = n,$$

we call Φ a k -nilpotent frame. Let S^j be the subbundle of T_X^* generated by $\{\sigma^1, \dots, \sigma^{r_j}\}$. We call

$$0 = S^0 \hookrightarrow S^1 \hookrightarrow \dots \hookrightarrow S^k = T_X^*,$$

the Φ -filtration of T_X^* .

Remark: We always have $1 \leq k \leq n$. For the Kodaira–Thurston manifold (see (5.1)), put

$$\sigma^1 = dx^4, \quad \sigma^2 = dx^3 - x^1 dx^2, \quad \sigma^3 = dx^2, \quad \sigma^4 = dx^1,$$

we have

$$\begin{cases} d\sigma^1 = 0, \\ d\sigma^2 = \sigma^3 \wedge \sigma^4, \\ d\sigma^3 = 0, \\ d\sigma^4 = 0. \end{cases}$$

Thus $k = 2$ and the Φ -filtration is

$$0 \hookrightarrow \text{Span}\{\sigma^1, \sigma^2\} \hookrightarrow \text{Span}\{\sigma^1, \sigma^2, \sigma^3, \sigma^4\} = T_X^*.$$

Put

$$T_v^* = \text{Span}\{\sigma^1, \sigma^2\}, \quad T_h^* = \text{Span}\{\sigma^3, \sigma^4\}.$$

We get the following vertical-horizontal decomposition of the T_X^*

$$T_X^* = T_v^* \oplus T_h^*.$$

with respect the Riemannian metric $\sum \sigma^j \otimes \sigma^j$. In general, we shall introduce the following definition

Definition 6.2. Denote by $T_{v_j}^*$ ($1 \leq j \leq k$) the subbundle of T_X^* generated by $\{\sigma^{r_{j-1}+1}, \dots, \sigma^{r_j}\}$. Put

$$T_{h_j}^* = \bigoplus_{l=j+1}^k T_{v_l}^*, \quad 0 \leq j \leq k-1, \quad T_{h_k}^* = 0.$$

We call

$$T_{h_{j-1}}^* := T_{v_j}^* \oplus T_{h_j}^*, \quad 1 \leq j \leq k,$$

the j -th vertical horizontal decomposition with respect to the following Riemannian metric

$$(6.3) \quad g_X := \sum_{j=1}^n \sigma^j \otimes \sigma^j,$$

on X associated to the nilpotent frame Φ .

Remark: The first vertical horizontal decomposition of T^*X gives

$$d := d^{h_0} = d^{v_1} + d^{h_1} + R_d^1,$$

where d^{v_1} (resp. d^{h_1}) increases the first vertical (resp. horizontal) degree by one and R_d^1 is the remaining term. In general, the j -th vertical horizontal decomposition gives

$$d^{h_{j-1}} = d^{v_j} + d^{h_j} + R_d^j, \quad 1 \leq j \leq k.$$

Notice that the k -th vertical horizontal decomposition reduces to

$$d^{h_{k-1}} = d^{v_k}, \quad R_d^k = 0, \quad d^{h_k} = 0.$$

We shall use our fundamental theorem (see Theorem 4.1) to prove the following result.

Theorem 6.3. With respect to the notation above, we have

$$(6.4) \quad \square_d = \square_{d^{v_1}} + \square_{d^{h_1} + R_d^1},$$

on the space of smooth forms on X . Moreover, if $k \geq 2$ then for every $2 \leq j \leq k$, we have

$$(6.5) \quad \square_{d^{h_{j-1}}} = \square_{d^{v_j}} + \square_{d^{h_j} + R_d^j},$$

on the space of smooth forms in $\ker \square_{d^{v_1}} \cap \dots \cap \ker \square_{d^{v_{j-1}}}$.

Proof. Denote by $\{T^{h_j}, T^{v_k}\}$ the subbundles of T_X that are dual to $\{T_{h_j}^*, T_{v_k}^*\}$. From Definition 6.1, we know that the distribution \mathcal{V}^1 associated to T^{v_1} is integrable and \mathcal{V}^1 defines a nilpotent foliation with respect to the Riemannian metric g_X in (6.3). Thus Theorem 4.1 gives (6.4).

Now let us prove (6.5) for $j = 2$. Let us write a smooth form u on X as

$$(6.6) \quad u = \sum f_{p,q} u_{h_1}^p \wedge u_{v_1}^q,$$

where $\{u_{h_1}^p\}$ (resp. $\{u_{v_1}^q\}$) denotes a basis of the exterior algebra generated by $\{\sigma^{r_1+1}, \dots, \sigma^n\}$ (resp. $\{\sigma^1, \dots, \sigma^{r_1}\}$) respectively. Denote by $\{V_l\}$ the frame of T_X that is dual to $\{\sigma^l\}$. Put

$$\varphi_{v_1}^l = \sigma^l, \quad \varphi_{h_1}^m = \sigma^{r_1+m}, \quad X_l^{v_1} = V_l, \quad X_m^{h_1} = X_{r_1+m}.$$

By Lemma 4.3 and Lemma 4.10, we have

$$\square_{d^{v_1}} = - \sum (X_l^{v_1})(X_l^{v_1}),$$

which gives

$$\square_{d^{v_1}} u = \sum (\square_{d^{v_1}} f_{p,q}) u_{h_1}^p \wedge u_{v_1}^q.$$

Thus $\square_{d^{v_1}} u = 0$ is equivalent to

$$\square_{d^{v_1}} f_{p,q} \equiv 0,$$

for all p, q . Since $f_{p,q}$ are globally defined smooth functions on X , we know that $\square_{d^{v_1}} f_{p,q} \equiv 0$ is equivalent to that $d^{v_1} f_{p,q} \equiv 0$. Thus we get the following lemma.

Lemma 6.4. $\square_{d^{v_1}} u = 0$ is equivalent to that all $f_{p,q}$ are constants on leaves of \mathcal{V}^1 .

By the above lemma and Definition 6.1, we know that the distribution \mathcal{V}^2 associated to T^{v_2} is integrable on the space of smooth forms in $\ker \square_{d^{v_1}}$. Moreover, on the space of smooth forms in $\ker \square_{d^{v_1}}$, \mathcal{V}^2 defines a nilpotent foliation with respect to the following decomposition

$$T_{h_1}^* = T_{v_2}^* \oplus T_{h_2}^*.$$

Thus the proof of Theorem 4.1 gives (6.5) for $j = 2$. The general case follows by induction on j . \square

6.2. Complex case. Let X be a compact complex manifold with smoothly trivial holomorphic cotangent bundle $\wedge^{1,0} T_X^*$. Let

$$\Psi := \{\xi^1, \dots, \xi^n\}$$

be a global smooth frame of $\wedge^{1,0} T_X^*$. Recall that Ψ is a *complex nilpotent frame* if

$$(6.7) \quad d\xi^j = \sum_{k,l>j} B_{kl}^j \xi^k \wedge \xi^l + \sum_{k,l>j} B_{k\bar{l}}^j \xi^k \wedge \bar{\xi}^l,$$

where B_{kl}^j and $B_{k\bar{l}}^j$ are *complex constants*.

Definition 6.5. Let Ψ be a complex nilpotent frame. Put $r_0 = 0$ and define r_j ($j \geq 1$) inductively by

$$r_j + 1 = \min U_j, \quad U_j := \cup_{l \geq r_{j-1} + 1} \{k, l : B_{kl}^j \text{ or } B_{k\bar{l}}^j \neq 0\},$$

where $r_j := n$ if U_j is empty. Fix k such that

$$0 = r_0 < r_1 < \dots < r_{k-1} < r_k = n,$$

we call Ψ a *complex k -nilpotent frame*. Let S^j be the subbundle of $\wedge^{1,0} T_X^*$ generated by $\{\xi^1, \dots, \xi^{r_j}\}$. We call

$$0 = S^0 \hookrightarrow S^1 \hookrightarrow \dots \hookrightarrow S^k = \wedge^{1,0} T_X^*,$$

the Ψ -filtration of $\wedge^{1,0} T_X^*$.

Similar as the real case, we have

Definition 6.6. Denote by $T_{v_j}^*$ ($1 \leq j \leq k$) the (smooth, may not be holomorphic) subbundle of $\wedge^{1,0} T_X^*$ generated by $\{\xi^{r_{j-1}+1}, \dots, \xi^{r_j}\}$. Put

$$T_{h_j}^* = \oplus_{l=j+1}^k T_{v_l}^*, \quad 0 \leq j \leq k-1, \quad T_{h_k}^* = 0.$$

We call

$$T_{h_{j-1}}^* := T_{v_j}^* \oplus T_{h_j}^*, \quad 1 \leq j \leq k,$$

the j -th vertical horizontal decomposition with respect to the following Hermitian form

$$(6.8) \quad \omega_X := i \sum_{j=1}^n \xi^j \wedge \bar{\xi}^j,$$

on X associated to the complex nilpotent frame Ψ .

Remark: The first vertical horizontal decomposition of T^*X gives

$$\bar{\partial} := \bar{\partial}^{h_0} = \bar{\partial}^{v_1} + \bar{\partial}^{h_1} + R_{\bar{\partial}}^1,$$

where $\bar{\partial}^{v_1}$ (resp. $\bar{\partial}^{h_1}$) increases the first vertical (resp. horizontal) degree by $(0, 1)$ and $R_{\bar{\partial}}^1$ is the remaining term. In general, the j -th vertical horizontal decomposition gives

$$\bar{\partial}^{h_{j-1}} = \bar{\partial}^{v_j} + \bar{\partial}^{h_j} + R_{\bar{\partial}}^j, \quad 1 \leq j \leq k.$$

Notice that the k -th vertical horizontal decomposition reduces to

$$\bar{\partial}^{h_{k-1}} = \bar{\partial}^{v_k}, \quad R_{\bar{\partial}}^k = 0, \quad \bar{\partial}^{h_k} = 0.$$

Similar as the real case, Theorem 4.1 implies the following result.

Theorem 6.7. *With respect to the notation above, we have*

$$(6.9) \quad \square_{\bar{\partial}} = \square_{\bar{\partial}^{v_1}} + \square_{\bar{\partial}^{h_1} + R_{\bar{\partial}}^1}, \quad \square_{\bar{\partial}^{v_1}} = \square_{\partial^{v_1}},$$

on the space of smooth forms on X . Moreover, if $k \geq 2$ then for every $2 \leq j \leq k$, we have

$$(6.10) \quad \square_{\bar{\partial}^{h_{j-1}}} = \square_{\bar{\partial}^{v_j}} + \square_{\bar{\partial}^{h_j} + R_{\bar{\partial}}^j}, \quad \square_{\bar{\partial}^{v_j}} = \square_{\partial^{v_j}},$$

on the space of smooth forms in $\ker \square_{\bar{\partial}^{v_1}} \cap \cdots \cap \ker \square_{\bar{\partial}^{v_{j-1}}}$.

7. PROOF OF THE MAIN THEOREM

7.1. Cohomology description of Theorem 4.1. Let X be a compact smooth manifold. By the de Rham theorem, we have the following isomorphism

$$H_d^* \simeq \mathcal{H}_d^* := \{u \in \mathcal{A}^* : \square_d u = 0\},$$

where H_d^* denotes the de Rham cohomology group of X , \mathcal{A}^* denotes the space of smooth forms on X . With the assumption in Theorem 4.1 (real case), every $u \in \mathcal{A}^*$ must satisfies

$$\square_{d^v} u = 0,$$

which gives, by Lemma 6.4, that all coefficients of u are constants on leaves of \mathcal{V} . Denote by \mathcal{A}_1^* the space of smooth forms on X whose coefficients are constants on leaves of \mathcal{V} , the usual d operator reduces to $d^h + R_d$ on \mathcal{A}_1^* , which suggests to look at the following cohomology (notice that $(d^h + R_d)^2 = 0$ on \mathcal{A}_1^* , since the coefficients of R_d are constants on leaves of \mathcal{V})

$$H_{d^h + R_d}^* := \frac{\{u \in \mathcal{A}_1^* : (d^h + R_d)u = 0\}}{(d^h + R_d)(\mathcal{A}_1^*)}.$$

A representative of a class in $H_{d^h + R_d}^*$ is minimal if and only if it lies in $\ker \square_{d^h + R_d}$. Thus Theorem 4.1 implies that

$$\mathcal{H}_d^* \simeq H_{d^h + R_d}^*.$$

A similar argument also works in the complex case, to summarize, we get

Theorem 7.1. *Theorem 4.1 (real case) implies*

$$H_d^* \simeq H_{d^h + R_d}^*.$$

Theorem 4.1 (real case) implies

$$H_{\bar{\partial}}^{*,*} \simeq H_{\bar{\partial}^h + R_{\bar{\partial}}}^{*,*}.$$

Remark: In the setting of Theorem 6.7 and Theorem 6.3, one may continue to get a sequence of isomorphisms. In fact, with the notation in Theorem 6.3, denote by \mathcal{V}^j ($1 \leq j \leq k$) the distribution associated to $T^{v_1} \oplus \cdots \oplus T^{v_j}$, let \mathcal{A}_j^* be the space of smooth forms on X whose coefficients are constants on leaves of \mathcal{V}^j , then one may verify that

$$(d^{h_j} + R_d^j)^2 = 0, \quad \text{on } \mathcal{A}_j^*.$$

Put

$$H_{d^{h_j} + R_d^j}^* := \frac{\{u \in \mathcal{A}_j^* : (d^{h_j} + R_d^j)u = 0\}}{(d^{h_j} + R_d^j)(\mathcal{A}_j^*)}, \quad H_{d^{h_j}}^* := \frac{\{u \in \mathcal{A}_j^* : d^{h_j}u = 0\}}{d^{h_j}(\mathcal{A}_j^*)}$$

we have

Theorem 7.2. *Theorem 6.3 implies*

$$H_d^* \simeq H_{d^{h_1} + R_d^1}^*, \quad H_{d^{h_{j-1}}}^* \simeq H_{d^{h_j} + R_d^j}^*, \quad \forall 2 \leq j \leq k.$$

Theorem 6.7 implies

$$H_{\partial}^{*,*} \simeq H_{\partial^{h_1} + R_{\partial}^1}^{*,*}, \quad H_{\partial^{h_{j-1}}}^{*,*} \simeq H_{\partial^{h_j} + R_{\partial}^j}^{*,*}, \quad \forall 2 \leq j \leq k.$$

Remark: In the proper fibration case, the above theorem is essentially equivalent to the use of the E_2 term in the Leray–Serre spectral sequence.

In order to apply the above theorem, one has to study the relation between $H_{d^{h_j} + R_d^j}^*$ and $H_{d^{h_j}}^*$; in the next section, we shall prove that, with respect to the spectral sequence of the double complex $d^{h_j} + R_d^j$, the E_2 term is equal to $H_{d^{h_j}}^*$ and the E_{∞} term is $H_{d^{h_j} + R_d^j}^*$.

7.2. Spectral sequence for double complex. We shall use the notation in [14]. We look at the complex

$$d^{h_j} + R_d^j : \mathcal{A}_j^* \rightarrow \mathcal{A}_j^*,$$

denote its kernel by Z , image by B and put

$$H := \frac{Z}{B} = H_{d^{h_j} + R_d^j}^*.$$

With the notation in [14], we denote by pT as the space of forms in \mathcal{A}_j^* whose j -th horizontal degree is no less than p . By Definition 6.1, the maximal j -th horizontal degree is $n - r_j$, thus we have

$$\mathcal{A}_j^* = {}^0T \supset {}^1T \supset \cdots \supset {}^{n-r_j}T \supset {}^{n-r_j+1}T = \{0\}.$$

Denote by T^m the space of degree- m forms in \mathcal{A}_j^* . We have

$$\mathcal{A}_j^* = \bigoplus_{m=0}^n T^m.$$

Put

$${}^pT^m := {}^pT \cap T^m,$$

we know that ${}^pT^m$ is the space of degree- m forms in \mathcal{A}_j^* whose horizontal degree is no less than p . One may easily verify the following compatibility conditions in [14]

$$(d^{h_j} + R_d^j)T^m \subset T^{m+1}, \quad (d^{h_j} + R_d^j){}^pT \subset {}^pT.$$

In fact, notice that $d^{h_j}{}^pT \subset {}^{p+1}T$ and $R_d^j{}^pT \subset {}^{p+2}T$, the following stronger compatibility condition holds

$$(d^{h_j} + R_d^j){}^pT \subset {}^{p+1}T.$$

Moreover, we shall introduce

$${}^pZ^m := Z \cap {}^pT^n, \quad {}^pB^m := B \cap {}^pT^n, \quad {}^pH^m := \frac{{}^pZ^m}{{}^pB^m}.$$

The fundamental definition is the following

Definition 7.3 (Spectral sequence of a filtered complex [14]). *The modules of spectral sequence are*

$$E_r^{p,q} := \frac{{}^pT_r^{p+q}}{{}^pT_{r-1}^{p+q} + {}^pB_{r-1}^{p+q}}, \quad E_\infty^{p,q} := \frac{{}^pZ^{p+q}}{{}^pZ^{p+q} + {}^pB^{p+q}}, \quad r = 1, 2, \dots,$$

where

$${}^pT_r^m := \{u \in {}^pT^m : (d^{h_j} + R_d^j)u \in {}^{p+r}T\}, \quad {}^pB_r^m := \{(d^{h_j} + R_d^j)u \in {}^pT^m : u \in {}^{p-r}T\}.$$

One may verify that $d^{h_j} + R_d^j$ induces the following d_r complex

$$E_r^{p-r, q+r-1} \xrightarrow{d_r} E_r^{p,q} \xrightarrow{d_r} E_r^{p+r, q-r+1}.$$

As general properties of the spectral sequence, we have

Proposition 7.4. *The associated d_r cohomology gives the E_{r+1} term, more precisely*

$$E_{r+1}^{p,q} \simeq \frac{\ker d_r \cap E_r^{p,q}}{\text{Im } d_r \cap E_r^{p,q}}.$$

Moreover, we have

$$E_\infty^{p,q} \simeq \frac{{}^pH^{p+q}}{{}^{p+1}H^{p+q}}, \quad E_r^{p,q} = E_\infty^{p,q}, \quad \forall r \geq n - r_j + 1.$$

Remark: For the double complex $d = \partial + \bar{\partial}$ (also called Frölicher spectral sequence), it is well known that [14]

$$E_1^{p,q} \simeq H_{\bar{\partial}}^{p,q}.$$

In our case, we obtain the following

Lemma 7.5. $\bigoplus_{p+q=m} E_2^{p,q} \simeq H_{d^{h_j}}^m$.

Proof. Recall that

$$E_2^{p,q} := \frac{{}^pT_2^{p+q}}{{}^pT_1^{p+q} + {}^pB_1^{p+q}}.$$

Let us write $u \in {}^pT^m$ as

$$u = u^{p|m-p} + u^{p+1|m-p-1} + \dots,$$

where $u^{\alpha|\beta}$ denotes the component of u with horizontal degree α and vertical degree β . Notice that: $(d^{h_j} + R_d^j)u$ has no horizontal degree $p+1$ component if and only if

$$d^{h_j} u^{p|m-p} = 0.$$

Thus u lies in ${}^pT_2^{p+q}$ if and only if $d^{h_j} u^{p|m-p} = 0$. A similar discussion gives that u lies in ${}^pT_1^{p+q} + {}^pB_1^{p+q}$ if and only if $u^{p|m-p} \in \text{Im } d^{h_j}$. Thus the $(p|q)$ -component of $H_{d^{h_j}}^m$ is isomorphic to $E_2^{p,q}$. Hence the lemma follows. \square

7.3. The final proof. Let us introduce the following notation: let D be a complex on \mathcal{A}_j^* , denote its associated cohomology by H_D^* . Recall that \mathcal{A}_j^* means the space of smooth forms on X whose coefficients are constants on leaves of \mathcal{V}^j , in particular

$$A^* := \mathcal{A}_k^*$$

is the finite dimensional real vector space spanned by wedge products of the frame Φ . Assume that D maps A^* to itself. Put

$$H_{D,\Phi}^* := \frac{\ker D|_{A^*}}{\operatorname{Im} D|_{A^*}},$$

we say that H_D^* is *simple* if it reduces to $H_{D,\Phi}^*$, i.e.

$$H_D^* \simeq H_{D,\Phi}^*.$$

The real case of our main theorem is of course equivalent to that H_d^* is simple. By Theorem 7.2, H_d^* is simple if and only if $H_{d^{h_1}+R_d^1}$ is simple. Now let us look at the double complex (d^{h_1}, R_d^1) , by Lemma 7.5, its E_2 term is $H_{d^{h_1}}^*$. Thus in order to prove that $H_{d^{h_1}+R_d^1}^*$ is simple, it suffices to show that $H_{d^{h_1}}^*$ is simple. Apply Theorem 7.2 again, we know that $H_{d^{h_1}}^*$ is simple if and only if $H_{d^{h_2}+R_d^2}^*$ is simple. Now again Lemma 7.5 ensures that it is enough to prove simplicity of $H_{d^{h_2}}^*$. Repeat the above argument, we know it suffices to show that $H_{d^{h_k}}^*$ is simple, which holds trivially since $\mathcal{A}_k^* = A^*$. The complex case follows by a similar argument.

Remark: Notice that the Lie algebra cohomology $H_{d,\Phi}^*$ has the following “harmonic” representative

$$H_{d,\Phi}^* \simeq \mathcal{H}_{d,\Phi}^* := \{u \in A^* : \square_{R_d^1+\dots+R_d^{k-1}} u = 0\}.$$

Since

$$\mathcal{H}_{d,\Phi}^* \in \ker \square_d,$$

our main theorem actually also gives a formula for the following d -harmonic space

$$\mathcal{H}_d^* := \ker \square_d = \mathcal{H}_{d,\Phi}^*.$$

A more careful study of $\mathcal{H}_{d,\Phi}^*$ will be given in section 8.

7.4. Proof of the strong version of Cordero-Fernández-Gray-Ugarte’s theorem.

Let us choose an appropriate basis of left-invariant vector fields for the nilpotent Lie group and a left-invariant metric making this a global orthonormal frame. Then we know that our main theorem applies.

Remark: The above argument also applies to the original version of Nomizu’s theorem.

8. AN EXPLICIT VERSION OF OUR MAIN THEOREM

In case $k = 2$, the remark at the end of section 7.3 gives

$$(8.1) \quad \ker \square_d = \{u \in A^* : \square_{R_d^1} u = 0\}.$$

The above formula can also be proved using the E_3 term of the spectral sequence of the double complex (d^{h_1}, R_d^1) . In fact, by Lemma 7.5, we know that the E_3 term is defined by the following complex

$$R_d^1 : A^* \rightarrow A^*.$$

The key point here is that both R_d^1 and its adjoint send A^* to itself, from which we know that the above spectral sequence degenerates at E_3 .

Example 1: By the example in section 5, if X is the Kodaira–Thurston manifold, then $k = 2$ and

$$R_{\bar{\partial}}^1 = -\frac{i}{2}\varphi_h \wedge \overline{\varphi_h} \wedge (X^v \rfloor), \quad (R_{\bar{\partial}}^1)^* = \frac{i}{2}\varphi_v \wedge (\overline{X^h} \rfloor)(X^h \rfloor).$$

The complex version of (8.1) gives

$$\mathcal{H}_{\bar{\partial}}^{*,*} = \{u \in A^{*,*} : R_{\bar{\partial}}^1 u = (R_{\bar{\partial}}^1)^* u = 0\}.$$

Example 2: Let \mathfrak{g} be the 3-dimensional 2-step complex nilpotent Lie algebra admitting a complex $(1, 0)$ -coframe $\{\xi^1, \xi^2, \xi^3\}$ satisfying

$$d\xi^1 = \xi^2 \wedge \overline{\xi^2} + D\xi^3 \wedge \overline{\xi^3}, \quad d\xi^2 = 0, \quad d\xi^3 = 0,$$

where D is a complex parameter having non-negative imaginary part. Let G be the simply-connected Lie group having \mathfrak{g} as Lie algebra. Let X be any compact quotient of G . Then, $k = 2$, the decomposition of $\wedge^{1,0} T_X^*$ into vertical and horizontal subbundles is given by

$$\wedge^{1,0} T_X^* = \langle \xi^1 \rangle \oplus \langle \xi^2, \xi^3 \rangle$$

and

$$R_{\bar{\partial}}^1 = (\xi^2 \wedge \overline{\xi^2} + D\xi^3 \wedge \overline{\xi^3}) \wedge (\xi_1 \rfloor), \quad (R_{\bar{\partial}}^1)^* = \xi^1 \wedge [(\overline{\xi_2} \rfloor)(\xi_2 \rfloor) + \overline{D}(\overline{\xi_3} \rfloor)(\xi_3 \rfloor)],$$

where $\{\xi_1, \xi_2, \xi_3\}$ satisfies $\xi^r(\xi_s) = \delta_s^r$. Also in this case, the complex version of (8.1) gives

$$\mathcal{H}_{\bar{\partial}}^{*,*} = \{u \in A^{*,*} : R_{\bar{\partial}}^1 u = (R_{\bar{\partial}}^1)^* u = 0\}.$$

For general k , we shall introduce the following:

Definition 8.1. *With the notation in our main theorem. In the real case, the nilpotent frame Φ is said to be admissible if $k \geq 2$ or*

$$R_d^{k-j-1} \left(\square_{R_d^{k-j}} + \cdots + \square_{R_d^{k-1}} \right) = \left(\square_{R_d^{k-j}} + \cdots + \square_{R_d^{k-1}} \right) R_d^{k-j-1}, \quad \forall 1 \leq j \leq k-2,$$

when $k \geq 3$. In the complex case, the nilpotent frame Ψ is said to be admissible if $k \geq 2$ or

$$R_{\bar{\partial}}^{k-j-1} \left(\square_{R_{\bar{\partial}}^{k-j}} + \cdots + \square_{R_{\bar{\partial}}^{k-1}} \right) = \left(\square_{R_{\bar{\partial}}^{k-j}} + \cdots + \square_{R_{\bar{\partial}}^{k-1}} \right) R_{\bar{\partial}}^{k-j-1}, \quad \forall 1 \leq j \leq k-2,$$

when $k \geq 3$.

We obtain the following result:

Theorem 8.2. *With the assumptions in our main theorem. Assume further that the associated frame is admissible. Then:*

(1) *In the real case, we have*

$$\mathcal{H}_d^* = \{u \in A^* : R_d^j u = (R_d^j)^* u = 0, \forall 1 \leq j \leq k-1\}.$$

(2) *In the complex case, we have*

$$\mathcal{H}_{\bar{\partial}}^{*,*} = \{u \in A^{*,*} : R_{\bar{\partial}}^j u = (R_{\bar{\partial}}^j)^* u = 0, \forall 1 \leq j \leq k-1\}.$$

Proof. Notice that

$$R_d^{k-2} \left(\square_{R_d^{k-1}} \right) = \left(\square_{R_d^{k-1}} \right) R_d^{k-2}$$

implies (notice that $\mathcal{H}_{d^{h_{k-2}}}^* = \ker \square_{R_d^{k-1}}$)

$$R_d^{k-2}(\mathcal{H}_{d^{h_{k-2}}}^*) \subset \mathcal{H}_{d^{h_{k-2}}}^*,$$

from which we know that the following complex

$$R_d^{k-2} : H_{d^{h_{k-2}}}^* \rightarrow H_{d^{h_{k-2}}}^*,$$

is also well defined on the corresponding harmonic space $\mathcal{H}_{d^{h_{k-2}}}^*$. Thus the spectral sequence $(d^{h_{k-2}}, R_d^{k-2})$ degenerates at E_3 and

$$\mathcal{H}_{d^{h_{k-2}}+R_d^{k-2}}^* = \mathcal{H}_{d^{h_{k-2}}}^* \cap \ker \square_{R_d^{k-2}},$$

which is equivalent to

$$\mathcal{H}_{d^{h_{k-3}}}^* = \ker \square_{R_d^{k-2}+R_d^{k-1}} = \ker \square_{R_d^{k-2}} \cap \square_{R_d^{k-1}}.$$

Repeating the above argument gives the proof of the real case, the complex case follows by a similar argument. \square

Remark: Not all nilpotent frames are admissible, we shall give a counterexample in the next section.

9. FURTHER EXAMPLES

9.1. A nilmanifold not satisfying Theorem 8.2. Let \mathfrak{h}_7 be the real 6-dimensional nilpotent Lie algebra with basis $\{e_1, \dots, e_6\}$ such that

$$[e_1, e_2] = -e_4, \quad [e_1, e_3] = -e_5, \quad [e_2, e_3] = -e_6,$$

the other brackets vanishing. Then, denoting by $\{e^1, \dots, e^6\}$ the dual basis of $\{e_1, \dots, e_6\}$, we obtain the following structure equations

$$de^1 = 0, \quad de^2 = 0, \quad de^3 = 0, \quad de^4 = e^1 \wedge e^2, \quad de^5 = e^1 \wedge e^3, \quad de^6 = e^2 \wedge e^3.$$

Then the simply connected Lie group H_7 whose Lie algebra is \mathfrak{h}_7 admits compact quotients $M = \Gamma \backslash H_7$. Define an almost complex structure J_0 on M by the following complex $(1, 0)$ -coframe

$$\omega^1 = e^1 + ie^2, \quad \omega^2 = -2(e^3 + ie^4), \quad \omega^3 = -4(e^5 + ie^6).$$

Then

$$d\omega^1 = 0, \quad d\omega^2 = \omega^1 \wedge \overline{\omega^1}, \quad d\omega^3 = \omega^1 \wedge \omega^2 + \omega^1 \wedge \overline{\omega^2},$$

that is J_0 is in fact integrable and by Ugarte [29] (see also [23]) any other complex structure on M is equivalent to J_0 . Set

$$\omega^1 = \xi^3, \quad \omega^2 = \xi^2, \quad \omega^3 = \xi^1.$$

Then,

$$(9.1) \quad d\xi^1 = -\xi^2 \wedge \xi^3 + \xi^3 \wedge \overline{\xi^2}, \quad d\xi^2 = \xi^3 \wedge \overline{\xi^3}, \quad d\xi^3 = 0$$

and $k = 3$. The decomposition of $\wedge^{1,0} T_M^*$ into vertical and horizontal subbundles reads as

$$\wedge^{1,0} T_M^* = \langle \xi^1 \rangle \oplus \langle \xi^2, \xi^3 \rangle.$$

A direct computation taking into account (9.1) gives

$$\begin{aligned} R_{\overline{\partial}}^1 &= -\overline{\xi^2} \wedge \overline{\xi^3} \wedge (\overline{\xi^1} \rfloor) + \xi^3 \wedge \overline{\xi^2} \wedge (\xi^1 \rfloor), & R_{\overline{\partial}}^2 &= \xi^3 \wedge \overline{\xi^3} \wedge (\xi_2 \rfloor), \\ (R_{\overline{\partial}}^1)^* &= -\overline{\xi^1} \wedge [(\overline{\xi^3} \rfloor)(\overline{\xi^2} \rfloor)] + \xi^1 \wedge [(\overline{\xi^2} \rfloor)(\xi_3 \rfloor)], & (R_{\overline{\partial}}^2)^* &= \xi^2 \wedge [(\overline{\xi^3} \rfloor)(\xi_3 \rfloor)], \end{aligned}$$

where $\{\xi_1, \xi_2, \xi_3\}$ satisfies $\xi^r(\xi_s) = \delta_s^r$.

Theorem 9.1. *The Dolbeault cohomology group of $\Gamma \backslash H_7$ satisfies*

$$(9.2) \quad \mathcal{H}_{\bar{\partial}}^{1,1} \simeq \text{Span}_{\mathbb{C}} \langle \xi^{1\bar{2}}, \xi^{2\bar{3}}, \xi^{2\bar{2}} + \xi^{3\bar{1}}, \xi^{2\bar{2}} + \xi^{1\bar{3}} \rangle,$$

where $\xi^{i\bar{j}} := \xi^i \wedge \bar{\xi}^j$. In particular, $\mathcal{H}_{\bar{\partial}}^{1,1}$ does not satisfy Theorem 8.2.

Proof. Let

$$u = a_1 \xi^{1\bar{1}} + b_1 \xi^{2\bar{1}} + c_1 \xi^{3\bar{1}} + a_2 \xi^{1\bar{2}} + b_2 \xi^{2\bar{2}} + c_2 \xi^{3\bar{2}} + a_3 \xi^{1\bar{3}} + b_3 \xi^{2\bar{3}} + c_3 \xi^{3\bar{3}}$$

be a harmonic (1,1)-form. By our main theorem, one may assume that all the above coefficients are constants. Then u is harmonic if and only if

$$\bar{\partial}u = \left(R_{\bar{\partial}}^1 + R_{\bar{\partial}}^2 \right) u = \bar{\partial}^* u = \left((R_{\bar{\partial}}^1)^* + (R_{\bar{\partial}}^2)^* \right) u = 0,$$

which gives (9.2). Notice that $\mathcal{H}_{\bar{\partial}}^{1,1}$ satisfies Theorem 8.2 if and only if

$$\mathcal{H}_{\bar{\partial}}^{1,1} \subset \ker R_{\bar{\partial}}^1 \cap \ker (R_{\bar{\partial}}^1)^* \cap \ker R_{\bar{\partial}}^2 \cap \ker (R_{\bar{\partial}}^2)^*$$

But obviously

$$R_{\bar{\partial}}^2(\xi^{2\bar{1}} + \xi^{3\bar{1}}) \neq 0,$$

thus $\mathcal{H}_{\bar{\partial}}^{1,1}$ does not satisfy Theorem 8.2. \square

9.2. Nilpotent frame in a non-nilpotent Lie group. Let us look at the following example from [16].

Example: Consider the following group structure

$$(a, \lambda) * (t, z) = (a + t, \lambda + e^{2\pi i a} z),$$

on $\mathbb{R} \times \mathbb{C}$. Then we know that

$$G := (\mathbb{R} \times \mathbb{C}, *),$$

is a Lie group. It is clear that

$$\Gamma := \mathbb{Z} \times (\mathbb{Z} + i\mathbb{Z})$$

is a discrete subgroup of G such that the quotient space (with respect to the left action)

$$X := \Gamma \backslash G,$$

is a compact smooth manifold. It is clear that

$$dt, e^{-2\pi i t} dz,$$

are G -invariant with respect to the left action of G . Put

$$e^1 = dt, e^2 = \text{Re}(e^{-2\pi i t} dz), e^3 = \text{Im}(e^{-2\pi i t} dz).$$

we know that $\{e^1, e^2, e^3\}$ is a basis of the dual Lie algebra \mathfrak{g}^* of G . Notice that, if $z = x + iy$ then

$$e^2 = (\cos 2\pi t) dx + (\sin 2\pi t) dy,$$

and

$$e^3 = -(\sin 2\pi t) dx + (\cos 2\pi t) dy.$$

Thus

$$de^1 = 0, de^2 = 2\pi e^1 \wedge e^3, de^3 = -2\pi e^1 \wedge e^2,$$

which gives

$$[\mathfrak{g}, [\mathfrak{g}, \mathfrak{g}]] = [\mathfrak{g}, \mathfrak{g}], [[\mathfrak{g}, \mathfrak{g}], [\mathfrak{g}, \mathfrak{g}]] = 0.$$

Thus \mathfrak{g} is solvable but not nilpotent. But notice that

$$\sigma^1 := dt, \quad \sigma^2 := dx, \quad \sigma^3 := dy,$$

give a smooth frame of T_X^* such that

$$d\sigma^j = 0, \quad j = 1, 2, 3.$$

Thus

$$\Phi := \{\sigma^1, \sigma^2, \sigma^3\},$$

is a nilpotent frame of X and our main theorem implies that the de Rham cohomology of X is isomorphic to the exterior algebra generated by Φ (in fact it is easy to see that X is diffeomorphic to a real torus).

REFERENCES

- [1] J. A. Alvarez López, A finiteness theorem for the spectral sequence of a Riemannian foliation, *Illinois J. Math.* **33** (1989), pp. 79–92.
- [2] J. A. Alvarez López, A decomposition theorem for the spectral sequence of Lie foliations. *Trans. Amer. Math. Soc.*, **329** (1992), pp. 173–184.
- [3] J. A. Alvarez López, P. Tondeur, Hodge decomposition along the leaves of a Riemannian foliation. *J. Funct. Anal.*, **99** (1991), pp. 443–458.
- [4] B. Berndtsson, M. Păun, X. Wang, *Algebraic fiber spaces and curvature of higher direct images*, arXiv:1704.02279.
- [5] A. Blanchard, Sur les variétés analytiques complexes, *Ann. Scient. Ecole Norm. Sup.*, **73** (1956) 157–202.
- [6] S. Console, A. Fino, On the de Rham cohomology of solvmanifolds, *Ann. Scuola Norm. Sup.*, **10** (2011), 801–818.
- [7] S. Console, A. Fino, H. Kasuya, On de Rham and Dolbeault cohomology of solvmanifolds, *Transf. Groups*, **21** (2016).pp. 653–680.
- [8] L. A. Cordero, M. Fernández, A. Gray, L. Ugarte, Nilpotent complex structures on compact nilmanifolds, *Rend. Circolo Mat. Palermo*, **49** (1997), 83–100.
- [9] L. A. Cordero, M. Fernández, A. Gray, L. Ugarte, Compact nilmanifolds with nilpotent complex structures: Dolbeault cohomology, *Trans. Amer. Math. Soc.*, **352** (2000) pp. 5405–5433.
- [10] L. J. Corwin and F. P. Greenleaf, *Representations of nilpotent Lie groups and their applications. Part I*, Cambridge Studies in Advanced Mathematics, vol. 18. Cambridge University Press, Cambridge (1990).
- [11] P. Deligne, Théorème de Lefschetz et critères de dégénérescence de suites spectrales, *Publications Mathématiques de l’Institut des Hautes Études Scientifiques*, **35** (1968) pp. 107–126.
- [12] J. P. Demailly, Complex analytic and differential geometry, Université de Grenoble I Institut Fourier, available in Demailly’s homepage.
- [13] A. Fino, S. Rollenske, J. Ruppenthal, Dolbeault cohomology of complex nilmanifolds foliated in toroidal groups, arXiv:1808.08090 [math.DG]
- [14] A. Frölicher, *Relations between the cohomology groups of Dolbeault and topological invariants*, Proc. Natl. Acad. Sci. USA **41** (1955) pp. 641–644.
- [15] P. A. Griffiths, The extension problem in complex analysis II; embeddings with positive normal bundle, *Amer. J. Math.* **88** (1966) pp. 366–446.
- [16] H. Kasuya, An extension of Nomizu’s Theorem—A user’s guide—, *Complex Manifolds* (2016) pp. 231–238.
- [17] K. Kodaira, On the structure of compact complex analytic surfaces. I, *Amer. J. Math.* **86** (1964) pp. 751–798.
- [18] Y. A. Kordyukov, Adiabatic limits and spectral geometry of foliations, *Math. Ann.* **313** (1999) pp. 763–783.
- [19] F. W. Kamber, P. Tondeur, de Rham–Hodge theory for Riemannian foliations, *Math. Ann.* **277** (1987) pp. 415–431.
- [20] K. Nomizu, On the cohomology of compact homogeneous spaces of nilpotent Lie groups, *Ann. of Math.* **59** (1954), pp. 531–538.
- [21] B. L. Reinhart, Harmonic Integrals on Foliated Manifolds, *Amer. J. Math.*, **81**, (1959), pp. 529–536.

- [22] S. Rollenske. Geometry of nilmanifolds with left-invariant complex structure and deformations in the large. *Proc. Lond. Math. Soc. (3)*, 99(2):425–460, 2009.
- [23] S. Rollenske. Dolbeault cohomology of nilmanifolds with left-invariant complex structure. In *Complex and differential geometry*, volume 8 of *Springer Proc. Math.*, pages 369–392. Springer, Heidelberg, 2011.
- [24] Y. Sakane, *On compact complex parallelisable solvmanifolds*, *Osaka J. Math.* **13** (1976), pp. 187–212,
- [25] Simon M. Salamon. Complex structures on nilpotent Lie algebras. *J. Pure Appl. Algebra*, 157(2-3):311–333, 2001.
- [26] R. Takagi and S. Yorozy, Notes on the Laplace–Beltrami operator on a foliated Riemannian manifold with a bundle-like metric, *Nihonkai Math. J.*, **1** (1990) pp. 89–106.
- [27] A. Tomassini, X. Wang, Some results on the Hard Lefschetz Condition, *Internat. J. Math.* **29** (2018), 1850095 (30 pp.).
- [28] W. P. Thurston, Some simple examples of symplectic manifolds, *Proc. Amer. Math. Soc.* **55** (1976), pp. 467–468.
- [29] L. Ugarte, Hermitian structures on six-dimensional nilmanifolds, *Transf. Groups*, **12** (2007) pp. 175–202.

FB 12 / MATHEMATIK UND INFORMATIK, PHILIPPS-UNIVERSITÄT MARBURG, HANS-MEERWEIN-STR. 6, 35032 MARBURG, GERMANY

Email address: `rollenske@mathematik.uni-marburg.de`

DIPARTIMENTO DI SCIENZE MATEMATICHE, FISICHE E INFORMATICHE, UNITÀ DI MATEMATICA E INFORMATICA, UNIVERSITÀ DEGLI STUDI DI PARMA, PARCO AREA DELLE SCIENZE 53/A, 43124, PARMA, ITALY

Email address: `adriano.tomassini@unipr.it`

DEPARTMENT OF MATHEMATICAL SCIENCES, NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, NO-7491 TRONDHEIM, NORWAY

Email address: `xu.wang@ntnu.no`