

# Chiral Phonon Transport Induced by Topological Magnons

Even Thingstad, Akashdeep Kamra, Arne Brataas, and Asle Sudbø  
*Center for Quantum Spintronics, Department of Physics,  
Norwegian University of Science and Technology, NO-7491 Trondheim, Norway*

The plethora of recent discoveries in the field of topological electronic insulators has inspired a search for boson systems with similar properties. There are predictions that ferromagnets on a two-dimensional honeycomb lattice may host chiral edge magnons. In such systems, we theoretically study how magnons and phonons couple. We find topological magnon-polarons around the avoided crossings between phonons and topological magnons. Exploiting this feature along with our finding of Rayleigh-like edge phonons in armchair ribbons, we demonstrate the existence of chiral edge modes with a phononic character. We predict that these modes mediate a chirality in the coherent phonon response and suggest measuring this effect via elastic transducers. These findings reveal a possible approach towards heat management in future devices.

*Introduction.*— Topological electronic insulators [1–5] are characterized by an insulating bulk with conducting ‘chiral’ edge states. The unidirectional propagation of these chiral modes is ‘topologically protected’ against defects at low temperatures when we can disregard inelastic scattering from phonons [5]. This has led to the development of a wide range of essential concepts, including Majorana modes [6–9], topological quantum computation [10, 11], and chiral transport. Inspired by these findings, there has been an upsurge of efforts towards finding similar states in other systems [12] with an emphasis on bosonic excitations [13–19]. There are predictions of topological magnons [15–17] in honeycomb ferromagnets with an engineered Dzyaloshinskii-Moriya interaction [20, 21] that induces the necessary band gap. In contrast to fermionic systems with Fermi energy within this band gap, the bulk is not necessarily insulating in bosonic systems [22].

The field of magnonics [23–26] focuses on pure spin transport mediated by magnons [27]. It is possible to exploit the low-dissipation and wave-like nature of these excitations in information processing [28, 29]. The coherent pumping of chiral surface spin wave (Damon-Eshbach) modes induces cooling via incoherent magnon-phonon scattering [30]. Besides application oriented properties, the bosonic nature of magnons, combined with spintronic manipulation techniques [24, 31], allows for intriguing physics [32–35]. The coupling [36] between magnons and phonons fundamentally differs from the electron-phonon interaction and results in a coherent hybridization of the modes [37], in addition to the temperature dependent incoherent effects [30, 38] discussed above. The direct influence of the hybridization between magnons and phonons, known as magnon-polarons [39, 40], has been observed in spin and energy transport in magnetic systems [41–46].

In this Letter, we address the robustness of the topological magnons in a honeycomb ferromagnet [15–17] against their coupling with the lattice vibrations. In contrast to the case of electron-phonon coupling, where phonons can be disregarded at low temperatures, the magnon dispersion may undergo significant changes with

new states emerging in the band gap [45, 46]. We find that in the honeycomb ferromagnet with spins oriented orthogonal to the lattice plane, only transverse phonon modes with out-of-plane displacement couple to spin. To understand the eigenmodes, we evaluate and analyze the coupled spin and out-of-plane phonon modes for an infinitely large plane as well as for a finite ribbon geometry. We quantify the effect of the magneto-elastic coupling on the magnon Hall conductivity and find a non-monotonic dependence on the coupling strength. Our analysis of the finite ribbons shows that topological magnons hybridize with bulk phonons around the avoided crossings in their coupled dispersion, forming magnon-polarons with topological chiral properties. Hence, while their edge localization is weakened, the magneto-elastic coupling does not completely remove the topological magnons. Furthermore, we find that armchair edges support Rayleigh-like edge phonon modes in sharp contrast to the zigzag edges. When topological magnons hybridize with these edge phonons, edge magnon-polarons with almost undiminished chirality are formed. We suggest a setup which utilizes this induced chirality in coherent phonon transport. Such systems enable the observation of the topological physics and serve as a prototype for a unidirectional heat pump. This offers a highly feasible alternative to producing topological phonon diodes [47–49].

*Model.*— We consider a ferromagnetic material with localized spins on a two-dimensional honeycomb lattice, allow for out-of-plane vibrations of the lattice sites, and assume there is magneto-elastic coupling. This system can be modelled by a Hamiltonian of the form  $H = H_m + H_{\text{ph}} + H_{\text{me}}$ , where  $H_m$  is the magnetic Hamiltonian,  $H_{\text{ph}}$  describes the phonons, and  $H_{\text{me}}$  represents the magneto-elastic coupling.

The Hamiltonian we consider is inspired by the Haldane model [1] given by [15–17]

$$H_m = -J \sum_{\langle i,j \rangle} \mathbf{S}_i \cdot \mathbf{S}_j + \mathcal{D} \sum_{\langle\langle i,j \rangle\rangle} \nu_{ij} \hat{z} \cdot \mathbf{S}_i \times \mathbf{S}_j - \mathcal{B} \sum_i S_i^z. \quad (1)$$

The first term describes the ferromagnetic exchange coupling between nearest neighbour sites, while the sec-

ond accounts for the Dzyaloshinskii-Moriya interaction [20, 21] between next-to-nearest neighbours [50]. The Haldane sign  $\nu_{ij} = \pm 1$  depends on the relative orientation of the next-to-nearest neighbours as shown in Fig. 1(a), and is the root of non-trivial topological properties. We let the nearest neighbour distance be  $d$  and the next-to-nearest neighbour distance be  $a$ . Refs. [16, 17] discuss the dispersion relation and Berry curvature of this spin model in linear spin wave theory.

For the phonon Hamiltonian, we consider only the out-of-plane degrees of freedom since only these modes couple to the spin to lowest order in the linear spin wave expansion. We assume nearest-neighbour interactions with elastic constant  $C$ , let the mass associated with the spins on the lattice sites be  $m$ , and disregard substrate coupling. Introducing  $S_k = \sum_{\beta} \cos(\mathbf{k} \cdot \boldsymbol{\beta})$ , where the sum is over the three next-to-nearest neighbour vectors  $\boldsymbol{\beta}$  of Fig. 1(a), we obtain the dispersion relation

$$\omega_{\pm}^{\text{ph}}(\mathbf{k}) = \sqrt{\frac{C}{m}} \sqrt{3 \pm \sqrt{3 + 2S_k}} \quad (2)$$

for the free phonon modes.

Motivated by the continuum limit description [36, 37], we write down the lattice magneto-elastic coupling to linear order in the magnon amplitude, obtaining

$$H_{\text{me}} = \kappa \sum_D \sum_{i \in D} \sum_{\boldsymbol{\alpha}_D} \mathbf{S}_i \cdot \boldsymbol{\alpha}_D (u_i^z - u_{i+\boldsymbol{\alpha}_D}^z) \quad (3)$$

where  $\kappa$  parametrizes the strength of the magnon-phonon coupling,  $\sum_D$  denotes the sum over sublattices,  $\sum_{i \in D}$  is the sum over the lattice sites on the  $D$  sublattice, and  $\boldsymbol{\alpha}_D$  are the corresponding nearest neighbour vectors. The out-of-plane deviation for lattice site  $i$  is denoted by  $u_i^z$ .

*Bulk spectrum.*—We introduce the Holstein-Primakoff representation of spins and use linear spin wave theory in the spin- and magneto-elastic terms [27]. Within the rotating wave approximation [51], the resulting Hamiltonian describing the phonon and magnon modes of the system is obtained as  $H = \sum_k \psi_k^\dagger M_k \psi_k$ , where  $\psi_k^\dagger = (a_k^\dagger, b_k^\dagger, c_{k-}^\dagger, c_{k+}^\dagger)$ . Here,  $a_k$  and  $b_k$  are annihilation operators for the sublattice magnon modes on the  $A$  and  $B$  sublattices, while  $c_{k\pm}$  are the annihilation operators for the phonon modes. The matrix  $M_k$  takes the form

$$M_k = \begin{pmatrix} A + h^z & h^- & g_{A-} & g_{A+} \\ h^+ & A - h^z & g_{B-} & g_{B+} \\ g_{A-}^* & g_{B-}^* & \omega_{k-}^{\text{ph}} & 0 \\ g_{A+}^* & g_{B+}^* & 0 & \omega_{k+}^{\text{ph}} \end{pmatrix}, \quad (4)$$

where  $A = 3JS + \mathcal{B}$ ,  $h^z(\mathbf{k}) = 2DS \sum_{\beta} \sin(\mathbf{k} \cdot \boldsymbol{\beta})$ ,  $h^-(\mathbf{k}) = -JS \sum_{\alpha} \exp(-i\mathbf{k} \cdot \boldsymbol{\alpha})$ , and  $h^+ = (h^-)^*$ . The coupling between the  $D$ -sublattice magnons and the phonon branch  $\pm$  is captured by  $g_{D\pm}$ , which is proportional to the dimensionless coupling strength  $\tilde{\kappa} =$




FIG. 1. (a) Lattice geometry showing the nearest neighbour vectors  $\boldsymbol{\alpha}$ , next-to-nearest neighbour vectors  $\boldsymbol{\beta}$ , and the Haldane sign  $\nu_{ij} = \pm 1$ . (b) The first Brillouin zone in reciprocal space, including the paths along which we plot the dispersion relation in figure (c). The parameter values used are  $\mathcal{D} = 0.1J$ ,  $\mathcal{B} = 0.4JS$ ,  $\sqrt{C/m} = 1.5JS$ , and rescaled coupling strength  $\tilde{\kappa} = 0.03$  (see main text). The magnon (yellow) and phonon (purple) character of the modes is indicated with colors. The modes are significantly affected by the magneto-elastic coupling only close to avoided crossings.

$(\kappa d/JS)^4 \sqrt{\hbar^2 S^2 / 16m^2 (C/m)}$ . The spectrum obtained by diagonalizing this matrix is plotted in Fig. 1(c) along the paths displayed in Fig. 1(b).

*Hall conductivity.*—The topological nature of the spin model is manifested in the magnon Hall conductivity that arises because of the time-reversal symmetry breaking caused by the Dzyaloshinskii-Moriya interaction.

The spin current operator  $J_\gamma$  may be found from a continuity equation or magnon group velocity approach [52], both yielding

$$J_\gamma = \sum_k \begin{pmatrix} a_k^\dagger & b_k^\dagger \end{pmatrix} \left( \frac{\partial H_m(\mathbf{k})}{\partial k_\gamma} \right) \begin{pmatrix} a_k \\ b_k \end{pmatrix} \quad (5)$$

along the Cartesian direction  $\gamma$ . Here,  $H_m(\mathbf{k})$  is the matrix representation of the magnon Hamiltonian. Assuming we apply a weak in-plane magnetic field gradient  $\nabla \mathcal{B}$ , we are interested in the current  $\mathbf{j} = \sigma \nabla \mathcal{B}$ , which is determined by the conductivity tensor  $\sigma$  [52]. The Hall conductivity can be calculated using the Kubo formula, giving

$$\sigma_{xy} = \sum_k \sum_{\alpha, \beta \neq \alpha} n_B(E_{k\alpha}) C_{\alpha\beta}(\mathbf{k}), \quad (6)$$

where  $E_{k\alpha}$  is the energy eigenvalue of band  $\alpha$  and  $n_B(E_{k\alpha})$  is the corresponding Bose factor. The curvature-tensor  $C_{\alpha\beta}$  is given by

$$C_{\alpha\beta}(\mathbf{k}) = i \frac{J_y^{\alpha\beta}(\mathbf{k})J_x^{\beta\alpha}(\mathbf{k}) - J_x^{\alpha\beta}(\mathbf{k})J_y^{\beta\alpha}(\mathbf{k})}{(E_{k\alpha} - E_{k\beta})^2}, \quad (7)$$

where  $(\alpha, \beta)$  are band-indices, and  $J_\gamma^{\alpha\beta}(\mathbf{k})$  are the energy eigenstate matrix elements of the current operator at quasimomentum  $\mathbf{k}$ . Disregarding the magneto-elastic coupling, the band-curvature  $C_\alpha = \sum_{\beta \neq \alpha} C_{\alpha\beta}$  can be identified as the Berry curvature.

Expressing the sublattice magnon operators in terms of the eigenmode operators, one may identify the current matrix elements  $J_\gamma^{\alpha\beta}$  and integrate the curvature over the Brillouin zone to obtain the Hall conductivity. We are particularly interested in the effect of the magneto-elastic coupling, and therefore present the dependence of the Hall conductivity on the dimensionless coupling  $\tilde{\kappa}$  in Fig 2.

To understand this dependence, we consider the curvature-tensor  $C_{\alpha\beta}$ . When the bands  $\alpha$  and  $\beta$  both have a predominant magnon content, the topological nature of the underlying magnons gives a finite curvature. This magnon curvature is largest close to the Dirac points [16, 17]. Close to an avoided crossing, the magneto-elastic coupling changes the spectrum and causes transfer of band-curvature between the relevant bands  $\alpha$  and  $\beta$ . The latter can be seen by plotting the curvature-tensor element  $C_{\alpha\beta}$  for the band-pairs with avoided crossings, as shown in the insets of Fig 2. The resulting change in Hall conductivity is given by these curvature-tensor elements weighted with the difference between the Bose factors of the relevant bands. This follows from the anti-symmetry property of the curvature-tensor. The two band-pairs in the insets contribute oppositely to the Hall conductivity, and the competition between their curvature transfer explains the non-monotonic behaviour of the Hall conductivity.

*Ribbon geometry and coherent transport.*—Due to the topological nature of the magnon model under consideration and the bulk-boundary correspondence, there are gapless magnon edge states in a finite sample [5, 15–17]. Considering an armchair ribbon with finite width, the one-dimensional projection of the energy spectrum is plotted in Fig. 3. The corresponding spectrum for the zigzag edge ribbon is given in the Supplemental Material [53], where also Refs. [54–58] appear. Magnon and phonon modes hybridize in regions with an avoided crossing. When the upper phonon band lies within the bandgap of the pure magnon spectrum, there are modes with a mixed content of chiral magnon edge states and phonons. Although the spectra look qualitatively similar, there is a crucial distinction between the two cases. For the zigzag edge configuration, all the phonon modes




FIG. 2. Dependence of the Hall conductivity on the magneto-elastic coupling strength  $\tilde{\kappa}$  for parameter values  $\mathcal{D} = 0.1J$ ,  $\mathcal{B} = 0.4JS$ , and  $\sqrt{C}/m = 1.5JS$  for different temperatures  $T$  in units of  $JS$ . The insets show the quasimomentum dependence of the curvature-tensor at  $\tilde{\kappa} = 0.03$  for band-pairs (1,2) and (2,3), where the bands are labelled according to their energy and band 1 is the lowest band. The dominant contribution in these band-pairs comes from the regions with avoided crossings of the respective bands.

are delocalized throughout the sample, while the armchair edges host “Rayleigh-like” edge phonon modes. In direct analogy with Rayleigh modes on the surface of a three-dimensional material, the localization length of these modes is directly proportional to their wavelength, as shown in the Supplemental Material [53]. These edge phonon modes are supported by the half-hexagon protrusions of the armchair edge that can pivot around the bonds parallel to the edges connecting these protrusions, see Fig. 3. No such parallel bonds exist for the zigzag edge.

The Hall conductivity is a hallmark of topological electronic properties and motivates a similar role for the Hall conductivity mediated by topological magnons. However, in contrast to electrons, the bosonic nature of the magnons results in the lack of a general proportionality between the magnon Hall conductivity and the Chern number [52]. Furthermore, the observation of a magnon planar Hall effect [59] in a cubic, non-topological magnet suggests that this Hall conductivity may not be regarded as a smoking-gun signature for topological properties. Thus, we suggest a complementary approach to observe the topological nature of the underlying magnons by elastically probing the chirality of the magneto-elastic hybrid modes.

We propose to observe coherent chiral phonon propagation in the experimental setup of Fig. 4(b) by utilizing the edge modes, as depicted schematically in Fig. 4(a) [60], on the upper armchair edge of the sample. Taking inspiration from previous related experiments [41, 61], we suggest to inject elastic energy into

the sample middle at the upper edge using a nano-scale variant of the interdigital transducer design [62, 63], elaborated further in the Supplemental Material [53]. For a given transducer design, modes are excited with fixed wavevectors  $\pm k_x$  and a tunable frequency. Similar transducers can be used to detect the elastic response  $p_{L/R}$  on the left (L) and right (R) edges of the sample. Here,  $p_{L/R}$  is the elastic power detected at the transducers.

Figure 4(a) schematically depicts the dispersion for the magnetoelastic modes localized on an armchair edge. Disregarding magnetoelastic coupling, the edge hosts two counterpropagating Rayleigh-like edge phonons and a single chiral edge magnon. There is thus no chirality in the phononic response. Due to magnetoelastic coupling, the Rayleigh-like phonon with wavevector  $-k_x$  hybridizes with the chiral magnon to form a magnon-polaron while the other phonon remains unchanged. This breaks the symmetry between the counterpropagating phononic modes and the result is non-zero chirality  $\chi = (p_R - p_L)/(p_R + p_L)$ . Furthermore, as shown in Fig. 4(a), the hybridization with the magnon mode reverses the group velocity direction of the participating phonon mode. In principle, this gives perfectly chiral phonon transport.

The wavevector location of the avoided crossing can be tuned via the Zeeman shift in the magnon dispersion. Performing a frequency integrated measurement over an energy range of the same order as the magnetoelastic coupling, one obtains a peaked chirality when the magnetic field is such that the wavevector of the avoided crossing coincides with the wavevector of the transducer, obtaining a chirality as shown in Fig. 4(b). Performing a similar transport experiment on the zigzag edge does not give chiral phonon transport since the delocalized phonons hybridize with counterpropagating magnons on both the edges, thereby destroying the overall chirality. In addition, the size of the avoided crossing is smaller due to the smaller overlap with the localized chiral magnon. The armchair edge is therefore crucial for obtaining the chirality.

*Summary.*— We have examined the robustness of topological magnons in a honeycomb ferromagnet against their interaction with phonons. Their topological properties, albeit weakened, survive the magneto-elastic coupling. The magnon Hall conductivity of the system is found to depend on the magneto-elastic coupling strength in a non-monotonic, temperature-sensitive manner. Exploiting the Rayleigh-like edge phonons in armchair ribbons, we predict the existence of topological magnon-polarons confined to the boundary. We have suggested an experimental setup capable of probing the chiral nature of the topological magnon-polarons by elastic means, which thus serves as a platform for chiral coherent phononic transport.

*Acknowledgments.*— We acknowledge support from the Research Council of Norway Grant Nos. 262633 “Cen-




FIG. 3. One-dimensional projection of the dispersion relation for the magnetoelastic modes on a honeycomb ribbon with armchair edges. In addition to the bulk bands, there are two topological edge magnon states crossing the magnon band gap, as well as Rayleigh-like edge phonons. The inset shows the avoided crossing of a topological magnon edge mode with the two quasi-degenerate edge phonon modes. The parameter values are  $B = 0$ ,  $\sqrt{C/m} = 1.37JS$ ,  $\mathcal{D} = 0.1J$  and  $\tilde{\kappa} = 0.03$ .

ter of Excellence on Quantum Spintronics”, and 250985, “Fundamentals of Low-dissipative Topological Matter”.

- 
- [1] F. D. M. Haldane, “Model for a quantum hall effect without landau levels: Condensed-matter realization of the “parity anomaly”,” *Phys. Rev. Lett.* **61**, 2015–2018 (1988).
  - [2] C. L. Kane and E. J. Mele, “ $Z_2$  topological order and the quantum spin hall effect,” *Phys. Rev. Lett.* **95**, 146802 (2005).
  - [3] C. L. Kane and E. J. Mele, “Quantum spin hall effect in graphene,” *Phys. Rev. Lett.* **95**, 226801 (2005).
  - [4] B. Bernevig, T. L. Hughes, and S.-C. Zhang, “Quantum spin hall insulator state in hgte quantum wells,” *Science* **314**, 1757–1761 (2006).
  - [5] M. Z. Hasan and C. L. Kane, “Colloquium: Topological insulators,” *Rev. Mod. Phys.* **82**, 3045–3067 (2010).
  - [6] C.W.J. Beenakker, “Search for majorana fermions in superconductors,” *Annual Review of Condensed Matter Physics* **4**, 113–136 (2013), <https://doi.org/10.1146/annurev-conmatphys-030212-184337>.
  - [7] Jason Alicea, “New directions in the pursuit of majorana fermions in solid state systems,” *Reports on Progress in Physics* **75**, 076501 (2012).
  - [8] V. Mourik, K. Zuo, S. M. Frolov, S. R. Plissard, E. P. A. M. Bakkers, and L. P. Kouwenhoven, “Signatures of majorana fermions in hybrid superconductor-semiconductor nanowire devices,” *Science* **336**, 1003–1007 (2012), <http://science.sciencemag.org/content/336/6084/1003.full.pdf>.
  - [9] Liang Fu and C. L. Kane, “Superconducting proximity




FIG. 4. (a) Schematic spectrum for the coupled Rayleigh-like edge phonons and the topological edge magnon on the upper armchair sample edge. The phonon at quasimomentum  $-k_x$  hybridizes with the chiral magnon, while the phonon at quasimomentum  $+k_x$  is unaffected due to the lack of a magnon with matching wavevector at this edge. At the avoided crossing, there is propagation direction reversal for the modes with a phononic content. The color of the dispersion represents its nature with purple representing phononic and yellow magnonic character. (b) Proposed experimental setup for detecting coherent chiral transport through excitation of phononic modes. Elastic energy is injected in the sample middle on the upper armchair edge and detected at the left (L) and right (R) edges using wavevector and frequency resolved elastic transducers (purple). By exciting modes at the avoided crossing in (a), only the elastic excitations at one of the two quasimomenta  $\pm k_x$  are converted into hybridized modes (green arrows). This gives a chiral response, and the chirality is peaked when the wavevector of the avoided crossing coincides with the fixed wavevector of the transducer.

effect and majorana fermions at the surface of a topological insulator,” *Phys. Rev. Lett.* **100**, 096407 (2008).

- [10] Chetan Nayak, Steven H. Simon, Ady Stern, Michael Freedman, and Sankar Das Sarma, “Non-abelian anyons and topological quantum computation,” *Rev. Mod. Phys.* **80**, 1083–1159 (2008).
- [11] Sankar Das Sarma, Michael Freedman, and Chetan Nayak, “Topologically protected qubits from a possible non-abelian fractional quantum hall state,” *Phys. Rev. Lett.* **94**, 166802 (2005).
- [12] Gregor Jotzu, Michael Messer, Rémi Desbuquois, Martin Lebrat, Thomas Uehlinger, Daniel Greif, and Tilman Esslinger, “Experimental realization of the topological haldane model with ultracold fermions,” *Nature* **515**, 237 EP – (2014).
- [13] Sylvain Lannebère and Mário G. Silveirinha, “Link between the photonic and electronic topological phases in

artificial graphene,” *Phys. Rev. B* **97**, 165128 (2018).

- [14] Yizhou Liu, Yong Xu, Shou-Cheng Zhang, and Wenhui Duan, “Model for topological phononics and phonon diode,” *Phys. Rev. B* **96**, 064106 (2017).
- [15] S A Owerre, “A first theoretical realization of honeycomb topological magnon insulator,” *Journal of Physics: Condensed Matter* **28**, 386001 (2016).
- [16] S. A. Owerre, “Topological honeycomb magnon hall effect: A calculation of thermal hall conductivity of magnetic spin excitations,” *Journal of Applied Physics* **120**, 043903 (2016), <https://doi.org/10.1063/1.4959815>.
- [17] Se Kwon Kim, Héctor Ochoa, Ricardo Zarzuela, and Yaroslav Tserkovnyak, “Realization of the haldane-kane-mele model in a system of localized spins,” *Phys. Rev. Lett.* **117**, 227201 (2016).
- [18] S A Owerre and J Nsofini, “Squeezed dirac and topological magnons in a bosonic honeycomb optical lattice,” *Journal of Physics: Condensed Matter* **29**, 455802 (2017).
- [19] Heung-Sik Kim and Hae-Young Kee, “Realizing haldane model in fe-based honeycomb ferromagnetic insulators,” *npj Quantum Materials* **2**, 20 (2017).
- [20] I. Dzyaloshinsky, “A thermodynamic theory of weak ferromagnetism of antiferromagnetics,” *Journal of Physics and Chemistry of Solids* **4**, 241 – 255 (1958).
- [21] Tôru Moriya, “Anisotropic superexchange interaction and weak ferromagnetism,” *Phys. Rev.* **120**, 91–98 (1960).
- [22] Andreas Rückriegel, Arne Brataas, and Rembert A. Duine, “Bulk and edge spin transport in topological magnon insulators,” *Phys. Rev. B* **97**, 081106 (2018).
- [23] V V Kruglyak, S O Demokritov, and D Grundler, “Magnonics,” *Journal of Physics D: Applied Physics* **43**, 264001 (2010).
- [24] A. V. Chumak, V. I. Vasyuchka, A. A. Serga, and B. Hillebrands, “Magnon spintronics,” *Nat Phys* **11**, 453 (2015).
- [25] Gerrit E. W. Bauer, Eiji Saitoh, and Bart J. van Wees, “Spin caloritronics,” *Nat Mater* **11**, 391 (2012).
- [26] K. Uchida, J. Xiao, H. Adachi, J. Ohe, S. Takahashi, J. Ieda, T. Ota, Y. Kajiwara, H. Umezawa, H. Kawai, G. E. W. Bauer, S. Maekawa, and E. Saitoh, “Spin seebeck insulator,” *Nat Mater* **9**, 894–897 (2010).
- [27] A.I. Akhiezer, V.G. Bar’iakhhtar, and S.V. Peletminski, *Spin waves* (North-Holland Publishing Company, Amsterdam, 1968).
- [28] Andrii V. Chumak, Alexander A. Serga, and Burkard Hillebrands, “Magnon transistor for all-magnon data processing,” *Nature Communications* **5**, 4700 (2014).
- [29] Kathrin Ganzhorn, Stefan Klingler, Tobias Wimmer, Stephan Geprgs, Rudolf Gross, Hans Huebl, and Sebastian T. B. Goennenwein, “Magnon-based logic in a multi-terminal yig/pt nanostructure,” *Applied Physics Letters* **109**, 022405 (2016), <https://doi.org/10.1063/1.4958893>.
- [30] T. An, V. I. Vasyuchka, K. Uchida, A. V. Chumak, K. Yamaguchi, K. Harii, J. Ohe, M. B. Jungfleisch, Y. Kajiwara, H. Adachi, B. Hillebrands, S. Maekawa, and E. Saitoh, “Unidirectional spin-wave heat conveyer,” *Nature Materials* **12**, 549 (2013).
- [31] E. Saitoh, M. Ueda, H. Miyajima, and G. Tatara, “Conversion of spin current into charge current at room temperature: Inverse spin-hall effect,” *Applied Physics Letters* **88**, 182509 (2006).
- [32] E.B. Sonin, “Spin currents and spin superfluidity,” *Advances in Physics* **59**, 181–255 (2010),

- <http://dx.doi.org/10.1080/00018731003739943>.
- [33] So Takei, Bertrand I. Halperin, Amir Yacoby, and Yaroslav Tserkovnyak, “Superfluid spin transport through antiferromagnetic insulators,” *Phys. Rev. B* **90**, 094408 (2014).
- [34] Akashdeep Kamra and Wolfgang Belzig, “Superpoissonian shot noise of squeezed-magnon mediated spin transport,” *Phys. Rev. Lett.* **116**, 146601 (2016).
- [35] Akashdeep Kamra, Utkarsh Agrawal, and Wolfgang Belzig, “Noninteger-spin magnonic excitations in untextured magnets,” *Phys. Rev. B* **96**, 020411 (2017).
- [36] Charles Kittel, “Physical theory of ferromagnetic domains,” *Rev. Mod. Phys.* **21**, 541–583 (1949).
- [37] C. Kittel, “Interaction of spin waves and ultrasonic waves in ferromagnetic crystals,” *Phys. Rev.* **110**, 836–841 (1958).
- [38] K. Uchida, H. Adachi, T. An, T. Ota, M. Toda, B. Hillebrands, S. Maekawa, and E. Saitoh, “Long-range spin seebeck effect and acoustic spinpumping,” *Nat Mater* **10**, 737 (2011).
- [39] Akashdeep Kamra, Hedyeh Keshtgar, Peng Yan, and Gerrit E. W. Bauer, “Coherent elastic excitation of spin waves,” *Phys. Rev. B* **91**, 104409 (2015).
- [40] Akashdeep Kamra and Gerrit E.W. Bauer, “Actuation, propagation, and detection of transverse magnetoelastic waves in ferromagnets,” *Solid State Communications* **198**, 35 – 39 (2014), SI: Spin Mechanics.
- [41] M. Weiler, H. Huebl, F. S. Goerg, F. D. Czeschka, R. Gross, and S. T. B. Goennenwein, “Spin pumping with coherent elastic waves,” *Phys. Rev. Lett.* **108**, 176601 (2012).
- [42] L. Dreher, M. Weiler, M. Pernpointner, H. Huebl, R. Gross, M. S. Brandt, and S. T. B. Goennenwein, “Surface acoustic wave driven ferromagnetic resonance in nickel thin films: Theory and experiment,” *Phys. Rev. B* **86**, 134415 (2012).
- [43] Benedetta Flebus, Ka Shen, Takashi Kikkawa, Ken-ichi Uchida, Zhiyong Qiu, Eiji Saitoh, Rembert A. Duine, and Gerrit E. W. Bauer, “Magnon-polaron transport in magnetic insulators,” *Phys. Rev. B* **95**, 144420 (2017).
- [44] Takashi Kikkawa, Ka Shen, Benedetta Flebus, Rembert A. Duine, Ken-ichi Uchida, Zhiyong Qiu, Gerrit E. W. Bauer, and Eiji Saitoh, “Magnon polarons in the spin seebeck effect,” *Phys. Rev. Lett.* **117**, 207203 (2016).
- [45] Dmytro A. Bozhko, Peter Clausen, Gennadii A. Melkov, Victor S. L’vov, Anna Pomyalov, Vitaliy I. Vasyuchka, Andrii V. Chumak, Burkard Hillebrands, and Alexander A. Serga, “Bottleneck accumulation of hybrid magnetoelastic bosons,” *Phys. Rev. Lett.* **118**, 237201 (2017).
- [46] Andreas Rückriegel, Peter Kopietz, Dmytro A. Bozhko, Alexander A. Serga, and Burkard Hillebrands, “Magnetoelastic modes and lifetime of magnons in thin yttrium iron garnet films,” *Phys. Rev. B* **89**, 184413 (2014).
- [47] Nianbei Li, Jie Ren, Lei Wang, Gang Zhang, Peter Hänggi, and Baowen Li, “Colloquium: Phononics: Manipulating heat flow with electronic analogs and beyond,” *Rev. Mod. Phys.* **84**, 1045–1066 (2012).
- [48] Martin Maldovan, “Sound and heat revolutions in phononics,” *Nature* **503**, 209 (2013).
- [49] Yizhou Liu, Yong Xu, Shou-Cheng Zhang, and Wenhui Duan, “Model for topological phononics and phonon diode,” *Phys. Rev. B* **96**, 064106 (2017).
- [50] The demagnetization energy is disregarded since it only causes minor shifts in the dispersion [35].
- [51] Marlan O. Scully and M. Suhail Zubairy, *Quantum optics* (Cambridge University Press, Cambridge, 1997).
- [52] Kouki Nakata, Jelena Klinovaja, and Daniel Loss, “Magnonic quantum hall effect and wiedemann-franz law,” *Phys. Rev. B* **95**, 125429 (2017).
- [53] See supplemental material.
- [54] C. Kittel, *Quantum Theory of Solids* (John Wiley & Sons, 1987).
- [55] G.S. Kino, *Acoustic Waves: Devices, Imaging, and Analog Signal Processing*, Prentice-Hall Contemporary Topics in Accounting Series (Prentice-Hall, 1987).
- [56] A. A. Maradudin and G. I Stegeman, “Surface acoustic waves,” in *Surface phonons*, edited by F. W. de Wette W. Kress (Springer-Verlag, Berlin-Heidelberg, 1991) Chap. 2, pp. 5–36.
- [57] B. Andrei Bernevig and Taylor L. Hughes, *Topological insulators and topological superconductors* (Princeton University Press).
- [58] Pascal Ruello and Vitaliy E Gusev, “Physical mechanisms of coherent acoustic phonons generation by ultrafast laser action,” *Ultrasonics* **56**, 21–35 (2015).
- [59] J. Liu, L. J. Cornelissen, J. Shan, T. Kuschel, and B. J. van Wees, “Magnon planar hall effect and anisotropic magnetoresistance in a magnetic insulator,” *Phys. Rev. B* **95**, 140402 (2017).
- [60] The inset of Fig. 3(b) shows two phonon modes. One is irrelevant since it is localized on the opposite edge.
- [61] P. G. Gowtham, T. Moriyama, D. C. Ralph, and R. A. Buhrman, “Traveling surface spin-wave resonance spectroscopy using surface acoustic waves,” *Journal of Applied Physics* **118**, 233910 (2015).
- [62] S. Datta, *Surface acoustic wave devices* (Prentice-Hall, 1986).
- [63] Alexander V. Mamishev, Kishore Sundara-Rajan, Fumin Yang, Yanqing Du, and Markus Zahn, “Interdigital sensors and transducers,” *Proceedings of the IEEE* **92**, 808 (2004).