

1 **Article: Discoveries**

2 **Rapid Parallel Adaptation to Anthropogenic Heavy Metal Pollution**

3 Alexander S.T. Papadopoulos^{1,2*}, Andrew J. Helmstetter^{2,3}, Owen G. Osborne¹, Aaron A. Comeault¹, Daniel P.
4 Wood¹, Edward A. Straw^{2,4}, Laurence Mason², Michael F. Fay^{2,5}, Joe Parker^{2,6}, Luke T. Dunning⁷, Andrew D.
5 Foote^{1,8}, Rhian J. Smith² & Jackie Lighten⁹

6

7 ¹Molecular Ecology and Evolution Bangor, Environment Centre Wales, School of Natural Sciences, Bangor
8 University, Bangor, LL57 2UW, UK

9 ²Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3AB, UK

10 ³FRB-CESAB, Institut Bouisson Bertrand, Rue de l'École de Médecine, 34000 Montpellier

11 ⁴Centre for Ecology, Evolution & Behaviour, Department of Biological Sciences, School for Life Sciences and
12 the Environment, Royal Holloway University of London, Egham, Surrey, TW20 0EX, UK

13 ⁵School of Plant Biology, University of Western Australia, Crawley, WA 6009, Australia

14 ⁶National Biofilms Innovation Centre, Department of Biological Sciences, University of Southampton, SO14
15 1BJ, UK

16 ⁷Department of Animal and Plant Sciences, University of Sheffield, Sheffield, UK

17 ⁸Department of Natural History, Norwegian University of Science and Technology, Trondheim, Norway

18 ⁹Biosciences, University of Exeter, Stocker Road, Exeter EX4 4PY, UK

19 *Correspondence to: a.papadopoulos@bangor.ac.uk

20

21

22

23

1 **Abstract**

2 The impact of human-mediated environmental change on the evolutionary trajectories of wild organisms is
3 poorly understood. In particular, species' capacities to adapt rapidly (in hundreds of generations or less),
4 reproducibly and predictably to extreme environmental change is unclear. *Silene uniflora* is predominantly a
5 coastal species, but it has also colonised isolated, disused mines with phytotoxic, zinc-contaminated soils. To
6 test whether rapid, parallel adaptation to anthropogenic pollution has taken place, we used reduced
7 representation sequencing (ddRAD) to reconstruct the evolutionary history of geographically proximate mine
8 and coastal population pairs and found largely independent colonisation of mines from different coastal sites.
9 Furthermore, our results show that parallel evolution of zinc tolerance has occurred without gene flow
10 spreading adaptive alleles between mine populations. In genomic regions where signatures of selection were
11 detected across multiple mine-coast pairs, we identified genes with functions linked to physiological
12 differences between the putative ecotypes, although genetic differentiation at specific loci is only partially
13 shared between mine populations. Our results are consistent with a complex, polygenic genetic architecture
14 underpinning rapid adaptation. This shows that even under a scenario of strong selection and rapid adaptation,
15 evolutionary responses to human activities (and other environmental challenges) may be idiosyncratic at the
16 genetic level and, therefore, difficult to predict from genomic data.

17 **Introduction**

18 Modification of the natural environment by humans has significant implications for biodiversity (Urban 2015;
19 Ceballos et al. 2017; Helmstetter et al. 2020). Rapid habitat loss or environmental change can drive species to
20 the brink of extinction, but also presents opportunities for adaptation and speciation (Johnson and Munshi-
21 South 2017; Otto 2018; Ravinet et al. 2018; Szulkin, M., Munshi-South, J., & Charmantier 2020). The ability
22 of species to adapt to human-modified landscapes or activities is a key determinant of their viability in the
23 Anthropocene (McNeilly and Bradshaw 1968; Antonovics and Bradshaw 1970; Wu and Bradshaw 1972;
24 Macnair 1979; Hof et al. 2016; Reid et al. 2016; Bosse et al. 2017). Thus, a key question in evolutionary
25 ecology is how repeatable and predictable adaptation is to human-altered habitats (Bay et al. 2018; Fitzpatrick
26 et al. 2018; Therkildsen et al. 2019; Van Etten et al. 2020; Santangelo et al. 2020). To demonstrate that local
27 adaptation has driven the evolution of distinct ecotypes, it is necessary to establish an association between
28 fitness differences of populations and specific habitats. However, we can investigate genomic processes that
29 might contribute to adaptation by examining the sequence-based signatures of selection associated with local
30 adaptation. This can be accomplished even when reduced representation sequencing methods are used (Lowry
31 et al. 2017). In such cases, examples of parallel colonization of habitats with novel selection pressures can
32 support the hypothesis that specific genetic loci underpin local adaptation (Rundle et al. 2000; Jones et al.
33 2012; Ravinet et al. 2016; Nosil et al. 2018). A genomic approach can also discriminate between single or
34 parallel origins of populations adapted to a specific habitat or selection pressure. Local gene flow between
35 differentiated populations can obscure the true evolutionary relationships between them and lead to false
36 inferences (Ravinet et al. 2016; James et al. 2020). Promising cases of rapid parallel adaptation do exist (e.g.,

1 Lescak et al. 2015; Marques et al. 2016; Alves et al. 2019), but few have ruled out the possibility of local gene
2 flow creating the false impression of independent origins (Roda et al. 2013; James et al. 2020).

3 Instances where the same toxic chemicals and contaminants have been repeatedly introduced into the
4 environment by humans in isolated locations can generate novel selection regimes that have the potential to
5 promote parallel adaptation. Strong selection, caused by herbicides, pesticides and heavy metals that
6 contaminate soils and water bodies, is capable of producing extremely rapid adaptive responses (Antonovics
7 and Bradshaw 1970; Wu and Bradshaw 1972; Macnair 1979; Hartley et al. 2006; Van Etten et al. 2020) and
8 trade-offs (Xie and Klerks 2004), and may be particularly prone to triggering parallel responses as a result
9 (MacPherson and Nuismer 2017). Indeed, there is evidence for rapid parallel adaptation from ‘ancient’
10 standing genetic variation during adaptation to copper mine contamination in two populations of *Mimulus*
11 *guttatus* (Wright et al. 2015; Lee and Coop 2017). In the Atlantic killifish, *Fundulus heteroclitus*, tolerance to
12 marine pollution has evolved in four populations (Reid et al. 2016). The mutations underlying this resistance
13 have evolved on at least two occasions, but migration between three of the four populations may have
14 contributed to the spread of tolerance (Lee and Coop 2017). Convergent herbicide resistance across species is
15 well documented, but there is more limited support for parallel origins within single species and the spread of
16 resistance by gene flow has been harder to rule out (Kreiner et al. 2019; Van Etten et al. 2020).

17 Here, we present evidence for multiple recent and independent origins of heavy metal tolerance in the
18 predominantly coastal plant *Silene uniflora* (sea campion). In Great Britain and Ireland, metal mining activities
19 had largely ceased by the early 20th century, but the legacy of spoil heaps and soils contaminated with heavy
20 metals forms a patchwork of highly localised and drastically altered environments across the landscape (Baker
21 et al. 2010). Heavy metals, such as zinc, copper, cadmium and lead, are highly toxic to plants, triggering
22 oxidative stress, inhibition of growth and photosynthesis, and death (Küpper and Andresen 2016). As a result,
23 many of these abandoned sites remain barren for hundreds of years after the mining itself has ceased (Baker
24 1974; Baker et al. 2010). Despite its largely linear coastal distribution, *S. uniflora* has managed to colonise a
25 number of isolated inland mine spoils in various regions of the UK and Ireland – although only a small
26 proportion of the >10,800 non-ferrous mines in Great Britain harbour the species (Baker 1974; Baker 1978;
27 Baker and Dalby 1980; Gill 2018). A common feature of the mines that it inhabits is an elevated level of zinc.
28 Experiments in the 1970s demonstrated that: (i) mine populations are more zinc tolerant than coastal
29 populations; (ii) mine plants exclude zinc from their shoots and (iii) zinc tolerance in each population is tightly
30 correlated with the concentration of zinc found in local soils (Baker 1978). Furthermore, in a common garden
31 experiment using zinc-enriched slag from a population in Morrision in Swansea, Baker (1974) demonstrated
32 that the local mine plants grew and produced flowers normally, while coastal plants remained in a dwarfed
33 state, developed chlorosis (yellowing due to lack of chlorophyll) and did not produce any flowers - even in
34 slag that had been heavily diluted with sandy soil. The link between the zinc tolerance phenotype, local levels
35 of environmental zinc, and reduced fitness of coastal plants in zinc-contaminated soils suggests that mine
36 populations are locally adapted to their environment.

1 Given the generally coastal distribution and the isolated nature of the colonised mines, we hypothesised that
2 the mine populations have independently adapted from the nearest coastal populations. Across four local mine-
3 coast population pairs, we used growth experiments to determine whether mine plants are more tolerant to zinc
4 toxicity than their nearest coastal counterparts. We combined a newly sequenced draft genome with reduced
5 representation genotypes for 216 individuals, conducting population genetic analyses to establish the
6 relationships between the populations and test the hypothesis that the mine populations had evolved
7 independently multiple times, following dispersal from their physically closest coastal populations. Finally,
8 we used these data to explore the extent to which evolution of the mine populations is controlled by a
9 parallel/convergent molecular basis.

11 Results and discussion

12 **Anthropogenic adaptation to heavy metal contamination.** Populations of *S. uniflora* were sampled from
13 four derelict mines and the nearest coastal population to each across the UK and Ireland (Fig 1A). Previous
14 research has shown that the contaminated mine sites all have elevated, toxic levels of zinc in the soil (2,410-
15 48,075ppm, Table S1) relative to typical coastal and inland sites (UK mean = 81.3ppm; Ross et al. 2007). Lead
16 levels were also high at all mine sites (>10,000ppm, Table S1; UK mean = 52.6; Ross et al. 2007), but only
17 the South Wales (SWA-M) and Irish (IRE-M) mines were heavily contaminated with copper (>10,000 ppm,
18 Table S1; UK mean = 20.6; Ross et al. 2007). We used root elongation experiments with wild collected seed
19 to determine whether mine populations were more tolerant of zinc and copper than the most geographically
20 proximate coastal population. In all cases, mine populations were significantly more zinc tolerant than the local
21 coastal population (Welch's t-test, two-sided, $p < 0.005$ for all four pairs; Fig 1B). Deep water culture
22 experiments with cuttings from individuals grown in standard conditions also confirmed that plants from mine
23 populations were more zinc tolerant than coastal populations: i.e., root growth continued in mine plants at
24 600 μ M ZnSO₄, but not in coastal plants (see Methods). However, only the Irish mine population was
25 significantly more copper tolerant than the respective local coastal population (Welch's t-test, two-sided, p
26 < 0.001 , Fig 1C). The lack of clear copper tolerance in SWA-M may be due to the relatively high copper
27 concentration used in the experiment, possibly beyond levels that can be tolerated by this population. It is
28 notable that both mine and coastal populations from Wales were more copper tolerant than the English
29 populations (Fig 1C), suggesting that SWA-M may be able to cope with high copper levels due to constitutive
30 copper tolerance in Welsh *S. uniflora*. High intraspecific variation in copper tolerance has been observed in
31 other species - even within a single mine (e.g., *Scopelophila cataractae*) - as has constitutive tolerance in non-
32 mine specialists (e.g., *Ceratodon purpureus*; Boquete et al. 2021). Overall, these results corroborate earlier
33 findings of zinc and copper tolerance in mine populations of *S. uniflora* (Baker 1978).

34 Although our experiments do not provide a direct measure of fitness in the wild, given the association between
35 zinc tolerance, levels of zinc contamination in soil, vegetative growth and flower production in *S. uniflora*
36 (Baker 1974; Baker 1978), our results indicate that all of the sampled mine populations are adapted to zinc
37 contamination. Due to the strong selection that heavy metal toxicity exerts, tolerance can evolve in plants

1 within as little as a single generation if there is sufficient genetic variation (Wu and Bradshaw 1972). Although
2 limited mining activity existed at some of these sites as far back as the bronze age, the most intensive working
3 took place between the 18th and 19th centuries (see Methods) and so it is likely that these anthropogenic mine
4 habitats only became available for colonisation once active excavation ceased at mining sites within the last
5 250 years (Baker 1974). Therefore, populations of zinc-tolerant *S. uniflora* studied here are likely to have
6 evolved since the 18th century (i.e., < 250 generations).

7 **Independent, parallel origins of the mine populations.** In total, 216 individuals (n per population; WWA-
8 M = 25, WWA-C = 28, SWA-M = 28, SWA-C = 27, ENG-M = 26, ENG-C = 27, IRE-M = 28, IRE-C = 27)
9 were genotyped at 74,064 SNPs. On average “local” mine and coastal populations were 20.8km apart (WWA
10 = 16.1km, SWA = 14.8km, ENG = 25.6km, IRE = 26.8km). Genetic differentiation between populations was
11 high (mean F_{ST} = 0.36; Table S2), reflecting the relatively poor dispersal capabilities and fragmented
12 distribution of the species (Baker 1974; Runyeon and Prentice 1997). Differentiation was substantially higher
13 between mine populations (mean F_{ST} = 0.45) than between coastal populations (mean F_{ST} = 0.25). Mine
14 populations were also substantially differentiated from their local coastal population (mean F_{ST} = 0.36),
15 suggestive of very limited geneflow between differentially adapted populations at the local level. In support
16 of this, analysis of molecular variance (AMOVA; Table S3) shows that most of the variation is partitioned
17 within and among individuals (~65%), but a large proportion of variation was among populations which were
18 grouped by either habitat (34%) or region (33%). Partitioning of genetic variation was low between habitats
19 (1.5%) and fractionally larger between regions (2.0%), reflecting the very high differentiation between mines
20 and greater degree of shared variation between local mine and coastal populations. Genetic diversity (π) was
21 also significantly higher in the coastal populations versus the mine populations (0.065 and 0.044, respectively;
22 Welch’s t-test, two-sided, $p < 0.036$, Table S4). Tajima’s D was slightly positive across all populations (mean
23 = 0.24, Table S4), but not significantly different between the mine and coastal populations. As Tajima’s D is
24 close to zero, the drop in diversity is unlikely to result from a population bottleneck, but this pattern matches
25 expectations for multiple soft selective sweeps taking place across the genome (Pennings and Hermisson 2006)
26 - as might be expected when colonising a new environment in the face of a strong selection pressure with
27 limited time for new adaptive mutations to evolve.

28 In the context of recent colonisation, relatively high differentiation and limited gene flow between populations,
29 we predicted that different colonisation scenarios would produce differing patterns of isolation by distance
30 among mine versus coastal habitats (IBD; Wright 1943; James et al. 2020) - specifically that a scenario of
31 independent origins of the mine populations would be distinguishable from a single origin. In a multiple origin
32 scenario, IBD among mine populations should be accentuated relative to the pattern across coastal populations,
33 whereas, in a single origin scenario, IBD among mine populations should be minimal. To test these predictions,
34 we conducted forward-in-time simulations in SLiM v3 (Haller and Messer 2019) and estimated within-habitat
35 IBD under ‘multiple-origin’ and ‘single-origin’ colonization scenarios (Fig 2A & B, See Methods). As
36 expected, the strength of IBD was significantly higher among the mine populations than among the coastal
37 populations for the multiple origin scenario (paired t-test, two-sided, $p < 0.001$; Fig 2A) and the reverse was

1 true for the single origin scenario (paired t-test, two-sided, $p < 0.001$; Fig 2B). The observed IBD in the sampled
2 populations (Fig 2C) closely matches the expectations for a parallel origin of mine populations, supporting the
3 hypothesis that the mine habitat has been colonised independently.

4 Phylogenetic reconstruction of evolutionary relationships between the *S. uniflora* populations based on 7,037
5 linkage disequilibrium pruned SNPs (Fig 3A) and principal component analysis (PCA) of genetic structure
6 from the full set of 74,064 genome aligned SNPs (Fig 3B), clearly indicate three independent origins of zinc-
7 tolerant mine populations; one in Ireland, one in England and one in Wales. The PCA highlights the much
8 higher genetic similarity between coastal populations than between mine populations, which occupy extremely
9 divergent areas of genotype space, suggesting that they may be on different evolutionary trajectories at the
10 genetic level, despite adapting to similar selection pressures. The two Welsh mines are genetically similar (Fig
11 3B and S1) and although we cannot rule out independent origins from unsampled non-tolerant populations, it
12 is likely that transport of workers, machinery or ore between Welsh mines dispersed zinc tolerant plants
13 between sites. In fact, records of mine ownership from 1758 indicate that human-mediated dispersal is possible
14 between West Wales and Swansea and it was common practice to transport ore mined elsewhere to be refined
15 in Swansea (Hughes 2000). There are at least 14 further records of *S. uniflora* growing on contaminated mine
16 spoil in the UK and Ireland (pers. obs. & Baker 1974), so our discovery of three independent origins is likely
17 to be a lower bound on the true number of independent origins for zinc-tolerant populations.

18 Three origins of zinc tolerant populations were further supported when modelling shared genetic drift among
19 populations (Treemix analysis; Fig 3C). This analysis also provided evidence of migration between the Welsh
20 coastal populations (WWA-C and SWA-C) and very weak migration between the Irish, English and Welsh
21 populations. To assess the significance of admixture in the evolution of the mine populations, we examined
22 genetic relationships across all population quartets using the less-parameterised f_4 statistics (Fig 4). The f_4
23 statistic quantifies shared drift between pairs of populations in a four-taxon tree - significant deviation of the
24 f_4 statistic from zero for the tested topology demonstrates that the relationships are not perfectly described by
25 a bifurcating tree. This is indicative of some shared drift between populations that conflicts with the topology,
26 for example due to admixture (Reich et al. 2009; Foote and Morin 2016; Peter 2016; Lipson 2020). The f_4
27 statistic for the tree containing all four mine populations (type 2; Fig 4) indicates that there has been no
28 admixture between mines (i.e., f_4 does not deviate from zero; $f_4 = 1.31 \times 10^{-5}$, s.d. = 7.75×10^{-5} , $p = 1.00$),
29 whereas f_4 for the coastal population quartet (type 1; Fig 4) demonstrates that admixture between coastal
30 populations has taken place (i.e., f_4 is significantly different from zero; $f_4 = -3.95 \times 10^{-4}$, s.d. = 7.57×10^{-5} , p
31 = 3.68×10^{-5}). Comparisons of quartets with three mine populations and one coastal population (type 4; Fig 4)
32 provide an additional test of the independent origins of the mine populations, in each case demonstrating that
33 there was no correlated drift between the mine outgroups and the mine-coast pair of more closely related
34 populations. On the other hand, the *three coastal : one mine* comparisons (type 3; Fig 4) provide further
35 confirmation of gene flow from coastal outgroups into more closely related mine-coast pairs in three quartets
36 and support the significance of migration edges between SWA-C and WWA-C, and IRE-C and ENG-C.

1 Overall, our results provide firm support for recent parallel evolution of mine populations, with migration
2 restricted to coastal sites.

3 **Evidence for molecular convergence/parallelism.** To investigate the genetic basis of mine-coast
4 differentiation and degree of molecular convergence in adaptation, we conducted pairwise F_{ST} -based genome
5 scans for each mine-coast pair and identified outlier loci potentially under divergent selection. Due to the
6 relatively sparse sampling of our ddRAD dataset and the highly fragmented draft genome (Table S5; N50 =
7 4,660bp, length = 0.77Gb), we designated genomic scaffolds containing at least one outlier SNP as an outlier
8 scaffold for each comparison (the number of outlier SNPs was not significantly associated with scaffold length;
9 Tukey's test Fig S2). Across the local mine-coast pairs, the number of outlier scaffolds ranged from 779-1,216
10 and the number of outlier SNPs varied from 1,346-2,261 - the degree of overlap between all sets of outlier
11 scaffolds (Fig 5A) and SNPs (Fig 5B) was significantly higher than expected by chance as assessed by Super
12 Exact Test (an extension of Fisher's Exact Test for multiple sets; Wang et al. 2015). In total, 34 scaffolds were
13 identified as outliers across all pairwise comparisons, while 187 and 756 outlier scaffolds were found across
14 the sets of three and two comparisons, respectively (Fig 5A). There was substantially less overlap at the level
15 of SNPs (Fig 5B), with four shared across all four sets, 85 shared by three sets and 870 shared by two sets.
16 This pattern suggests a highly polygenic basis to mine-coast differentiation, with a substantial proportion of
17 shared targets of selection found in three or fewer pairs. However, we are unable to rule out the possibility that
18 the shared scaffolds are physically close to each other in the genome, although linkage disequilibrium between
19 the scaffolds is low (mean $r^2 = 0.021$).

20 It is currently unclear whether the adaptive variation that underpins tolerance and colonisation of the mine
21 habitat has arisen through new independent mutations in each population (as in *F. heteroclitus*; Reid et al.
22 2016), has been drawn from standing variation (as in *M. guttatus*; Lee and Coop 2017), or has been obtained
23 through adaptive introgression from close relatives (as in *Fundulus grandis*; Oziolor et al. 2019). Despite this
24 limitation, the lack of parallelism at the SNP level provides some indication that introgression is unlikely to be
25 the source of adaptive alleles. Dramatically greater overlap between the two Welsh comparisons (WWA and
26 SWA) and a bias towards shared outlier SNPs rather than scaffolds, further supports the single origin of the
27 Welsh mine populations and provides a clear contrast with the degree of outlier overlap with mine populations
28 that evolved in other regions. It is possible that the difference in distribution of overlap between scaffolds and
29 SNPs is due to a limited role of parallelism at the level of individual nucleotides, but greater convergence at
30 the genic level (Conte et al. 2012). However, the sparse sampling inherent to the ddRAD approach may mean
31 that the specific adaptive sites are not captured in the analysis (Lowry et al. 2017) and there may be more
32 substantial sharing and parallelism of adaptive SNPs across independently derived mine populations.

33 A polygenic basis to differentiation in *S. uniflora* is at odds with previous investigations of heavy metal
34 tolerance in *Silene*. Using controlled crosses and hydroponic experiments, these studies indicated that both
35 zinc and copper tolerance have relatively simple genetic bases and are not controlled by the same molecular
36 mechanisms (Schat et al. 1996; Schat and Vooijs 1997). The simple architecture for copper tolerance in *S.*
37 *vulgaris* is also supported by the recent discovery of two related ATPase copper transporters which additively

1 contribute to copper tolerance (Li et al. 2017). The potential for polygenic convergence in *S. uniflora* is further
2 supported by gene ontology enrichment analysis of the subset of genes found on the 34 scaffolds which were
3 outliers in all 4 pairwise comparisons. This group was significantly enriched for genes involved in metabolism
4 of reactive oxygen species and the regulation of salicylic acid (Table S6), which are critical in responses to
5 cold, salt, drought and heavy metal stresses (Khan et al. 2015). Further systematic investigation of gene
6 functions revealed that 15 genes have well-supported roles in processes that are relevant to differentiation
7 between coastal and mine plants: eight associated with salt stress, eight with heavy metal stress and four with
8 root development and morphology (Table S7). This points to a potential trade-off in the molecular processes
9 which govern mine-coast differentiation, with selection against salt tolerance alleles in mines and against metal
10 tolerance alleles in coastal environments. Alternatively, some alleles for genes that contribute to metal
11 tolerance may be conditionally neutral in coastal plants and under positive selection in the mine environment.
12 In this latter scenario, we might expect a higher incidence of metal tolerance among coastal population, but
13 further work is needed to establish which model underlies local adaptation.

14 The exact mechanism of zinc tolerance in *Silene* is not well understood. However, hydroponic experiments
15 with mine and coastal *S. uniflora* demonstrated that mine plants grown in zinc-contaminated media accumulate
16 a higher proportion of absorbed zinc in the roots relative to their shoots whereas the reverse is true for coastal
17 plants (Baker 1978). Additional research in *S. vulgaris* indicates that zinc uptake into tonoplast vesicles of
18 zinc-tolerant *S. vulgaris* is higher than in non-tolerant plants (Chardonens et al. 1999). In our study, three
19 genes on outlier scaffolds (*PSD2*, *WRKY23* and *RWPI*) have direct links to these physiological differences
20 between tolerant and non-tolerant *Silene*: (i) *PSD2* encodes a form of phosphatidylserine decarboxylase which
21 is located in the tonoplast (Nerlich et al. 2007), confers cadmium tolerance in *Saccharomyces cerevisiae*
22 (Gulshan et al. 2009) and produces phosphatidylethanolamine, which is involved in zinc homeostasis in
23 *Pseudomonas fluorescens* (Appanna et al. 1995); (ii) *WRKY23* is a transcription factor that regulates root
24 development by altering auxin distribution through the control of flavanol biosynthesis in *Arabidopsis thaliana*
25 - overexpression of *WRKY23* increases quercetin root concentrations (Grunewald et al. 2012). Quercetin is a
26 very efficient chelator of heavy metals (i.e., a molecule that binds metal ions) and supplementation of wild
27 type *A. thaliana* with quercetin stimulates root growth in the presence of zinc ions (Keilig and Ludwig-Müller
28 2009); and (iii) *RWPI* is required for the production of the cell wall polymer suberin. In *A. thaliana*, *RWPI*
29 mutants lack suberin and have increased root permeability for NaCl (Gou et al. 2009). Furthermore, *Esb1*
30 mutants have increased levels of root suberin, which both decreases accumulation of cadmium, manganese
31 and zinc in the shoots and increases accumulation of sodium in the shoots (Baxter et al. 2009).

32 Parallel evolution is expected to be facilitated in spatially structured environments when loci have large,
33 spatially antagonistic fitness effects (Chevin et al. 2010). Evidence of such trade-offs in wild plants is lacking,
34 with loci displaying conditional neutrality more commonly detected (Lowry et al. 2009; Hall et al. 2010;
35 Anderson et al. 2011). The dual effect of high suberin levels on restriction of zinc ions to the roots and exposure
36 of the shoots to sodium raises the possibility of a direct trade-off in suberin production and opens the possibility
37 of antagonistic pleiotropy at *RWPI* influencing the parallel evolution of zinc tolerant populations. Of the three

1 genes, only the scaffold containing *RWPI* had consistently lower genetic diversity in the mine populations
2 (paired t-test, two-sided, $p = 0.030$), whereas for *WRKY23* and *PSD2* diversity was only lower in the mines
3 from West Wales and Ireland (Table S4). These findings further support the polygenic nature of parallel
4 adaptation in *S. uniflora* and the potential importance of antagonistic pleiotropy in the rapid evolution of
5 differentially adapted populations.

6 In a rapidly changing world, the adaptability of species will be critical for their long-term persistence. This
7 study shows that some species will be capable of responding quickly to strong selection pressures across their
8 range. We argue that plant species with sufficient genetic variation may adapt quickly to a single physiological
9 stress repeatedly in different places, while using subtly different genetic mechanisms. As in *S. uniflora*, those
10 species that evolved to survive in environments with natural sources of high abiotic stress, but which do not
11 compete well in low-abiotic stress/high-biotic competition environments, may be particularly well suited to
12 cope with the ongoing human modification of the planet. Alongside evidence of widespread local adaptation
13 to different environmental conditions in other species (Fournier-Level et al. 2011; Papadopoulos et al. 2014),
14 our findings indicate that while it may be possible to predict which species will adapt to specific environments,
15 the underlying genetic basis to that adaptation may be considerably more variable than is currently understood
16 from the limited number of well-studied examples (Bay et al. 2018; Fitzpatrick et al. 2018; Oomen et al. 2020).
17 In order to be accurate, predictions of evolutionary responses to environmental change from genomic data will
18 need to account for the possibility that multiple genetic architectures can produce similar phenotypic responses.

19

20 **Materials and Methods**

21 **Sample collection:** Four focal mine sites where *S. uniflora* was known to occur were selected for sampling;
22 Grogwynion (West Wales; WWA; worked 1588 – 1889 C.E.; Hartley 2009), White Rock (Swansea, South
23 Wales; SWA; 1736 – 1928; Hughes 2000), Priddy Pools (Somerset, South-West England; ENG; 1850 – 1908,
24 evidence of Roman mining; Gough 1967) and Ross Island (Co. Kerry, South-West Ireland; IRE; 1707-1829,
25 evidence of Bronze Age mining; O’Brien 2020). For three of these sites (WWA, ENG and IRE), metal
26 tolerance has previously been tested (Baker 1978; Schat et al. 1996). White Rock was also located near a
27 previously tested population in Morryston, Swansea (Baker 1978) that no longer exists. The BSBI Database
28 was used to identify the nearest accessible coastal populations to each mine. See Table S8 for population co-
29 ordinates. At each of the eight populations, leaf tissue was sampled from 30 individuals and preserved for
30 DNA extraction in fine mesh silica gel. Individuals were sampled at least one metre apart and samples were
31 collected at even intervals across the extent of each population. At each site, we collected seeds from a
32 minimum of twelve individuals, which were then dried and stored separately with silica gel. For assembly of
33 a draft genome, cuttings from a single coastal individual were collected in Tresaith (West Wales), propagated
34 and self-fertilised to produce an inbred F1 (SUTF1P) with reduced heterozygosity.

35 **Phenotyping:** Root elongation experiments were conducted to determine the level of zinc and copper tolerance
36 in each population (Baker 1978). Seeds were germinated in groups of eight (one seed per population) on $\frac{1}{4}x$

1 Murashige-Skoog media in 1% Agar with no supplemental heavy metals (control treatment), 24 μ M copper
2 sulphate (copper treatment) or 459 μ M zinc sulphate (zinc treatment). Twenty graduated plates were prepared
3 per treatment and the positions of populations within plates was determined using a random seed. Plates were
4 placed upright in a germination cabinet with a 12-hour light/dark cycle for 10 days and then photographed
5 using a digital camera. Radicle length of all seedlings with emerged cotyledons was measured using ImageJ
6 v1.8.0. Zinc and copper tolerance were calculated as the radicle length in the treatment divided by the mean
7 length in the control for each population. Six individuals per population germinated on control media were
8 grown into adults and zinc tolerance was assessed using deep water culture. To do this, cuttings from each
9 individual were rooted in a mist propagator for two weeks before being transferred to a deep-water culture set
10 up with 1/10x Hoagland's solution. After acclimatisation for one week, the plant roots were stained using a
11 suspension of activated charcoal and rinsed with ddH₂O, the solution was refreshed and 600 μ M Zinc sulphate
12 was added. After a further two days root growth was inspected by eye - the presence of unstained root tips
13 (i.e., ongoing root growth) was taken as confirmation of zinc tolerance (Schat et al. 1996; Bratteler et al.
14 2006a).

15 **Genome assembly:** DNA was extracted from silica dried leaf tissue using Qiagen DNeasy Plant tissue kits.
16 DNA quality was assessed using agarose gel electrophoresis and DNA was quantified using a Promega
17 Quantus fluorometer with Quantifluor dsDNA kits. For draft genome assembly, four NEBnext Ultra II libraries
18 were prepared for SUTFP and each was sequenced using illumina MiSeq v3 600bp PE cartridges. Adapter
19 and quality trimming were performed using cutadapt v2.1 (Martin 2011) and Trimmomatic v0.36 (Bolger et
20 al. 2014) (minimum quality = 15, minimum length = 64). Overlapping read pairs were merged using Abyss-
21 mergepairs (Jackman et al. 2017) and non-overlapping pairs merged using konnector v2.0 (Vandervalk et al.
22 2015) with a bloom filter containing merged and unmerged reads for all libraries (kmer length=96, bloom filter
23 FPR = 1.01%). illumina reads were assembled into contigs using Abyss v2.0 (Jackman et al. 2017) with a kmer
24 length=241 – selected after estimation with kmergenie v1.7048 and Abyss runs with kmers = 96/127/151. To
25 scaffold the assembly, the same individual was sequenced using an Oxford Nanopore MinION (Three R9 flow
26 cells and one R9.4 flow cell with SQK-NSK007 kits). Nanopore reads were corrected with Proovread v2.12
27 (Hackl et al. 2014) using the processed illumina reads. Redundans v0.14a (Pryszcz and Gabaldón 2016) was
28 used to reduce contig redundancy caused by heterozygosity (minimum identity 95%) and scaffold contigs
29 using the corrected nanopore data. Abyss-sealer (Paulino et al. 2015) was used to fill gaps in the scaffolded
30 assembly (kmers = 94/89/84) and completeness was assessed with BUSCO v3 (Benchmarking Universal
31 Single-Copy Orthologs; complete and fragmented = 78.5%, Table S5). Augustus (Stanke et al. 2006) was used
32 to predict genes in the genomic scaffolds using the annotation training files from *Solanum lycopersicum*. The
33 resulting predicted amino acid sequences were BLASTp-searched (Camacho et al. 2009) against the
34 *Arabidopsis thaliana* proteome (Araport11) and only the best scoring hit from each predicted amino acid
35 sequence was retained.

36 **Genotyping:** Double-digest RAD sequencing was performed following a modified protocol of Peterson et al
37 (2012) detailed in Papadopoulos et al (2019) and restriction was performed using *EcoRI*-HF and *MspI*. For this

1 study, size selection was conducted with a pippin prep (468-546bp) and one pool of 230 uniquely barcoded
2 individuals was sequenced on five lanes of an illumina HiSeq 2500 (100bp, PE) at the Earlham Institute. Raw
3 reads were demultiplexed, trimmed to 90bp and low-quality reads were discarded, resulting in an average of
4 4.76M reads per sample (s.d. 2.01M). Reads were mapped to the draft genome using bowtie v2.3.4 (Langmead
5 and Salzberg 2012) in end-to-end mode and excluding reads with low mapping quality ($Q < 20$). SNPS were
6 called from the resulting BAM files using gstacks v2.0b (Rochette et al. 2019), 14 samples were excluded from
7 further analysis due to low coverage. Genotypes for SNPS with less than 20% missing data were extracted in
8 VCF and RADpainter format using Populations v2.0b (Rochette et al. 2019). In total, 216 individuals were
9 genotyped at 74,064 SNPs.

10 **Evolutionary genetics:** Population genetic structure across *S. uniflora* was assessed using principal
11 components analysis implemented in adegenet v2.1.3 (Jombart 2008) in R and genetic co-ancestry was
12 estimated using the haplotype-based inference method of fineRADstructure v0.3.2 (Malinsky et al. 2018).
13 Analysis of Molecular Variance (AMOVA) was conducted in Arlequin v3.5.2.2 (Excoffier and Lischer 2010)
14 To assess patterns of isolation by distance, pairwise genetic differentiation between the sampled populations
15 (Weir and Cockerham's F_{ST}) was calculated using Arlequin v3.5.2.2 (Excoffier and Lischer 2010), pairwise
16 geographic distances between populations were calculated with the distm function in the geosphere R package
17 and isolation by distance estimated in R using linear regression. Tajima's D was calculated for 20kb sliding
18 windows in VCFtools v0.1.16 (Danecek et al. 2011) and averaged over the subset of windows for which D
19 could be calculated in all populations. To identify the isolation by distance signature expected from parallel vs
20 single origins of the mine populations, we conducted simulations in SLiM v3.3.2 under two scenarios:
21 independent colonisation of mines from the nearest coastal population and non-independent colonisation of
22 mines from the same individual coastal population. In the latter case, the 'founding' coastal population was
23 randomly chosen in each independent iteration of the simulation. All simulations were initiated with a burn-in
24 period of 100,000 generations and a population size of 10,000 individuals. Each individual in the population
25 was diploid and hermaphroditic, and generations were non-overlapping (i.e. Wright-Fisher simulations). To
26 track genetic relationships among populations, we simulated a single chromosome that was 50,000 bp long
27 with a uniform mutation rate of 7.5×10^{-9} - based on estimates for *S. latifolia* (Krasovec et al. 2018) - and a
28 recombination rate of 4.0×10^{-9} - based on the genetic map length (446cM; Bratteler et al. 2006b) and genome
29 size (1.13Gb) of *S. vulgaris* (Pellicer and Leitch 2020). In the 100,000th generation, two populations (p1 and
30 p2) were colonised with 500 individuals each from the ancestral population. These two populations represented
31 those that initially colonised Ireland and the west coast of England/Wales at the end of the Last Glacial
32 Maximum. Subsequent stepwise colonisation of populations (i.e. p2 -> p3 -> p4), representing coastal
33 populations, occurred every 20 generations until there were four coastal populations in the 100,040th
34 generation. Coastal populations were always founded with 500 individuals and population sizes increased to
35 1,000 individuals ten generations after a population was initially founded. After colonisation, p1 and p2
36 exchanged migrants at a rate of 0.00001 per generation, p2 and p3 at a rate of 0.0001, and p3 and p4 at a rate
37 of 0.0001. P1 through p4 were therefore effectively arranged along a line and migration rates between non-
38 adjacent populations were equivalent to the product of migration rates connecting them. Ten-thousand

1 generations after the coastal populations were founded, 100 individuals were used to found each of four
2 populations meant to reflect those found in mine environments. After founding the mine populations, these
3 populations exchanged migrants with the nearest coastal population at a rate of 0.0002. All populations then
4 evolved for an additional 100 generations. At the end of the simulations (i.e. at generation 110,150), we
5 calculated and output F_{ST} between each of the four coastal populations (all pairwise comparisons) and each of
6 the four mine populations. We ran 100 independent replicates for each of the three colonisation scenarios
7 described above.

8 To further establish the evolutionary relationships between the populations, the dataset was pruned to 7,037
9 SNPs using a linkage disequilibrium threshold of 0.1 and minor allele frequency threshold of 0.05, and the
10 phylogenetic tree estimated with 1,000 bootstrap replicates using the maximum likelihood approach
11 implemented in SNPhylo v2 (Lee et al. 2014). This reduced dataset was then used to explore the possibility of
12 migration and introgression between the populations using Treemix v0.1.15 (Pickrell and Pritchard 2012). For
13 the maximum likelihood estimation of the tree in Treemix, one to ten migration edges were fitted and the
14 number of edges that explained 99.8% of the variance selected as the best model. Using the fourpop function
15 in Treemix, f_4 statistics (Reich et al. 2009) were calculated for all population quartets to assess whether
16 relationships between the populations deviated significantly (after Dunn-Bonferroni correction) from tree-
17 likeness. The premise of the f_4 statistic and our test is that for any four populations there are three possible
18 trees [((A,B),(C,D)); ((A,C),(B,D)); and ((A,D),(B,C))]. If ((A,B),(C,D)) is the correct tree, the allele frequency
19 difference between A and B will not be correlated with the frequency difference between C and D, i.e., the
20 correlation in frequency differences (f_4) would not deviate from zero (Reich et al. 2009). For each quartet of
21 populations in our sample, we determined the correct tree based on Fig 3A and tested whether f_4 significantly
22 deviated from zero using the z -score.

23 To investigate the level of parallel evolution at the molecular level, we calculated Weir and Cockerham's F_{ST}
24 at all variable sites in pairwise comparisons between the geographically proximate mine-coast pairs using
25 VCFtools v0.1.16. SNPs falling in the upper 95% percentile of values in each pairwise comparison were
26 designated as outlier loci and scaffolds containing one or more outlier SNPs were designated as outlier
27 scaffolds. Overlap of outlier SNPs and scaffolds was visualised using upsetR v1.4.0 (Conway et al. 2017) and
28 significance of overlap was assessed using SuperExactTest v1.0.7 (Wang et al. 2015). To investigate the
29 possible functions of genes in outlier regions, all genes on the outlier scaffolds that were in common across
30 the four pairwise mine-coast comparisons were subjected to gene ontology enrichment analysis performed in
31 topGo v3.11 (Alexa and Rahnenfuhrer 2020) using the "elim" algorithm and Fisher's Exact tests to assess
32 significance. Further assessments of gene functions were made from The *Arabidopsis* Information Resource
33 (TAIR) descriptions and associated references. Systematic searches were performed using gene names with
34 and without the terms "stress" and "heavy metal" using Google Scholar.

35

36 **References**

- 1 Bratteler M, Lexer C, Widmer A. 2006a. Genetic architecture of traits associated with serpentine adaptation
2 of *Silene vulgaris*. *J. Evol. Biol.* 19:1149–1156.
- 3 Bratteler M, Lexer C, Widmer A. 2006b. A genetic linkage map of *Silene vulgaris* based on AFLP markers.
4 *Genome* 49:320–327.
- 5 Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST plus:
6 architecture and applications. *BMC Bioinformatics* 10:1.
- 7 Ceballos G, Ehrlich PR, Dirzo R. 2017. Biological annihilation via the ongoing sixth mass extinction
8 signaled by vertebrate population losses and declines. *Proc. Natl. Acad. Sci. U. S. A* 114:6089–6096.
- 9 Chardonnens AN, Koevoets PLM, van Zanten A, Schat H, Verkleij J a. C. 1999. Properties of enhanced
10 tonoplast zinc transport in naturally selected zinc-tolerant *Silene vulgaris*. *Plant Physiol.* 120:779–786.
- 11 Chevin LM, Martin G, Lenormand T. 2010. Fisher’s model and the genomics of adaptation: Restricted
12 pleiotropy, heterogenous mutation, and parallel evolution. *Evolution.* 64:3213–3231.
- 13 Conte GL, Arnegard ME, Peichel CL, Schluter D. 2012. The probability of genetic parallelism and
14 convergence in natural populations. *Proc. R. Soc. B Biol. Sci.* 279:5039–5047.
- 15 Conway JR, Lex A, Gehlenborg N. 2017. UpSetR: An R package for the visualization of intersecting sets
16 and their properties. *Bioinformatics* 33:2938–2940.
- 17 Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT,
18 Sherry ST, et al. 2011. The variant call format and VCFtools. *Bioinformatics* 27:2156–2158.
- 19 Van Etten M, Lee KM, Chang SM, Baucom RS. 2020. Parallel and nonparallel genomic responses contribute
20 to herbicide resistance in *Ipomoea purpurea*, a common agricultural weed. *PLoS Genet.* 16:e1008593.
- 21 Excoffier L, Lischer HEL. 2010. Arlequin suite ver 3.5: a new series of programs to perform population
22 genetics analyses under Linux and Windows. *Mol. Ecol. Resour.* 10:564–567.
- 23 Fitzpatrick MC, Keller SR, Lotterhos KE. 2018. Comment on “Genomic signals of selection predict climate-
24 driven population declines in a migratory bird.” *Science.* 361:2–4.
- 25 Foote AD, Morin PA. 2016. Genome-wide SNP data suggest complex ancestry of sympatric North Pacific
26 killer whale ecotypes. *Heredity.* 117:316–325.
- 27 Fournier-Level A, Korte A, Cooper MD, Nordborg M, Schmitt J, Wilczek AM. 2011. A map of local
28 adaptation in *Arabidopsis thaliana*. *Science.* 334:86–89.
- 29 Gill M. 2018. Gazetteer of British Miscellaneous Mines. Available from: <https://www.nmrs.org.uk/>
- 30 Gou JY, Yu XH, Liu CJ. 2009. A hydroxycinnamoyltransferase responsible for synthesizing suberin
31 aromatics in *Arabidopsis*. *Proc. Natl. Acad. Sci. U. S. A.* 106:18855–18860.

- 1 Gough JW. 1967. The Mines of Mendip. Newton Abbot: David & Charles
- 2 Grunewald W, De Smet I, Lewis DR, Löffke C, Jansen L, Goeminne G, Vanden Bossche R, Karimi M, De
3 Rybel B, Vanholme B, et al. 2012. Transcription factor *WRKY23* assists auxin distribution patterns
4 during *Arabidopsis* root development through local control on flavonol biosynthesis. *Proc. Natl. Acad.*
5 *Sci. U. S. A.* 109:1554–1559.
- 6 Gulshan K, Shahi P, Moyer-Rowley WS. 2009. Compartment-specific synthesis of phosphatidylethanolamine
7 is required for normal heavy metal resistance. *Mol. Biol. Cell* 21:443–455.
- 8 Hackl T, Hedrich R, Schultz J, Förster F. 2014. Proovread: Large-scale high-accuracy PacBio correction
9 through iterative short read consensus. *Bioinformatics* 30:3004–3011.
- 10 Hall MC, Lowry DB, Willis JH. 2010. Is local adaptation in *Mimulus guttatus* caused by trade-offs at
11 individual loci? *Mol. Ecol.* 19:2739–2753.
- 12 Haller BC, Messer PW. 2019. SLiM 3: Forward genetic simulations beyond the Wright-Fisher model. *Mol.*
13 *Biol. Evol.* 36:632–637.
- 14 Hartley CJ, Newcomb RD, Russell RJ, Yong CG, Stevens JR, Yeates DK, La Salle J, Oakeshott JG. 2006.
15 Amplification of DNA from preserved specimens shows blowflies were preadapted for the rapid
16 evolution of insecticide resistance. *Proc. Natl. Acad. Sci. U. S. A.* 103:8757–8762.
- 17 Hartley S. 2009. Remediation of abandoned metal mine drainage using dealginated seaweed. PhD.
- 18 Helmstetter AJ, Cable S, Rakotonasolo F, Rabarijaona R, Rakotoarinivo M, Eiserhardt WL, Baker WJ,
19 Papadopulos AST. 2020. The demographic history of micro-endemics: have rare species always been
20 rare? *bioRxiv*. Available from: <https://doi.org/10.1101/2020.03.10.985853>
- 21 Hof AEV t., Campagne P, Rigden DJ, Yung CJ, Lingley J, Quail MA, Hall N, Darby AC, Saccheri IJ. 2016.
22 The industrial melanism mutation in British peppered moths is a transposable element. *Nature*
23 534:102–105.
- 24 Hughes S. 2000. Copperopolis: Landscapes of the Early Industrial Period in Swansea. Royal Commission on
25 the Ancient and Historical Monuments of Wales.
- 26 Jackman SD, Vandervalk BP, Mohamadi H, Chu J, Yeo S, Hammond SA, Jahesh G, Khan H, Coombe L,
27 Warren RL, et al. 2017. ABySS 2.0: Resource-efficient assembly of large genomes using a Bloom
28 filter. *Genome Res.* 27:768–777.
- 29 James ME, Arenas-Castro H, Groh JS, Engelstaedter J, Ortiz-Barrientos D. 2020. Highly replicated evolution
30 of parapatric ecotypes. *bioRxiv* Available from: <https://doi.org/10.1101/2020.02.05.936401>
- 31 Johnson MTJ, Munshi-South J. 2017. Evolution of life in urban environments. *Science.* 358.
- 32 Jombart T. 2008. Adegnet: A R package for the multivariate analysis of genetic markers. *Bioinformatics*

- 1 24:1403–1405.
- 2 Jones FC, Grabherr MG, Chan YF, Russell P, Mauceli E, Johnson J, Swofford R, Pirun M, Zody MC, White
3 S, et al. 2012. The genomic basis of adaptive evolution in threespine sticklebacks. *Nature* 484:55–61.
- 4 Keilig K, Ludwig-Müller J. 2009. Effect of flavonoids on heavy metal tolerance in *Arabidopsis thaliana*
5 seedlings. *Bot. Stud.* 50:311–318.
- 6 Khan MIR, Fatma M, Per TS, Anjum NA, Khan NA. 2015. Salicylic acid-induced abiotic stress tolerance
7 and underlying mechanisms in plants. *Front. Plant Sci.* 6:1–17.
- 8 Krasovec M, Chester M, Ridout K, Filatov DA. 2018. The mutation rate and the age of the sex chromosomes
9 in *Silene latifolia*. *Curr. Biol.*
- 10 Kreiner JM, Giacomini DA, Bemm F, Waithaka B, Regalado J, Lanz C, Hildebrandt J, Sikkema PH, Tranel
11 PJ, Weigel D, et al. 2019. Multiple modes of convergent adaptation in the spread of glyphosate-
12 resistant *Amaranthus tuberculatus*. *Proc. Natl. Acad. Sci. U. S. A.* 116:23363.
- 13 Küpper H, Andresen E. 2016. Mechanisms of metal toxicity in plants. *Metallomics* 8:269–285.
- 14 Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* 9:357–359.
- 15 Lee KM, Coop G. 2017. Distinguishing among modes of convergent adaptation using population genomic
16 data. *Genetics* 207:1591–1619.
- 17 Lee TH, Guo H, Wang X, Kim C, Paterson AH. 2014. SNPhylo: A pipeline to construct a phylogenetic tree
18 from huge SNP data. *BMC Genomics* 15:1–6.
- 19 Lescak EA, Bassham SL, Catchen J, Gelmond O, Sherbick ML, von Hippel FA, Cresko WA. 2015.
20 Evolution of stickleback in 50 years on earthquake-uplifted islands. *Proc. Natl. Acad. Sci.* 201512020.
- 21 Li Y, Iqbal M, Zhang Q, Spelt C, Bliet M, Hakvoort HWJ, Quattrocchio FM, Koes R, Schat H. 2017. Two
22 *Silene vulgaris* copper transporters residing in different cellular compartments confer copper
23 hypertolerance by distinct mechanisms when expressed in *Arabidopsis thaliana*. *New Phytol.*
24 215:1102–1114.
- 25 Lipson M. 2020. Interpreting f -statistics and admixture graphs : theory and examples. *Preprints*:1–18.
- 26 Lowry DB, Hall MC, Salt DE, Willis JH. 2009. Genetic and physiological basis of adaptive salt tolerance
27 divergence between coastal and inland *Mimulus guttatus*. *New Phytol.* 183:776–788.
- 28 Lowry DB, Hoban S, Kelley JL, Lotterhos KE, Reed LK, Antolin MF, Storfer A. 2017. Breaking RAD: an
29 evaluation of the utility of restriction site-associated DNA sequencing for genome scans of adaptation.
30 *Mol. Ecol. Resour.* 17:142–152.
- 31 Macnair MR. 1979. The genetics of copper tolerance in the yellow monkey flower, *Mimulus guttatus*. I.
32 Crosses to nontolerants. *Genetics* 91:553–563.

- 1 MacPherson A, Nuismer SL. 2017. The probability of parallel genetic evolution from standing genetic
2 variation. *J. Evol. Biol.* 30:326–337.
- 3 Malinsky M, Trucchi E, Lawson DJ, Falush D. 2018. RADpainter and fineRADstructure: Population
4 Inference from RADseq Data. *Mol. Biol. Evol.* 35:1284–1290.
- 5 Marques DA, Lucek K, Meier JI, Mwaiko S, Wagner CE, Excoffier L, Seehausen O. 2016. Genomics of
6 rapid incipient speciation in sympatric threespine stickleback. *PLoS Genet.* 12:1–34.
- 7 Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
8 *EMBnet.journal* 17.
- 9 McNeilly T, Bradshaw AD. 1968. Evolutionary processes in populations of copper tolerant *Agrostis tenuis*.
10 *Evolution.* 22:108–118.
- 11 Nerlich A, Von Orlow M, Rontein D, Hanson AD, Dörmann P. 2007. Deficiency in phosphatidylserine
12 decarboxylase activity in the psd1 psd2 psd3 triple mutant of *Arabidopsis* affects
13 phosphatidylethanolamine accumulation in mitochondria. *Plant Physiol.* 144:904–914.
- 14 Nosil P, Villoutreix R, De Carvalho CF, Farkas TE, Soria-Carrasco V, Feder JL, Crespi BJ, Gompert Z.
15 2018. Natural selection and the predictability of evolution in timema stick insects. *Science.* 359:765–
16 770.
- 17 O’Brien W. 2020. Ross Island Mine Heritage. *Dep. Archaeol. NUI Galw.* Available from:
18 http://www.nuigalway.ie/ross_island/index.htm
- 19 Oomen RA, Kuparinen A, Hutchings JA. 2020. Consequences of single-locus and tightly linked genomic
20 architectures for evolutionary responses to environmental change. *J. Hered.* esaa020.
- 21 Otto SP. 2018. Adaptation, speciation and extinction in the Anthropocene. *Proc. R. Soc. B Biol. Sci.* 285.
- 22 Oziolor E, Reid N, Yair S, KM L, VerPloeg S, Bruns P, Shaw J, Whitehead A, Matson C. 2019. Adaptive
23 introgression enables evolutionary rescue from extreme environmental pollution. *Science.* 364:455–
24 457.
- 25 Papadopulos AST, Igea J, Smith TP, Hutton I, Baker WJ, Butlin RK, Savolainen V. 2019. Ecological
26 speciation in sympatric palms: 4. Demographic analyses support speciation of *Howea* in the face of
27 high gene flow. *Evolution.* 73:1996–2002.
- 28 Papadopulos AST, Kaye M, Devaux C, Hipperson H, Lighten J, Dunning LT, Hutton I, Baker WJ, Butlin
29 RK, Savolainen V. 2014. Evaluation of genetic isolation within an island flora reveals unusually
30 widespread local adaptation and supports sympatric speciation. *Philos. Trans. R. Soc. B Biol. Sci.*
31 369:20130342.
- 32 Paulino D, Warren RL, Vandervalk BP, Raymond A, Jackman SD, Birol I. 2015. Sealer: A scalable gap-
33 closing application for finishing draft genomes. *BMC Bioinformatics* 16:1–8.

- 1 Pellicer J, Leitch IJ. 2020. The Plant DNA C-values database (release 7.1): an updated online repository of
2 plant genome size data for comparative studies. *New Phytol.* 226:301–305.
- 3 Pennings PS, Hermisson J. 2006. Soft sweeps III: The signature of positive selection from recurrent
4 mutation. *PLoS Genet.* 2:1998–2012.
- 5 Peter BM. 2016. Admixture, population structure, and f-statistics. *Genetics* 202:1485–1501.
- 6 Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE. 2012. Double digest RADseq: an inexpensive
7 method for de novo SNP discovery and genotyping in model and non-model species. *PLoS One*
8 7:e37135.
- 9 Pickrell JK, Pritchard JK. 2012. Inference of population splits and mixtures from genome-wide allele
10 frequency data. *PLoS Genet.* 8.
- 11 Prysycz LP, Gabaldón T. 2016. Redundans: An assembly pipeline for highly heterozygous genomes. *Nucleic*
12 *Acids Res.* 44:e113.
- 13 Ravinet M, Elgvin TO, Trier C, Aliabadian M, Gavrilov A, Sætre GP. 2018. Signatures of human-
14 commensalism in the house sparrow genome. *Proc. R. Soc. B Biol. Sci.* 285.
- 15 Ravinet M, Westram A, Johannesson K, Butlin R, André C, Panova M. 2016. Shared and nonshared genomic
16 divergence in parallel ecotypes of *Littorina saxatilis* at a local scale. *Mol. Ecol.* 25:287–305.
- 17 Reich D, Thangaraj K, Patterson N, Price AL, Singh L. 2009. Reconstructing Indian population history.
18 *Nature* 461:489–494.
- 19 Reid NM, Proestou DA, Clark BW, Warren WC, Colbourne JK, Shaw JR, Karchner SI, Hahn ME, Nacci D,
20 Oleksiak MF, et al. 2016. The genomic landscape of rapid repeated evolutionary adaptation to toxic
21 pollution in wild fish. *Science.* 354:1305–1308.
- 22 Rochette NC, Rivera-Colón AG, Catchen JM. 2019. Stacks 2: Analytical methods for paired-end sequencing
23 improve RADseq-based population genomics. *Mol. Ecol.* 28:4737–4754.
- 24 Roda F, Ambrose L, Walter GM, Liu HL, Schaul A, Lowe A, Pelsler PB, Prentis P. 2013. Genomic evidence
25 for the parallel evolution of coastal forms in the *Senecio lautus* complex. *Mol. Ecol.* 22:2941–2952.
- 26 Ross SM, Wood MD, Copplestone D, Warriner M, Crook P. 2007. UK Soil and Herbage - Pollutant Survey
27 Environmental Concentrations of Heavy Metals in UK Soil and Herbage.
- 28 Rundle HD, Nagel L, Wenrick Boughman J, Schluter D. 2000. Natural selection and parallel speciation in
29 sympatric sticklebacks. *Science.* 287:306–308.
- 30 Runyeon H, Prentice HC. 1997. Genetic differentiation in the Bladder champions, *Silene vulgaris* and *S.*
31 *uniflora* (Caryophyllaceae), in Sweden. *Biol. J. Linn. Soc.* 61:559–584.
- 32 Santangelo JS, Thompson KA, Cohan B, Syed J, Ness RW, Johnson MTJ. 2020. Predicting the strength of

- 1 urban-rural clines in a Mendelian polymorphism along a latitudinal gradient. *Evol. Lett.* 4:212–225.
- 2 Schat H, Vooijs R. 1997. Multiple tolerance and co-tolerance to heavy metals in *Silene vulgaris*: A co-
3 segregation analysis. *New Phytol.* 136:489–496.
- 4 Schat H, Vooijs R, Kuiper E. 1996. Identical major gene loci for heavy metal tolerances that have
5 independently evolved in different local populations and subspecies of *Silene vulgaris*. *Evolution.*
6 50:1888–1895.
- 7 Stanke M, Schöffmann O, Morgenstern B, Waack S. 2006. Gene prediction in eukaryotes with a generalized
8 hidden Markov model that uses hints from external sources. *BMC Bioinformatics* 7:62.
- 9 Szulkin, M., Munshi-South, J., & Charmantier A. 2020. Urban Evolutionary Biology. Oxford University
10 Press.
- 11 Therkildsen NO, Wilder AP, Conover DO, Munch SB, Baumann H, Palumbi SR. 2019. Contrasting genomic
12 shifts underlie parallel phenotypic evolution in response to fishing. *Science.* 490:487–490.
- 13 Urban M. 2015. Accelerating extinction risk from climate change. *Science.* 348.
- 14 Vandervalk BP, Yang C, Xue Z, Raghavan K, Chu J, Mohamadi H, Jackman SD, Chiu R, Warren RL, Birol
15 I. 2015. Konnector v2.0: Pseudo-long reads from paired-end sequencing data. *BMC Med. Genomics*
16 8:S1.
- 17 Wang M, Zhao Y, Zhang B. 2015. Efficient test and visualization of multi-set intersections. *Sci. Rep.* 5:1–12.
- 18 Wright KM, Hellsten U, Xu C, Jeong AL, Sreedasyam A, Chapman JA, Schmutz J, Coop G, Rokhsar DS,
19 Willis JH. 2015. Adaptation to heavy-metal contaminated environments proceeds via selection on pre-
20 existing genetic variation. *bioRxiv* Available from: <http://biorxiv.org/lookup/doi/10.1101/029900>
- 21 Wright S. 1943. Isolation by distance. *Genetics* 28:114.
- 22 Wu L, Bradshaw AD. 1972. Aerial pollution and the rapid evolution of copper tolerance. *Nature* 238:167–
23 169.
- 24 Xie L, Klerks PL. 2004. Fitness cost of resistance to cadmium in the least killifish (*Heterandria formosa*).
25 *Environ. Toxicol. Chem.* 23:1499–1503.
- 26


27 **Acknowledgements:** This work was supported by the Natural Environment Research Council (NERC); and
28 the Royal Society We thank Alan Baker, Roger Butlin, Andrew Leitch & Steve Rossiter for encouragement
29 and discussion; Robyn Cowan, Wendy Grail & Jonathan Kendon for laboratory support; and the Botanical
30 Society of the British Isles & Mike Gill for access to databases.

31 **Author Contributions:** ASTP conceived and designed the research with contributions from all co-authors.
32 ASTP, RJS, JL & ES conducted fieldwork. AJH conducted ddRAD lab work. ASTP, ES & LM conducted

1 tolerance experiments. ASTP analysed the data, with contributions from OGO, AF, AC & JL. AAC conducted
2 simulations. ASTP wrote the manuscript and all authors commented on the final version.


3 **Data Availability:** DNA sequence data that support the findings of this study have been deposited in the NCBI
4 Short Read Archive and are accessible through accession number PRJNA699303. The draft genome has been
5 deposited in the GenBank repository under accession number JAGPOY010000000. Custom code for
6 simulations is included as the supplementary material.

7




8

9 **Figure 1. Differential heavy metal tolerance between local mine and coastal populations.** (A) Map of
10 population sampling locations. Fill colours denote habitat type (mine - orange, coastal - blue). Outline colours
11 denote local populations (West Wales - WWA; South Wales - SWA; South-West England – ENG; South-West
12 Ireland – IRE). The same colour scheme is used throughout. (B) Zinc and (C) copper tolerance for each mine-
13 coast pair (centre line, median; box limits, upper and lower quartiles; whiskers, 1.5x interquartile range; points,
14 outliers; Zinc treatment left to right $n = 15/17/14/14/14/16/15/19$; Copper treatment left to right $n =$
15 $17/17/16/17/15/18/16/18$). Local mine and coastal populations have significantly different zinc tolerance, but
16 only the Irish pair have significantly different copper tolerance.




1

2 **Figure 2. Isolation by distance (IBD) patterns arising from multiple or single origins of mine**
 3 **populations.** (A) Under a simulated multiple independent origin model, the correlation between F_{ST} and
 4 migration between mine populations (orange) is steeper (i.e., IBD is stronger) and has a higher intercept than
 5 isolation by distance between coastal populations (blue). (B) In contrast, under a single origin model the
 6 relationship between genetic differentiation and geography breaks down between mine populations – the slope
 7 is not significantly different from zero and the intercept is lower than between coastal populations (centre line,
 8 median; box limits, upper and lower quartiles; whiskers, 1.5x interquartile range; points, outliers). (C) The
 9 observed IBD relationships in *S. uniflora* conform to the patterns expected from multiple origins of the mine
 10 populations. IBD between mine and coastal populations in green.




1
 2 **Figure 3. Evidence for three independent origins of zinc tolerant populations in *S. uniflora*.** (A)
 3 Phylogenetic reconstruction (mine populations in orange and coastal populations in blue). Nodes with greater
 4 than 90% bootstrap support are denoted by black circles. (B) Principal components analysis points to three,
 5 well supported, independent origins of zinc tolerant populations. Variance explained by PC1 = 12.3% and PC2
 6 = 9.0%. Points are coloured by region as Fig. 1. All points from a specific population are surrounded by a
 7 single ellipse which is coloured by habitat type (mine - orange, coast - blue). (C) Treemix analysis with four
 8 migration edges. Points are coloured as in Fig. 1. The topology is almost identical to that produced by the
 9 SNPhylo analysis - the relationship of ENG-C and SWA-C to ENG-M is reversed. Colour scale indicates
 10 migration edge weight. Only migration between coastal populations was supported by f_4 statistics (see Fig. 4).



1

2 **Figure 4. Evidence for admixture between coastal populations but not between mines.** z -scores of f_4
 3 statistics for the six different permutations of four taxon trees (Types 1-6), with all of the different
 4 combinations of mine (orange) and coastal (blue) populations based on relationships in Fig 3A. The red line
 5 denotes the z -score at which the f_4 statistic is significantly different from zero at the 5% level after Dunn-
 6 Bonferroni correction for multiple tests ($z = 3.67$). There is evidence of admixture in the *four coast tree* (Type
 7 1; $z = 5.23$) and *three coast : one mine trees* (Type 3; $z > 3.67$ for three of the quartets), which is also reflected
 8 in the four Type 5 quartets with z -scores exceeding 3.67. On the other hand, the *four mine* (Type 2; $z = 0.17$)
 9 and *three mine : one coast trees* (Type 4; $z = 0.14 - 2.35$) demonstrate that there has not been introgression
 10 between mine sites.

11



1

2 **Figure 5. Molecular convergence and divergence across regional mine-coast pairs.** Upset plots of the
 3 shared (A) outlier scaffolds and (B) individual SNPs across the four regional mine-coast pairs. Filled points
 4 below bars denote which regional sets are intersected for each bar (e.g., the leftmost bar in each plot represents
 5 the set including all four mine-coast comparisons). Inset scatterplots show observed overlap (y-axis) vs
 6 expected overlap (x-axis) across combinations of regional sets, with line at 1:1. Black bars denote outliers
 7 found in a single geographic region. The remaining bars are coloured by super exact test p-value (all < 0.001)
 8 with darker green denoting smaller p-values and purple denoting extremely small values ($< 10^{-150}$).