

# Three-dimensional surface gravity waves of a broad bandwidth on deep water

Yan Li<sup>1,2†</sup>

<sup>1</sup>Department of Energy and Process Engineering, Norwegian University of Science and Technology, N-7491 Trondheim, Norway

<sup>2</sup>Department of Engineering Science, University of Oxford, Parks Road, Oxford OX1 3PJ, UK

(Received xx; revised xx; accepted xx)

A new nonlinear Schrödinger equation (NLSE) is presented for ocean surface waves. Earlier derivations of NLSEs that describe the evolution of deepwater waves have been limited to a narrow bandwidth, for which the bound waves at second order in wave steepness are described in leading-order approximations. This work generalizes these earlier works to allow for deepwater waves of a broad bandwidth with large directional spreading. The new NLSE permits simple numerical implementations and can be extended in a straightforward manner in order to account for waves on water of finite depth. For the description of second-order waves, this paper proposes a semi-analytical approach that can provide accurate and computationally efficient predictions. With a leading-order approximation to the new NLSE, the instability region and energy growth rate of Stokes waves are investigated. Compared with the exact results based on McLean (*J. Fluid Mech.*, vol. 511, 1982), predictions by the new NLSE show better agreement than by Trulsen *et al.* (*Phys. Fluids*, vol. 12, 2000). With numerical implementations of the new NLSE, the effects of wave directionality are investigated by examining the evolution of a directionally spread focused wave group. A downward shift of the spectral peak is observed, owing to the asymmetry in the change rate of energy in a more complex manner than that for uniform Stokes waves. Rapid oblique energy transfers near the group at linear focus are observed, likely arising from the instability of uniform Stokes waves appearing in a narrow spectrum subject to oblique sideband disturbances.

## 1. Introduction

An important task in the study of surface gravity waves is the development of the theoretical description of flow fields. Theoretical models of surface gravity waves are essential in a wide range of fields, such as in engineering practices and for research purposes. A theoretical study has obvious advantages in elucidating the underlying physics, thereby advancing the understanding of realistic wave problems, compared with other approaches such as experiments, field studies, and direct numerical computations. Theoretical findings have contributed to providing possible explanations to the formation mechanism of extremely large waves that appear suddenly of much larger amplitude than their surroundings, known also as ‘rogue’ or ‘freak’ waves. A few examples of the possible mechanisms proposed are modulational or Benjamin-Feir instability in deep water (Onorato *et al.* 2009), refraction by ambient currents or bathymetry (Janssen & Herbers 2009; Onorato *et al.* 2011), and effects on weakly nonlinear waves of a depth change in shallow or intermediate water depth (Trulsen *et al.* 2020; Li *et al.* 2021*a,b*).

† Email address for correspondence: yan.li@ntnu.no

TABLE 1. A summary of nonlinear Schrödinger equations (NLSEs) for waves on deep water and the regimes of applicability which refer to the accuracy of lower-order wave fields considered in a NLSE for the dynamic evolution of the (potential or elevation) envelope of the first harmonic, in addition to the NLSEs for an arbitrary depth;  $\epsilon$  denotes the dimensionless wave steepness,  $\delta_x$  and  $\delta_y$  denote the dimensionless bandwidth in the main propagation direction and in the direction normal to the main, respectively.  $\delta_{x,y}^\infty$  denotes arbitrary order in bandwidth.

	Order in wave steepness			Overall order of accuracy
	first order	second order	third order	
Dysthe (1979) ( $\mathcal{O}(\delta_{x,y}) \sim \mathcal{O}(\epsilon)$ )	$\mathcal{O}(\epsilon \delta_x^n \delta_y^{3-n})$	$\mathcal{O}(\epsilon^2 \delta_{x,y})$	$\mathcal{O}(\epsilon^3 \delta_{x,y})$	$\mathcal{O}(\epsilon^3 \delta_{x,y}, \epsilon \delta_x^n \delta_y^{3-n})$ ( $n = 0, 1, 2, 3$ )
Trulsen & Dysthe (1996) ( $\mathcal{O}(\delta_{x,y}) \sim \mathcal{O}(\sqrt{\epsilon})$ )	$\mathcal{O}(\epsilon \delta_x^n \delta_y^{5-n})$	$\mathcal{O}(\epsilon^2 \delta_{x,y})$	$\mathcal{O}(\epsilon^3 \delta_{x,y})$	$\mathcal{O}(\epsilon^3 \delta_{x,y}, \epsilon \delta_x^n \delta_y^{5-n})$ ( $n = 0, 1, \dots, 5$ )
Trulsen <i>et al.</i> (2000) ( $\delta_{x,y} < 1$ )	$\mathcal{O}(\epsilon \delta_{x,y}^\infty)$	$\mathcal{O}(\epsilon^2 \delta_{x,y})$	$\mathcal{O}(\epsilon^3 \delta_{x,y})$	$\mathcal{O}(\epsilon^3 \delta_{x,y}, \mathcal{O}(\epsilon \delta_{x,y}^\infty))$
This paper ( $\delta_{x,y} \sim \mathcal{O}(1)$ )	$\mathcal{O}(\epsilon)$	$\mathcal{O}(\epsilon^2)$	$\mathcal{O}(\epsilon^3)$	$\mathcal{O}(\epsilon^3)$

The modulational instability of weakly nonlinear Stokes waves subject to sideband disturbances was discovered by Benjamin & Feir (1967). Theoretical advances made in the late 1960s in this perspective include contributions by Lighthill (1965); Benjamin (1967); Whitham (1967); Zakharov (1968) and Benney & Roskes (1969). Since the discovery of the instability of a train of Stokes waves, progress has been made both numerically (Janssen 1983; Lo & Mei 1985; Trulsen & Dysthe 1997; Dysthe *et al.* 2003) and experimentally (Lake *et al.* 1977; Melville 1982; Su 1982; Chabchoub *et al.* 2011) in understanding the evolution properties of surface waves on deep water. For smaller values of wave steepness, symmetrical upper and lower sidebands tend to grow equally in the initial stage. This is due to the degenerate resonant interaction in the prestigious ‘figure of eight’ quartet resonance loop (Phillips 1960, 1967; Longuet-Higgins 1976; Lake *et al.* 1977; McLean 1982), which corresponds to the special cases where two out of the four wavenumbers obeying the ‘figure of eight’ loop are identical. As nonlinearity increases, the sidebands appear to grow unequally, with the lower sideband growing faster than the upper sideband (Lake *et al.* 1977; Lo & Mei 1985), reaching a maximum larger than the minimum to which the spectrum peak drops. This unequal growth in energy, as a result of a combination of dissipation, wave breaking, and nonlinear wave evolution, likely causes the downward shift of the spectral peak of wind waves (Lake *et al.* 1977; Lo & Mei 1985; Trulsen & Dysthe 1997) .

The so-called nonlinear Schrödinger equation (NLSE) has been widely known as a convenient approach for analysing the instability of Stokes waves. For deepwater waves, Zakharov (1968) found that the wave envelope satisfies the NLSE which is cubic in wave slope  $\epsilon$  (or wave steepness). The NLSE is derived based on the assumption of a small wave slope and the so-called narrow-banded assumption that restricts the modulation of the wave envelope to be slow in both space and time relative to a rapidly varying wave phase.

Many attempts have been made to expand the applicability scope of the NLSE for deepwater waves through adding higher-order terms, with a primary focus on reducing the restrictions due to bandwidth, as shown in table 1. Table 1 indicates the lower-order wave fields considered in a NLSE for the nonlinear and dynamic evolution of the (potential or elevation) envelope of the first harmonic. Specifically, the second-order wave fields do not appear alone in a NLSE but in combination with the first-order wave fields.

Let  $\delta_{x,y}$  be a small non-dimensional parameter as a measure of the bandwidth in the main propagation ( $\delta_x$ ) and the transverse ( $\delta_y$ ) direction, respectively. With one step further, Dysthe (1979) has obtained a NLSE that is correct to  $\mathcal{O}(\epsilon^3 \delta_{x,y}, \epsilon \delta_x^n \delta_y^{3-n})$  ( $n = 0, 1, 2, 3$ ), known as Dysthe's equation, or the fourth-order NLSE as  $\mathcal{O}(\delta_{x,y}) \sim \mathcal{O}(\epsilon)$  is implied. With the fourth-order NLSE, Dysthe (1979) finds that second-order mean flows play a significant role in the instability of Stokes waves. It is shown by Stiassnie (1984) that Dysthe's equation can be derived from Zakharov's integral equation (Zakharov 1968) through a narrow-banded assumption. Higher-order extensions in bandwidth are derived by Trulsen & Dysthe (1996) with a modified NLSE correct to  $\mathcal{O}(\epsilon \delta_x^n \delta_y^{5-n}, \epsilon^3 \delta_{x,y})$  ( $n = 0, 1, \dots, 5$ ) and by Trulsen *et al.* (2000) in which the linear evolution of the envelope of linear wave is described by the exact linear dispersion relation, giving a NLSE correct to  $\mathcal{O}(\epsilon \delta_{x,y}^\infty, \epsilon^3 \delta_{x,y})$ . Using some of the earlier versions of NLSEs, Trulsen & Dysthe (1997) and Dysthe *et al.* (2003) find that the NLSEs can recover a downward shift of the spectral peak in a three- and two-dimensional narrow spectrum, respectively. The latter occurs after the spectrum has reached a quasi-steady state and hence, is unlikely due to the modulational instability which results in symmetrical growth of sidebands. Strong frequency dependence developed in the temporal evolution of directional waves initially of no frequency dependence is observed in NLSE-based numerical simulations and field data (Simanesev *et al.* 2016). Simanesev *et al.* (2017) find the modified NLSE of Trulsen *et al.* (2000) gives less accurate predictions for short crest waves with larger directional spread.

Recent progress has also been made in generalising a NLSE in order to take into account the interaction of waves with ambient environments, e.g. dissipation and forcing effects due to wind actions and turbulence (cf. Wu *et al.* (2006); Dias *et al.* (2008); Kharif *et al.* (2010); Slunyaev *et al.* (2015)) and wave current interaction (Dysthe & Das 1981; Stocker & Peregrine 1999; Hjelmervik & Trulsen 2009; Curtis *et al.* 2018). In addition to the NLSEs for deepwater waves, coupled NLSEs are derived to consider crossing sea states, see Gramstad & Trulsen (2011); Trulsen *et al.* (2015) and references therein. A generalisation to water of finite uniform depth is done by Davey & Stewartson (1974) and higher-order corrections are added by Johnson (1977); Brinch-Nielsen & Jonsson (1986); Slunyaev (2005); Gramstad (2014) among others. Modified cubic equations are obtained that allow for waves in water of a finite, varying depth due to a mildly-sloping seabed (Djordjević & Redekopp 1978; Kirby & Dalrymple 1983).

Many other approaches have been developed as alternative tools to the NLSE for understanding the properties of surface deepwater waves; such as the Zakharov integral equation and its leading order approximations (see Zakharov (1968); Crawford *et al.* (1980, 1981); Krasitskii (1994)), numerical wave tanks (e.g., Ducrozet *et al.* (2012); Bateman *et al.* (2001)) based on the High-Order Spectral (HOS) method (Dommermuth & Yue 1987; West *et al.* 1987), and direct numerical solutions of fully nonlinear potential-flow equations (Engsig-Karup *et al.* 2009; Bihs *et al.* 2020; Zheng *et al.* 2020). Through studying the temporal evolution of three-dimensional steep focused wave groups, rapid energy changes in a wave spectrum are reported by earlier works e.g. Gibbs & Taylor (2005); Gibson & Swan (2007). They attribute this phenomenon to the third-order resonant interactions, which are considered as the cause for the formation of the so-called 'wing waves' that appear in the transverse direction after the group at nonlinear focus reported in Barratt *et al.* (2020). Direct numerical solutions were used by Barratt *et al.* (2020) for the evolution of a steep focused wave group with directional spread. In three dimensions, oblique energy transfer in a wave spectrum and a downward shift of the spectral peak are observed in previous works, e.g. Trulsen & Dysthe (1997) and Barratt *et al.* (2021).

There is still room and need for further development of the models based on a NLSE, with two aspects addressed here. Firstly, ocean surface waves are generally three dimensional and broad banded. Nevertheless, it should be noted that the NLSEs mentioned above, e.g. Dysthe (1979); Trulsen *et al.* (2000), are only correct to order  $(\epsilon^3 \delta_{x,y})$ , as shown in table 1. This suggests the need for extending the bandwidth at third order in wave steepness. Secondly, surface waves interact with ambient environments such as sub-surface currents, turbulence, and wind actions in the atmosphere, which often require explicit and accurate vertical structures of flow fields below the surface to allow for coupling. In contrast, it is known that a NLSE is based on the description of the envelope of the surface displacement, accompanied by mostly lowest-order approximations to the vertical profiles of flow fields at second order in wave steepness, as shown in table 1.

This paper aims to make attempts to fill in gaps in the aforementioned two aspects, through the development of a new framework that would allow for the study of three-dimensional waves of a broad bandwidth and would take into account the structures of flow fields below the surface without compromising much the computational efficiency of a NLSE-based model. The objective of this paper is threefold. First, a new NLSE for the evolution of three-dimensional deepwater surface waves is derived in §2–§4 that does not rely on the assumption of a narrow bandwidth as aforementioned NLSEs. Specially, the newly developed NLSE should allow for  $\delta_{x,y} > 1$  since it has relaxed the narrowband assumption for wave fields at second order, as shown in table 1. Simple numerical implementations of the new NLSE as performed with earlier versions of NLSEs are explained. Second, a semi-analytical approach for the description of wave fields for bound waves at second order in wave steepness is proposed in §3 which allows for simpler and more efficient numerical implementations than by Dalzell (1999). Finally, a study of the instability region and energy growth rate of Stokes waves and the temporal evolution of a directionally-spread focused wave group are presented in §6, where comparisons with Longuet-Higgins (1978), Dysthe (1979), Crawford *et al.* (1981), McLean (1982), and Trulsen *et al.* (2000) are made.

## 2. Mathematical formulation and methodology

### 2.1. Problem definition

We consider ocean surface waves propagating on deep water in the framework of potential-flow theory, thereby assuming incompressible inviscid flows, irrotational fluid motions, and negligible effects of surface tension. A Cartesian coordinate system is chosen with the undisturbed water surface located at  $z = 0$ . The system can be described as a boundary value problem governed by the Laplace equation:

$$\nabla_3^2 \Phi = 0 \quad \text{for} \quad -\infty < z < \zeta(\mathbf{x}, t), \quad (2.1)$$

where  $\Phi(\mathbf{x}, z, t)$  denotes the velocity potential,  $\zeta(\mathbf{x}, t)$  is the free surface elevation,  $\mathbf{x}$  is the position vector in the horizontal plane,  $t$  is the time, and  $\nabla_3 = (\nabla, \partial_z)$  with  $\nabla = (\partial_x, \partial_y)$  denoting the gradient in the horizontal plane. Equation (2.1) should be solved subject to the nonlinear kinematic and dynamic boundary conditions (cf. Davey & Stewartson (1974)) at the free water surface  $z = \zeta(\mathbf{x}, t)$ :

$$\partial_t \zeta + \nabla \Phi \cdot \nabla \zeta - \partial_z \Phi = 0 \quad \text{and} \quad \Gamma \Phi + \partial_t (\nabla_3 \Phi)^2 + \frac{1}{2} \nabla_3 \Phi \cdot \nabla_3 (\nabla_3 \Phi)^2 = 0 \quad (2.2a,b)$$



where the operator  $\Gamma$  is defined as  $\Gamma = \partial_{tt} + g\partial_z$  with  $g$  the gravitational acceleration; A deepwater boundary condition:

$$\partial_z \Phi = 0 \quad \text{for } z \rightarrow -\infty. \quad (2.3)$$

## 2.2. Stokes expansion and separation of harmonics

In order to solve the boundary value problem (2.1)-(2.3), we seek the solutions of unknown  $\Phi$  and  $\zeta$  in a form of power series in wave steepness defined as  $\epsilon = k_0 A_0$  (a so-called Stokes expansion), with  $k_0$  and  $A_0$  denoting the characteristic wavenumber and wave amplitude, respectively,

$$\Phi = \epsilon \Phi^{(1)} + \epsilon^2 \Phi^{(2)} + \epsilon^3 \Phi^{(3)} + \mathcal{O}(\epsilon^4) \quad \text{and} \quad \zeta = \epsilon \zeta^{(1)} + \epsilon^2 \zeta^{(2)} + \epsilon^3 \zeta^{(3)} + \mathcal{O}(\epsilon^4), \quad (2.4a,b)$$

where we consider the first three orders and the superscripts denote the order in  $\epsilon$ . Substituting (2.4) into the boundary value problem (2.1)-(2.3) leads to the decomposition of the fully nonlinear system into different problems through a collection of the terms at the same order in  $\epsilon$ . The decomposed problems can be solved successively from the lowest to higher orders, as presented in the following from the first (§2.3) to third order (§5).

Let linear surface elevation  $\zeta^{(1)}$  be expressed in two equivalent forms as follows,

$$\zeta^{(1)}(\mathbf{x}, t) = \frac{1}{2} A(\mathbf{x}, t) e^{i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)} + \text{c.c.} \quad \text{or} \quad (2.5a)$$

$$\zeta^{(1)}(\mathbf{x}, t) = \frac{1}{2} \int_{-\infty}^{\infty} \hat{\zeta}(\mathbf{k}) e^{i(\mathbf{k} \cdot \mathbf{x} - \omega(\mathbf{k})t)} d\mathbf{k}^2 + \text{c.c.}, \quad (2.5b)$$

in which c.c. denotes the complex conjugates,  $A$  denotes the complex wave envelope of the surface displacement of the carrier wave with  $\mathbf{k}_0 = (k_0, 0)$  the wavenumber vector chosen in the  $x$  direction,  $\omega_0$  denotes the angular frequency of the carrier wave that satisfies the dispersion relation  $\omega_0 = \omega(\mathbf{k}_0)$  given by  $\omega(\mathbf{k}) = \sqrt{gk}$  (where  $k = |\mathbf{k}|$ );  $\mathbf{k} = (k_x, k_y)$  denotes a wave vector in the horizontal plane. (2.5a) denotes the first-order elevation being expressed in an envelope-type form and (2.5b) a form through linear superposition of monochromatic waves in the Fourier  $\mathbf{k}$  plane. This paper focuses on the former which leads to a nonlinear evolution equation for  $A$  that allows for the relaxation of a narrow banded assumption (cf. Chu & Mei (1971) and Davey & Stewartson (1974)).

Equating (2.5a) and (2.5b) we obtain a relation between  $A$  and  $\hat{\zeta}$  as follows

$$A(\mathbf{x}, t) = \int_{-\infty}^{\infty} \hat{\zeta}(\boldsymbol{\kappa} + \mathbf{k}_0) e^{i[\boldsymbol{\kappa} \cdot \mathbf{x} - (\omega(\boldsymbol{\kappa} + \mathbf{k}_0) - \omega_0)t]} d\boldsymbol{\kappa}^2, \quad (2.6)$$

where the integral variable was replaced with  $\boldsymbol{\kappa}$  through  $\mathbf{k} = \boldsymbol{\kappa} + \mathbf{k}_0$ . Introducing a Fourier transform for  $A$  defined as follows

$$A(\mathbf{x}, t) = \int_{-\infty}^{\infty} \hat{A}(\mathbf{k}, t) e^{i\mathbf{k} \cdot \mathbf{x}} d\mathbf{k}, \quad \text{we obtain } \hat{A}(\mathbf{k}, t) = \hat{\zeta}(\mathbf{k} + \mathbf{k}_0) e^{-i(\omega(\mathbf{k} + \mathbf{k}_0) - \omega_0)t}. \quad (2.7a,b)$$

A linear evolution equation for  $A$  for waves of a broad bandwidth is derived by Trulsen

*et al.* (2000) in two equivalent forms as follows

$$\partial_t A + i\omega_0 \mathcal{L}A = 0 \quad \text{with} \quad (2.8a)$$

$$\mathcal{L}(\partial_x, \partial_y) = \left[ \left( 1 - \frac{i\partial_x}{k_0} \right)^2 + \frac{(i\partial_y)^2}{k_0^2} \right]^{1/4} - 1 \quad \text{and} \quad (2.8b)$$

$$\partial_t \hat{A} + i(\omega(\mathbf{k} + \mathbf{k}_0) - \omega_0) \hat{A} = 0, \quad (2.8c)$$

where (2.8a) is used for constructing the solutions in §3, §4, and §6.2; equation (2.8c) is in a form convenient for numerical implementations, as explained in §4.4. It is clear that a narrow-banded assumption implies  $k_0 \mathcal{L}A \ll \mathcal{O}(\epsilon)$ , which, again, this paper aims to drop. For convenience and later reference, we introduce a dimensionless bandwidth parameter defined as follows

$$\delta_{x,y} \sim \frac{\Delta k_{x,y}}{k_0} \quad \text{or} \quad \delta_{x,y} \sim \mathcal{O}\left(\frac{\partial_{x,y} A}{\epsilon}\right), \quad (2.9)$$

where  $\delta_x$  ( $\Delta k_x$ ) and  $\delta_y$  ( $\Delta k_y$ ) denote the dimensionless (dimensional) bandwidth in the  $x$  and  $y$  direction, respectively. The bandwidth is not assumed small, in contrast to conventional, reduced-form evolution equations for  $A$  by earlier works e.g. Dysthe (1979); Stiassnie (1984). Specifically,  $\delta_{x,y} > 1$  are permitted in this paper. Similar to (2.5a),  $\zeta$  and  $\Phi$  are expressed in an envelope-type form through the separation of wave harmonics as follows

$$\begin{aligned} \Phi(\mathbf{x}, z, t) = & \frac{1}{2}\epsilon(B + \epsilon^2 B^{(31)})e^{k_0 z} e^{i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)} + \text{c.c.} + \epsilon^2 [\Phi^{(20)}(\mathbf{x}, z, t) + (\Phi^{(22)}(\mathbf{x}, z, t) + \text{c.c.})] \\ & + \epsilon^3 (\Phi^{(33)}(\mathbf{x}, z, t) + \text{c.c.}), \end{aligned} \quad (2.10a)$$

$$\begin{aligned} \zeta(\mathbf{x}, t) = & \frac{1}{2}\epsilon(A + \epsilon^2 A^{(31)})e^{i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)} + \text{c.c.} + \epsilon^2 \left[ \zeta^{(20)} + \left( \frac{1}{2} A^{(22)} e^{2i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)} + \text{c.c.} \right) \right] + \\ & \frac{1}{2}(\epsilon^3 A^{(33)} e^{3i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)} + \text{c.c.}), \end{aligned} \quad (2.10b)$$

in which the superscripts  $(ij)$  denote  $\mathcal{O}(\epsilon^i)$  and  $j$ -th harmonic,  $A^{(ij)} = A^{(ij)}(\mathbf{x}, t)$  and  $B^{(ij)} = B^{(ij)}(\mathbf{x}, z, t)$  are the modulated amplitude and potential, respectively, the potentials at second and third harmonics are given by, respectively

$$\Phi^{(22)}(\mathbf{x}, z, t) = \frac{1}{2}B^{(22)}e^{2k_0 z} e^{2i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)} \quad \text{and} \quad \Phi^{(33)}(\mathbf{x}, z, t) = \frac{1}{2}B^{(33)}e^{3k_0 z} e^{3i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)}. \quad (2.11a,b)$$

It is worth noting, although (2.10a,b) are in a form similar to the so-called harmonic expansion, the wave fields at second and third order in such a form are introduced for convenience. They are due to the separation of wave harmonics arising from mainly the forcing equations (e.g. (3.1b) and (4.1)) at still water surface presented in §3 at second order and in §4 at third order.

### 2.3. Linear wave fields

Solutions to linearised boundary value problem from (2.1)-(2.3) are obtained for linear wave fields. They are given here without detailed derivations (see §13 in Mei *et al.* (2005)),

expressed in terms of  $\hat{A}(\mathbf{k})$  as follows

$$\mathbf{V}^{(1)} \equiv [\mathbf{u}^{(1)}, w^{(1)}] = \frac{1}{2} \bar{\mathbf{V}}(\mathbf{x}, z, t) e^{k_0 z} e^{i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)} + \text{c.c.}, \text{ with} \quad (2.12a)$$

$$\bar{\mathbf{V}} = [\bar{\mathbf{u}}(\mathbf{x}, z, t), \bar{w}(\mathbf{x}, z, t)] \text{ and} \quad (2.12b)$$

$$\begin{bmatrix} B \\ \bar{\mathbf{u}} \\ \bar{w} \end{bmatrix} = -i \int_{-\infty}^{\infty} \begin{bmatrix} 1 \\ i(\mathbf{k} + \mathbf{k}_0) \\ |\mathbf{k} + \mathbf{k}_0| \end{bmatrix} c_p(\mathbf{k} + \mathbf{k}_0) \hat{A}(\mathbf{k}, t) e^{(|\mathbf{k} + \mathbf{k}_0| - k_0)z} e^{i\mathbf{k} \cdot \mathbf{x}} d\mathbf{k}^2, \quad (2.12c)$$

in which  $\mathbf{V}$  denotes linear velocity vector and  $\bar{\mathbf{V}}$  its magnitude in the envelope-type form,  $\mathbf{u}^{(1)}(\mathbf{x}, z, t)$  and  $w^{(1)}(\mathbf{x}, z, t)$  denote linear velocity vector in the horizontal plane and vertical component, respectively,  $\bar{\mathbf{u}}$  and  $\bar{w}$  are their corresponding magnitude in the envelope-type form, and  $c_p(\mathbf{k}) = \sqrt{g/|\mathbf{k}|}$  denotes the phase velocity of wave  $\mathbf{k}$ . It is worth noting that (2.12c) is fundamental to the theory presented in this paper due to the following two aspects; it allows for a nonlinear evolution equation for  $A$  being expressed in terms of  $\bar{\mathbf{V}}$  and it also facilitates simple numerical implementations of the evolution equation for  $A$  presented in §4.

Using the definition of linear velocity potential  $\Phi^{(1)}$  and following similar procedures that lead to the first-order evolution equation (2.8a), it is understood the following identities hold (to the first order in wave steepness)

$$\bar{\mathbf{V}} = \nabla_3 B + \mathbf{k}_0^{(3d)} B \quad \text{and} \quad \partial_t \begin{bmatrix} B \\ \bar{\mathbf{u}} \\ \bar{w} \end{bmatrix} + i\omega_0 \mathcal{L} \begin{bmatrix} B \\ \bar{\mathbf{u}} \\ \bar{w} \end{bmatrix} = 0, \quad (2.13a,b)$$

in which  $\mathbf{k}_0^{(3d)} = [i\mathbf{k}_0, k_0]$  is introduced for convenience.

#### 2.4. Quadratic property of linear waves: velocity head

Before presenting the wave solutions at second and third order in  $\epsilon$ , we introduce a leading-order approximation to the wave velocity head,  $H_v(\mathbf{x}, z, t)$ , defined as

$$H_v(\mathbf{x}, z, t) = \frac{1}{2g} |\nabla_3 \Phi^{(1)}|^2, \quad (2.14)$$

which is correct to  $\mathcal{O}(\epsilon^2)$ . The wave velocity head plays an essential role in the forcing of bound waves at second order and, thus, the nonlinear evolution equation for  $A$  presented in §4. Inserting the envelope-type expression for  $\Phi^{(1)}$  into (2.14) and using (2.12b), we obtain

$$H_v = (H^{(22)} + \text{c.c.}) + H^{(20)} \text{ with} \quad (2.15a)$$

$$H_v^{(22)} = \frac{1}{8g} \bar{\mathbf{V}} \cdot \bar{\mathbf{V}} e^{2k_0 z} e^{2i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)} \quad \text{and} \quad H_v^{(20)} = \frac{1}{4g} \bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^* e^{2k_0 z} \quad (2.15b)$$

in which the asterisk “\*” denotes the complex conjugates, and we note that  $\bar{\mathbf{V}} \cdot \bar{\mathbf{V}} = \bar{u}^2 + \bar{v}^2 + \bar{w}^2$  with  $\bar{u}$  and  $\bar{v}$  denoting the velocity component of  $\bar{\mathbf{u}}$  ( $\bar{\mathbf{u}} = [\bar{u}, \bar{v}]$ ) in the  $x$  and  $y$  direction, respectively. It is noticeable that  $H_v^{(22)}$  varies with the superharmonic factor  $\exp(2i\mathbf{k}_0 \cdot \mathbf{x} - 2i\omega_0 t)$  whereas the mean velocity head  $H_v^{(20)}$  is independent of the carrier wave phase  $\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t$ . Especially for a quasi-monochromatic wave group (or a monochromatic wave) that admits  $\nabla A \ll \mathcal{O}(1)$  (or  $\nabla A = 0$ ), we obtain a leading order approximation to the mean velocity head  $H_v^{(20)}$  given by

$$H_v^{(20)}(\mathbf{x}, z, t) = \frac{1}{g} U_{SD} c_{g0} \quad \text{with} \quad U_{SD} = k_0 \omega_0 |A|^2 e^{2kz} \quad \text{and} \quad c_{g0} = \frac{\omega_0}{2k_0}, \quad (2.16)$$

where  $U_{SD}$  and  $c_{g0}$  denote Stokes drift and the group velocity, respectively. We understand from (2.16) that the mean velocity head  $H_v^{(20)}$  maintains the energy for the propagation of the Stokes drift at group velocity  $c_{g0}$  for a quasi-monochromatic wave group. We show in §3.3.2 that it is also responsible for the generation of the Eulerian return flow at second order.

In contrast, the rapidly varying velocity head, associated with the factor of second harmonic  $\exp(2i\mathbf{k}_0 \cdot \mathbf{x} - 2i\omega_0 t)$ , has been rarely investigated as it is zero for unidirectional waves. In particular, we note

$$\frac{1}{4} \bar{\mathbf{V}} \cdot \bar{\mathbf{V}} e^{2k_0 z} e^{2i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)} + \text{c.c.} = 0, \quad (2.17)$$

which holds for unidirectional waves of a broad bandwidth, as can also be inferred from earlier works e.g. equations (23) and (25) by Dalzell (1999). Due to this, we examine the effects of the rapidly varying velocity head in §3.3.1 and §6.3.1 for multi-directional waves.

### 3. Second-order solutions $\mathcal{O}(\epsilon^2)$

In this section, the solutions for the second-order super- and sub-harmonic waves forced by linear surface waves are presented. Compared with Dalzell (1999), the solutions in this section are as exact but expressed in an envelope-type form. In contrast to Chu & Mei (1971); Davey & Stewartson (1974); Dysthe (1979), the derivations do not rely on a narrow-banded assumption except for §3.3 where leading-order approximations are presented.

#### 3.1. Governing equation and boundary conditions

The velocity potential at second order in wave steepness,  $\Phi^{(2)} = \Phi^{(22)} + \text{c.c.} + \Phi^{(20)}$ , are described by the following boundary value problem (Dalzell 1999; Li *et al.* 2021c)

$$\nabla_3 \Phi^{(2)} = 0 \text{ for } -\infty < z < 0, \quad (3.1a)$$

$$(\partial_{tt} + g\partial_z)\Phi^{(2)} = \underbrace{-\zeta^{(1)} \Gamma_z \Phi^{(1)}}_{=0} - \partial_t (|\nabla_3 \Phi^{(1)}|^2) \text{ for } z = 0, \quad (3.1b)$$

$$\partial_z \Phi^{(2)} = 0 \text{ for } z \rightarrow -\infty. \quad (3.1c)$$

in which  $\Gamma_z$  is defined as  $\Gamma_z = \partial_z \Gamma$ . As highlighted in (3.1b), the first term on the right hand side does not contribute to the forcing on still water surface as it vanishes and, thus, only the second term needs to be considered. Using the second term and (2.15b) we define a forcing term  $S$  for  $z = 0$  as follows

$$S(\mathbf{x}, t) = \left[ -\partial_t (|\nabla_3 \Phi^{(1)}|^2) \right]_{z=0}, \text{ which leads to } S = (S^{(22)} e^{2i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)} + \text{c.c.}) + S^{(20)} \quad (3.2a,b)$$

with

$$S^{(22)} = -2g\partial_t H_v^{(22)}(\mathbf{x}, z, t) e^{-2i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)} \text{ and } S^{(20)} = -2g\partial_t H_v^{(20)}(\mathbf{x}, z, t) \text{ for } z = 0. \quad (3.3a,b)$$

The forcing term  $S$  (or the velocity head on still water surface) is responsible for the forcing of the second-order bound waves. The negative sign on the right hand side of (3.3a,b) implies a flow generated below still water surface, propagating in an opposite direction to the change of the velocity head in time at  $z = 0$ ; a so-called return flow which is known for second-order subharmonic waves.

In order to obtain the solution to boundary value problem (3.1a)–(3.1c) for  $\Phi^{(2)}$ , two different and novel approaches are proposed in this paper and presented in §3.2 and §3.3, respectively. In particular, the first approach is applicable to three-dimensional waves of a broad bandwidth, and the second is an approximate method based on the assumption of a narrow bandwidth with  $\delta_{x,y} \ll 1$ .

### 3.2. Approach I: semi-analytical approach for $\Phi^{(2)}$

Approach I, referred to as a semi-analytical approach, proposes solving boundary value problem (3.1a)–(3.1c) in the Fourier  $\mathbf{k}$  plane by using a pseudospectral and a finite difference method. The boundary value problem for  $\Phi^{(2j)}$  from (3.1a)–(3.1c) is given by

$$\nabla^2 \Phi^{(2j)} = 0 \text{ for } -\infty < z < 0, \Gamma \Phi^{(2j)} = S^{(2j)} \text{ for } z = 0 \text{ and } \partial_z \Phi^{(2j)} = 0 \text{ for } z \rightarrow -\infty, \quad (3.4a,b,c)$$

in which  $j = 0$  and  $j = 2$  denote the subharmonic and superharmonic wave, respectively. The solutions of (3.4a,b,c) can be expressed in a form as follows

$$\Phi^{(2j)}(\mathbf{x}, z, t) = e^{j i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)} \int_{-\infty}^{\infty} \hat{B}^{(2j)}(\mathbf{k}, t) \exp e^{i \mathbf{k} \cdot \mathbf{x} + |\mathbf{k} + j \mathbf{k}_0| z} d\mathbf{k}^2, \quad (3.5)$$

for  $j = 0$  and  $j = 2$ . Substituting (3.5) into (3.4b) leads to the second-order differential equation for  $\hat{B}^{(2j)}$ :

$$(\partial_t - i j \omega_0)^2 \hat{B}^{(2j)} + g |\mathbf{k} + j \mathbf{k}_0| \hat{B}^{(2j)} = 2^{j/2} \hat{S}^{(2j)}(\mathbf{k}, t), \quad (3.6)$$

where  $\hat{S}^{(2j)}$  denotes the Fourier transform for  $S^{(2j)}$  for  $j = 0$  and  $j = 2$ . It is understood that the second-order waves are bound which do not admit homogeneous solution of (3.6) (Phillips 1960; Hasselmann 1962). As a result, equation (3.6) for  $\hat{B}^{(2j)}$  can be solved numerically by a *semi-analytical approach* that proposes evaluating  $S^{(2j)}(\mathbf{x}, t)$  by using a pseudospectral method and obtaining the numerical solution of (3.6) by available time marching methods. The numerical procedures are explained in §4.4. For numerical implementations, prescribed conditions at an initial instant are required for  $\Phi^{(2)}(\mathbf{x}, 0, t_0)$  and  $\partial_t \Phi^{(2)}(\mathbf{x}, 0, t_0)$ . In practice, multiple choices are available to this end, depending on the purpose of wave predictions. For instance, a second-order wavemaker theory based on Schäffer (1996), periodic boundary conditions as in Dommermuth & Yue (1987), the framework by Bonnefoy *et al.* (2006) for the waves generation in a numerical wave tank, stationary waves based on Dalzell (1999), and approximate initial conditions as presented in §3.3. If the waves generation is based on linear wave theory,  $\Phi^{(2)}(\mathbf{x}, 0, t_0) = 0$  and  $\partial_t \Phi^{(2)}(\mathbf{x}, 0, t_0) = 0$  are implied, leading to inconsistency and additional generation of spurious waves (Schäffer 1996). The semi-analytical approach leads to improved computational efficiency compared with the analytical method by Dalzell (1999) based on Fourier integrals. For  $N$  Fourier modes, the semi-analytical approach requires computational operations at  $\mathcal{O}(N \ln(N))$ , whereas Dalzell (1999) requires  $\mathcal{O}(N^2)$ . The validation of the semi-analytical approach is presented in §6.1.

### 3.3. Approach II: an approximate method for $\Phi^{(2)}$

With a narrow-banded assumption, this section presents an approximate method for  $\Phi^{(22)}$  (§3.3.1) and  $\Phi^{(20)}$  (§3.3.2). For identifying the correct order in bandwidth, we assume small and non-dimensional bandwidth scaling parameters  $\delta_x$  and  $\delta_y$  that denote the bandwidth in the  $x$  and  $y$  direction, respectively. The order of accuracy of the approximate solutions is indicated by the product of  $\epsilon^2$  and  $\delta_{x,y}^j$  with  $j$  non-zero integers.

With a narrow-banded assumption, we seek the solutions of (3.1a)–(3.1c) for unknown  $B^{(22)}$  and  $\Phi^{(20)}$  in a form of power series in bandwidth as follows,

$$B^{(22)} = \delta_y B_{\approx,1}^{(22)} + \delta_{x,y}^2 B_{\approx,2}^{(22)} + \dots \text{ and } \Phi^{(20)} = \delta_x \Phi_{\approx,1}^{(20)} + \delta_{x,y}^2 \Phi_{\approx,2}^{(20)} + \dots \quad (3.7a,b)$$

in which the subscripts ‘ $\approx, j$ ’ denote an approximation at  $\mathcal{O}(\epsilon^2 \delta_{x,y}^j)$ , and we consider leading-order approximations up to  $j = 2$ .

### 3.3.1. Forcing of second-order superharmonic waves by multi-directional waves

It is understood that the forcing term on the right hand side of (3.1b) (see also (2.17)) for  $\Phi^{(22)}$  for unidirectional waves is zero, which suggests  $\Phi^{(22)}(\mathbf{x}, z, t) = 0$ . Hence, the derivations for  $\Phi^{(22)}$  are only needed for multi-directional waves. Based on the boundary value problem described by (3.1a)–(3.1c), the forcing equation of the superharmonic bound waves for  $B^{(22)}$  on still water surface reads

$$(\partial_{tt} - 4i\omega_0 \partial_t - 4\omega_0^2 + g\partial_z + 2gk_0)B^{(22)} = i\omega_0 \bar{\mathbf{V}} \cdot \bar{\mathbf{V}} - \bar{\mathbf{V}} \cdot \partial_t \bar{\mathbf{V}} \text{ for } z = 0, \quad (3.8)$$

in which the first term on the right hand side is at  $\mathcal{O}(\epsilon^2 \delta_y)$  and the second term at  $\mathcal{O}(\epsilon^2 \delta_y \delta_x)$ . Inserting the expanded form (3.7a) for  $B^{(22)}$  into (3.8) and the Laplace equation for  $\Phi^{(22)}$  and following the conventional procedures for perturbed solutions, we obtain

$$B_{\approx,1}^{(22)} = \int_{-\infty}^{\infty} \frac{i\omega_0 \hat{V}_{sq}^{(22)}(\mathbf{k}, t)}{g|\mathbf{k} + 2\mathbf{k}_0| - 4\omega_0^2} e^{i\mathbf{k} \cdot \mathbf{x} + (|\mathbf{k} + 2\mathbf{k}_0| - 2k_0)z} d\mathbf{k}^2 \text{ and} \quad (3.9a)$$

$$B_{\approx,2}^{(22)} = \int_{-\infty}^{\infty} \frac{-4\omega_0^2 \partial_t \hat{V}_{sq}^{(22)} + (4\omega_0^2 - g|\mathbf{k} + 2\mathbf{k}_0|) \hat{V}_{sq,t}^{(22)}(\mathbf{k}, t)}{(g|\mathbf{k} + 2\mathbf{k}_0| - 4\omega_0^2)^2} e^{i\mathbf{k} \cdot \mathbf{x} + (|\mathbf{k} + 2\mathbf{k}_0| - 2k_0)z} d\mathbf{k}^2, \quad (3.9b)$$

where  $\hat{V}_{sq}^{(22)}$  and  $\hat{V}_{sq,t}^{(22)}$  denote the Fourier transform for  $\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}$  and  $\bar{\mathbf{V}} \cdot \partial_t \bar{\mathbf{V}}$  with respect to  $\mathbf{x}$  for  $z = 0$ , respectively.

### 3.3.2. Potential for the second-order mean flows in two and three dimensions

Similar to the second-order superharmonic waves, the forcing equation of the subharmonic bound waves at second order for  $\Phi^{(20)}$  on still water surface (cf. (3.1b)) reads

$$\partial_{tt} \Phi^{(20)} + g\partial_z \Phi^{(20)} = S^{(20)} \text{ for } z = 0, \quad (3.10)$$

where  $S^{(20)}$  is defined by (3.3b) and  $S^{(20)} \sim \mathcal{O}(\epsilon^2 \delta_x)$ . With a narrow-banded assumption and following the same procedures as for  $B^{(22)}$ , we obtain

$$\Phi_{\approx,1}^{(20)} = \int_{-\infty}^{\infty} \frac{\hat{S}^{(20)}}{g|\mathbf{k}|} e^{i\mathbf{k} \cdot \mathbf{x} + |\mathbf{k}|z} d\mathbf{k}^2 \text{ and } \Phi_{\approx,2}^{(20)} = \int_{-\infty}^{\infty} \frac{-2\hat{S}_t^{(20)}}{(g|\mathbf{k}|)^2} e^{i\mathbf{k} \cdot \mathbf{x} + |\mathbf{k}|z} d\mathbf{k}^2, \quad (3.11a,b)$$

in which  $\hat{S}^{(20)}(\mathbf{k}, t)$  and  $\hat{S}_t^{(20)}(\mathbf{k}, t)$  denote the Fourier transform for  $S^{(20)}$  and  $\partial_t S^{(20)}$  with respect to  $\mathbf{x}$ , respectively.

## 3.4. Wave elevation $\zeta^{(2)}$ and velocity $\mathbf{V}^{(2)}$ at second order

With second-order potential  $\Phi^{(2)}$  given by the semi-analytical approach presented in §3.2 or the approximate method presented in §3.3, the surface elevation at second order is obtained from

$$\zeta^{(2)}(\mathbf{x}, t) = -\frac{1}{g} \left[ \partial_t \Phi^{(2)} + \zeta^{(1)} \partial_{tz} \Phi^{(1)} + \frac{1}{2} |\nabla_3 \Phi^{(1)}|^2 \right] \text{ for } z = 0, \quad (3.12)$$

where  $\Phi^{(2)} = \Phi^{(20)}$  for unidirectional waves and  $\Phi^{(2)} = \Phi^{(20)} + \Phi^{(22)} + \text{c.c.}$  for multidirectional waves. Inserting the envelope-type expression for  $\Phi^{(1)}$  and  $\Phi^{(2)}$ , we arrive at

$$\zeta^{(2)} = \zeta^{(20)} + (\zeta^{(22)} + \text{c.c.}) \quad \text{with} \quad (3.13a)$$

$$\zeta^{(22)} = -\frac{1}{g} \left[ \partial_t \Phi^{(22)} + \frac{1}{4} \left( A(\partial_t - i\omega_0)\bar{w} + \frac{1}{2} \bar{\mathbf{V}} \cdot \bar{\mathbf{V}} \right) e^{2i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)} \right] \quad \text{and} \quad (3.13b)$$

$$\zeta^{(20)} = -\frac{1}{g} \left[ \partial_t \Phi^{(20)} + \frac{1}{4} \bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^* + \frac{1}{4} (A^*(\partial_t - i\omega_0)\bar{w} + \text{c.c.}) \right] \quad \text{for } z = 0. \quad (3.13c)$$

With  $\Phi^{(22)} = 0$  in mind for unidirectional waves, (3.13b) gives an exact expression for the elevation for second-order superharmonic waves that depend only on linear modulational parameters e.g.  $A$  and  $\bar{\mathbf{V}}$ .

Similarly, the velocity at second order,  $\mathbf{V}^{(2)}(\mathbf{x}, z, t)$ , is given by

$$\mathbf{V}^{(2)} = \mathbf{V}^{(20)}(\mathbf{x}, z, t) + [\bar{\mathbf{V}}^{(22)}(\mathbf{x}, z, t) e^{2k_0 z} e^{2i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)} + \text{c.c.}] \quad \text{where} \quad (3.14a)$$

$$\mathbf{V}^{(20)} = \nabla \Phi^{(20)} \quad \text{and} \quad \bar{\mathbf{V}}^{(22)} = \frac{1}{2} (2\mathbf{k}_0^{(3d)} + \nabla_3) B^{(22)}. \quad (3.14b)$$

#### 4. A new nonlinear Schrödinger equation for $A$ ( $\mathcal{O}(\epsilon^3)$ )

In this section, a new NLSE for linear wave envelope  $A$  is presented in §4.1 for multidirectional waves of a broad bandwidth. Based on a narrow-banded assumption, leading order approximations to the new NLSE are derived in §4.2 and how this new evolution equation recovers existing NLSEs is explained in §4.3. The numerical implementations of the new NLSE are presented in §4.4.

##### 4.1. An evolution equation correct to $\mathcal{O}(\epsilon^3)$ for waves of a broad bandwidth

Collecting the terms at third order in wave steepness in (2.2), we obtain

$$(\partial_{tt} + g\partial_z)\Phi^{(3)} = -Q^{(3)}(\mathbf{x}, z, t) \quad \text{for } z = 0, \quad (4.1)$$

where the forcing term  $Q^{(3)}$  is given by

$$Q^{(3)} = \underbrace{\zeta^{(2)} \Gamma_z \Phi^{(1)} + \frac{1}{2} (\zeta^{(1)})^2 \Gamma_{zz} \Phi^{(1)}}_{=0} + \zeta^{(1)} \partial_z \left[ \Gamma \Phi^{(2)} + \partial_t (|\nabla_3 \Phi^{(1)}|^2) \right] + 2\partial_t (\nabla_3 \Phi^{(1)} \cdot \nabla_3 \Phi^{(2)}) + \frac{1}{2} \nabla_3 \Phi^{(1)} \cdot \nabla_3 (|\nabla_3 \Phi^{(1)}|^2), \quad (4.2)$$

with  $\Gamma_{zz} = \partial_z \Gamma_z$ . As indicated in (4.2), one would show that  $\Gamma_z \Phi^{(1)} = 0$  and  $\Gamma_{zz} \Phi^{(1)} = 0$ , which are not necessarily so for a finite depth. The terms in the square bracket on the right hand side of (4.2) can be simplified due to (3.1b). In particular, it is noticed that the following identity holds

$$\partial_z \left[ \Gamma \Phi^{(2)} + \partial_t (|\nabla_3 \Phi^{(1)}|^2) \right] = k_0 \partial_t (\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*) \quad \text{for } z = 0. \quad (4.3)$$

Inserting the harmonic expansion for  $\Phi^{(1)}$  and  $\Phi^{(2)}$  into (4.2) leads to  $Q^{(3)}$  being expressed in a form as follows

$$Q^{(3)}(\mathbf{x}, t) = Q^{(31)}(\mathbf{x}, t) + Q^{(33)}(\mathbf{x}, t), \quad (4.4)$$

where  $Q^{(31)}$  denotes the forcing term with a factor  $\exp(i\mathbf{k}_0 \cdot \mathbf{x} - i\omega_0 t)$  and  $Q^{(33)}$  the term with a factor  $\exp(3i\mathbf{k}_0 \cdot \mathbf{x} - 3i\omega_0 t)$ . Leaving the expression for  $Q^{(33)}$  in §5 and focusing

on  $Q^{(31)}$  in this section, we obtain (see details in Appendix A)

$$Q^{(31)} = \bar{Q}^{(31)}(\mathbf{x}, z, t)e^{i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)} + \text{c.c. for } z = 0, \text{ with } \bar{Q}^{(31)} = \bar{Q}_{\text{all}}^{(31)} + \bar{Q}_{\text{dir}}^{(31)}, \quad (4.5a)$$

$$\bar{Q}_{\text{all}}^{(31)} = (-i\omega_0 + \partial_t) \left[ \nabla_3 \Phi^{(20)} \cdot \bar{\mathbf{V}} \right] + \frac{1}{8} \left[ \nabla_3 (\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*) \right] \cdot \bar{\mathbf{V}} + \frac{1}{4} (k_0 \bar{w} + 2k_0 A \partial_t) (\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*), \quad (4.5b)$$

$$\bar{Q}_{\text{dir}}^{(31)} = (-i\omega_0 + \partial_t) \left\{ \frac{1}{2} \left[ (2\mathbf{k}_0^{(3d)} + \nabla_3) B^{(22)} \right] \cdot \bar{\mathbf{V}}^* \right\} + \frac{1}{16} \left[ (2\mathbf{k}_0^{(3d)} + \nabla_3) (|\bar{\mathbf{V}}|^2) \right] \cdot \bar{\mathbf{V}}^*, \quad (4.5c)$$

in which the subscripts ‘all’ and ‘dir’ are used to distinguish two different cases; the former means the parameter expressed in terms that are non-constant for both unidirectional and multi-unidirectional waves, and ‘dir’ means the parameter composed of terms that are non-zero only for directional waves. Hence,  $\bar{Q}^{(31)}$  admits a much simpler expression for waves considered in two dimensions or for long-crested waves due to  $\bar{Q}_{\text{dir}}^{(31)} = 0$ .

Physically,  $Q^{(31)}$  would lead to secular solutions that should be removed (see the discussion on page 376 in Madsen & Fuhrman (2006)). In order to achieve this, a conventional approach is to introduce a nonlinear amplitude-dependent frequency  $\omega_2$  ( $\sim \mathcal{O}(\epsilon^2)$ ) for the first-order wave fields through replacing factor  $\exp[i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)]$  in (2.5a) and (2.12) with  $\exp[i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t) - i\epsilon^2 \omega_2 t]$ . Thereby, it leads to additional contribution of the updated first-order potential  $\Phi^{(1)}$  to (4.1) at  $\mathcal{O}(\epsilon^3)$ . Especially arising from  $\partial_{tt} \Phi^{(1)}$ , we obtain an additional term associated with first harmonic at  $\mathcal{O}(\epsilon^3)$  expressed as  $-i\omega_2 (\partial_t - i\omega_0) B \exp(i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t) - i\epsilon^2 \omega_2 t) + \text{c.c.}$ . Mathematically, the secular solutions can be removed if the sum of this additional term and  $\bar{Q}^{(31)}$  equal zero. Hence, we arrive at

$$i\omega_2 (\partial_t - i\omega_0) B = \bar{Q}^{(31)} \text{ for } z = 0. \quad (4.6)$$

Noticing that  $\zeta^{(1)} = -\partial_t \Phi^{(1)} / g$  for  $z = 0$  which gives  $(\partial_t - i\omega_0) B = -gA$ , substituting this expression for  $B$  into (4.6) leads to the equation for  $A$  as follows

$$-ig\omega_2 A = \bar{Q}^{(31)} \text{ for } z = 0. \quad (4.7)$$

The evolution equation for  $A'$  with  $A' = A \exp(-i\omega_2 t)$ , correct to third order in wave steepness, can now be expressed as

$$\partial_t A' + i\omega_0 \mathcal{L} A' + i\omega_2 A' = 0, \quad (4.8)$$

Inserting (4.7) into (4.8) and omitting the prime leads to

$$\partial_t A + i\omega_0 \mathcal{L} A + \mathcal{N}(\mathbf{x}, z, t) = 0 \text{ for } z = 0, \text{ with} \quad (4.9)$$

where  $\mathcal{N} = -\bar{Q}^{(31)} / g$  denotes the nonlinear term, given by

$$\mathcal{N} = \mathcal{N}_{\text{all}} + \mathcal{N}_{\text{dir}} \text{ with} \quad (4.10a)$$

$$\mathcal{N}_{\text{all}} = \frac{1}{g} (i\omega_0 - \partial_t) \left[ \mathbf{V}^{(20)} \cdot \bar{\mathbf{V}} \right] - \frac{1}{8g} \left[ 2k_0 \bar{w} + \bar{\mathbf{V}} \cdot \nabla_3 + 4k_0 A \partial_t \right] (\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*), \quad (4.10b)$$

$$\mathcal{N}_{\text{dir}} = \frac{1}{g} (i\omega_0 - \partial_t) \left( \bar{\mathbf{V}}^{(22)} \cdot \bar{\mathbf{V}}^* \right) - \frac{1}{16g} \left[ (2\mathbf{k}_0^{(3d)} + \nabla_3) (|\bar{\mathbf{V}}|^2) \right] \cdot \bar{\mathbf{V}}^*. \quad (4.10c)$$

Equation (4.9) denotes the *new nonlinear Schrödinger equation* that describes the evolution of linear wave envelope  $A$  in time and space. In addition to the linear term associated with the time derivative  $\partial_t(\dots)$ , equation (4.9) is composed of a term that denotes linear dispersion relation due to  $i\omega_0 \mathcal{L}$  and a nonlinear term denoted by  $\mathcal{N}$  that is correct to



$\mathcal{O}(\epsilon^3)$ . For (4.9), no assumptions associated with the bandwidth have been introduced. Thereby, it has relaxed the limitation arising from a narrow-banded assumption on which most NLSEs are based, such as the NLSEs by Dysthe (1979), Trulsen & Dysthe (1996), and Trulsen *et al.* (2000); i.e., the new NLSE should permit the prediction of waves with bandwidth  $\delta_{x,y} > 1$ . The implementation of the new NLSE for the wave envelope,  $A$ , of the first-harmonic elevation requires the evaluation of linear term  $\partial_t A$  in the Fourier plane based on (2.8c) and the evaluation of the linear envelope (vector) of both the linear and second-order velocity for  $\mathcal{N}$ , which will be explained in detail in §4.4.

Extensions of the new NLSE (i.e. (4.9)) to more general cases are simple and straightforward. For example, the new NLSE can be extended to consider a finite water depth following the derivations in this paper, except that one would expect that the nonlinear term  $\mathcal{N}$  may contain a few more terms that introduce a negligible additional computational cost. Following Dias *et al.* (2008) and Kharif *et al.* (2010), the new NLSE would allow for weakly damped/forced free surface flows through adding the viscous effects in the dynamic and kinematic boundary condition (cf. Eqs. (35, 37) by Dias *et al.* (2008)) at the water surface. Moreover, despite the new NLSE is formulated in the framework of potential flow theory for simplicity, it is essentially an equation associated with linear envelope  $A$ , linear velocity  $\bar{\mathbf{V}}$ , and second-order velocity  $\bar{\mathbf{V}}^{(2)}$ . This suggests the weak effects due to viscosity and vorticity in a sub-surface layer can be easily incorporated, allowing for waves interacting with ambient environments e.g. turbulence and background rotational flows (cf. McWilliams *et al.* (2004)). These aspects will be considered in future work.

#### 4.2. Leading-order approximations to the new NLSE

Approximations to (4.9) correct to  $\mathcal{O}(\epsilon^3 \delta_{x,y}^j)$ , with  $j = 0, 1, 2$ , and  $j = 3$ , are presented in this section with a narrow-banded assumption, i.e.  $\mathcal{O}(\delta) \ll 1$ . In particular, we assume that the bandwidths both in the longitudinal ( $x$ ) direction and the transverse direction to the propagation of the carrier wave are small and that  $\mathcal{O}(\delta_y) \sim \mathcal{O}(\delta_x^2)$ . Small directionally spread is, hence, implied. Based on previous papers (e.g. Chu & Mei (1971); Li *et al.* (2021c)), the order of magnitude associated with the relevant terms in (4.5) is assumed as follows

$$[\bar{\mathbf{V}}, \bar{\mathbf{V}}^*] \sim \mathcal{O}(\epsilon), \quad [|\bar{\mathbf{V}}|^2, \bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*] \sim \mathcal{O}(\epsilon^2 \delta_y^2, \epsilon^2), \quad \Phi^{(20)} \sim \mathcal{O}(\epsilon^2 \delta_x), \quad \text{and} \quad B^{(22)} \sim \mathcal{O}(\epsilon^2 \delta_y^2). \quad (4.11a,b,c,d)$$

Moreover, it is understood that the derivative operators ( $\partial_t, \nabla_3$ ) on the terms in (4.11) would lead to an order of magnitude higher in bandwidth  $\delta_{x,y}$ . With the right orders of magnitude in mind, collecting the terms up to a particular order in  $\epsilon$  and  $\delta$  based on (4.5) and (4.9b) and thus, leading order approximations to (4.9) are obtained. In particular, they are expressed in a form as follows

$$\partial_t A + i\omega_0 \mathcal{L}A + \mathcal{N}_{\approx}^{(3,j)} = \mathcal{O}(\epsilon^3 \delta_{x,y}^{j+1}) \quad \text{for } z = 0, \quad (4.12)$$

where  $\mathcal{N}_{\approx}^{(3,j)}$  denotes an approximation to  $\mathcal{N}$  correct to  $\mathcal{O}(\epsilon^3 \delta_{x,y}^j)$ , with  $j = 0, 1, 2$ , and 3, given by,

$$\mathcal{N}_{\approx}^{(3,0)} = -\frac{1}{4g}(\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*)k_0\bar{w}, \quad (4.13a)$$

$$\mathcal{N}_{\approx}^{(3,1)} = \mathcal{N}_{\approx}^{(3,0)} - \frac{1}{8g}[\nabla_3(\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*)] \cdot \bar{\mathbf{V}} - \frac{k_0}{2g}A\partial_t(\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*), \quad (4.13b)$$

$$\mathcal{N}_{\approx}^{(3,2)} = \mathcal{N}_{\approx}^{(3,1)} + \frac{i\omega_0}{g}(\nabla_3\Phi_{\approx,1}^{(20)} \cdot \bar{\mathbf{V}}) + \mathcal{N}_{\approx,\text{dir}}^{(3,2)}, \quad (4.13c)$$

$$\mathcal{N}_{\approx}^{(3,3)} = \mathcal{N}_{\approx}^{(3,2)} + \frac{i\omega_0}{g}(\nabla_3\Phi_{\approx,2}^{(20)} \cdot \bar{\mathbf{V}}) - \frac{1}{g}\partial_t(\nabla_3\Phi_{\approx,1}^{(20)} \cdot \bar{\mathbf{V}}) + \mathcal{N}_{\approx,\text{dir}}^{(3,3)}, \quad (4.13d)$$

$$\mathcal{N}_{\approx}^{(3,4)} = \mathcal{N}, \quad (4.13e)$$

where (4.13e) denotes that  $\mathcal{N}^{(3,4)}$  has an identical expression as  $\mathcal{N}$ . As noted earlier, the subscript ‘dir’ means the parameter being composed of nonlinear terms that are non-zero only for multi-directional waves, given by

$$\mathcal{N}_{\approx,\text{dir}}^{(3,2)} = \frac{1}{g}\left(B_{\approx,1}^{(22)} - \frac{1}{8}\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}\right)(i\mathbf{k}_0 \cdot \bar{\mathbf{u}}^* + k_0\bar{w}^*), \quad (4.13f)$$

$$\begin{aligned} \mathcal{N}_{\approx,\text{dir}}^{(3,3)} = & -\frac{1}{16g}[\nabla_3(|\bar{\mathbf{V}}|^2)] \cdot \bar{\mathbf{V}}^* - \frac{1}{g}\partial_t[(i\mathbf{k}_0 \cdot \bar{\mathbf{u}}^* + k_0\bar{w}^*)B_{\approx,1}^{(22)}] + \\ & \frac{1}{g}B_{\approx,2}^{(22)}(i\mathbf{k}_0 \cdot \bar{\mathbf{u}}^* + k_0\bar{w}^*). \end{aligned} \quad (4.13g)$$

Equation (4.13b) denotes that a leading order approximation to  $\mathcal{N}$ , correct to  $\mathcal{O}(\epsilon^3\delta)$ , does not require an evaluation of the second-order mean potential. Equation (4.13c) suggests that the approximations to  $\Phi^{(20)}$  and  $B^{(22)}$  that are presented in §3.3 are sufficient to give a leading order approximation to (4.9), correct up to  $\mathcal{O}(\epsilon^3\delta^3)$ . Any approximations correct to an order higher than  $\mathcal{O}(\epsilon^3\delta^2)$  would require the same computational cost as  $\mathcal{N}$  because the approximations  $\Phi_{\approx,2}^{(20)}$  and  $B_{\approx,2}^{(22)}$  do not lead to enhanced computational efficiency, compared with solving for  $\Phi^{(2)}$  by the semi-analytical method.

### 4.3. Comparisons to two reduced-form equations for $A$

Two limiting cases are considered in this section in order to demonstrate that the new NLSE can recover two reduced-form equations for  $A$ , including the NLSE for the evolution of a train of Stokes wave (§4.3.1) and the NLSEs by Dysthe (1979); Trulsen & Dysthe (1996) and Trulsen *et al.* (2000) with a narrow-banded assumption (§4.3.2).

#### 4.3.1. Uniform Stokes waves

For uniform Stokes waves on an infinite depth, that denote  $A$  and  $\bar{\mathbf{V}}$  having no slow modulation in  $\mathbf{x}$  and  $z$ , the new NLSE leads to

$$\mathcal{N} = \mathcal{N}^{(3,0)} \text{ and hence, } \mathcal{N} = \frac{1}{2}ik_0^2\omega_0|A|^2A. \quad (4.14)$$

Introducing  $\epsilon = k_0^2|A|^2$  and  $\alpha_1 = \frac{1}{2}$  for Stokes waves, we obtain the evolution equation for  $A$  as follows

$$\partial_t A + i\alpha_1\epsilon^2\omega_0 A = 0, \quad (4.15)$$

which agrees with the classic NLSE for a train of Stokes waves as presented in many papers e.g. Benjamin & Feir (1967); Zakharov (1968). Equation (4.15) will be used for the analysis of sideband instability of Stokes waves presented in §6.2.

4.3.2. Waves of a narrow bandwidth ( $\mathcal{O}(\epsilon^3\delta_{x,y})$ )

In order to recover the previous versions of NLSEs by Dysthe (1979); Trulsen & Dysthe (1996); Trulsen *et al.* (2000) from the NLSE (4.9), two aspects are addressed. First, the term associated with linear dispersion relation  $i\omega_0\mathcal{L}$  is identical to Eqs. (12) and (13) by Trulsen *et al.* (2000) using the dimensional versions. It has been demonstrated in Trulsen *et al.* (2000) how this term can recover the terms associated with linear dispersion relation derived by Trulsen & Dysthe (1996) and Dysthe (1979). Second, the nonlinear term in the NLSEs by Trulsen *et al.* (2000) and Trulsen & Dysthe (1996) are directly obtained from the Dysthe equation by Dysthe (1979). Hence, we only have to demonstrate that  $\mathcal{N}$  can recover the nonlinear terms in the Dysthe equation. To this end, we start with (4.9b) for unidirectional waves as the Dysthe equation has neglected the contribution from  $\Phi^{(22)}$  (which is non-zero only for multi-directional waves). Moreover, a narrow-banded assumption as in Dysthe (1979) is introduced; i.e.  $\mathcal{O}(\delta_{x,y}) \sim \mathcal{O}(\epsilon)$  is assumed in this section.

With a narrow-banded assumption, the velocity (magnitude)  $\bar{\mathbf{V}}$  correct to  $\mathcal{O}(\epsilon\delta_{x,y})$  reads (cf. Eqs. (13.2.21) and (13.2.30) by Mei *et al.* (2005) for deepwater waves)

$$\bar{\mathbf{V}}(\mathbf{x}, z, t) = \frac{-ig}{\omega_0} \left[ \mathbf{k}_0^{(3d)} A + \begin{pmatrix} -i(\mathbf{k}_0 z + \mathbf{e}_x) \nabla A \\ -i(k_0 z + 1) \partial_x A \end{pmatrix} \right] + \mathcal{O}(\epsilon\delta^2), \quad (4.16)$$

in which  $\mathbf{e}_x = [1, 0]$  denotes a unit vector along the  $x$  direction in the horizontal plane. Inserting (4.16) into (4.9b) leads to a leading order approximation to  $g\mathcal{N}$  as follows (see Appendix A.2 for details)

$$\Gamma_{\text{Dysthe}} C \equiv -g\mathcal{N} = 4k_0^4 C|C|^2 + 8ik_0^3 C(C\partial_x C^* - C^*\partial_x C) - 4ik_0 C\partial_x(C^*C) + 2k_0\omega_0 C(\partial_x\Phi^{(20)} - \partial_z\Phi^{(20)}), \quad (4.17)$$

where  $C$  is defined as  $C \equiv B/2 = -i\omega_0 A/(2k_0)$ . The subscript ‘Dysthe’ means that the parameter corresponds to the ‘ $\Gamma$ ’ by Dysthe (1979); i.e. (4.17) recovers ‘ $\Gamma$ ’ on the right hand side of equation (2.17) by Dysthe (1979), while observing the following two aspects. Firstly, the notation  $C$  defined here is denoted by ‘ $A$ ’ in Dysthe (1979). Secondly, there is one misprint in the equation for ‘ $\Gamma$ ’, as also pointed out by Brinch-Nielsen & Jonsson (1986) (cf. page 461) and Janssen (1983); i.e. the second term in the expression ‘ $3ik(\dots)$ ’ for ‘ $\Gamma$ ’ should read  $8ik(\dots)$ . For completeness, the detailed derivations for (4.17) are presented in appendix A.2. The modified NLSE by Trulsen *et al.* (2000) (i.e. equations (19-22) therein) is essentially a leading order approximation to (4.9), correct to  $\mathcal{O}(\epsilon^3\delta_{x,y})$ .

## 4.4. Numerical implementation

Figure 1 shows the detailed procedures of the numerical implementation of the new NLSE (4.9) for  $A$ , including the solution of the second-order equation, (3.6), for  $B^{(2j)}$  and the evaluation of  $\mathcal{N}$  by a pseudospectral method and a finite-difference method. We remark here that it is conventional to solve a NLSE numerically for  $A$  in a moving coordinate system through the transforms  $\xi = x - c_{g0}t$  and  $\eta = t$ , which would also follow similar numerical procedures as follows but for  $A'(\xi, y, \eta)$  with  $A'(\xi, y, \eta) = A(\mathbf{x}, t)$  instead.

At any temporal step  $t_n = t_0 + n\Delta t$  with  $t_0$  denoting the initial time, it is understood that the value of  $A(\mathbf{x}, t_n)$  for all  $\mathbf{x}_s < \mathbf{x} < \mathbf{x}_e$ , with  $\mathbf{x}_s$  and  $\mathbf{x}_e$  denoting the start and end position vector of a chosen spatial domain, respectively, is given from earlier computations. The spatial domain is discretised by  $2N_x \times 2N_y$  points at which  $A(\mathbf{x}, t_n)$  is known, with  $2N_x$  and  $2N_y$  denoting the total number of points in  $x$  and  $y$  direction, respectively. Transforming  $A$  to the discretised Fourier space using a fast Fourier transform (FFT)

Numerical implementation of the extended NLS equation, Eq. (4.9), for  $A(\mathbf{x}, t)$

$$\begin{aligned}
 t_n &= t_0 + n \Delta t & \hat{A}(\mathbf{k}, t) &= F\{A(\mathbf{x}, t)\} \\
 t_{n+0.5} &= t_0 + (n + \frac{1}{2})\Delta t & \widetilde{A}(\mathbf{x}, t_{n+0.5}) &= A(\mathbf{x}, t_n) - \frac{1}{2} \Delta t \mathcal{N}(A_{t_n}, \bar{\mathbf{V}}_{t_n}, \bar{\mathbf{V}}_{t_n}^{(20)}, \bar{\mathbf{V}}_{t_n}^{(22)}) \\
 t_{n+1} &= t_0 + (n+1)\Delta t & \widetilde{\widetilde{A}}(\mathbf{x}, t_{n+1}) &= \widetilde{A}(\mathbf{x}, t_{n+0.5}) - \Delta t \mathcal{N}(A_{t_{n+0.5}}, \bar{\mathbf{V}}_{t_{n+0.5}}, \bar{\mathbf{V}}_{t_{n+0.5}}^{(20)}, \bar{\mathbf{V}}_{t_{n+0.5}}^{(22)}) \\
 & & A(\mathbf{x}, t_{n+1}) &= F^{-1}\{F[\widetilde{\widetilde{A}}(\mathbf{x}, t_{n+1})]\exp[-i(\omega(\mathbf{k}+\mathbf{k}_0) - \omega_0)\Delta t]\}
 \end{aligned}$$

How to evaluate  $\mathcal{N}$  is organised in a form of three consecutive steps highlighted with three different colors

Step I: obtaining linear terms based on  $\hat{A}(\mathbf{k}, t)$  using iFFTs based on Eq. (4.19)

$$[\partial_t \bar{\mathbf{V}}, \partial_x \bar{\mathbf{V}}, \partial_y \bar{\mathbf{V}}, \partial_z \bar{\mathbf{V}}] = F^{-1}\{-i(\omega(\mathbf{k}+\mathbf{k}_0) - \omega_0)\hat{\mathbf{V}}, ik_x \hat{\mathbf{V}}, ik_y \hat{\mathbf{V}}, (|\mathbf{k}+\mathbf{k}_0| - k_0)\hat{\mathbf{V}}\}$$

Step II: solving the second-order equations for second-order sub- and super-harmonic potentials numerically

$$\begin{aligned}
 &>> \text{The midpoint method based on Eq. (3.6) for } \hat{B}_{t_{n+0.5}}^{(2j)}, \hat{B}_{t_{n+0.5}}^{(2j)}, \hat{B}_{t_{n+1}}^{(2j)} \text{ and } \hat{B}_{t_{n+1}}^{(2j)} \text{ for } j = 0 \text{ and } 2; \\
 &>> [\partial_t \hat{B}_s^{(2j)}, \partial_x \hat{B}_s^{(2j)}, \partial_y \hat{B}_s^{(2j)}, \partial_z \hat{B}_s^{(2j)}] = F^{-1}\{\hat{B}^{(2j)}, ik_x \hat{B}^{(2j)}, ik_y \hat{B}^{(2j)}, (|\mathbf{k}+j\mathbf{k}_0| - jk_0)\hat{B}^{(2j)}\} \\
 &>> \mathbf{V}^{(2j)} = \nabla_s \hat{B}^{(2j)} + j\mathbf{k}_0^{(3d)} \hat{B}^{(2j)}
 \end{aligned}$$

Step III: evaluation of  $\mathcal{N}_{t_n}$  and  $\mathcal{N}_{t_{n+1}}$  by a pseudospectral method based on Eq. (4.10)

FIGURE 1. Flow diagram of the numerical implementation of the new NLSE for  $A$ , and of the numerical solution of (3.6) for  $B^{(2j)}(\mathbf{x}, z, t)$ . Dot denotes the derivative with respect to time.

(Frigo & Johnson 1998), we obtain

$$\hat{A}(\mathbf{k}, t_n) = \mathcal{F}\{A(\mathbf{x}, t)\}, \quad \text{with } \mathbf{k} = (k_x, k_y), \quad (4.18a)$$

$$k_x = (0, \pm 1, \pm 2, \dots, \pm N_x) \times dk_x, \quad k_y = (0, \pm 1, \pm 2, \dots, \pm N_y) \times dk_y, \quad (4.18b)$$

where  $dk_x$  and  $dk_y$  denote the interval between two adjacent points in the Fourier  $k_x$  and  $k_y$  direction, respectively;  $\mathcal{F}$  denotes a FFT with respect to  $\mathbf{x}$ .

As highlighted in grey in figure 1 with the known value of  $\hat{A}(\mathbf{k}, t_n)$ , the new NLSE is solved numerically for by using a split-step Fourier method (Tappert 1974). The main difference between the numerical procedures presented here and these presented in Lo & Mei (1985), lie in the fact that the evaluation of  $\mathcal{N}$  per time step is obtained differently as  $\mathcal{N}$  has different expressions. The numerical procedures for  $\mathcal{N}$  at  $t = t_n$  are decomposed into three consecutive steps, as explained in the following.

(*Step I*) The first step is to use the known value of  $\hat{A}(\mathbf{k}, t_n)$  to evaluate  $\bar{\mathbf{V}}(\mathbf{x}, 0, t_n)$  and its derivative with respect to  $t$ ,  $x$ ,  $y$ , and  $z$  using an inverse FFT, as highlighted in figure 1 in blue. Let  $\hat{\mathbf{V}}$  be the transformed linear velocity,  $\bar{\mathbf{V}}$ , in the Fourier  $\mathbf{k}$  space. We understand from (2.12) that

$$\hat{\mathbf{V}}(\mathbf{k}, z) = \begin{bmatrix} \mathbf{k} + \mathbf{k}_0 \\ -i|\mathbf{k} + \mathbf{k}_0| \end{bmatrix} c_p(\mathbf{k} + \mathbf{k}_0) \hat{A} e^{|\mathbf{k}_0 + \mathbf{k}|z - k_0 z}, \quad \text{and thus} \quad (4.19a)$$

$$\mathcal{F}\{[\partial_t, \partial_x, \partial_y, \partial_z]\bar{\mathbf{V}}\} = i[-\omega(\mathbf{k} + \mathbf{k}_0) - \omega_0, k_x + k_0, k_y, |\mathbf{k}_0 + \mathbf{k}| - k_0]\hat{\mathbf{V}}(\mathbf{k}, z). \quad (4.19b)$$

Hence,  $\bar{\mathbf{V}}$  and also its derivatives  $[\partial_t, \partial_x, \partial_y, \partial_z]\bar{\mathbf{V}}$  are evaluated for all spatial points through an inverse FFT using (4.19) for  $z = 0$ .

(*Step II*) The second step (highlighted in green in figure 1) is to solve the second-order equations (3.6) for  $B^{(2j)}$  with  $j = 0$  and  $j = 2$  by the semi-analytical approach presented in §3.2. The forcing terms  $S^{(2j)}$  for  $j = 0$  and  $j = 2$  are computed based on (3.3) in the physical plane using  $\partial_t \bar{\mathbf{V}}$ ,  $\bar{\mathbf{V}}$ , and  $\bar{\mathbf{V}}^*$  evaluated in *Step I*. The second-order potentials  $\hat{B}^{(2j)}$  and their derivatives with respect to time,  $\partial_t \hat{B}^{(2j)}$  for  $j = 0$  and  $j = 2$ , with prescribed initial conditions, can be evaluated based on (3.6) by using a time marching

method e.g. the midpoint method as indicated in figure 1. With the potentials computed per time step, second-order velocity at different harmonics,  $\bar{\mathbf{V}}^{(2j)}$  for  $j = 0$  and  $j = 2$ , and their derivatives with respect to  $t$ ,  $x$ ,  $y$ , and  $z$  can be readily obtained by an inverse FFT.

(*Step III*) The third step, as highlighted in purple in figure 1, is to evaluate  $\mathcal{N}$  in the physical space based on (4.9b) using the terms evaluated in *Step I* and *Step II*.

Some remarks on the numerical efficiency of the new NLSE are explained here for two dimensional waves, which allow for a fair comparison with the efficiency of existing NLSEs e.g. the Dysthe equation as explained in Lo & Mei (1985). In the pseudospectral method, the nonlinear terms are evaluated through FFT at each  $\mathbf{x}$  and  $t_n$  before the integration in time is performed. The evaluation requires 20 FFTs (cf. the areas highlighted in blue and green in figure 1). The third-order nonlinear terms are evaluated in the physical domain, and additional eight FFT computations arise from the terms associated with  $\partial_t(\partial_x, \partial_z)B^{(2j)}$  and  $(\partial_x, \partial_z)B^{(2j)}$  due to an inverse FFT (cf. the areas highlighted in green and purple in figure 1). Two FFTs are needed in the linear equation (i.e. the equation in the last line highlighted in grey in figure 1). If  $N$  Fourier modes are used, a total of  $30N \ln(N)$  operations are needed for advancing the solution by one step in time whereas  $10N \ln(N)$  operations are required with NLSEs e.g. the Dysthe equation and modified NLSE (Dysthe 1979; Lo & Mei 1985; Trulsen *et al.* 2000). Although a slightly increased computational cost relative to the previous versions are inevitable as expected, the new NLSE offers much better computational efficiency compared with other methods e.g. solving the Zakharov integral equation that requires roughly  $\mathcal{O}(N^3)$  operations when an explicit time-differencing scheme is used. Moreover, it is worthy noting that the kinematics of the system, i.e. the elevation and velocities at the first and second orders, are obtained at almost no additional cost throughout the numerical implementations of the NLSE.

In contract to the High-Order Spectral (HOS) method which is also based on efficient FFTs, a pseudospectral method and numerical methods for time marching (cf. Dommermuth & Yue (1987); West *et al.* (1987)), the new NLSE has the following main features. First, as aforementioned, the new NLSE allows for a computational domain chosen based on the scaling of the wave envelope (or the maximum of the side bandwidth of a spectrum), in a manner similar to other NLSE-based models. This suggests a smaller number of discrete points than the HOS method can be chosen in a computational domain. It is mainly achieved by the semi-analytical approach for second-order superharmonic bound waves, which are expressed in an envelope-type form. It is equivalent to shifting the spectrum of the second-order superharmonic bound waves by  $2k_0$  towards the origin of the Fourier plane. In practice, the choice of the carrier wave (i.e.  $k_0$ ) allows for flexibility and, therefore, can contribute to reducing the computational cost, in particular for ocean spectra of a long tail. With a reasonable selection of  $k_0$ , the side bandwidth of an asymmetrical spectrum can be much reduced and therefore, a less fine mesh would be allowed for the implementation of the new NLSE. For example, a numerical test for a JONSWAP spectrum (see appendix C) was carried out with  $k_0 = 4k_p$  chosen for computations, where  $k_p$  denotes the spectrum peak. Secondly, the new NLSE separately obtains the wave fields due to free waves and second-order bound waves at still water surface. Last, with proper separation of the perturbed solutions and multiple scales in time (in terms of wave steepness), the author conjectures that one would show that the solution of the third-order HOS method following Onorato *et al.* (2007) (i.e. Eq. (13) and (14) in the paper) can recover the new NLSE for the envelope of the first-harmonic wave elevation. This will be addressed in future work.

## 5. Solutions at third order $\mathcal{O}(\epsilon^3)$

For completeness, this section presents the solutions at third order for both potential  $\Phi^{(33)}$  given in §5.1 and elevation  $\zeta^{(3)}$  in §5.2.

### 5.1. Potential $\Phi^{(33)}$ at third harmonic

Collecting the terms at third order in wave steepness in (2.1)–(2.3) and keeping the terms that have a factor  $\exp(3i\mathbf{k}_0 \cdot \mathbf{x} - 3i\omega_0 t)$ , we obtain the following equations for  $\Phi^{(33)}$

$$\nabla_3^2 \Phi^{(33)} = 0 \text{ for } -\infty < z < 0, \Gamma \Phi^{(33)} = Q^{(33)} \text{ for } z = 0 \text{ and } \partial_z \Phi^{(33)} = 0 \text{ for } z \rightarrow -\infty, \quad (5.1)$$

where the third-order term  $Q^{(33)}(\mathbf{x}, t)$  denotes the forcing of bound waves at third harmonic on still water surface, as noted in §4, expressed as

$$Q^{(33)} = \overline{Q}^{(33)}(\mathbf{x}, z, t) e^{3i\mathbf{k}_0 \cdot \mathbf{x} - 3i\omega_0 t} \text{ for } z = 0 \text{ and} \quad (5.2a)$$

$$\overline{Q}^{(33)} = \frac{1}{2}(\partial_t - 3i\omega_0)[\overline{\mathbf{V}} \cdot (\nabla_3 + 2i\mathbf{k}_0^{(3d)})B^{(22)}] + \frac{1}{16}\overline{\mathbf{V}} \cdot [(\nabla_3 + 2i\mathbf{k}_0^{(3d)})(\overline{\mathbf{V}} \cdot \overline{\mathbf{V}})]. \quad (5.2b)$$

For unidirectional waves, it is understood that both  $B^{(22)}$  and  $(\overline{\mathbf{V}} \cdot \overline{\mathbf{V}})$  have no contribution to the third-order nonlinear terms as they vanish, meaning  $\overline{Q}^{(33)} = 0$  and thus,  $\Phi^{(33)} = 0$ .

The solution of equations (5.1a,b,c) for  $\Phi^{(33)}$  can be obtained in a way similar to the semi-analytical approach for  $\Phi^{(2)}$ , i.e the solution can be obtained by a finite-difference method in the Fourier  $\mathbf{k}$  space based on the forcing equation on still water surface as follows

$$(\partial_t - 3i\omega_0)^2 \hat{B}^{(33)}(\mathbf{k}, z, t) + g|\mathbf{k} + 3\mathbf{k}_0| \hat{B}^{(33)}(\mathbf{k}, z, t) = 2\hat{Q}^{(33)}(\mathbf{k}, z, t) \text{ for } z = 0, \quad (5.3)$$

where, as defined in previous sections,  $(\hat{\dots})$  denotes a Fourier transform to the  $\mathbf{k}$  plane with respect to  $\mathbf{x}$ . Thereby,  $\Phi^{(33)}$  is expressed as

$$\Phi^{(33)} = \frac{1}{2}B^{(33)}e^{3k_0 z}e^{3i\mathbf{k} \cdot \mathbf{x} - 3i\omega_0 t} \text{ with } B^{(33)} = \int_{-\infty}^{\infty} \hat{B}^{(33)}(\mathbf{k}, 0, t)e^{|\mathbf{k} + 3\mathbf{k}_0|z - 3k_0 z}e^{i\mathbf{k} \cdot \mathbf{x}}d\mathbf{k}^2. \quad (5.4a,b)$$

### 5.2. Elevation $\zeta^{(3)}$ at third order

Based on the boundary conditions (2.2a,b) it is understood that the wave elevation at third order can be obtained from (cf. (13.2.3) by Mei *et al.* (2005))

$$\begin{aligned} \zeta^{(3)} = & -\frac{1}{g} \left[ \partial_t \Phi^{(3)} + \zeta^{(1)} \partial_{tz} \Phi^{(2)} + \zeta^{(2)} \partial_{tz} \Phi^{(1)} + \nabla_3 \Phi^{(1)} \cdot \nabla_3 \Phi^{(2)} \right. \\ & \left. + \frac{1}{2}(\zeta^{(1)})^2 \partial_{tzz} \Phi^{(1)} + \frac{1}{2} \zeta^{(1)} \partial_z (\nabla_3 \Phi^{(1)} \cdot \nabla_3 \Phi^{(1)}) \right]. \end{aligned} \quad (5.5)$$

Based on (5.5), it is understood  $\zeta^{(3)}$  can be expressed in a form as follows

$$\zeta^{(3)} = \left( \frac{1}{2}A^{(31)}(\mathbf{x}, t)e^{i(\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)} + \text{c.c.} \right) + \left( \frac{1}{2}A^{(33)}(\mathbf{x}, t)e^{3i\mathbf{k} \cdot \mathbf{x} - 3i\omega_0 t} + \text{c.c.} \right), \quad (5.6)$$

where amplitudes  $A^{(31)}(\mathbf{x}, t)$  and  $A^{(33)}(\mathbf{x}, t)$  are in a form given by, respectively,

$$\begin{aligned}
 A^{(31)} = & -\frac{1}{2g}[(\partial_t - 2i\omega_0)(2k_0 + \partial_z)B^{(22)}]A^* - \frac{1}{g}A\partial_{tz}\Phi^{(20)} - \frac{1}{2g}A^{(22)}[(\partial_t + i\omega_0)\bar{w}^*] \\
 & - \frac{1}{g}\zeta^{(20)}[(\partial_t - i\omega_0)\bar{w}] - \frac{1}{2g}\bar{\mathbf{V}}^* \cdot [(2\mathbf{k}_0^{(3d)} + \nabla_3)B^{(22)}] - \frac{A^2}{8g}(\partial_t + i\omega_0)(k_0 + \partial_z)\bar{w}^* \\
 & - \frac{1}{4g}AA^*(\partial_t - i\omega_0)(k_0 + \partial_z)\bar{w} - \frac{1}{8g}A^*[(\partial_z + 2k_0)(\bar{\mathbf{V}} \cdot \bar{\mathbf{V}})] \\
 & - \frac{1}{4g}A(\partial_z + 2k_0)(\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*) - \frac{1}{g}(\partial_t + i\omega_0\mathcal{L})B, \quad \text{for } z = 0, \tag{5.7}
 \end{aligned}$$

$$\begin{aligned}
 A^{(33)} = & -\frac{1}{g}(\partial_t - 3i\omega_0)B^{(33)} - \frac{1}{2g}[(\partial_t - 2i\omega_0)(2k_0 + \partial_z)B^{(22)}]A - \frac{1}{2g}A^{(22)}[(\partial_t + i\omega_0)\bar{w}] \\
 & - \frac{1}{2g}\bar{\mathbf{V}} \cdot [(2\mathbf{k}_0^{(3d)} + \nabla_3)B^{(22)}] - \frac{A^2}{8g}(\partial_t - i\omega_0)(k_0 + \partial_z)\bar{w} - \frac{1}{8g}A[(\partial_z + 2k_0)(\bar{\mathbf{V}} \cdot \bar{\mathbf{V}})], \tag{5.8}
 \end{aligned}$$

in which the nonlinear terms are obtained in the physical plane using a pseudospectral method. As noted in §2.4 and §3.3.1, both  $B^{(22)}$  and  $\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}$  vanishes for unidirectional waves, under which circumstances the amplitudes at third order embrace a much simpler expressions given by,

$$\begin{aligned}
 A_{\text{uni}}^{(31)} = & -\frac{1}{g}A\partial_{tz}\Phi^{(20)} - \frac{1}{2g}A^{(22)}(\partial_t + i\omega_0)\bar{w}^* - \frac{1}{g}\zeta^{(20)}(\partial_t - i\omega_0)\bar{w} - \frac{A^2}{8g}(\partial_t + i\omega_0)(k_0 + \partial_z)\bar{w}^* \\
 & - \frac{1}{4g}AA^*(\partial_t - i\omega_0)(k_0 + \partial_z)\bar{w} - \frac{1}{4g}A[(\partial_z + 2k_0)(\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*)] - \frac{1}{g}(\partial_t + i\omega_0\mathcal{L})B, \tag{5.9a}
 \end{aligned}$$

$$A_{\text{uni}}^{(33)} = -\frac{A^2}{8g}(\partial_t - i\omega_0)(k_0 + \partial_z)\bar{w} - \frac{1}{2g}A^{(22)}(\partial_t - i\omega_0)\bar{w}. \tag{5.9b}$$

in which the subscript ‘uni’ refers to the cases for unidirectional waves. With a narrow-banded assumption  $\mathcal{O}(\delta) \ll 1$ , we readily obtain from (5.9) (through omitting terms associated with the derivatives with respect to  $t$  and  $z$  and the second-order mean fields and noting  $\bar{w} = -i\omega_0 A + \mathcal{O}(\epsilon\delta)$ ):

$$A_{\text{uni},\approx}^{(31)} \approx -\frac{3}{8}k_0^2|A|^2A + \mathcal{O}(\epsilon^3\delta) \quad \text{and} \quad A_{\text{uni},\approx}^{(33)} \approx \frac{3}{8}k_0^2A^3 + \mathcal{O}(\epsilon^3\delta), \tag{5.10a,b}$$

in which the subscript ‘ $\approx$ ’ denotes an approximate expression due to a narrow-banded assumption. Inserting (5.10a,b) into (5.6) leads to a leading order approximation for  $\zeta^{(3)}$  that agrees with (2.12) in Lo & Mei (1985).

## 6. Results

This section focuses on three aspects in order to explore and validate the theory presented, including the second-order solutions presented in §3, sideband instability of Stokes waves, and the roles of wave directionality. They are examined in §6.1, §6.2, and §6.3, respectively.

For numerical implementations, a Gaussian amplitude spectrum ( $\Xi(k)$ ) in wavenumber and a (unnormalised) directional distribution ( $D(\theta)$ ) were chosen to generate a short




FIGURE 2. A comparison of second-order superharmonic wave elevation  $\zeta^{(22)}$  for a unidirectional wave group at different times between the predictions by (3.13b) (black solid line), Dalzell (1999) (red dashed line), and (6.4)(blue dot-dashed line) based on a second-order Stokes wave theory. The wave group started to propagate at an initial time  $t_0 = -15T_0$  with  $T_0$  the period of the peak wave; it focuses linearly at  $x_f = -25\lambda_0$  with  $\lambda_0 = 2\pi/k_p$ ,  $k_0 = k_p$ , and  $t_f = -10T_0$ . At  $t = t_0$ , an asymmetrical Gaussian spectrum, with  $k_w = 0.27k_0$  for  $k < k_0$  and  $k_w = 1.62k_0$  for  $k > k_0$  and  $\epsilon_0 = 0.3$ , was chosen for  $\zeta^{(1)}$  based on (6.2). The top panels show for  $t = -10T_0$ , the middle panes for  $t = 20T_p$ , and the lower panels for  $t = 40T_0$ . The right panels show the area highlighted by the thick blue box in the left panels, respectively.

(focused) wave group, given by, respectively

$$\Xi(k) = \exp\left[-\frac{(k - k_p)^2}{2k_w^2}\right] \quad \text{and} \quad D(\theta) = \exp\left[-\frac{(\theta - \theta_p)^2}{2\theta_w^2}\right], \quad (6.1a,b)$$

where  $k_p$  denotes the peak wavenumber and  $\theta_p$  denotes the propagation direction of the peak wave at linear focus,  $k_w$  and  $\theta_w$  denote the standard deviation of the spectrum in wavenumber and direction, respectively. An asymmetrical Gaussian spectrum is obtained from (6.1a) through using two different values of  $k_w$  for the upper sideband  $k > k_p$  and lower sideband  $k \leq k_p$ , respectively. For the relevant cases computed in this section, we used  $\theta_p = 0$  and  $k_p = 0.02769 \text{ m}^{-1}$  which is typical of the North Sea (Barratt *et al.* 2021). Using the spectra described by (6.1a,b), linear elevation  $\zeta^{(1)}$  at  $t = t_0$  is given as input by

$$\zeta^{(1)}(x, t) = \frac{\epsilon_0}{k_0 \Xi_k} \sum_{n=1}^{N_k} \Xi(k_n) \cos[k_n(x - x_f) - \sqrt{gk_n}(t_0 - t_f) + \psi_f] \quad \text{with} \quad \Xi_k = \sum_{n=1}^{N_k} \Xi(k_n) \quad (6.2a,b)$$

for unidirectional waves, where  $\epsilon_0$  denotes the dimensionless steepness of a wave group at linear focus,  $k_n$  and  $N_k$  denote the evenly spaced discrete wavenumbers and the total number chosen for numerical implementations, respectively; the subscript ‘f’ denotes the position or time for the wave group at linear focus;  $\psi_f$  denotes the wave phase at linear



focus; for multi-directional waves:

$$\zeta^{(1)}(\mathbf{x}, t) = \frac{\epsilon_0}{k_0 \Xi_{k,\theta}} \sum_{n=1}^{N_k} \sum_{m=1}^{M_\theta} \Xi(k_n) D(\theta_m) \cos[\mathbf{k}_{n,m} \cdot (\mathbf{x} - \mathbf{x}_f) - \omega(\mathbf{k}_{n,m})(t_0 - t_f) + \psi_f] \quad \text{with} \quad (6.3a)$$

$$\Xi_{k,\theta} = \sum_{n=1}^{N_k} \sum_{m=1}^{M_\theta} \Xi(k_n) D(\theta_m) \quad \text{and} \quad \mathbf{k}_{n,m} = [k_n \cos \theta_m, k_n \sin \theta_m], \quad (6.3b)$$

where  $\theta_m$  and  $M_\theta$  denote the evenly spaced discrete wave directions in the range of  $(-\pi, \pi)$  and the total number chosen for numerical implementations, respectively. Despite that the directional distribution described by (6.1b) is in an unnormalised form, (6.3a,b) suggest that the integral unity of the directions of the linear multi-directional waves generated has implicitly been satisfied due to the additional scaling factor,  $\Xi_{k,\theta}$ , defined in (6.3b).

### 6.1. Second-order solutions

We validate the second-order solutions presented in §3 in this section through comparisons with the analytical method by Dalzell (1999) and with existing approximate methods. We focus on the temporal-spatial evolution of a wave group. For simplicity, the effects at third order are neglected in this section as they do not affect qualitatively the discussion presented here. Let a wave group start to propagate at  $t = t_0$  at which the linear elevation  $\zeta^{(1)}(\mathbf{x}, t_0)$  of a group is given in space by (6.2) for a unidirectional group and by (6.3a) for a directionally spread focused wave group. Inserting  $\zeta^{(1)}(\mathbf{x}, t_0)$  into (2.6) leads to an expression for  $A(\mathbf{x}, t)$ . For a fair comparison, the input at  $t = t_0$  for second-order wave fields is based on the results by Dalzell (1999). Following the numerical procedures explained in §4.4,  $\zeta^{(22)}$  and  $\zeta^{(20)}$  are readily obtained from (3.13b) and (3.13c), respectively.

#### 6.1.1. Unidirectional and multi-directional wave group

Figure 2 shows a comparison of the second-order superharmonic wave elevation for a unidirectional wave group at two different times (noting that  $\Phi^{(22)}(x, t) = 0$ ), between the results predicted by (3.13b), Dalzell (1999), and a leading order approximation based on a second-order Stokes theory given by

$$\zeta_{\text{Stokes}}^{(22)} = \frac{1}{2} k_0 A^2(\mathbf{x}, t) \cos[k_0(x - x_f) - \omega_0(t - t_f) + \psi_0]. \quad (6.4)$$

Due to an asymmetrical Gaussian amplitude spectrum chosen in figure 2, a good estimate of the side bandwidth can be given by  $\delta = 3k_w/k_p$  for  $k > k_p$ , which measures from the wavenumber (i.e.  $k = k_p + 3k_w$ ) dropping by 99% in magnitude relative to the spectrum peak. This gives  $\delta = 4.86$  for the case examined in figure 2 and 4. The good agreement between (3.13b) and Dalzell (1999) is evident in figure 2, whereas the less satisfactory agreement between the second-order superharmonic elevation (6.4) and the other two methods is also shown. For completeness, the expressions for both  $\zeta^{(22)}$  and  $\zeta^{(20)}$  based on Dalzell (1999) are presented in appendix B. It is understood Dalzell (1999) is capable of providing exact predictions of second order wave fields based on Fourier integrals. The agreement with Dalzell (1999) in figure 2 clearly demonstrates the semi-analytical approach presented in §3.2 can also work for this purpose. Similar observations are also shown in figure 3 for  $\zeta^{(22)}$  predicted for the evolution of a directionally spread focused wave group at different times.




FIGURE 3. Second-order superharmonic wave elevation  $\zeta^{(22)}$  for a directionally spread focused wave group at different times; the left panels show the results predicted by (3.13b); the right panels show a comparison of the prediction by (3.13b) (blue dot-dashed line) and Dalzell (1999) (red solid line) for  $y = 0$ . The wave group started to propagate at an initial time  $t_0 = -15T_0$  with  $T_0$  denoting the period of the peak wave  $k_p = k_0$ ; it focuses linearly at  $x_f = 0$  and  $t_f = 0$ . At  $t = t_0$ , a symmetrical Gaussian spectrum with  $k_w = 0.3k_0$ ,  $\epsilon_0 = 0.3$ ,  $\theta_w = 15$  deg was chosen for  $\zeta^{(1)}$  based on (6.3a). In the figure  $\lambda_0 = 2\pi/k_0$  denotes the wavelength of the carrier wave of the group.




FIGURE 4. A comparison of surface wave elevation  $\zeta^{(20)}$  for the second-order subharmonic waves forced by a unidirectional wave group at different times between the prediction by (3.13c) (black solid line), Dalzell (1999) (red dashed line), and Trulsen *et al.* (2000) (blue dot-dashed line). In the figure, the top panels show for  $t = -10T_0$ , the middle panels for  $t = 20T_0$ , and the bottom panels  $t = 40T_0$ ; and the parameters were chosen the same as in figure 2.

Figure 4 shows a comparison of the elevation for second-order subharmonic waves forced by a unidirectional linear wave group on still water surface between the results predicted by (3.13c), Dalzell (1999) (cf. appendix B), and Trulsen *et al.* (2000) (cf. appendix D). It is clearly seen in figure 4 that (3.13c) can predict  $\zeta^{(20)}$  as good as Dalzell (1999), whereas Trulsen *et al.* (2000) leads to less satisfactory agreement with Dalzell (1999), owing to a narrow banded assumption.

### 6.2. Sideband instability of Stokes waves

In this section, we examine the capabilities of the new NLSE by investigating the sideband instability of Stokes waves through comparisons with earlier works e.g. Dysthe (1979); Crawford *et al.* (1981); McLean (1982); Trulsen *et al.* (2000). The approximations to  $\mathcal{N}$  presented in §4.2 are used in this section as they are sufficient to this end. The contribution due to  $\mathcal{N}_{\approx, \text{dir}}$  are neglected for simplicity. Based on (4.15), the stability of Stokes waves to sideband perturbations can be investigated by assuming small disturbances in amplitude and phase for  $A$  in a form given by (cf. Trulsen *et al.* (2000))

$$A = a_0(1 + a + i\theta)e^{-i\alpha_1\epsilon_0^2\omega_0 t} \quad \text{with} \quad [a, \theta] = 2[\hat{a}, \hat{\theta}] \cos \psi(\mathbf{x}, t), \quad (6.5)$$

in which  $a_0$  denotes the amplitude of a train of Stokes wave,  $\epsilon_0 = k_0 a_0$  denotes the dimensionless steepness of the Stokes wave,  $\hat{a}$  and  $\hat{\theta}$  are infinitesimal real parameters (i.e.  $\ll \mathcal{O}(1)$ ) that denote small perturbations in amplitude and phase, respectively,  $\psi(\mathbf{x}, t) = k_0[(1 + \delta_x)x + \delta_y y] - \omega_0 \delta_\Omega t$  denotes the modulated phase, with  $k_0$  and  $\omega_0$  the wavenumber and angular frequency of the Stokes wave, respectively,  $\delta_x$  and  $\delta_y$  the dimensionless sideband in wavenumber in the longitudinal ( $k_x$ ) and transverse/spanwise ( $k_y$ ) direction, respectively, and  $\delta_\Omega$  the dimensionless sideband in frequency; as defined in §4.3.1,  $\alpha_1$  is defined as  $\alpha_1 = 1/2$ .

Treating the expression (6.5) for  $A$  as the modulated amplitude of a plane wave  $\mathbf{k}_0$

and inserting (6.5) into (2.12) leads to  $\bar{\mathbf{V}}(\mathbf{x}, z, t)$  given by

$$\bar{\mathbf{V}} = -i \frac{\omega_0}{k_0} \epsilon_0 \left\{ \begin{bmatrix} i \\ 0 \\ 1 \end{bmatrix} + \begin{bmatrix} (i\hat{a} - \hat{\theta})V_{x,+}(z) \cos \psi + (\hat{a} + i\hat{\theta})V_{x,-}(z) \sin \psi \\ (i\hat{a} - \hat{\theta})V_{y,+}(z) \cos \psi + (\hat{a} + i\hat{\theta})V_{y,-}(z) \sin \psi \\ (\hat{a} + i\hat{\theta})V_{z,+}(z) \cos \psi + (i\hat{a} - \hat{\theta})V_{z,-}(z) \sin \psi \end{bmatrix} \right\} e^{-i\alpha_1 |a_0|^2 t}, \quad (6.6)$$

where the  $z$  dependent coefficients  $V_{i,\pm}(z)$  are

$$V_{x,\pm}(z) = \frac{\sqrt{1 + \delta_{z,+}}}{\delta_{z,+}} (1 + \delta_x) e^{k_0 z \delta_{z,+}} \pm \frac{\sqrt{1 + \delta_{z,-}}}{\delta_{z,-}} (1 - \delta_x) e^{k_0 z \delta_{z,-}}, \quad (6.7a)$$

$$V_{y,\pm}(z) = \frac{\sqrt{1 + \delta_{z,+}}}{\delta_{z,+}} \delta_y e^{k_0 z \delta_{z,+}} \pm \frac{\sqrt{1 + \delta_{z,-}}}{\delta_{z,-}} (-\delta_y) e^{k_0 z \delta_{z,-}}, \text{ and} \quad (6.7b)$$

$$V_{y,\pm}(z) = \sqrt{1 + \delta_{z,+}} e^{k_0 z \delta_{z,+}} \pm \sqrt{1 + \delta_{z,-}} e^{k_0 z \delta_{z,-}}, \text{ with} \quad (6.7c)$$

$$\delta_{z,\pm} = \sqrt{(1 \pm \delta_x)^2 + \delta_y^2} - 1. \quad (6.7d)$$

The full solution for  $\bar{\mathbf{V}}$  in a form as (6.6) allows for  $\min(|\delta_x|, |\delta_y|) > 1$ . It should be noted that, time dependant disturbances (small) for  $\bar{\mathbf{V}}$  except for these due to linear dispersion relation are neglected for simplicity. Or otherwise, the derivations following Benjamin (1967) (e.g. starting from Eq. (11) and (21) in the paper) are needed that would lead to numerical analysis for the sideband instability. The insertion of (6.6) into (4.12) for  $j = 3$  leads to an eigenvalues equation for  $\delta_\Omega$ , with the terms at  $\mathcal{O}(\epsilon_0^2 \delta_y^2)$  and higher neglected. In particular, the eigenvalues equation for  $\delta_\Omega$  reads

$$(-\delta_\Omega + \mathcal{L}_I)^2 + (-\delta_\Omega + \mathcal{L}_I)(\alpha_{11} + \alpha_{22}) + \alpha_{12}\alpha_{21} = 0, \quad (6.8)$$

where  $\mathcal{L}_r = \frac{1}{2}[\mathcal{L}(ik_0\delta_x, ik_0\delta_x) + \mathcal{L}(-ik_0\delta_x, -ik_0\delta_x)]$  and  $\mathcal{L}_I = \frac{1}{2}[\mathcal{L}(ik_0\delta_x, ik_0\delta_y) - \mathcal{L}(-ik_0\delta_x, -ik_0\delta_y)]$  denote the contribution of the linear dispersion, and coefficients  $\alpha_{ij}$  are given by

$$\alpha_{11} = \frac{1}{4}V_{z,-}\epsilon_0^2 + (V_{x,+} + V_{z,+}) \left[ -\frac{1}{8}\delta_x + \left( \frac{1}{2}\mathcal{L}_2 + \frac{\mathcal{L}_2(1 + \epsilon_0^2)\delta_{20} - \delta_x\mathcal{L}_2^2}{\delta_{20} - \mathcal{L}_2^2} \right) \right] \epsilon_0^2, \quad (6.9a)$$

$$\alpha_{12} = \alpha_1\epsilon_0^2 - \mathcal{L}_r - \frac{1}{4}V_{z,+}\epsilon_0^2 - (V_{x,-} - V_{z,-}) \left[ \frac{1}{8}\delta_x - \frac{1}{2} \left( \mathcal{L}_2 + \frac{2\mathcal{L}_2(1 + \epsilon_0^2)\delta_{20} - 2\delta_x(\mathcal{L}_2)^2}{\delta_{20} - (\mathcal{L}_2)^2} \right) \right] \epsilon_0^2, \quad (6.9b)$$

$$\alpha_{22} = \left[ -\frac{1}{4}(V_{x,-} - 2V_{z,-}) - \frac{1}{8}\partial_z(V_{x,-} - V_{z,-}) + (V_{x,-} - V_{z,-}) \frac{\delta_x(1 + \epsilon_0^2)\mathcal{L}_2 - \delta_{20}\mathcal{L}_2^2}{\delta_{20} - \mathcal{L}_2^2} \right] \epsilon_0^2, \quad (6.9c)$$

$$\alpha_{21} = \mathcal{L}_r - \alpha_1\epsilon_0^2 + \frac{1}{4}(V_{x,+} + 2V_{z,+})\epsilon_0^2 + \frac{1}{8}\partial_z(V_{x,+} + V_{z,+})\epsilon_0^2 - \epsilon_0^2(V_{x,+} + V_{z,+}) \times \\ [\delta_x(1 + \epsilon_0^2)\mathcal{L}_2 - \delta_{20}\mathcal{L}_2^2] / (\delta_{20} - \mathcal{L}_2^2), \quad (6.9d)$$

in which the  $z$  dependent variables associated with  $V_{i,\pm}(z)$  are evaluated for  $z = 0$ ,  $\delta_{20} = \sqrt{\delta_x^2 + \delta_y^2}$  denotes the dimensionless magnitude of the sidebands away from the wavenumber of the Stokes wave, and the approximation  $\mathcal{L}_2 \approx \frac{1}{2}(\mathcal{L}_I + \mathcal{L}_r)$  (cf. equations (11) and (15) by Crawford *et al.* (1981) for similar discussions) was used due to the terms associated with  $\partial_t(\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*)$  and, therefore,  $\mathbf{V}^{(20)}$ , arising from the symmetry of the disturbances to the Stokes wave.

It is understood that the instability occurs if (6.8), a quadratic equation in  $\delta_\Omega$ , has a




FIGURE 5. A comparison of the instability region predicted by Trulsen *et al.* (2000), the exact computations based on McLean (1982), and the present theory based on (6.10) for three different values of wave steepness defined as  $\epsilon_0 = k_0 a_0$ ; (a,d)  $\epsilon_0 = 0.0995$ , (b,e)  $\epsilon_0 = 0.197$ , and (c,f)  $\epsilon_0 = 0.327$ . (a-f) the exact results based on McLean (1982) (indicated by black dashed lines) are computed; (a-c) computations are based on equation (18) by Trulsen *et al.* (2000) (blue dots) and a leading-order approximation based on (4.12) and (4.13a) (red solid); both approximations are correct to  $\mathcal{O}(\epsilon_0^3 \delta_0^0)$  with the exact linear dispersion relation; (d-f) the approximations based on equations (19-22) by Trulsen *et al.* (2000) (blue dot-dashed), correct to  $\mathcal{O}(\epsilon_0^2 \delta_x^2, \epsilon_0^2 \delta_y^0)$ , and (6.10) derived in this paper (red dot-dashed).

positive imaginary root for  $\delta_\Omega$ . Thereby, the instability occurs if the following inequality holds

$$R_\Omega \equiv \frac{1}{4}(\alpha_{11} + \alpha_{22})^2 - \alpha_{12}\alpha_{21} < 0, \quad (6.10)$$

in which  $R_\Omega$  is defined for later reference.

Figure 5 shows a comparison of the instability region predicted by (6.10) and its lower order expression, equations (18-22) by Trulsen *et al.* (2000), and the exact computations by McLean (1982) for three different values of wave steepness. The equations by Trulsen *et al.* (2000) are presented in appendix D for reference. In the cases chosen by McLean (1982) the value of a different steepness, defined as half the crest to trough height of Stokes waves and referred to as  $\epsilon_M$  here, is given. For a direct comparison, the identity derived by Trulsen & Dysthe (1996) (i.e. equation (30) therein) was used to evaluate  $\epsilon_0$  as follows

$$\epsilon_0 = \epsilon_M - \frac{1}{2}\epsilon_M^3. \quad (6.11)$$

It is seen from figure 5(a-c) that, compared with McLean (1982), both approximations behave similar; good agreement is shown for small  $\delta_x$  and small steepness  $\epsilon_0$ , whereas

the agreement becomes less satisfactory for larger  $\epsilon_0$ . As approximations of higher-order accuracy are used, better agreement with McLean (1982) is clearly seen in figure 5(d-f) compared with figure 5(a-c). In particular, as seen in figure 5(d-f), equation (6.10) agrees with McLean (1982) in a nearly perfect manner for  $\epsilon_0 \approx 0.1$  and  $\epsilon_0 \approx 0.2$  in the range  $\delta_x \lesssim 1$ , while the predictions by (6.10) are also satisfactory for  $\epsilon_0 \approx 0.33$ . Comparing the results by Trulsen *et al.* (2000) and (6.10) with McLean (1982) shown in 5(d-f), it is clear that (6.10) provides better prediction for all three cases. In particular, equation (6.10) is capable of providing more accurate predictions for the instability of Stokes waves subject to sidebands in the transverse direction, as clearly shown in figure 5(e,f). Focusing on the instability due to transverse sidebands ( $\delta_y$ ), figure 5 shows that equations (19-22) by Trulsen *et al.* (2000) fail to improve the accuracy of the lowest-order approximation (i.e. equation (18) by Trulsen *et al.* (2000)) for small  $\delta_x$  for all three cases. The reason to this is explained as follows. Equations (19-22) by Trulsen *et al.* (2000), with the notation in this paper, lead to the instability region described by the inequality as follows

$$R_T \equiv \mathcal{L}_r \left[ \mathcal{L}_r + \epsilon_0^2 \left( 1 - \frac{\delta_x^2}{\delta_{20}^2} \right) \right] + \frac{\epsilon_0^4 \delta_x^2}{16} < 0. \quad (6.12)$$

where  $R_T$  is introduced for later reference. Equation (6.12) gives the instability region based on equation (18) by Trulsen *et al.* (2000) by setting  $\delta_x$  to zero. For small  $\epsilon_0$  ( $\epsilon_0 \lesssim 0.5$ ), the last term in  $R_T$  is negligible, which gives two conditions for instability to occur

$$\mathcal{L}_r < 0 \quad \text{or} \quad \mathcal{L}_r + \epsilon_0^2 \left( 1 - \frac{\delta_x^2}{\delta_{20}^2} \right) < 0. \quad (6.13)$$

Noting that  $\delta_x^2/\delta_{20}^2 < 1$  for  $\delta_x \lesssim 1$ , the first condition in (6.13) dominates for small  $\delta_x$ . Using a leading order approximation to  $\mathcal{L}_r$ , i.e.  $\mathcal{L}_r \approx (2\delta_y^2 - \delta_x^2)/8$ , the first condition is approximately

$$2\delta_y^2 - \delta_x^2 < 0, \quad (6.14)$$

which corresponds to an (neutral) instability slope near the origin defined by  $\delta_y/\delta_x = \pm 1/\sqrt{2}$  (corresponding to  $\sim \pm 35.26^\circ$  to the carrier wave  $\mathbf{k}_0$ ), as pointed out by Longuet-Higgins (1976) for the analysis of resonant energy transfer in a narrow spectrum and clearly shown by the predictions based on Trulsen *et al.* (2000) in figure 5 (see also figure 12).

In contrast, equation (6.10) is capable of providing accurate prediction of the instability region of Stokes waves subject to span-wise ( $y$  direction) sideband perturbations for moderate values of wave steepness,  $\epsilon_0 \lesssim 0.2$ , as can be clearly seen in figure 5(d, e). This is due to that (6.6) has taken into account the transverse sidebands  $\delta_y$  in  $\delta_{z,\pm}$  which denotes the effects on wave kinematics (i.e.  $\bar{\mathbf{V}}$ ) in addition to the wave envelope  $A$ . For  $\epsilon_0 \approx 0.33$ , the difference between (6.10) and McLean (1982) is likely due to that the terms described by  $\mathcal{N}_{\approx, \text{dir}}$  presented in §4.2 at  $\mathcal{O}(\epsilon_0^2 \delta_y^2)$  were neglected in the computations.

Proceeding to examining the energy growth rate of Stokes waves, which can be defined as the positive imaginary component of the roots of (6.8) as follows

$$\text{Im}(\delta_\Omega) = \text{Im}(\sqrt{R_\Omega}), \quad (6.15)$$

where  $\text{Im}$  denotes the imaginary component. When instability occurs, the rate of amplitude growth/decline behaves like  $\exp[\frac{1}{2}\text{Im}(\delta_\Omega)\omega_0 t]$  based on (6.5). This will be further examined in §6.3.2.

Figure 6 shows the energy growth rate,  $\text{Im}(\delta_\Omega)$ , varying with the sideband  $\delta_x$  for five values of  $\epsilon_0$  and varying with  $\epsilon_0$  for two values of  $\delta_x$ . In addition to the experimental




FIGURE 6. The growth rate  $\text{Im}(\delta_\Omega)$  varying with the longitudinal bandwidth  $\delta_x$  (left panel) and  $\epsilon_0 = k_0 a_0$  (right panel) when the instability of Stokes waves occurs. In the figure, the predictions are computed based on Crawford *et al.* (1981), (6.15), and equations (19-22) by Trulsen *et al.* (2000) for three different values of steepness (left panel) and two different values of  $\delta_x$  (right panel). The right panel shows, in addition, experimental measurements by Benjamin & Feir (1967) and Lake *et al.* (1977).




FIGURE 7. Stability boundary for growth of unstable perturbations for two-dimensional wave trains predicted based on (6.15), compared with the results of Benjamin & Feir (1967); Longuet-Higgins (1978), and the Zakharov equation due to Crawford *et al.* (1981).

measurements and the predictions of (6.15), figure 6 also presents the results due to Crawford *et al.* (1981), for which small corrections following Krasitskii (1994) and Janssen (2004) (cf. §4.11) were used. Crawford *et al.* (1981) is based on the Zakharov integral equation for the study of a uniform wave train and the results are correct to third order in wave steepness with all higher-order dispersion effects included. It is seen in figure 6 that (6.15) shows better agreement with Crawford *et al.* (1981) than Trulsen *et al.* (2000) (i.e. predicted by  $\text{Im}\sqrt{R_T}$ ), which is more so for  $0.15 \lesssim \epsilon_0$  and  $0.4 \lesssim \delta_x$ .

Figure 7 shows the instability boundary for growth of unstable perturbations for two dimensional wave trains, predicted due to Benjamin & Feir (1967); Longuet-Higgins (1978); Crawford *et al.* (1981), and by this paper based on (6.15). Compared with the exact numerical results due to Longuet-Higgins (1976), (6.15) is in qualitative good agreement whereas the quantitative discrepancy is  $\sim 23\%$  for the restabilization of steeper




FIGURE 8. The nondimensional superharmonic velocity head  $k_0(H_v^{(22)} + \text{c.c.})$  for the evolution of a focused wave group at different times for  $z = 0$ . The parameters are the same as figure 3 except that  $\theta_w$  differs: (a-c)  $\theta_w = 15$  deg; (d-f)  $\theta_w = 30$  deg; (g-i)  $\theta_w = 45$  deg; (a,d,g)  $t = -15 \times T_0$ ; (b,e,h)  $t = 0$  for the groups at linear focus with  $\epsilon_0 = 0.3$ ; (c,f,i)  $t = 15 \times T_0$  after focus.

waves perturbed by long waves with  $\delta_x \rightarrow 0$ . Although the results based on (6.15) in figure 7 show slightly worse performance than the Zakharov equation, it can obviously yield restabilization for larger wave steepness, being an improvement over earlier versions of NLSEs.

For large values of wave steepness, i.e.  $0.2 \lesssim \epsilon_0$ , as shown in figure 6(a,b) and 7, the disagreement between (6.15) and Crawford *et al.* (1981) can be observed, which likely arises from that the additional time-dependant (small) disturbances are neglected in (6.15) to the linear velocity envelope (vector),  $\bar{\mathbf{V}}$ . The analysis in this section shows the results due to (6.15) are satisfactory.

Figures 5, 6, and 7 suggest that the new NLSE described by (4.9) should work for  $\epsilon_0 \lesssim 0.25$  and  $\delta_{x,y} > 1$  for the evolution of surface waves with directional spread.

### 6.3. Effects of wave directionality

The presented theory permits us to investigate the effects of wave directionality based on an order in wave steepness. Thereby, we focus the wave-directionality effects on waves at second order (§6.3.1) and on nonlinear energy transfer in a narrow wave spectrum by implementing the new NLSE numerically in §6.3.2.

#### 6.3.1. Waves at second order

As explained in §2.4, one of the major effects of wave directionality arises from that the superharmonic velocity head  $H_v^{(22)}$  throughout the water columns becomes non-zero for directionally spread waves. This leads to both non-zero  $\Phi^{(22)}$ , additional contributions






FIGURE 9. The dimensionless velocity head  $k_0 H_v^{(20)}$  at different times for  $z = 0$ . See the caption of figure 8 for the specific parameters.

to elevation  $\zeta^{(22)}$  due to both  $\Phi^{(22)}$  and  $H_v^{(22)}$ . These second-order effects on the spatial-temporal evolution of potential or elevation envelope of the first harmonic are considered negligible in most models based on an earlier version of NLSE e.g. the NLSE by Dysthe (1979); Stiassnie (1984); Trulsen *et al.* (2000). We focus on examining the velocity head  $H_v^{(22)}$  and  $H_v^{(20)}$  by studying the evolution of a directionally spread focused wave group at different times. For simplicity, third order effects were neglected in the computations as they do not affect the discussions here in a qualitative manner.

Figures 8 and 9 show the dimensionless superharmonic and subharmonic velocity head defined in §2.4, respectively. It is seen from figure 8 that, except for the group at linear focus (figure 8(b,e,h)), larger values of directional spread lead to smaller magnitudes in superharmonic velocity head, owing to that, as the total wave energy is kept the same, the wave energy distributes in larger areas for larger values of directional spread. The magnitude of the superharmonic velocity head is shown to be  $\lesssim 10 \sim 30\% \times \epsilon_0$  in figure 8. Similar observations can be found in figure 9 for the subharmonic velocity head which is of a similar order of magnitude as the superharmonic velocity head shown in figure 8. This indicates both components can be equally important for waves with directional spread.

### 6.3.2. Nonlinear energy transfer in a narrow wave spectrum

In order to examine the nonlinear energy transfer in a narrow wave spectrum, the temporal evolution of a ‘steep’ wave group is investigated numerically with the implementation of the new NLSE and equations (19-22) by Trulsen *et al.* (2000). Figure 10 shows the evolution of the discrete amplitude spectra,  $\hat{A}(\mathbf{k}, t)$ , in the first quadrant of




FIGURE 10. Discrete dimensionless amplitude spectra,  $k_0|\hat{A}(\mathbf{k}, t)|$ , in the Fourier space at different times in the first quadrant for the spatial evolution of a wave group with  $\epsilon_0 = 0.2$ . The group starts to propagate at  $t = -15T_0$  and focuses linearly at  $\mathbf{x}_f = 0$  and  $t_f = 0$ . (a-c): the approximate NLSE with (4.13b) was used for computations; (d-f): equations (19-22) by Trulsen *et al.* (2000) were implemented. Contour levels (black solid line) are evenly distributed between 0.005 and 0.12 m in intervals of 0.005 m. (a) the red circles denote the wavenumbers chosen to investigate the growth rate of  $|\hat{A}(\mathbf{k}, t)|$  in figure 11. The longitudinal and transverse sidebands are defined as  $\delta_x = (k_x - k_0)/k_0$  and  $\delta_y = k_y/k_0$ , respectively.

the Fourier space for a wave group at different times with  $\epsilon_0 = 0.2$ . It is seen in figure 10 that the results based on (4.9) and Trulsen *et al.* (2000) do not show differences in a qualitative manner and the energy conservation described by

$$\int_{\mathbf{x}_s}^{\mathbf{x}_e} |A(\mathbf{x}, t)|^2 d\mathbf{x} = \text{constant} \quad (6.16)$$

were verified (not shown) to be within 0.01% for both models. The dimensionless sidebands in the longitudinal and transverse direction are defined as, respectively,

$$\delta_x = \frac{k_x - k_0}{k_0} \text{ and } \delta_y = \frac{k_y}{k_0}, \quad (6.17a,b)$$

in which  $k_0$  was chosen to be the peak wavenumber of the initial wave spectrum,  $\delta_x > 0$  and  $\delta_x < 0$  denote the region of upper and lower sidebands in the longitudinal direction, respectively.

Compared with the initial time shown in figure 10(a,d), figure 10(b,e) shows broadened spectral amplitude in both the lower and higher sidebands in the longitudinal direction but narrowed in the region of larger sidebands in the transverse ( $k_y$ ) direction, suggesting energy transfers from the latter to the former.

From  $t = 0$  to  $t = 24 \times T_0$  with  $T_0$  denoting the wave period of the spectral peak, two main features can be observed in figure 10. Firstly, oblique energy transfers are clear and the change of the spectral amplitude in the regions of the lower ( $k_x/k_0 < 1$  or

$\delta_x < 0$ ) and upper ( $k_x/k_0 > 1$  or  $\delta_x < 0$ ) sidebands behaves differently. This indicates asymmetrical change rate of the spectral amplitude in sidebands. Particularly, in the region of the lower sidebands described by  $0 < \delta_x = (k_x - k_0)/k_0 < -\sqrt{2}k_y/k_0$ , a decline in the amplitude is indicated whereas an energy expansion is obvious in the region of upper sidebands, especially for  $\delta_y > \delta_x/\sqrt{2}$ . Secondly, relative to the original spectral peak at  $k_x/k_0 = 1$ , a small downward shift of the spectral peak can be observed in figure 10(c,f), as is clearly shown by the nearest contour in the vicinity of (1, 0) that is no longer symmetrical relative to  $k_x/k_0 = 1$ .

In order to examine the change rate of the spectral amplitude in time,  $|\hat{A}(\mathbf{k}, t)|$  were calculated based on (4.9) at different times and  $|\hat{A}(\mathbf{k}, t)|$  at 35 different wavenumbers were chosen and shown in figure 11. Figure 11 shows the nondimensional spectral amplitude as a function of the dimensionless time  $\frac{1}{2}\epsilon_0^2\omega_0 t$ . As is seen in figure 11, the change rate of  $|\hat{A}(\mathbf{k}, t)|$  behaves differently at different wavenumbers. The most interesting range is obviously for  $\frac{1}{2}\epsilon_0^2\omega_0(t - t_0)$  in  $\sim [1, 3]$  in which the amplitude changes rapidly. This range is in the vicinity of  $t = t_f$  at which the wave group focuses linearly. We refer to  $t \lesssim t_f$  and  $t_f \lesssim t$  in this time range as the pre-focus and post-focus phase, respectively.

Focusing on the pre-focus phase first, figure 11 shows the amplitude grows for the wavenumbers in the range  $\delta_y \lesssim 0.1$  and wavenumbers in the upper sidebands with  $\delta_x = 0.36$  for  $\delta_y \lesssim 0.35$ . In contrast, the amplitudes for larger sidebands in the transverse direction, i.e. for  $0.18 \lesssim \delta_y$ , experience a decline except for  $\delta_x = 0.36$ . In the post-focus phase, the amplitudes that have experienced a growth (decline) in the pre-focus phase tend to decrease (grow) exponentially. Especially, for the amplitudes at the wavenumbers in small transverse sidebands, i.e.  $\delta_y \lesssim 0.09$ , the rate of decline is larger than that of increase. The asymmetry in the rate of amplitude change in different regions is likely the cause for the downward shift of the spectral peak shown in figure 10(c,f).

The change rate of  $\ln(k_0|\hat{A}|)$  in the range for  $\frac{1}{2}\epsilon_0^2\omega_0(t - t_0)$  in  $\sim [1, 3]$  in figure 11 shows that  $\ln(k_0|\hat{A}|)$  decreases/grows with  $\frac{1}{2}\epsilon_0^2\omega_0(t - t_0)$  (approximately) linearly with absolute slopes  $\lesssim 1$ . This suggests that  $|\hat{A}(\mathbf{k}, t)|$  changes exponentially with the dimensionless time  $\frac{1}{2}\epsilon_0^2\omega_0(t - t_0)$ , unlikely arising from resonant energy transfers as resonance would lead to the change rate to be linear in time. The rapid change of the spectral amplitudes shown in figure 11 may be due to instability. This paper argues that it is likely due to a mechanism similar to oblique sideband instability of Stokes waves. To illustrate this, the amplitude change rate in the instability region of Stokes waves is shown in figure 12 for five different values of wave steepness. The results in figure 12 were computed based on (6.15) and equations (19-22) by Trulsen *et al.* (2000) (i.e.  $\text{Im}\sqrt{R_T}$ ). As noted in §6.2, it is understood that, when instability occurs, the amplitude behaves like

$$|A| \sim a_0 e^{\text{Im}(\delta_\Omega)\omega_0(t-t_0)} \quad \text{and thus, } R_c \equiv \frac{\ln(k_0|A|)}{\frac{1}{2}\omega_0\epsilon_0^2(t-t_0)} \sim \ln(\epsilon_0) \frac{\text{Im}(\delta_\Omega)}{\frac{1}{2}\epsilon_0^2}, \quad (6.18a,b)$$

where  $R_c$  denotes the change rate of amplitude in the logarithmic scale, introduced for convenience. It also corresponds to the gradient of  $\ln(k_0|A|)$  with respect to  $\frac{1}{2}\omega_0\epsilon_0^2(t-t_0)$  shown in figure 11.

Based on (6.18b), it is clear that  $R_c$  depends on both  $\epsilon_0$  and the magnitude of  $\text{Im}(\delta_\Omega)$ . Due to  $\epsilon_0 < 0.5$  in general which means  $\ln(\epsilon_0) < 0$ , whether or not the amplitude grows in time depends on the sign of  $\text{Im}(\delta_\Omega)$ ; a positive sign leads to a decline whereas a negative sign a growth. In practice  $0.1 \lesssim \epsilon_0 \lesssim 0.3$  would give a good estimate of the steepness range, yielding  $-2.3 \lesssim \ln(\epsilon_0) \lesssim -1.2$ . Figure 12 shows  $\delta_\Omega/(\frac{1}{2}\epsilon_0^2)$  varying with transverse and longitudinal sidebands at five different values of  $\epsilon_0$  varying from 0.1 to 0.3. The contours of large growth rates shown in Figure 12 suggest that oblique energy transfers




FIGURE 11. Amplitude spectra  $k_0|\hat{A}(\mathbf{k}, t)|$  at 35 wavenumbers in the first quadrant of the Fourier  $\mathbf{k}$  space as a function of the dimensionless time  $\frac{1}{2}\epsilon_0^2\omega_0(t-t_0)$ . The results are based on the new NLSE and the parameters were chosen the same as figure 10. In the figure,  $\delta_x = (k_x - k_0)/k_0$  and  $\delta_y = k_y/k_0$  denote the dimensionless sidebands in the longitudinal and transverse/span-wise direction, respectively. Short vertical lines near  $\frac{1}{2}\epsilon_0^2\omega_0(t-t_0) = 2$  indicates for  $t = t_f$  at which the wave group focuses linearly.

would be favoured when instability occurs, although the maximum  $\delta_\Omega$  always lies on the  $\delta_x$  axis for deepwater Stokes waves. The magnitude of  $\delta_\Omega/(\frac{1}{2}\epsilon_0^2)$  predicted by both the present theory and Trulsen *et al.* (2000) is  $\lesssim 1$ . Obvious ‘coincidences’ between figures 11 and 12 are that both indicate the same magnitude for  $R_c$ , i.e.  $\lesssim 1$ , and that the wave amplitude in both cases changes exponentially. Due to the two points, this paper argues the instability of Stokes waves subject to sidebands disturbances in oblique directions would be a possible cause for the rapid change of wave amplitude shown in figure 11 in the neighbourhood of the linear focused time  $t = t_f$ .

Examining figures 10, 11, and 12 together, the following features are highlighted.

(i) The rapid energy transfers between upper and lower sidebands shown in figure 10 are likely due to the instability of Stokes waves in addition to the degenerate resonant interaction in the ‘figure of eight’ quartet resonant loop (Phillips 1967), as a result of that the change rate of the amplitude shown in figure 11 is of the same order of magnitude as those in figure 12 when instability occurs. It is well understood that the instability of Stokes waves is mostly accompanied by the degenerate resonant interaction, as pointed out by Phillips (1967) where it is shown that the neutral instability modes (i.e.  $\text{Im}(\delta_\Omega) = 0$ ) of Stokes waves can be directly derived from the degenerate resonant interaction (Hammack & Henderson 1993) and, most importantly, the instability of Stokes waves subject to oblique wavetrain disturbances is likely to occur in a narrow spectrum.

(ii) A downward shift of the spectral peak arises likely from the asymmetrical change rates of energy between the different regions of sidebands. This shift is consistent with the finding of Trulsen & Dysthe (1997). When a downward shift of spectral peak is observed, rapid energy transfers are shown between lower and upper sidebands in the longitudinal direction, and between the transverse and longitudinal sidebands.

(iii) Oblique energy transfers towards large upper sidebands ( $0.3 \lesssim \delta_x$  or  $0.27 \lesssim \delta_y$ )




FIGURE 12. Instability growth rate of Stokes waves, defined as  $\text{Im}(\delta_\Omega)/(\epsilon_0^2/2)$ , subject to oblique sideband perturbations for five values of wave steepness  $\epsilon_0 = k_0 a_0$ , obtained using the prediction by (6.8) (a-e) and equations (19-22) (f-j) by Trulsen *et al.* (2000) for five different values of steepness .

are obviously seen in figure 10(c,f), likely due to the different change rates within the instability region of Stokes waves; the contour levels of large change rates of amplitude have demonstrated that the change of energy favours more in oblique directions. This observation is consistent with Trulsen & Dysthe (1997) and the rapid energy transfer reported in Barratt *et al.* (2021). Barratt *et al.* (2021) focus on a numerical study of the propagation of a steep wave group based on direct numerical solutions of fully nonlinear potential flow equations. Barratt *et al.* (2021) reports an angle along  $\text{ang}(\delta_y/\delta_x) = \pm 55^\circ$  to the spectral peak in the region of upper sidebands. This paper argues that the angle as such  $\pm 55^\circ$  may not exist. It is likely a coincidence to the effect shown in figure 11(d,e) that larger upper sidebands ( $0.3 \lesssim \delta_x$  or  $0.27 \lesssim \delta_y$ ) tend to grow exponentially at a rate among the largest. Moreover, Barratt *et al.* (2021) attribute the rapid oblique energy transfers, similar to those shown in the region (i.e.  $\delta_y > \delta_x/\sqrt{2}$  for  $0.3 \lesssim \delta_x$ ) of the upper sidebands in figure 10(c,f), to non-degenerate resonant interaction. This argument is different from what is shown in figures 10, 11, and 12; i.e. that the amplitude changes exponentially is unlikely a result of resonant interactions.

## 7. Concluding remarks

Using a perturbation expansion, this paper has derived a new nonlinear Schrödinger equation (NLSE) for the linear envelope evolution of three-dimensional surface gravity waves without the assumption of a narrow bandwidth. The new NLSE can be extended for more general cases e.g. waves on water of finite uniform or slowly varying depth. Simple numerical implementations of the NLSE for the wave envelope using a pseudospectral, a split-step, and a finite-difference method are demonstrated. The computational cost is in the same order as the numerical implementations of an existing NLSE e.g. the Dysthe equation by Dysthe (1979) and the modified NLSE by Trulsen *et al.* (2000). With leading-order approximations to the NLSE, its capabilities for studying the instability region and energy growth rate of Stokes waves are demonstrated. The approximation is shown to be capable of providing satisfactory predictions of the instability region, compared with the exact computations by McLean (1982) for dimensionless wave steepness equal to as large as 0.33. It can also provide good estimates of the energy growth rate and stability boundary of unstable perturbations for two dimensional wave trains, compared with the results due to Crawford *et al.* (1981) and the exact numerical results based on Longuet-Higgins (1978). In these comparisons, the approximation has been demonstrated of better performance than the modified NLSE by Trulsen *et al.* (2000), particularly for sideband disturbances of waves in an oblique direction to a train of Stokes wave.

This paper has also proposed a semi-analytical approach for the description of wave fields at second order in wave steepness, based on a pseudospectral and a finite-difference method. It is demonstrated that the semi-analytical approach is capable of providing exact predictions as that by Dalzell (1999). Compared with Dalzell (1999), the semi-analytical approach can significantly improve the computational efficiency; the numerical implementations of Dalzell (1999) require computational operations at  $\mathcal{O}(N^2)$  and the semi-analytical approach requires  $\mathcal{O}(N \ln(N))$  if  $N$  Fourier modes are used. In analysis of velocity head at second order in wave steepness, it is suggested that the super- and sub-harmonic waves may be equally important in directional sea states due to the same order of magnitude.

Through a study of the temporal evolution of a steep focused wave group, energy transfers in a narrow spectrum have been investigated. For the time range in the vicinity of the group at linear focus, rapid oblique energy transfers between different regions of sidebands are observed, likely arising from degenerate resonant interactions in the so-called ‘figure of eight’ quartet resonant loop (Phillips 1967) and the instability of Stokes waves subject to oblique sideband perturbations. The change rate of wave amplitude is found to be at the same order of magnitude as that of oblique instability of Stokes waves. A downward shift of the spectral peak is also observed in this paper, consistent with the finding in Trulsen & Dysthe (1997). This shift is observed in the course of rapid energy transfers between different sideband regions when oblique modulational instability occurs, in contrast to the temporal evolution of a unidirectional spectrum which is found to lead to a permanent shift of the spectrum peak only in a quasi-steady state in Dysthe *et al.* (2003). Rapid energy transfers towards the angle,  $\sim \pm 55^\circ$ , to the spectral peak in the region of upper sideband, that are reported in Barratt *et al.* (2021), are not supported by this study. Instead, the upper sideband range of large oblique directions ( $> 45^\circ$ ) to the spectral peak is found to be favoured the most for the energy growth, coinciding with the sideband region where the instability growth rate of three-dimensional Stokes waves is among the largest.

## Acknowledgement

The author acknowledges support from the Research Council of Norway through FRIPRO mobility project 287389. The author would like to thank Prof. Karsten Trulsen and Prof. Paul Taylor for insightful discussions, the kind hospitality of Associate Prof. Thomas A.A. Adcock and Associate Prof. Ton van den Bremer and their support as well as the support of Prof. Simen A. Å. Ellingsen to the project. The author is grateful to the anonymous reviewers for their constructive comments and suggestions which improved the quality of the paper.

## Declaration of interests

The author reports no conflict of interest.

## Appendix A. Third-order forcing term for the first harmonic

### A.1. Full expression of $Q^{(31)}$

The first term that has nonzero contribution to  $Q^{(31)}$  is denoted by  $Q_1^{(31)}$ , expressed as

$$Q_1^{(31)} = 2\partial_t(\nabla_3\Phi^{(1)} \cdot \nabla_3\Phi^{(2)}), \quad (\text{A } 1)$$

which leads to

$$Q_1^{(31)} = E(\mathbf{x}, t)(-i\omega_0 + \partial_t) \left\{ \frac{1}{2}[(2\mathbf{k}_0^{(3d)} + \nabla_3)B^{(22)}] \cdot \bar{\mathbf{V}}^* + \nabla_3\Phi^{(20)} \cdot \bar{\mathbf{V}} \right\} + \text{c.c.}, \quad (\text{A } 2)$$

where  $E(\mathbf{x}, t) = \exp(i\mathbf{k}_0 \cdot \mathbf{x} - \omega_0 t)$ . The second term that has nonzero contribution to  $Q^{(31)}$  is denoted by  $Q_2^{(31)}$  in a form as

$$Q_2^{(31)} = \frac{1}{2}\nabla_3\Phi^{(1)} \cdot \left[ \nabla_3(|\nabla_3\Phi^{(1)}|^2) \right], \quad (\text{A } 3)$$

yielding

$$Q_2^{(31)} = \left\{ \frac{1}{16} \left[ (2\mathbf{k}_0^{(3d)} + \nabla_3)|\bar{\mathbf{V}}|^2 \right] \cdot \bar{\mathbf{V}}^* + \frac{1}{4}\nabla_3 \left[ (\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*)e^{2k_0 z} \right] \cdot \frac{1}{2}\bar{\mathbf{V}} \right\} E(\mathbf{x}, t) + \text{c.c.} \quad (\text{A } 4)$$

The third term that contribute to  $Q^{(31)}$  is defined as

$$Q_3^{(31)} = k_0\zeta\partial_t(\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*), \quad (\text{A } 5)$$

which leads to

$$Q_3^{(31)} = \frac{1}{2}k_0A\partial_t(\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*)E(\mathbf{x}, t) + \text{c.c.} \quad (\text{A } 6)$$

The summation of the three terms for  $Q^{(31)}$  yields

$$\begin{aligned} \bar{Q}^{(31)} = & (-i\omega_0 + \partial_t) \left( \nabla_3\Phi^{(20)} \cdot \bar{\mathbf{V}} \right) + \frac{1}{8} \left[ \nabla_3(\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*) \right] \cdot \bar{\mathbf{V}} + \frac{1}{4}(k_0\bar{w} + 2k_0A\partial_t)(\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*) + \\ & (-i\omega_0 + \partial_t) \left\{ \frac{1}{2} \left[ (2\mathbf{k}_0^{(3d)} + \nabla_3)B^{(22)} \right] \cdot \bar{\mathbf{V}}^* \right\} + \frac{1}{16} \left[ (2\mathbf{k}_0^{(3d)} + \nabla_3)(|\bar{\mathbf{V}}|^2) \right] \cdot \bar{\mathbf{V}}^*, \\ & \text{for } z = 0. \end{aligned} \quad (\text{A } 7)$$

For unidirectional waves we note that

$$B^{(22)} = 0 \quad \text{and} \quad \bar{\mathbf{V}} \cdot \bar{\mathbf{V}} = 0, \quad (\text{A } 8)$$

which leads to

$$\bar{Q}_{\text{all}}^{(31)} = (-i\omega_0 + \partial_t) \left( \nabla_3 \Phi^{(20)} \cdot \bar{\mathbf{V}} \right) + \frac{1}{8} \left[ \nabla_3 (\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*) \right] \cdot \bar{\mathbf{V}} + \frac{1}{4} (k_0 \bar{w} + 2k_0 A \partial_t) (\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*). \quad (\text{A } 9)$$

Combining (A 7) and (A 9), (4.5) in the main is obtained.

### A.2. Details for $\Gamma_{\text{Dythe}}$ based on a narrow banded assumption

With a narrow-banded assumption, i.e.  $\delta \ll \mathcal{O}(1)$ ,  $\bar{\mathbf{V}}$  and  $\bar{\mathbf{V}}^*$  read, respectively

$$\bar{\mathbf{V}} = \frac{-i\omega_0}{k_0} \left\{ \begin{bmatrix} ik_0 \\ 0 \\ k_0 \end{bmatrix} A + \begin{bmatrix} i(k_0 z + 1) \\ 0 \\ 1 + k_0 z \end{bmatrix} (-i\partial_x A) \right\} + \mathcal{O}(\epsilon \delta^2), \quad (\text{A } 10a)$$

that leads to

$$\begin{aligned} \bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^* &= 2\omega_0^2 AA^* + \frac{\omega_0^2}{k_0} [iA\partial_x A^* (2 + 2k_0 z) - iA^* \partial_x A (2 + 2k_0 z)] \\ &\quad + \mathcal{O}(\epsilon^2 \delta^2), \end{aligned} \quad (\text{A } 11a)$$

$$\begin{aligned} \nabla_3 (\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^* e^{2k_0 z}) &= 2\omega_0^2 [\nabla, 2k_0] (AA^*) + \frac{\omega_0^2}{k_0} [0, 0, 6k_0] (iA\partial_x A^* - iA^* \partial_x A) \\ &\quad + \mathcal{O}(\epsilon^2 \delta^2) \quad \text{for } z = 0 \quad \text{and} \end{aligned} \quad (\text{A } 11b)$$

$$\partial_t (\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*) = -\frac{\omega_0^3}{k_0} (A^* \partial_x A + A \partial_x A^*) + \mathcal{O}(\delta^2 \epsilon^2), \quad (\text{A } 11c)$$

where the relations  $\partial_t A = -\omega_0 \partial_x A / (2k_0)$  and  $\partial_t \partial_x A^* = -\omega_0 A^* / (2k_0)$  were used in (A 11c). Thus, we obtain for unidirectional waves

$$\begin{aligned} \frac{1}{8} \nabla_3 [(\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*) e^{2k_0 z}] \cdot \bar{\mathbf{V}} + \frac{1}{2} k_0 A \partial_t (\bar{\mathbf{V}} \cdot \bar{\mathbf{V}}^*) &= -\frac{1}{2} i\omega_0^3 k_0 A^2 A^* - \frac{1}{2} \omega_0^3 A \partial_x (AA^*) + \\ &\quad \omega_0^3 (A^2 \partial_x A^* - AA^* \partial_x A) + \mathcal{O}(\epsilon^3 \delta^2) \quad \text{for } z = 0. \end{aligned} \quad (\text{A } 12)$$

Moreover, the second term in (4.5b) reads

$$(-i\omega_0 + \partial_t) \left[ \nabla_3 \Phi^{(20)} \cdot \bar{\mathbf{V}} \right] = -\frac{\omega_0^2}{k_0} A [ik_0 \partial_x \Phi^{(20)} + k_0 \partial_z \Phi^{(20)}] + \mathcal{O}(\epsilon^3 \delta^2). \quad (\text{A } 13)$$

Combing (A 12) and (A 13) leads to

$$\begin{aligned} \bar{Q}^{(31)} &= -\frac{1}{2} i\omega_0^3 k_0 A^2 A^* + \omega_0^3 (A^2 \partial_x A^* - AA^* \partial_x A) - \frac{1}{2} \omega_0^3 A \partial_x (AA^*) \\ &\quad - i\omega_0^2 A \partial_x \Phi^{(20)} - \omega_0^2 A \partial_z \Phi^{(20)} + \mathcal{O}(\epsilon^3 \delta^2). \end{aligned} \quad (\text{A } 14)$$

Introducing  $C = -\frac{i\omega_0 A}{2k_0}$  and inserting it for  $A$  into (A 14) yields (4.17) in the main.



## Appendix B. Wave elevation at second order by Dalzell (1999)

Based on Dalzell (1999),  $\zeta^{(20)}$  and  $\zeta^{(22)}$  are given by, respectively,

$$\zeta^{(20)}(\mathbf{x}, t) = \frac{1}{2} \sum_{n=1}^N \sum_{m=1}^M A_m^{(20)} |\hat{\zeta}^{(1)}(\mathbf{k}_n)| |\hat{\zeta}^{(1)}(\mathbf{k}_m)| \cos(\psi_n - \psi_m), \quad (\text{B } 1)$$

$$\zeta^{(22)}(\mathbf{x}, t) = \frac{1}{2} \sum_{n=1}^N \sum_{m=1}^M A_m^{(22)} |\hat{\zeta}^{(1)}(\mathbf{k}_n)| |\hat{\zeta}^{(1)}(\mathbf{k}_m)| \cos(\psi_n + \psi_m), \quad (\text{B } 2)$$

where  $\hat{\zeta}_n^{(1)}$  and  $\hat{\zeta}_m^{(1)}$  denotes the Fourier transform of  $\zeta^{(1)}(\mathbf{x}, t)$ ,  $\psi_n = \mathbf{k}_n \cdot \mathbf{x} + \text{ang}(\hat{\zeta}^{(1)}(\mathbf{k}_n))$ ,  $\psi_m = \mathbf{k}_m \cdot \mathbf{x} + \text{ang}(\hat{\zeta}^{(1)}(\mathbf{k}_m))$ ,  $\mathbf{k}_n = |\mathbf{k}_n|(\cos \theta_n, \sin \theta_n)$ ,  $\mathbf{k}_m = |\mathbf{k}_m|(\cos \theta_m, \sin \theta_m)$ ,

$$A_m^{(20)} = \frac{\omega_n^2 + \omega_m^2}{2g} + \frac{\omega_1 \omega_2}{2g} (1 + \cos(\theta_n - \theta_m)) \frac{(\omega_1 - \omega_2)^2 + g(|\mathbf{k}_n| - |\mathbf{k}_m|)}{(\omega_1 - \omega_2)^2 - g(|\mathbf{k}_n| - |\mathbf{k}_m|)}, \quad (\text{B } 3)$$

$$A_p^{(22)} = \frac{\omega_n^2 + \omega_m^2}{2g} - \frac{\omega_1 \omega_2}{2g} (1 - \cos(\theta_n - \theta_m)) \frac{(\omega_1 + \omega_2)^2 + g(|\mathbf{k}_n| + |\mathbf{k}_m|)}{(\omega_1 + \omega_2)^2 - g(|\mathbf{k}_n| + |\mathbf{k}_m|)}. \quad (\text{B } 4)$$

## Appendix C. Additional results

This section shows additional results for second-order superharmonic and subharmonic elevations predicted by the present paper, compared with these due to Dalzell (1999) and two other methods, as shown in figure 13 and figure 14, respectively. Figure 13 and figure 14 show the spatial distribution of the second-order elevations driven by a focused wave group at three instants, based on a JONSWAP (amplitude) spectrum with the peak enhancement factor of 3.3 in frequency. For the results based on the semi-analytical approach, the spectrum was truncated at the high wavenumber side up to  $k = 15k_p$  with  $k_p$  the spectrum peak wavenumber. Agreement between Dalzell (1999) and the semi-analytical approach is clearly seen in figure 13 and 14.

## Appendix D. The modified evolution equation for $A$ by Trulsen *et al.* (2000)

Using the notation herein, equations (19-22) by Trulsen *et al.* (2000) read

$$\partial_t A + i\omega_0 \mathcal{L}A + \frac{ik_0^2 \omega_0}{2} |A|^2 A + \frac{3k_0 \omega_0}{2} |A|^2 \partial_x A - \frac{k_0 \omega_0}{4} A^2 \partial_x A^* + ik_0 \partial_x \Phi^{(20)} = 0, \quad (\text{D } 1a)$$

$$\partial_z \Phi^{(20)} = \frac{\omega_0}{2} \partial_x |A|^2 \text{ for } z = 0, \quad (\text{D } 1b)$$

$$\partial_z \Phi^{(20)} = 0 \text{ for } z \rightarrow -\infty, \quad (\text{D } 1c)$$

it should be noted that the sign in front of the term that reads ' $k_0 \omega_0 A^2 \partial_x A^* / 4$ ' is positive in Trulsen *et al.* (2000) whereas the negative sign is used in this paper due to the different definition of  $A$  in this paper from that by Trulsen *et al.* (2000), consistent with equation (2.1) in Lo & Mei (1985) and equation (14) by Stiassnie (1984). In Trulsen *et al.* (2000),  $A$  is defined as the sum of the first- and third-order elevation envelope of the first harmonic.

## REFERENCES

- BARRATT, D., BINGHAM, H. B. & ADCOCK, THOMAS A. A. 2020 Nonlinear evolution of a steep, focusing wave group in deep water simulated with oceanwave3d. *J. Offshore Mech. Arct. Eng.* **142** (2).




FIGURE 13. A comparison of second-order superharmonic wave elevation  $\zeta^{(22)}$  for a unidirectional wave group at different times between the predictions by (3.13b) (black solid line), Dalzell (1999) (red dashed line), and (6.4) (blue dot-dashed line) that is based on a second-order Stokes wave theory. The values for  $k_p$ ,  $x_f$ ,  $t_f$ , and  $t_0$  are the same as figure 2 except that a JONSWAP (amplitude) spectrum with the peak enhancement factor of 3.3 in frequency was used in this figure. The top panels for  $t = -10T_0$  with  $T_0$  the peak wave period, the middle panels for  $t = 20T_p$ , and the bottom panels for  $t = 40T_0$ . The right panels show the area highlighted by the thick blue box in the left panels, respectively.




FIGURE 14. A comparison of elevation  $\zeta^{(20)}$  for the second-order subharmonic waves forced by a unidirectional wave group at different times between the predictions by (3.13c) (black solid line), Dalzell (1999) (red dashed line), and Trulsen *et al.* (2000) (blue dot-dashed line). The detailed parameters are the same as figure 13.

- BARRATT, D., BINGHAM, H. B., TAYLOR, P. H., VAN DEN BREMER, T. S. & ADCOCK, T. A. A 2021 Rapid spectral evolution of steep surface wave groups with directional spreading. *J. Fluid Mech.* **907**.
- BATEMAN, W. J. D., SWAN, C. & TAYLOR, P. H. 2001 On the efficient numerical simulation of directionally spread surface water waves. *Journal of Computational Physics* **174** (1), 277–305.
- BENJAMIN, T. B. 1967 Instability of periodic wavetrains in nonlinear dispersive systems. *Proc. R. Soc. London, Ser. A. Math. & Phys. Sci.* **299** (1456), 59–76.
- BENJAMIN, T. B. & FEIR, J. E. 1967 The disintegration of wave trains on deep water Part 1. Theory. *J. Fluid Mech.* **27** (3), 417–430.
- BENNEY, D. J. & ROSKES, G. J. 1969 Wave instabilities. *Stud. Appl. Math.* **48** (4), 377–385.
- BIHS, H., WANG, W., PAKOZDI, C. & KAMATH, A. 2020 REEF3D: FNPF–A flexible fully nonlinear potential flow solver. *J. Offshore Mech. Arct. Eng.* **142** (4).
- BONNEFOY, F., LE TOUZÉ, D. & FERRANT, P. 2006 A fully-spectral 3d time-domain model for second-order simulation of wavetank experiments. part a: Formulation, implementation and numerical properties. *Appl. Ocean Res.* **28** (1), 33–43.
- BRINCH-NIELSEN, U. & JONSSON, I. G. 1986 Fourth order evolution equations and stability analysis for Stokes waves on arbitrary water depth. *Wave Motion* **8** (5), 455–472.
- CHABCHOUB, A., HOFFMANN, N. P. & AKHMEDIEV, N. 2011 Rogue wave observation in a water wave tank. *Phys. Rev. Lett.* **106** (20), 204502.
- CHU, V. H. & MEI, C. C. 1971 The non-linear evolution of Stokes waves in deep water. *J. Fluid Mech.* **47** (2), 337–351.
- CRAWFORD, D. R., LAKE, B. M., SAFFMAN, P. G. & YUEN, H. C. 1981 Stability of weakly nonlinear deep-water waves in two and three dimensions. *J. Fluid Mech.* **105**, 177–191.
- CRAWFORD, D. R., SAFFMAN, P. G. & YUEN, H. C. 1980 Evolution of a random inhomogeneous field of nonlinear deep-water gravity waves. *Wave motion* **2** (1), 1–16.
- CURTIS, C. W., CARTER, J. D. & KALISCH, H. 2018 Particle paths in nonlinear Schrödinger models in the presence of linear shear currents. *J. Fluid Mech.* **855**.
- DALZELL, J. F. 1999 A note on finite depth second-order wave–wave interactions. *Appl. Ocean Res.* **21** (3), 105–111.
- DAVEY, A. & STEWARTSON, K. 1974 On three-dimensional packets of surface waves. *Proc. R. Soc. London, Ser. A. Math. & Phys. Sci.* **338** (1613), 101–110.
- DIAS, F., DYACHENKO, A. I. & ZAKHAROV, V. E. 2008 Theory of weakly damped free-surface flows: a new formulation based on potential flow solutions. *Phys. Lett. A* **372** (8), 1297–1302.
- DJORDJEVIĆ, V. D. & REDEKOPP, L. G. 1978 On the development of packets of surface gravity waves moving over an uneven bottom. *Zeitschrift für angewandte Mathematik und Physik ZAMP* **29** (6), 950–962.
- DOMMERMUTH, D. G. & YUE, D. K. P. 1987 A high-order spectral method for the study of nonlinear gravity waves. *J. Fluid Mech.* **184**, 267–288.
- DUCROZET, GUILLAUME, BONNEFOY, FÉLICIEN, LE TOUZÉ, DAVID & FERRANT, PIERRE 2012 A modified high-order spectral method for wavemaker modeling in a numerical wave tank. *Eur. J. Mech. B Fluids* **34**, 19–34.
- DYSTHE, K.B. & DAS, K.P. 1981 Coupling between a surface-wave spectrum and an internal wave: modulational interaction. *J. Fluid Mech.* **104**, 483–503.
- DYSTHE, K., TRULSEN, K., KROGSTAD, H. E. & SOCQUET-JUGLARD, H. 2003 Evolution of a narrow-band spectrum of random surface gravity waves. *J. Fluid Mech.* **478**, 1–10.
- DYSTHE, K. B. 1979 Note on a modification to the nonlinear Schrödinger equation for application to deep water waves. *Proc. R. Soc. London, Ser. A. Math. & Phys. Sci.* **369** (1736), 105–114.
- ENGSIG-KARUP, A. P., BINGHAM, H. B. & LINDBERG, O. 2009 An efficient flexible-order model for 3D nonlinear water waves. *J. Comput. Phys.* **228** (6), 2100–2118.
- FRIGO, M. & JOHNSON, S. G. 1998 FFTW: An adaptive software architecture for the FFT. In *Proceedings of the 1998 IEEE International Conference on Acoustics, Speech and Signal Processing, ICASSP'98 (Cat. No. 98CH36181)*, , vol. 3, pp. 1381–1384. IEEE.
- GIBBS, R. H. & TAYLOR, P. H. 2005 Formation of walls of water in ‘fully’nonlinear simulations. *Appl. Ocean Res.* **27** (3), 142–157.

- GIBSON, R. S. & SWAN, C. 2007 The evolution of large ocean waves: the role of local and rapid spectral changes. *Proc. R. Soc. London, Ser. A. Math. & Phys. Sci.* **463** (2077), 21–48.
- GRAMSTAD, O. 2014 The zakharov equation with separate mean flow and mean surface. *J. Fluid Mech.* **740**, 254.
- GRAMSTAD, O. & TRULSEN, K. 2011 Fourth-order coupled nonlinear Schrödinger equations for gravity waves on deep water. *Phys. Fluids* **23** (6), 062102.
- HAMMACK, J. L. & HENDERSON, D. M. 1993 Resonant interactions among surface water waves. *Annu. Rev. Fluid Mech.* **25** (1), 55–97.
- HASSELMANN, K. 1962 On the non-linear energy transfer in a gravity-wave spectrum. *J. Fluid Mech* **12** (15), 481–500.
- HJELMERVIK, K. B. & TRULSEN, K. 2009 Freak wave statistics on collinear currents. *J. Fluid Mech* **637**, 267.
- JANSSEN, P. A. E. M. 1983 On a fourth-order envelope equation for deep-water waves. *J. Fluid Mech.* **126**, 1–11.
- JANSSEN, P. A. E. M. 2004 *The interaction of ocean waves and wind*. Cambridge University Press.
- JANSSEN, T. T. & HERBERS, T. H. C. 2009 Nonlinear wave statistics in a focal zone. *J. Phys. Oceanogr.* **39** (8), 1948–1964.
- JOHNSON, R. S. 1977 On the modulation of water waves in the neighbourhood of  $kh \approx 1.363$ . *Proc. R. Soc. London, Ser. A. Math. & Phys. Sci.* **357** (1689), 131–141.
- KHARIF, C., KRAENKEL, R. A., MANNA, M. A. & THOMAS, R. 2010 The modulational instability in deep water under the action of wind and dissipation. *J. Fluid Mech.* pp. 138–149.
- KIRBY, J. T. & DALRYMPLE, R. A. 1983 A parabolic equation for the combined refraction diffraction of Stokes waves by mildly varying topography. *J. Fluid Mech.* **136**, 453–466.
- KRASITSKII, VLADIMIR P 1994 On reduced equations in the hamiltonian theory of weakly nonlinear surface waves. *J. Fluid Mech.* **272**, 1–20.
- LAKE, B. M., YUEN, H. C., RUNGALDIER, H. & FERGUSON, W. E. 1977 Nonlinear deep-water waves: theory and experiment. Part 2. Evolution of a continuous wave train. *J. Fluid Mech.* **83** (1), 49–74.
- LI, Y., DRAYCOTT, S., ADCKOCK, T. A. A. & VAN DEN BREMER, T. S. 2021a Surface wavepackets subject to an abrupt depth change. Part II: experimental analysis. *J. Fluid Mech.* **915** (A72).
- LI, Y., DRAYCOTT, S., ZHENG, Y., LIN, Z., ADCKOCK, T. A. A. & VAN DEN BREMER, T. S. 2021b Why rogue waves occur atop abrupt depth transitions. *J. Fluid Mech.* **919** (R2).
- LI, Y., ZHENG, Y.K., LIN, Z.L., ADCKOCK, T. A. A. & VAN DEN BREMER, T. S. 2021c Surface wavepackets subject to an abrupt depth change. Part I: second-order theory. *J. Fluid Mech.* **915** (A71).
- LIGHTHILL, MJ 1965 Contributions to the theory of waves in non-linear dispersive systems. *IMA J. Appl. Math.* **1** (3), 269–306.
- LO, E. & MEI, C. C. 1985 A numerical study of water-wave modulation based on a higher-order nonlinear Schrödinger equation. *J. Fluid Mech.* **150**, 395–416.
- LONGUET-HIGGINS, M. S. 1976 On the nonlinear transfer of energy in the peak of a gravity-wave spectrum: a simplified model. *Proc. R. Soc. London, Ser. A. Math. & Phys. Sci.* **347** (1650), 311–328.
- LONGUET-HIGGINS, M. S. 1978 The instabilities of gravity waves of finite amplitude in deep water ii. subharmonics. *Proc. R. Soc. London, Ser. A. Math. & Phys. Sci.* **360** (1703), 471–488.
- MADSEN, P. A & FUHRMAN, D. R 2006 Third-order theory for bichromatic bi-directional water waves. *J. Fluid Mech.* **557**, 369–397.
- MCLEAN, J. W. 1982 Instabilities of finite-amplitude water waves. *J. Fluid Mech.* **114**, 315–330.
- MCWILLIAMS, J. C, RESTREPO, J. M. & LANE, E. M. 2004 An asymptotic theory for the interaction of waves and currents in coastal waters. *J. Fluid Mech.* **511**, 135.
- MEI, C. C., STIASSNIE, M. & YUE, D. K. P. 2005 *Theory and applications of ocean surface waves: nonlinear aspects*, , vol. 23. World scientific.
- MELVILLE, W. K. 1982 The instability and breaking of deep-water waves. *J. Fluid Mech.* **115**, 165–185.

- ONORATO, MIGUEL, OSBORNE, ALFRED RICHARD & SERIO, MARINA 2007 On the relation between two numerical methods for the computation of random surface gravity waves. *European Journal of Mechanics-B/Fluids* **26** (1), 43–48.
- ONORATO, M., PROMENT, D. & TOFFOLI, A. 2011 Triggering rogue waves in opposing currents. *Phys. Rev. Lett.* **107** (18), 184502.
- ONORATO, M., WASEDA, T., TOFFOLI, A., CAVALERI, L., GRAMSTAD, O., JANSSEN, P. A. E. M., KINOSHITA, T., MONBALIU, J., MORI, N., OSBORNE, A. R. & TRULSEN, K. 2009 Statistical properties of directional ocean waves: the role of the modulational instability in the formation of extreme events. *Phys. Rev. Lett.* **102** (11), 114502.
- PHILLIPS, O. M. 1960 On the dynamics of unsteady gravity waves of finite amplitude part 1. the elementary interactions. *J. Fluid Mech.* **9** (2), 193–217.
- PHILLIPS, O. M. 1967 Theoretical and experimental studies of gravity wave interactions. *Proc. R. Soc. London, Ser. A. Math. & Phys. Sci.* **299** (1456), 104–119.
- SCHÄFFER, H. A. 1996 Second-order wavemaker theory for irregular waves. *Ocean Eng.* **23** (1), 47–88.
- SIMANESEW, A., KROGSTAD, H. E., TRULSEN, K. & BORGE, J. C. N 2016 Development of frequency-dependent ocean wave directional distributions. *Appl. Ocean Res.* **59**, 304–312.
- SIMANESEW, A., KROGSTAD, H. E., TRULSEN, K. & BORGE, J. C. N 2017 Surface wave predictions in weakly nonlinear directional seas. *Appl. Ocean Res.* **65**, 79–89.
- SLUNYAEV, A., SERGEEVA, A. & PELINOVSKY, E. 2015 Wave amplification in the framework of forced nonlinear Schrödinger equation: the rogue wave context. *Physica D: Nonlinear Phenomena* **303**, 18–27.
- SLUNYAEV, A. V. 2005 A high-order nonlinear envelope equation for gravity waves in finite-depth water. *Journal of Experimental and Theoretical Physics* **101** (5), 926–941.
- STIASSNIE, M. 1984 Note on the modified nonlinear Schrödinger equation for deep water waves. *Wave motion* **6** (4), 431–433.
- STOCKER, J.R. & PEREGRINE, D.H. 1999 The current-modified nonlinear schrödinger equation. *J. Fluid Mech.* **399**, 335–353.
- SU, M. Y. 1982 Evolution of groups of gravity waves with moderate to high steepness. *Phys. Fluids* **25** (12), 2167–2174.
- TAPPERT, F. 1974 Numerical solutions of the Korteweg-de Vries equation and its generalizations by the split-step Fourier method. *Nonlinear Wave Motion* **15**, 215–216.
- TRULSEN, K. & DYSTHE, K. B. 1996 A modified nonlinear Schrödinger equation for broader bandwidth gravity waves on deep water. *Wave Motion* **24** (3), 281–289.
- TRULSEN, K. & DYSTHE, K. B. 1997 Frequency downshift in three-dimensional wave trains in a deep basin. *J. Fluid Mech.* **352**, 359–373.
- TRULSEN, K., KLIAKHANDLER, I., DYSTHE, KRISTIAN B. & VELARDE, M. G. 2000 On weakly nonlinear modulation of waves on deep water. *Phys. Fluids* **12** (10), 2432–2437.
- TRULSEN, K., NIETO BORGE, J. C., GRAMSTAD, O., AOUF, L. & LEFÈVRE, J. 2015 Crossing sea state and rogue wave probability during the Prestige accident. *J. Geophys. Res. C: Oceans* **120** (10), 7113–7136.
- TRULSEN, K., RAUSTØL, A., JORDE, S. & RYE, L.B. 2020 Extreme wave statistics of long-crested irregular waves over a shoal. *J. Fluid Mech.* **882**.
- WEST, BRUCE J, BRUECKNER, KEITH A, JANDA, RALPH S, MILDER, D MICHAEL & MILTON, ROBERT L 1987 A new numerical method for surface hydrodynamics. *J. Geophys. Res.: Oceans* **92** (C11), 11803–11824.
- WHITHAM, G. B. 1967 Non-linear dispersion of water waves. *J. Fluid Mech.* **27** (2), 399–412.
- WU, G., LIU, Y. & YUE, D. K. P. 2006 A note on stabilizing the Benjamin-Feir instability. *J. Fluid Mech.* **556**, 45.
- ZAKHAROV, V. E. 1968 Stability of periodic waves of finite amplitude on the surface of a deep fluid. *J. Appl. Mech. Tech. Phys.* **9** (2), 190–194.
- ZHENG, Y., LIN, Z., LI, Y., ADCOCK, T.A.A., LI, Y. & VAN DEN BREMER, T.S. 2020 Fully nonlinear simulations of extreme waves provoked by strong depth transitions: the effect of slope. *Phys. Rev. Fluids* **5**, 064804.