



OPEN ACCESS

TOP2A gene copy number change in breast cancer

M J Engstrøm,¹ B Ytterhus,¹ L J Vatten,² S Opdahl,² A M Bofin¹

¹Department of Laboratory Medicine, Children's and Women's Health, Norwegian University of Science and Technology, Trondheim, Norway

²Department of Public Health and General Practice, Norwegian University of Science and Technology, Trondheim, Norway

Correspondence to

Dr Monica Jernberg Engstrøm, NTNU, Faculty of Medicine, Department for laboratory Medicine, Children's and Women's Health, Laboratoriseretret, St. Olav's Hospital, Erling Skjalgssons gate, Trondheim 7030, Norway; monica.j.engstrom@ntnu.no

Received 4 November 2013

Revised 3 December 2013

Accepted 9 December 2013

Published Online First

8 January 2014

ABSTRACT

Aims The clinical significance of *TOP2A* as a prognostic marker has not been clarified. The aims of this study were to investigate the frequency of *TOP2A* copy number change; to correlate *TOP2A* with *HER2* status, hormone receptor (HR) status and molecular subtype, and further to explore differences in breast cancer-specific survival according to *TOP2A* and *HER2*.

Methods In this study, *TOP2A*, *HER2* and chromosome 17 copy number were assessed in 670 cases of breast cancer using in situ hybridisation techniques. Gene to chromosome ratios ≥ 2 were classified as amplification. *TOP2A* deletion (gene to chromosome ratio ≤ 0.8) or monosomy (only one signal for both gene and chromosome in more than 75% of nuclei) were classified as gene loss.

Results A strong association between *TOP2A* change and HR and *HER2* status was found. During the first 5 years after diagnosis, the risk of death from breast cancer was significantly higher for cases with *HER2* amplification irrespective of *TOP2A* status.

Conclusions *TOP2A* copy number change was strongly associated with HR and *HER2* status and as a prognostic marker *TOP2A* is probably of limited value.

with high histopathological grade³ and high proliferation,⁴ but the clinical significance of *TOP2A* and its relationship to *HER2* have not been clarified.

The aims of this study were to investigate the frequency of *TOP2A* copy number change in a well-characterised cohort of women with breast cancer⁵ and to correlate *TOP2A* with *HER2* status, hormone receptor (HR) status and molecular subtype. A further objective was to explore differences in breast cancer-specific survival (BCSS) according to *TOP2A* and *HER2*.

MATERIALS AND METHODS**Study population**

A screening programme for early diagnosis of breast cancer was conducted by the Norwegian Cancer Registry between 1956 and 1959. The patients developed breast cancer in a time period with limited access to adjuvant treatment. None were treated with anthracyclines or trastuzumab. According to the guidelines at the time of diagnosis, 30.7% patients may have qualified for treatment with tamoxifen. The population has been described in detail previously.⁵⁻⁷ A total of 1393 women in the underlying population developed breast cancer in the follow-up period from 1961 to the end of 2008. Of these, 945 had tissue samples available at the Department of Pathology and Medical Genetics, St. Olav's Hospital, Trondheim, Norway, and 670 were suitable for assessment of *TOP2A* and *HER2* copy number. Survival data were generated after linkage between the Cause of Death Registry of Norway and the Norwegian Cancer Registry.

INTRODUCTION

The *HER2* gene has a well-established biological and clinical role in breast cancer, and the *HER2* amplicon on chromosome 17 harbours a number of genes involved in breast cancer pathophysiology. Copy number change among these genes is frequently observed though their significance remains to be clarified.¹

TOP2A is one of the genes close to *HER2* and its protein product, topoisomerase II α , is the molecular target of anthracycline treatment. *TOP2A* amplification status has been thought to be linked to response to treatment. However, data are conflicting and, as yet, unresolved.² *HER2* and *TOP2A* are associated

Specimen characteristics

All cases in this study have previously been classified according to histopathological type and grade and reclassified in molecular subtypes according to figure 1⁵ using oestrogen receptor (ER),




Figure 1 Classification algorithm for molecular subtyping.



Open Access
Scan to access more
free content

To cite: Engstrøm MJ, Ytterhus B, Vatten LJ, et al. *J Clin Pathol* 2014;**67**: 420-425.

Table 1 Descriptive statistics of the 670 breast cancer cases

	<i>TOP2A</i> normal	<i>TOP2A</i> amplified	<i>TOP2A</i> loss	<i>HER2</i> normal	<i>HER2</i> amplified	Co-amplified	<i>TOP2A</i> loss, <i>HER2</i> amplified	<i>TOP2A</i> amplified, <i>HER2</i> normal	<i>TOP2A</i> loss, <i>HER2</i> normal	Normal <i>TOP2A</i> and <i>HER2</i>	Total
Number (%)	604 (90.2)	41 (6.1)	25 (3.7)	560 (83.6)	110 (16.4)	32 (4.8)	6 (0.9)	9 (1.3)	19 (2.8)	532 (79.4)	670
Mean age at diagnosis (SD)	73.4 (9.7)	69.5 (9.6)	72.0 (12.5)	74.0 (9.3)	68.3 (11.0)	68.2 (10.1)	69.2 (13.6)	74.3 (5.6)	72.8 (12.3)	74.0 (9.2)	73.1 (9.8)
Median years of follow-up after diagnosis (IQR)	6.7 (9.4)	5.8 (11.9)	6.4 (5.9)	7.1 (9.1)	4.5 (10.6)	5.1 (12.6)	5.0 (7.9)	6.0 (10.0)	6.7 (8.1)	7.1 (9.2)	6.6 (9.4)
Tumour grade (%)											
1	71 (11.8)	1 (2.4)	0	71 (12.7)	1 (0.9)	0	0	1 (11.1)	0	70 (13.2)	72 (10.8)
2	319 (52.8)	16 (39.0)	16 (64.0)	318 (56.8)	33 (30.0)	11 (34.4)	2 (33.3)	5 (55.6)	14 (73.7)	299 (56.2)	351 (52.4)
3	214 (35.4)	24 (58.5)	9 (36.0)	171 (30.5)	76 (69.1)	21 (65.6)	4 (66.7)	3 (33.3)	5 (26.3)	163 (30.6)	247 (36.9)
Tumour size (%)											
<2	136 (22.5)	8 (19.5)	5 (20.0)	135 (24.1)	14 (12.7)	5 (15.6)	0	3 (33.3)	5 (26.3)	127 (23.9)	149 (22.2)
2-5	292 (48.3)	15 (36.6)	7 (28.0)	270 (48.2)	44 (40.0)	14 (43.8)	3 (50.0)	1 (11.1)	4 (21.1)	265 (49.8)	314 (46.9)
>5	41 (6.8)	3 (7.3)	5 (20.0)	33 (5.9)	16 (14.6)	2 (6.3)	1 (16.7)	1 (11.1)	4 (21.1)	28 (5.3)	49 (7.3)
Uncertain	135 (22.4)	15 (36.6)	8 (32.0)	122 (21.8)	36 (32.7)	11 (34.4)	2 (33.3)	4 (44.4)	6 (31.6)	112 (21.1)	158 (23.6)
Molecular subtypes (%)											
Luminal A	300 (49.7)	7 (17.1)	10 (40.0)	317 (56.6)	0	0	0	7 (77.8)	10 (52.6)	300 (56.4)	317 (47.3)
Luminal B (HER2-)	166 (27.5)	1 (2.4)	6 (24.0)	173 (30.9)	0	0	0	1 (11.1)	6 (31.6)	166 (31.2)	173 (25.8)
Luminal B (HER2+)	37 (6.1)	23 (56.1)	1 (4.0)	0	61 (55.5)	23 (71.9)	1 (16.7)	0	0	0	61 (9.1)
HER2 type	35 (5.8)	8 (19.5)	5 (20.0)	0	49 (44.6)	9 (28.1)	5 (83.3)	0	0	0	49 (7.3)
Five negative phenotype	22 (3.6)	0	0	22 (3.9)	0	0	0	0	0	22 (4.1)	22 (3.3)
Basal phenotype	44 (7.3)	2 (4.9)	3 (12.0)	48 (8.6)	0	0	0	1 (11.1)	3 (15.8)	44 (8.3)	48 (7.2)
Hormone receptor											
Positive	503 (83.3)	31 (75.6)	17 (68.0)	490 (87.5)	61 (55.5)	23 (71.9)	1 (16.7)	8 (88.9)	16 (84.2)	466 (87.6)	551 (82.2)
Negative	101 (16.7)	10 (24.4)	8 (32.0)	70 (12.5)	49 (44.5)	9 (28.1)	5 (83.3)	1 (11.1)	3 (15.8)	66 (12.4)	119 (17.8)

progesterone receptor (PR), Ki67, cytokeratin 5 and epithelial growth factor receptor (EGFR) 1 as surrogate markers for gene expression. *HER2* status was assessed using chromogenic in situ hybridisation (CISH).

Assay methods

For the present study, fluorescence in situ hybridisation (FISH) was employed for detection of *TOP2A* and chromosome 17 according to the manufacturer's guidelines. Pretreatment was done using Histology FISH Accessory Kit, code K5799 (Dako). The probe mix (VYSIS *TOP2A*/CEP 17 FISH Probe Kit, code 03N89-020 Abbott Molecular Inc) was applied and denatured at 73°C for 5 min before hybridisation at 37°C overnight. For *HER2* and chromosome 17, the *HER2* CISH pharmDx Kit, code 109 (Dako), was used and immunostaining for ER (ER SP1 Cell Marque 33 mg/mL 1:100) and PR (PR 16 Novocastra 360 mg/mL 1:400) was done in a DakoCytomation Autostainer Plus (Dako) using Dako REAL EnVision Detection System with Peroxidase/DAB+, Rabbit/Mouse, code K5007, as previously described.⁵

Scoring and reporting

TOP2A gene copy number was evaluated under a fluorescence microscope (Nikon Eclipse 90i) and *HER2* gene under a bright field microscope (Nikon Eclipse 80i) by three of the authors (AMB, BY and MJE). A minimum of 20 non-overlapping tumour cell nuclei with signals for both chromosome and gene were counted in each case. Gene to chromosome ratios ≥ 2 were classified as amplification.^{8–11} *TOP2A* was considered to be deleted when the gene to chromosome ratio was ≤ 0.8 .^{9–12} Cases with only one signal for both gene and chromosome in more than 75% of all nuclei were recorded as monosomy. In the analyses, deletion and monosomy were grouped together. ER and PR were classified as positive when $\geq 1\%$ of the tumour cells showed positive nuclear staining.

Statistical analyses

Follow-up was from breast cancer diagnosis to death from breast cancer, death from any other cause or to December 31, 2010, whichever occurred first. BCSS was estimated using the Kaplan–Meier method, and Cox proportional hazards models were used to estimate risk of death from breast cancer. HRs were calculated with 95% CIs using Stata V.12.1 IC for Windows (Stata Corp).

RESULTS

Description of breast cancer cases

Of the 670 cases, 251 (37.5%) died of breast cancer, 314 (46.9%) died of other causes, and at the end of the observation period, 105 (15.6%) were still alive. Mean age at diagnosis was 73.1 years (SD 9.8; range 41–96 years), and median follow-up was 6.6 years (IQR 9.42 years). Histopathological grade, tumour size and molecular subtypes are given in table 1.

Amplification and deletion

Table 2 shows amplification of *TOP2A* was found in 41 cases (6.1%) and monosomy or deletion in 25 (3.7%). *HER2* was amplified in 110 cases (16.4%) and co-amplified with *TOP2A* in 32 cases (4.8%). Of the 25 cases with *TOP2A* loss, 6 were amplified for *HER2*. The majority with *TOP2A* amplification (78.1%) were co-amplified with *HER2*, whereas 34.5% of the *HER2* amplified tumours were either *TOP2A* amplified or showed *TOP2A* loss. The proportion of HR+ tumours was higher among cases with *TOP2A* amplification (75.6%) and

Table 2 Number of positive and negative cases for each marker

IHC (%)	<i>TOP2A</i> normal	<i>TOP2A</i> amplified	<i>TOP2A</i> loss	Total
HER2+	72 (11.9)	32 (78.1)	6 (24.0)	110 (16.4)
HER2–	532 (88.1)	9 (21.9)	19 (76.0)	560 (83.6)
ER+	500 (82.8)	31 (75.6)	17 (68.0)	548 (81.8)
ER–	102 (16.9)	10 (24.4)	8 (32.0)	120 (17.9)
PR+	361 (59.8)	19 (46.3)	5 (20.0)	385 (57.5)
PR–	243 (40.2)	22 (53.7)	20 (80.0)	285 (42.5)
Ki67 >15%	270 (44.7)	24 (58.5)	13 (52.0)	307 (45.8)
Ki67 >15%	333 (55.1)	17 (41.5)	12 (48.0)	362 (54.0)
CK5+	115 (19.0)	9 (21.9)	5 (20.0)	129 (19.3)
CK5–	489 (81.0)	32 (78.1)	20 (80.0)	541 (80.8)
EGFR+	46 (7.6)	1 (2.4)	3 (12.0)	50 (7.5)
EGFR–	558 (92.4)	40 (97.6)	22 (88.0)	620 (92.5)
Total	604 (90.2)	41 (6.1)	25 (3.7)	670 (100.0)

ER, oestrogen receptor; PR, progesterone receptor.

TOP2A loss (68.0%) compared with *HER2* amplification (55.5%).

Amplification and loss according to molecular subtypes

With the exception of 5NP, *TOP2A* copy number aberrations were found in all subtypes and were associated with both HR and *HER2* status. A majority of 56.1% of *TOP2A* amplified cases were Luminal B (*HER2*+). Loss of *TOP2A* was found among the HR+ and *HER2* negative subtypes (Luminal A and Luminal B (*HER2*–)) (64.0%) or *HER2* subtype (20.0%). One of four *TOP2A* deleted case was Luminal B (*HER2*+).

BCSS, *TOP2A* and *HER2*

The Kaplan–Meier plots in figures 2 and 3 show BCSS according to *TOP2A* and *HER2*, respectively, and in figure 4 the BCSS according to the status of both genes. Loss of *TOP2A* in the absence of *HER2* amplification did not affect BCSS. The Kaplan–Meier plots show poorest survival in *HER2*-amplified cases and *TOP2A* aberrations did not affect this.




Figure 2 Kaplan–Meier plot. Breast cancer-specific survival (BCSS) according to *TOP2A*. p Value from log-rank test of differences in BCSS first 5 years after diagnosis was 0.02. After 5 years, the p value was 0.4.




Figure 3 Kaplan–Meier plot. Breast cancer-specific survival (BCSS) according to *HER2*. p Value from log-rank test of differences in BCSS first 5 years after diagnosis was <0.0001. After 5 years, the p value was 0.9.

Risk of death from breast cancer, *TOP2A*, *HER2* and HR status

During the first 5 years, risk of death from breast cancer appears to be significantly higher in cases with amplification of *TOP2A*

and *HER2* when analysed separately. When compared with no amplification for *TOP2A* and *HER2*, respectively, the HR for *TOP2A* amplification was 2.03 (95% CI 1.22 to 3.360) and for *HER2* was 2.77 (95% CI 1.97 to 3.89). Adjusting for age and stage did not change the results. For those who survived the first 5 years after diagnosis, there were no statistically significant differences in survival according to gene amplification status.

However, as shown in table 3 and figure 4, *TOP2A* did not exert an independent effect on prognosis. Adjusting for HR status in the Cox proportional hazards model did not change the results (data not shown). During the first 5 years after diagnosis, the risk of death from breast cancer was significantly higher for HR+ cases with *HER2* amplification irrespective of *TOP2A* status. Among the HR– cases, the numbers in each category were low and the results must be interpreted with caution.

DISCUSSION

TOP2A gene copy number change in breast cancer is an infrequent finding and its significance has been difficult to establish. In this study of 670 cases of breast cancer with long-term follow-up, the number of cases with *TOP2A* amplification or loss was far lower than the number of *HER2*-positive cases. However, there was a large proportion of co-amplification. In contrast to others who have found that amplification of one or both genes entails a poorer prognosis compared with cases with no amplification,^{11 13 14} this study demonstrates that

Table 3 Risk of death from breast cancer according to *TOP2A* and *HER2* amplification

	Number of cases	Deaths from breast cancer	Hazard ratio 95% CI unadjusted		Hazard ratio 95% CI adjusted for age		Hazard ratio 95% CI adjusted for stage	
<i>TOP2A</i>								
Follow-up first 5 years after diagnosis	604	132	1.00		1.00		1.00	
Not amplified	41	17	2.03	1.22 to 3.36	2.07	1.24 to 3.47	2.11	1.27 to 3.50
Amplified	25	5	0.91	0.37 to 2.21	0.82	0.33 to 2.01	0.70	0.29 to 1.73
Loss	670	154						
<i>TOP2A</i>								
Follow-up from 5 years after diagnosis*	359	87	1.00		1.00		1.00	
Not amplified	22	5	0.75	0.30 to 1.85	0.74	0.30 to 1.86	1.02	0.41 to 2.54
Amplified	15	5	1.63	0.66 to 4.03	1.93	0.77 to 4.84	1.41	0.56 to 3.52
Loss	396	97						
<i>HER2</i>								
Follow-up first 5 years after diagnosis	560	105	1.00		1.00		1.00	
Not amplified	110	49	2.77	1.97 to 3.89	2.81	1.95 to 4.04	2.66	1.89 to 3.75
Amplified	670	154						
<i>HER2</i>								
Follow-up from 5 years after diagnosis*	346	83	1.00		1.00		1.00	
Not amplified	50	14	0.95	0.54 to 1.67	0.95	0.52 to 1.73	1.04	0.60 to 1.86
Amplified	396	97						
<i>HER2 and TOP2A</i>								
Follow-up first 5 years after diagnosis	532	100	1.00		1.00		1.00	
Normal <i>TOP2A</i> and <i>HER2</i>	38	17	2.61	1.56 to 4.36	2.76	1.63 to 4.69	2.68	1.60 to 4.51
<i>TOP2A</i> change and <i>HER2</i> amplification	28	5	0.96	0.39 to 2.37	0.89	0.36 to 2.21	0.77	0.31 to 1.90
<i>TOP2A</i> change and <i>HER2</i> normal	72	32	2.86	1.92 to 4.26	2.81	1.84 to 4.29	2.59	1.74 to 3.87
Amplified <i>HER2</i> , <i>TOP2A</i> normal	670	154						
<i>HER2 and TOP2A</i>								
Follow-up from 5 years after diagnosis*	328	79	1.00		1.00		1.00	
Normal <i>TOP2A</i> and <i>HER2</i>	19	6	0.99	0.43 to 2.28	0.97	0.41 to 2.28	1.44	0.62 to 3.37
<i>TOP2A</i> change and <i>HER2</i> amplification	18	4	1.07	0.39 to 2.94	1.26	0.45 to 3.50	0.91	0.33 to 2.51
<i>TOP2A</i> change and <i>HER2</i> normal	31	8	0.92	0.45 to 1.91	0.95	0.44 to 2.04	0.84	0.40 to 1.79
Amplified <i>HER2</i> , <i>TOP2A</i> normal	396	97						

*Conditional on surviving the first 5 years CI.




Figure 4 Kaplan-Meier plot. Breast cancer-specific survival (BCSS) according to *TOP2A* and *HER2*. p Value from log-rank test of differences in BCSS first 5 years after diagnosis was <0.0001. After 5 years, the p value was 1.0.

associations between BCSS and *TOP2A* copy number change are not independent of *HER2* and HR status.

The most important finding in this study is the strong association between *TOP2A* copy number change and HR and *HER2* status. These markers are well established as prognostic and predictive factors, and are to a high degree decisive for treatment after surgery. To the best of our knowledge, few studies have been designed to examine the prognostic value of *TOP2A*, though it has been shown that *TOP2A* amplification affects BCSS and risk of death from breast cancer¹⁵ and that *TOP2A* may be a prognostic marker in ER+ breast cancer.^{14 16} However, when the analyses include HR and *HER2* status, the present study shows that *TOP2A* has no independent prognostic impact. *TOP2A* may still have some modulating effects on prognostication, but this is probably of limited benefit in clinical practice.

Twenty of twenty-five cases with *TOP2A* loss were PR-, and of these, 12 were ER+. PR negativeness is a predictor of poor prognosis and appears to be associated with *TOP2A* loss. However, in this study, survival tended to be better in PR- cases with loss of *TOP2A* compared with cases with normal or amplified *TOP2A* (data not shown).

The proportion of amplification and co-amplification of *TOP2A* and *HER2* in breast cancer varies between studies. *HER2* amplification is reported to be around 15%.² For *TOP2A*, amplification varies from 5% to 19%.^{3 17 18} In *HER2*-positive breast cancer, amplification of *TOP2A* varies from 25% to 42%.^{1 19} Both amplification and deletion of *TOP2A* in the absence of *HER2* amplification have been demonstrated.^{3 20} In the present study, 29.1% of the *HER2*-amplified cases were co-amplified with *TOP2A*. The proportion of *TOP2A* positive tumours in this study was lower than in other studies.² However, the frequency of *HER2* amplification is comparable with others, and this weighs against methodological problems. Furthermore, a short DNA probe for *TOP2A* was used to avoid overlap with *HER2*.²¹ This may in part account for the low number of *TOP2A*-amplified cases in this study compared with previous studies and may reflect the true frequency of this finding.

Assessment of loss should be carried out with caution in histopathological sections because nuclear truncation may lead

to a falsely low estimation of copy number. The cut-off for amplification is usually set at a gene/chromosome ratio of ≥ 2.0 , and for deletion the cut-off level ranges from 0.5 to 1.0.²¹ It is possible that monosomy may have an impact similar to loss of individual genes, but this is uncertain. In this study, only four cases showed deletion and monosomy and deletion were grouped together.

HER2-positive breast cancer has been shown to be more aggressive than *HER2*-negative breast cancer. Co-amplification with other genes, such as *STARD3* and *GRB7*, may contribute to and possibly strengthen this aggressive behaviour.² The proportion of amplification and co-amplification of *TOP2A* and *HER2* in breast cancer is low, and even in a series of 670 patients, the numbers are too low to draw reliable conclusions. As a prognostic marker, *TOP2A* is probably of limited value. *TOP2A* aberrations are strongly associated with HR and *HER2* status, and the importance of these markers in prognostication is still unchallenged.

Take-home messages

- ▶ *TOP2A* gene copy number change is an infrequent finding in breast cancer.
- ▶ There is a strong association between *TOP2A* copy number change and hormone receptor and *HER2* status.
- ▶ As a prognostic marker, *TOP2A* is probably of limited value, and hormone receptor and *HER2* status remain unchallenged.

Acknowledgements The authors thank the Department of Pathology and Medical Genetics, St. Olav's Hospital, for making the archives available for the study and the Cancer Registry of Norway for providing the patient data.

Contributors MJE contributed to interpretation of in situ markers, carried out statistical analyses, interpretation of the results and drafted the manuscript. BY participated in planning and performing the laboratory work, contributed to interpretation of in situ markers and to discussion and review of the manuscript. LJV contributed to discussion of the study and review of the manuscript. SO participated in acquisition of data and tissue blocks, discussion of the statistical analyses and reviewed the manuscript. AMB contributed to conception and design of the study, interpretation of the in situ markers, interpretation and analyses of the data, and draft and review of the manuscript. All authors read and approved the final manuscript.

Funding The study has received financial support from the Liaison Committee between the Central Norway Regional Health Authority and the Norwegian University of Science and Technology and the Cancer Fund, St. Olav's Hospital, Trondheim University Hospital, Norway.

Competing interests None.

Ethics approval Approval of the study and dispensation from the requirement of patient consent was granted by the Regional Committee for Medical and Health Sciences Research Ethics (REK, Midt-Norge, ref. nr: 836/2009).

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

REFERENCES

- 1 Fountzilias G, Dafni U, Bobos M, *et al*. Evaluation of the prognostic role of centromere 17 gain and *HER2*/topoisomerase II alpha gene status and protein expression in patients with breast cancer treated with anthracycline-containing adjuvant chemotherapy: pooled analysis of two Hellenic Cooperative Oncology Group (HeCOG) phase III trials. *BMC Cancer* 2013;13:163.

- 2 Jacot W, Fiche M, Zaman K, *et al*. The HER2 amplicon in breast cancer: Topoisomerase IIA and beyond. *Biochim Biophys Acta* 2013;1836:146–57.
- 3 Bofin AM, Ytterhus B, Hagmar BM. TOP2A and HER-2 gene amplification in fine needle aspirates from breast carcinomas. *Cytopathology* 2003;14:314–9.
- 4 Romero A, Martin M, Cheang MC, *et al*. Assessment of Topoisomerase II alpha status in breast cancer by quantitative PCR, gene expression microarrays, immunohistochemistry, and fluorescence in situ hybridization. *Am J Pathol* 2011;178:1453–60.
- 5 Engstrom MJ, Opdahl S, Hagen AI, *et al*. Molecular subtypes, histopathological grade and survival in a historic cohort of breast cancer patients. *Breast Cancer Res Treat* 2013;140:463–73.
- 6 Kvåle G, Heuch I, Eide G. A prospective study of reproductive factors and breast cancer. *Am J Epidemiol* 1987;126:831–41.
- 7 Opdahl S, Alsaker MD, Janszky I, *et al*. Joint effects of nulliparity and other breast cancer risk factors. *Br J Cancer* 2011;105:731–6.
- 8 Bartlett JM, Starczynski J, Atkey N, *et al*. HER2 testing in the UK: recommendations for breast and gastric in-situ hybridisation methods. *J Clin Pathol* 2011;64:649–53.
- 9 Di Leo A, Desmedt C, Bartlett JM, *et al*. HER2 and TOP2A as predictive markers for anthracycline-containing chemotherapy regimens as adjuvant treatment of breast cancer: a meta-analysis of individual patient data. *Lancet Oncol* 2011;12:1134–42.
- 10 Nielsen KV, Ejlersen B, Moller S, *et al*. Lack of independent prognostic and predictive value of centromere 17 copy number changes in breast cancer patients with known HER2 and TOP2A status. *Mol Oncol* 2012;6:88–97.
- 11 Sauter G, Lee J, Bartlett JM, *et al*. Guidelines for human epidermal growth factor receptor 2 testing: biologic and methodologic considerations. *J Clin Oncol* 2009;27:1323–33.
- 12 Olsen KE, Knudsen H, Rasmussen BB, *et al*. Amplification of HER2 and TOP2A and deletion of TOP2A genes in breast cancer investigated by new FISH probes. *Acta Oncol* 2004;43:35–42.
- 13 Starczynski J, Atkey N, Connelly Y, *et al*. HER2 gene amplification in breast cancer: a rogues' gallery of challenging diagnostic cases: UKNEQAS interpretation guidelines and research recommendations. *Am J Clin Pathol* 2012;137:595–605.
- 14 Rody A, Karn T, Ruckhaberle E, *et al*. Gene expression of topoisomerase II alpha (TOP2A) by microarray analysis is highly prognostic in estrogen receptor (ER) positive breast cancer. *Breast Cancer Res Treat* 2009;113:457–66.
- 15 Zaczek A, Markiewicz A, Supernat A, *et al*. Prognostic value of TOP2A gene amplification and chromosome 17 polysomy in early breast cancer. *Pathol Oncol Res* 2012;18:885–94.
- 16 Sparano JA, Goldstein LJ, Davidson NE, *et al*. TOP2A RNA expression and recurrence in estrogen receptor-positive breast cancer. *Breast Cancer Res Treat* 2012;134:751–7.
- 17 Jarvinen TA, Tanner M, Rantanen V, *et al*. Amplification and deletion of topoisomerase IIalpha associate with ErbB-2 amplification and affect sensitivity to topoisomerase II inhibitor doxorubicin in breast cancer. *Am J Pathol* 2000;156:839–47.
- 18 Park K, Kim J, Lim S, *et al*. Topoisomerase II-alpha (topoII) and HER2 amplification in breast cancers and response to preoperative doxorubicin chemotherapy. *Eur J Cancer* 2003;39:631–4.
- 19 Jarvinen TA, Tanner M, Barlund M, *et al*. Characterization of topoisomerase II alpha gene amplification and deletion in breast cancer. *Genes Chromosomes Cancer* 1999;26:142–50.
- 20 Hicks DG, Yoder BJ, Pettay J, *et al*. The incidence of topoisomerase II-alpha genomic alterations in adenocarcinoma of the breast and their relationship to human epidermal growth factor receptor-2 gene amplification: a fluorescence in situ hybridization study. *Hum Pathol* 2005;36:348–56.
- 21 Varga Z, Moelans CB, Zuerrer-Hardi U, *et al*. Topoisomerase 2A gene amplification in breast cancer. Critical evaluation of different FISH probes. *Breast Cancer Res Treat* 2012;133:929–35.



TOP2A gene copy number change in breast cancer

M J Engstrøm, B Ytterhus, L J Vatten, S Opdahl and A M Bofin

J Clin Pathol 2014 67: 420-425 originally published online January 8, 2014

doi: 10.1136/jclinpath-2013-202052

Updated information and services can be found at:
<http://jcp.bmj.com/content/67/5/420>

These include:

References

This article cites 21 articles, 4 of which you can access for free at:
<http://jcp.bmj.com/content/67/5/420#BIBL>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (75)
[Breast cancer](#) (472)
[Molecular genetics](#) (324)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>