

MODULES OF CONSTANT JORDAN TYPE
OVER QUANTUM COMPLETE INTERSECTIONS

PETTER ANDREAS BERGH, KARIN ERDMANN,
AND DAVID A. JORGENSEN

Received: October 24, 2019

Revised: August 16, 2020

Communicated by Julia Pevtsova

ABSTRACT. We initiate the study of modules of constant Jordan type for quantum complete intersections, and prove a range of basic properties. We then show that for these algebras, constant Jordan type is an invariant of Auslander-Reiten components. Finally, we classify modules with stable constant Jordan type $[1]$ or $[n - 1]$ in the 2-generator case.

2020 Mathematics Subject Classification: 16D50, 16E05, 16G70, 16S80, 16U80, 20C20

Keywords and Phrases: Constant Jordan type, quantum complete intersections

1 INTRODUCTION

Varieties for modular group representations were introduced and studied by Carlson and others in order to understand modules without having to rely on complete classifications. Many aspects in this approach are controlled by elementary abelian p -groups, making use of the fact that their group algebras are truncated polynomial algebras. In particular, the concept of rank variety has proved to be extremely powerful. These varieties control projectivity of finite-dimensional modules, and this is known as “Dade’s Lemma.”

Modules of constant Jordan type were introduced by Carlson, Friedlander and Pevtsova in [CFP]. By using the idea of π -points, they obtained versions not only for finite groups, but more generally for finite group schemes; see the book [Ben] for extensive discussions. It turns out that a substantial part of the theory relies on Dade's Lemma. Around the same time, in [BEH], rank varieties were introduced for certain q -complete intersections. It was proved that the analog of Dade's Lemma holds, namely that rank varieties control projectivity also in this case. It is therefore natural to study modules of constant Jordan type in this setting.

The aim of this paper is to start this investigation. We introduce the relevant algebras, and prove a range of general properties. For example, we show that constant Jordan type is preserved under taking direct sums and summands, and taking syzygies and cosyzygies. We prove that certain constant Jordan types are unachievable by modules. We also address the question of how abundant are modules of constant Jordan type.

We also show that in our setting, constant Jordan type is preserved under Auslander-Reiten translation, and moreover that constant Jordan type is an invariant of Auslander-Reiten components.

For the group algebra setting, it is known that modules with stable constant Jordan type $[1]$ or $[p - 1]$ are precisely the syzygies of the trivial module. One expects that this might generalize. In the last section, we prove a result in this direction. Namely, for the 2-generator case, with certain assumptions, we show that the modules with stable constant Jordan type $[1]$ or $[n - 1]$ are precisely the syzygies of the simple module.

For group representations and group schemes in general, the tools which were used are mostly based on the fact that the rings are Hopf algebras. This is not the case in our setting, with quantum complete intersections. This means that a lot of the machinery and techniques from the group scheme setting are not available, and consequently the proofs are more difficult. Also, it is not clear whether all the results in the group scheme setting actually generalize.

2 CONSTANT JORDAN TYPE

All modules we consider are assumed to be finitely generated left modules. Let k be a field, $n \geq 2$ a positive integer, and define n' by

$$n' = \begin{cases} n & \text{if } \text{char } k = 0 \\ n/p^r & \text{if } \text{char } k = p > 0 \text{ and } p^r \mid n \text{ but } p^{r+1} \nmid n \end{cases}$$

Furthermore, let $q \in k$ be a primitive n' th root of unity, and fix a positive integer $c \geq 2$. The algebra we shall consider is the quantum complete intersection

$$A_q^c = k\langle x_1, \dots, x_c \rangle / (x_i^n, x_i x_j - q x_j x_i (i < j))$$

which is a finite dimensional selfinjective algebra of dimension n^c .

Quantum complete intersections can be defined more generally, but the essential thing for us is that any linear form in the variables x_1, \dots, x_c is n -nilpotent. Namely, by [BEH, Lemma 2.3], given any nonzero c -tuple $\lambda = (\lambda_1, \dots, \lambda_c) \in k^c$, the element

$$u_\lambda = \lambda_1 x_1 + \dots + \lambda_c x_c$$

in A_q^c satisfies $u_\lambda^n = 0$ (and n is the smallest such power with $u_\lambda^n = 0$). Consequently, the subalgebra $k[u_\lambda]$ of A_q^c generated by u_λ is isomorphic to the n -dimensional truncated polynomial ring $k[x]/(x^n)$. Note that A_q^c is free both as a left and as a right module for this subalgebra.

Up to isomorphism, the truncated polynomial ring $k[x]/(x^n)$ admits n finitely generated indecomposable left modules, namely $k[x]/(x^i)$ for $1 \leq i \leq n$. Since $\dim_k k[x]/(x^i) = i$, these are uniquely determined by their dimensions. Now let M be an A_q^c -module. When we restrict the module to the subalgebra $k[u_\lambda]$, it decomposes into a direct sum of the n indecomposable $k[u_\lambda]$ -modules.

DEFINITION. Fix an A_q^c -module M .

(1) Let λ be a nonzero c -tuple in k^c , and M_i the indecomposable $k[u_\lambda]$ -module with $\dim_k M_i = i$. Then the *Jordan type* of M with respect to λ is

$$[1]^{d_1} [2]^{d_2} \dots [n]^{d_n}$$

if M as a $k[u_\lambda]$ -module decomposes as a direct sum $M \simeq M_1^{d_1} \oplus \dots \oplus M_n^{d_n}$. In this case, the *stable Jordan type* of M with respect to λ is

$$[1]^{d_1} \dots [n-1]^{d_{n-1}}$$

(2) The module M has *constant Jordan type* if its Jordan type is the same for all nonzero $\lambda \in k^c$.

The phrase *Jordan type* refers to the action of the truncated polynomial ring $k[x]/(x^n)$ on its n indecomposable modules M_1, \dots, M_n , where $M_i = k[x]/(x^i)$. For M_i has a k -vector space basis w_1, \dots, w_i , with $xw_j = w_{j+1}$ for $1 \leq j \leq i-1$, and so the action of x on M_i can be defined in terms of the transpose of the Jordan block

$$\begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \\ 0 & 0 & 0 & \dots & 0 \end{pmatrix}$$

of size i and with eigenvalue 0. Thus if M has Jordan type $[1]^{d_1} \dots [n]^{d_n}$ with respect to $\lambda = (\lambda_1, \dots, \lambda_c)$, then the action of $u_\lambda = \lambda_1 x_1 + \dots + \lambda_c x_c$ on M is given by the transpose of a Jordan matrix having d_i such Jordan blocks of size i for each $1 \leq i \leq n$.

To simplify the notation, it is convenient to write Jordan types in such a way that only the Jordan blocks that are actually involved appear. That is, if M is

a nonzero A_q^c -module and λ a nonzero c -tuple, then there are integers $1 \leq a_1 < \dots < a_t \leq n$ such that $M \simeq M_{a_1}^{d_{a_1}} \oplus \dots \oplus M_{a_t}^{d_{a_t}}$ when restricted to $k[u_\lambda]$ (where M_i is the indecomposable $k[u_\lambda]$ -module of dimension i and all the exponents d_{a_i} are nonzero). We shall then in most cases write the Jordan type of M with respect to λ as

$$[a_1]^{d_{a_1}} \dots [a_t]^{d_{a_t}}$$

instead of having to write

$$[1]^{d_1} \dots [n]^{d_n}$$

with $d_i = 0$ for $i \notin \{a_1, \dots, a_t\}$. Similarly we apply this short-hand notation also for the stable Jordan type. Furthermore, when $d_i = 1$ we just write $[i]$ instead of $[i]^{d_i}$.

EXAMPLE. Two trivial examples of indecomposable A_q^c -modules of constant Jordan type are k and A_q^c . For k , the Jordan type with respect to any nonzero λ is clearly $[1]$, whereas for A_q^c it is $[n]^{n^{c-1}}$ since A_q^c is free of rank n^{c-1} over $k[u_\lambda]$. More generally, denote the radical of A_q^c by \mathfrak{r} . The Loewy length of A_q^c is $(n-1)^c + 1$, and we claim that for every pair of integers $0 \leq s < t \leq (n-1)^c + 1$, the A_q^c -module $\mathfrak{r}^s/\mathfrak{r}^t$ has constant Jordan type (but may not be indecomposable, for example when $t = s + 1$). To see this, note that as a k -vector space, the module $\mathfrak{r}^s/\mathfrak{r}^t$ has a basis

$$\{x_1^{e_1} x_2^{e_2} \dots x_c^{e_c} \mid s \leq e_1 + \dots + e_c \leq t\}$$

Now let $\lambda = (\lambda_1, \dots, \lambda_c)$ be a nonzero c -tuple in k^c , and let i be the smallest index such that λ_i is nonzero. Using now that

$$x_i = \lambda_i^{-1} \left(u_\lambda - \sum_{j \neq i} \lambda_j x_j \right)$$

it is straightforward to check that

$$\{x_1^{e_1} \dots x_{i-1}^{e_{i-1}} u_\lambda^{e_i} x_{i+1}^{e_{i+1}} \dots x_c^{e_c} \mid s \leq e_1 + \dots + e_c \leq t\}$$

is another basis for the module $\mathfrak{r}^s/\mathfrak{r}^t$. Consequently, the module structure of $\mathfrak{r}^s/\mathfrak{r}^t$ when restricted to $k[u_\lambda]$ is the same as when restricted to $k[x_i]$. Moreover, it is easy to see that the Jordan type when we restrict to $k[x_i]$ is the same as when we restrict to $k[x_1]$. This shows that $\mathfrak{r}^s/\mathfrak{r}^t$ has constant Jordan type. As a specific example, take the indecomposable module A_q^c/\mathfrak{r}^3 , and assume first that $n \geq 3$ so that $x_i^2 \neq 0$. The monomials

$$\begin{array}{ccccccc} & & & & & & 1 \\ & & & & & & x_1 & \dots & x_c \\ & & & & & & x_1^2 & x_1 x_2 & \dots & x_1 x_c & x_i x_j \end{array}$$

form a basis for the module, where $2 \leq i \leq j \leq c$. Over the subalgebra $k[x_1]$, the subset $\{1, x_1, x_1^2\}$ is a basis for the indecomposable module of dimension 3, and the $c - 1$ subsets $\{x_i, x_1x_i\}$ for $2 \leq i \leq c$ form bases for indecomposable modules of dimension 2. Finally, the $c(c-1)/2$ subsets $\{x_ix_j\}$ for $2 \leq i < j \leq c$ form bases for indecomposable modules of dimension 1. Thus the module A_q^c/\mathfrak{r}^3 has constant Jordan type

$$[1]^{c(c-1)/2}[2]^{(c-1)}[3]$$

Note that if n were 2, then there would be no indecomposable $k[x_1]$ -module of dimension 3, but instead one more of dimension 2. Also, there would be $c - 1$ less indecomposable $k[x_1]$ -module of dimension 1. Therefore, in this case, the constant Jordan type of A_q^c/\mathfrak{r}^3 would be

$$[1]^{(c^2-3c+2)/2}[2]^c$$

The following result records some elementary properties on modules of constant Jordan type

PROPOSITION 2.1. (1) If M is an A_q^c -module of constant Jordan type $[1]^{d_1} \dots [n]^{d_n}$, then $\dim_k M = \sum_{i=1}^n id_i$. Moreover, the dual $\text{Hom}_k(M, k)$ has the same constant Jordan type as a right module.

(2) If M and N are A_q^c -modules of constant Jordan types $[1]^{d_1} \dots [n]^{d_n}$ and $[1]^{e_1} \dots [n]^{e_n}$, respectively, then the direct sum $M \oplus N$ has constant Jordan type $[1]^{d_1+e_1} \dots [n]^{d_n+e_n}$.

Proof. For the dual module, note that when dualizing an indecomposable module over $k[x]/(x^n)$, the result is an isomorphic indecomposable module. □

The modules of the form $\mathfrak{r}^s/\mathfrak{r}^t$ that we looked at in the example are well behaved and easy to deal with. In general, there does not seem to exist any effective method for determining whether or not an arbitrary indecomposable module has constant Jordan type. However, it turns out that such modules can be described in terms of certain open subsets of the affine space k^c , where we use the Zariski topology. We shall use this to prove the converse of the second part of Proposition 2.1, namely that the direct summand of a module of constant Jordan type also has constant Jordan type. An alternative method would be to adapt the arguments from [Ben, Sections 4.5 and 5.1], which are based on Carlson, Friedlander and Pevtsova’s original arguments from [CFP].

DEFINITION. For an A_q^c -module M and integer $1 \leq i \leq n - 1$, define

$$\mathcal{U}_M^i = \left\{ \lambda \in k^c \setminus \{0\} \mid \text{rank}(M \xrightarrow{u_\lambda^i} M) \geq \text{rank}(M \xrightarrow{u_\sigma^i} M) \text{ for all } \sigma \in k^c \right\}$$

In other words, the set \mathcal{U}_M^i is the collection of all nonzero λ for which the linear operator u_λ^i on M has maximal rank. Note that this is a nonempty set

by definition. Recall that for a truncated polynomial algebra $k[x]/(x^n)$, the rank of x^i as a linear operator on the t -dimensional module $M_t = k[x]/(x^t)$ is $\max\{0, t - i\}$. Consequently, if the Jordan type of M with respect to λ is $[1]^{d_1} [2]^{d_2} \cdots [n]^{d_n}$, that is, if M as a $k[u_\lambda]$ -module decomposes as $M_1^{d_1} \oplus \cdots \oplus M_n^{d_n}$, then the rank of the linear operator u_λ^i on M is

$$d_{i+1} + 2d_{i+2} + \cdots + (n - i)d_n$$

provided $i \leq n - 1$. We use this elementary fact in the proof of the following lemma, which shows that the sets we have just defined describe the modules of constant Jordan type.

LEMMA 2.2. *Let M be an A_q^c -module.*

(1) *If λ and σ belong to $\cap_{i=1}^{n-1} \mathcal{U}_M^i$, then the Jordan types of M with respect to λ and σ are the same.*

(2) *The module M has constant Jordan type if and only if $\cap_{i=1}^{n-1} \mathcal{U}_M^i = k^c \setminus \{0\}$.*

Proof. To prove (1), suppose that M decomposes as $M_1^{d_1} \oplus \cdots \oplus M_n^{d_n}$ over $k[u_\lambda]$, and as $M_1^{e_1} \oplus \cdots \oplus M_n^{e_n}$ over $k[u_\sigma]$. Since both λ and σ belong to $\cap_{i=1}^{n-1} \mathcal{U}_M^i$, we see that for each i , the rank of the linear operators u_λ^i and u_σ^i on M are the same. It therefore follows from the discussion preceding the lemma that there are equalities

$$\begin{aligned} d_2 + 2d_3 + \cdots + (n - 1)d_n &= e_2 + 2e_3 + \cdots + (n - 1)e_n \\ d_3 + 2d_4 + \cdots + (n - 2)d_n &= e_3 + 2e_4 + \cdots + (n - 2)e_n \\ &\vdots \\ d_{n-1} + 2d_n &= e_{n-1} + 2e_n \\ d_n &= e_n \end{aligned}$$

This gives $d_i = e_i$ for $2 \leq i \leq n$, and in turn also $d_1 = e_1$ by considering dimensions.

For (2), note that if M has constant Jordan type then trivially $\mathcal{U}_M^i = k^c \setminus \{0\}$ for all i . The converse is an immediate consequence of (1). □

The following lemma shows that the maximal rank sets that we have defined are open subsets of affine c -space. Moreover, when the ground field is infinite, then the sets corresponding to a direct sum of modules is the intersection of the sets corresponding to the summands.

LEMMA 2.3. *Assume that k is infinite. For every A_q^c -module M and integer $1 \leq i \leq n - 1$, the set \mathcal{U}_M^i is open in k^c . Moreover, $\mathcal{U}_{M \oplus N}^i = \mathcal{U}_M^i \cap \mathcal{U}_N^i$ for every A_q^c -module N .*

Proof. An argument similar to the proof of [Op1, Lemma 9] shows that the sets \mathcal{U}_M^i are open. Also, the intersection of two nonempty open sets in k^c is always nonempty, in particular $\mathcal{U}_M^i \cap \mathcal{U}_N^i \neq \emptyset$. Since the rank of u_λ^i on $M \oplus N$ is the sum of the ranks on M and N , this proves that $\mathcal{U}_{M \oplus N}^i = \mathcal{U}_M^i \cap \mathcal{U}_N^i$. □

We can now show that the converse of the second part of Proposition 2.1 also holds.

COROLLARY 2.4. *If the field k is infinite, then for every pair M, N of A_q^c -modules, the direct sum $M \oplus N$ has constant Jordan type if and only if both M and N have.*

Proof. If $M \oplus N$ has constant Jordan type, then from Lemma 2.2 we see that $\mathcal{U}_{M \oplus N}^i = k^c \setminus \{0\}$ for all i . By Lemma 2.3, this implies that both \mathcal{U}_M^i and \mathcal{U}_N^i equal $k^c \setminus \{0\}$ for all i , and so by Lemma 2.2 again both M and N have constant Jordan type. □

Having seen some examples and elementary properties of modules of constant Jordan type, it is natural to ask the following basic questions:

- QUESTIONS.** (1) Which indecomposable A_q^c -modules have constant Jordan type?
 (2) Which sequences (a_1, \dots, a_n) in \mathbb{Z}_+^n occur as the (exponents of the) Jordan type $[1]^{a_1} \dots [n]^{a_n}$ for some indecomposable A_q^c -module of constant Jordan type?
 (3) Which sequences (a_1, \dots, a_{n-1}) in \mathbb{Z}_+^{n-1} occur as the (exponents of the) stable Jordan type $[1]^{a_1} \dots [n-1]^{a_{n-1}}$ for some indecomposable A_q^c -module of constant Jordan type?

Regarding the second question, the following result shows that the sequences

$$(0, 1, 0, \dots, 0), (0, 0, 1, 0, \dots, 0), \dots, (0, \dots, 0, 1)$$

in \mathbb{Z}_+^n are not the exponents of the Jordan types of modules of constant Jordan type.

PROPOSITION 2.5. *For $2 \leq a \leq n$, there does not exist an A_q^c -module of constant Jordan type $[a]$.*

Proof. Suppose such a module M exists, and denote the radical of A_q^c by \mathfrak{r} . For every nonzero $\lambda \in k^c$, the restriction of M to $k[u_\lambda]$ is isomorphic to the indecomposable module $k[u_\lambda]/(u_\lambda^a)$ of dimension a . Therefore there are strict inclusions

$$M \supset u_\lambda M \supset \dots \supset u_\lambda^{a-1} M \supset 0$$

with one-dimensional quotients. Now consider the radical filtration

$$M \supseteq \mathfrak{r}M \supseteq \dots \supseteq \mathfrak{r}^{a-1}M \supseteq \dots$$

of M as an A_q^c -module. If $\mathfrak{r}^i M = \mathfrak{r}^{i+1} M$, then $\mathfrak{r}^i M = 0$ by Nakayama's lemma. Moreover, since $u_\lambda \in \mathfrak{r}$, the inclusion $u_\lambda^i M \subseteq \mathfrak{r}^i M$ holds for all i . Combining all this with the fact that $\dim_k M = a$, we see that the radical series of M is

$$M \supset \mathfrak{r}M \supset \dots \supset \mathfrak{r}^{a-1}M \supset 0$$

with one-dimensional quotients. Thus M has a k -vector space basis $\{m_1, \dots, m_a\}$ with $m_i \in \mathfrak{r}^{i-1}M \setminus \mathfrak{r}^iM$ for all i .

From the above, for every nonzero $\lambda \in k^c$ there is a nonzero element $\alpha \in k$ such that $u_\lambda m_{a-1} = \alpha m_a$. In particular, there are nonzero elements α_1 and α_2 such that $x_1 m_{a-1} = \alpha_1 m_a$ and $x_2 m_{a-1} = \alpha_2 m_a$. But then

$$(\alpha_2 x_1 - \alpha_1 x_2) m_{a-1} = 0$$

which means that $u_\lambda m_{a-1} = 0$ for $\lambda = (\alpha_2, -\alpha_1, 0, \dots, 0)$. This is a contradiction. \square

Next, we look at modules of constant Jordan type from a homological point of view. For a finite-dimensional algebra A , every module M has a minimal projective resolution

$$\dots \rightarrow P_2 \xrightarrow{\partial_2} P_1 \xrightarrow{\partial_1} P_0 \xrightarrow{\partial_0} M \rightarrow 0$$

with $\text{Im } \partial_i \subseteq \mathfrak{r}_A P_{i-1}$. The *complexity* of M is defined as

$$\text{cx } M = \inf\{m \geq 0 \mid \text{there exists } b \in \mathbb{R} \text{ with } \dim_k P_t \leq bt^{m-1} \text{ for all } t \geq 1\}$$

The complexity of a module might be infinite, and is at most the maximal complexity obtained by the simple A -modules. For our quantum complete intersection A_q^c , it follows from [BeO, Theorem 5.3] that the complexity of the simple module k is c , since it equals the rate of growth of $\text{Ext}_{A_q^c}^*(k, k)$. Consequently, the complexity of every A_q^c -module is at most c . Moreover, for every integer $0 \leq m \leq c$, there exists an A_q^c -module having complexity m . Namely, by [BeO, Theorem 5.5], the Hochschild cohomology ring $\text{HH}^*(A_q^c)$ of A_q^c is finitely generated, and $\text{Ext}_{A_q^c}^*(M, N)$ is a finitely generated $\text{HH}^*(A_q^c)$ -module for all A_q^c -modules M and N . The claim now follows from [EHSST, Theorem 2.5(c) and Theorem 4.4].

The following result shows that the non-free A_q^c -modules of constant Jordan type must have maximal complexity. Moreover, it shows that the property of having constant Jordan type is preserved under syzygies and cosyzygies.

THEOREM 2.6. (1) *If M is an A_q^c -module of constant Jordan type, then either M is free, or $\text{cx } M = c$.*

(2) *Given a short exact sequence*

$$0 \rightarrow M \rightarrow F \rightarrow N \rightarrow 0$$

of A_q^c -modules with F a free module, the module N has constant Jordan type if and only if M does. In fact, if N has constant Jordan type $[1]^{d_1} \dots [n]^{d_n}$, then M has constant Jordan type $[1]^{d_{n-1}} \dots [n-1]^{d_1} [n]^d$, where $d = rn^{c-1} - (d_1 + \dots + d_n)$ and r is the rank of the free A_q^c -module F . Conversely, if M has constant Jordan type $[1]^{d_1} \dots [n]^{d_n}$, then N has constant Jordan type $[1]^{d_{n-1}} \dots [n-1]^{d_1} [n]^d$.

Proof. (1) Suppose that M is a non-free module of constant Jordan type. By [BEH, Theorem 2.6], there exists a nonzero c -tuple $\lambda \in k^c$ with the property that M is not free as a $k[u_\lambda]$ -module. As M has constant Jordan type, the same must hold for every nonzero $\lambda \in k^c$. Thus the rank variety $V_{A_q^c}^r(M)$ of M , as defined in [BEH], must equal k^c . By [BE1, Corollary 3.7], the dimension of the rank variety of a module equals its complexity, hence $\text{cx } M = c$.
 (2) As before, let M_1, \dots, M_n be the indecomposable $k[x]/(x^n)$ -modules, with $M_i = k[x]/(x^i)$. For $1 \leq i \leq n - 1$, there are short exact sequences

$$0 \rightarrow M_{n-i} \rightarrow k[x]/(x^n) \rightarrow M_i \rightarrow 0$$

and so $\Omega_{k[x]/(x^n)}^1(M_i) \simeq M_{n-i} \simeq \Omega_{k[x]/(x^n)}^{-1}(M_i)$. Now suppose that N has Jordan type $[1]^{d_1} \dots [n]^{d_n}$ with respect to λ . Since the module F is free as a $k[u_\lambda]$ -module, the module M is isomorphic to $\Omega_{k[x]/(x^n)}^1(N) \oplus Q$ over $k[u_\lambda]$, where Q is some free $k[u_\lambda]$ -module. Thus the Jordan type of M with respect to λ must be

$$[1]^{d_{n-1}} \dots [n-1]^{d_1} [n]^d$$

for some $d \geq 0$. Comparing dimensions we obtain

$$\begin{aligned} d_{n-1} + 2d_{n-2} + \dots + (n-1)d_1 + nd &= \dim_k M \\ &= \dim_k F - \dim_k N \\ &= rn^c - (d_1 + 2d_2 + \dots + nd_n) \end{aligned}$$

which in turn gives

$$d = rn^{c-1} - (d_1 + \dots + d_n)$$

The converse is proved exactly the same way. □

COROLLARY 2.7. *Let $i \in \mathbb{Z}$ be any integer. An A_q^c -module M has constant Jordan type if and only if $\Omega_{A_q^c}^i(M)$ does. Moreover, if M has constant stable Jordan type $[1]^{d_1} \dots [n-1]^{d_{n-1}}$, then so does $\Omega_{A_q^c}^{2i}(M)$, whereas $\Omega_{A_q^c}^{2i+1}(M)$ has constant stable Jordan type $[1]^{d_{n-1}} \dots [n-1]^{d_1}$.*

A reasonable question to ask is how abundant are modules of constant Jordan type. The following discussion offers an answer to this question.

Let M be a finitely generated A_q^c -module of dimension d . We fix a k -basis of M . For $\lambda = (\lambda_1, \dots, \lambda_c) \in k^c$ and $u_\lambda = \lambda_1 x_1 + \dots + \lambda_c x_c$ we let $[u_\lambda]$ denote the matrix representing the linear operator $u_\lambda : M \rightarrow M$ with respect to the fixed basis. In particular, $[x_i]$ represents the linear operator $x_i : M \rightarrow M$, for $1 \leq i \leq c$. The matrices $[x_i]$ determine the A_q^c -module M in the following sense. Let N be another d -dimensional A_q^c -module. Then M and N are isomorphic as A_q^c -modules, with isomorphism $\alpha : M \rightarrow N$, if and only if there exists an invertible $d \times d$ matrix E such that $E[x_i] = [x_i]_N E$ for all $1 \leq i \leq c$, where $[u_\lambda]_N$ is the matrix of the linear operator $u_\lambda : N \rightarrow N$ with respect to the fixed basis of M and the corresponding (under α) basis of N . In this case, E

represents the A_q^c -linear isomorphism α , as a k -linear map. Note that we also have $E[u_\lambda] = [u_\lambda]_N E$ for all λ ,

We want to associate A_q^c -modules to points in a certain affine space. We will do this using the matrices $[x_i]$. For $1 \leq i \leq c$ we let $(X_{r,s}^i)$ be a $d \times d$ generic matrix of indeterminates. Consider the polynomial ring

$$k[X_{r,s}^i \mid 1 \leq r, s \leq d, 1 \leq i \leq c]$$

and let P be the homogeneous ideal generated by the entries of the matrices $(X_{r,s}^i)(X_{r,s}^j) - q(X_{r,s}^j)(X_{r,s}^i)$, $i < j$, and $(X_{r,s}^i)^n$, $1 \leq i \leq c$. Let V denote the affine variety of P . Then any point p in V corresponds to a A_q^c -module M_p of dimension d , in the sense that the underlying k -vector space of M_p is k^d and the matrix $(X_{r,s}^i)(p)$ (substitute the coordinates of p in for the $X_{r,s}^i$) represents the linear operator $x_i : M_p \rightarrow M_p$, with respect to the standard basis of k^d , for $1 \leq i \leq c$.

Before discussing which A_q^c -modules have constant Jordan type, we first consider a weaker condition, namely, that the linear operators $u_\lambda : M \rightarrow M$ have constant rank for all λ . For a matrix B we let $I_g(B)$ denote the ideal of $g \times g$ minors of B . Consider now the polynomial ring $k[\Lambda_i, X_{r,s}^i \mid 1 \leq i \leq c, 1 \leq r, s \leq d]$, in the additional indeterminates Λ_i , and the $d \times d$ matrix

$$U_\Lambda = \Lambda_1(X_{r,s}^1) + \cdots + \Lambda_c(X_{r,s}^c)$$

For $1 \leq g \leq d$ and $p \in V$, we let $U_\Lambda(p)$ denote the matrix, and $I_g(U_\Lambda)(p)$ the ideal of $k[\Lambda_1, \dots, \Lambda_c]$, obtained by substituting the coordinates of p in for the $X_{r,s}^i$ in U_Λ , and $I_g(U_\Lambda)$, respectively.

PROPOSITION 2.8. *For $p \in V$, the linear operators $u_\lambda : M_p \rightarrow M_p$ on the A_q^c -module M_p of dimension d and corresponding to p , have constant rank g if and only if the following conditions are satisfied.*

1. $I_{g+1}(U_\Lambda)(p) = 0$
2. $\sqrt{I_g(U_\Lambda)(p)} \supseteq (\Lambda_1, \dots, \Lambda_c)$

Proof. The first condition guarantees that the $(g+1) \times (g+1)$ minors of the matrix $U_\Lambda(p)$ vanish. This implies that the linear operators $u_\lambda : M_p \rightarrow M_p$ have rank at most g for all λ . If $g = 0$, then the linear operators $u_\lambda : M_p \rightarrow M_p$ automatically have constant rank 0, and by convention $I_0(U_\Lambda) = A_q^c$. Otherwise, the second condition says that the only possible way for $u_\lambda : M_p \rightarrow M_p$ to have rank less than g is if $\lambda = 0$. Thus for all nonzero λ , the linear operator $u_\lambda : M_p \rightarrow M_p$ has rank g . \square

Proposition 2.8 suggests that the linear operators $u_\lambda : M \rightarrow M$ on A_q^c -modules of large dimension relative to c tend to have constant rank. On the other hand, we have the following.

COROLLARY 2.9. *There do not exist linear operators $u_\lambda : M \rightarrow M$ of positive constant rank g on A_q^c -modules of dimension d when $\binom{d}{g}^2 < c$. Consequently, there are no A_q^c -modules of dimension d and of constant Jordan type of rank $g > 0$ when $\binom{d}{g}^2 < c$.*

Proof. Any A_q^c -module of dimension d is isomorphic to M_p for some point $p \in V$, and so it suffices to consider modules M_p .

There are $\binom{d}{g}^2$ many $g \times g$ minors of the matrix U_Λ , and so the ideal $I_g(U_\Lambda)$, being generated by $\binom{d}{g}^2$ elements, cannot possibly have radical $(\Lambda_1, \dots, \Lambda_c)$ when $\binom{d}{g}^2 < c$. Thus there are no A_q^c -modules of constant rank $g > 0$ and dimension d when $\binom{d}{g}^2 < c$, and so no such modules of constant Jordan type. □

For $p \in V$, let us define the A_q^c -module M_p to be of *generic rank g* if $I_{g+1}(U_\Lambda)(p) = 0$, but $I_g(U_\Lambda)(p) \neq 0$. Then Proposition 2.8 says that among the $p \in V$ for which the A_q^c -modules M_p have generic rank g , those p corresponding to modules M_p of constant rank g constitute a Zariski open set. Thus the A_q^c -modules M_p generically have constant rank. We want to have a similar statement for modules of constant Jordan type. This will involve an embellishment of the immediate discussion to the sets \mathcal{U}_M^i defined above. Consider the powers $U_\Lambda, U_\Lambda^2, \dots, U_\Lambda^{n-1}$ of the generic matrix U_Λ .

DEFINITION. For $p \in V$, we say that the corresponding A_q^c -module M_p has *generic rank $g = (g_1, \dots, g_{n-1})$* if $I_{g_i+1}(U_\Lambda^i)(p) = 0$ and $I_{g_i}(U_\Lambda^i)(p) \neq 0$ for $1 \leq i \leq n-1$.

THEOREM 2.10. *For $p \in V$, suppose that M_p has generic rank $g = (g_1, \dots, g_{n-1})$. Then M_p has constant Jordan type if and only if*

$$\sqrt{I_{g_i}(U_\Lambda^i)(p)} = (\Lambda_1, \dots, \Lambda_c)$$

for $1 \leq i \leq n-1$.

Proof. The condition shows that $u_\lambda^i : M_p \rightarrow M_p$ is of maximal rank for all nonzero λ , and all $1 \leq i \leq n-1$. Thus by Lemma 2.2, M_p has constant Jordan type. □

For fixed i , the condition that $p \in V$ satisfies $\sqrt{I_{g_i}(U_\Lambda^i)(p)} = (\Lambda_1, \dots, \Lambda_c)$ corresponds to a Zariski open set of V . Thus the condition of Theorem 2.10 corresponds to a finite intersection of Zariski open sets, and thus is open. The question remains when are these open sets nonempty. The answer is simple: they are nonempty if and only if $\binom{d}{g_i}^2 \geq c$. The upshot of Theorem 2.10 is therefore that the A_q^c -modules M_p generically have constant Jordan type when $d^2 \geq c$.

REMARKS 2.11. (1). The conditions $\sqrt{I_{g_i}(U_\Lambda^i)(p)} = (\Lambda_1, \dots, \Lambda_c)$ seem to depend on the point $p \in V$. However, if p' is another point of V such that the A_q^c -modules M_p and $M_{p'}$ are isomorphic, then there exists a $d \times d$ invertible matrix E such that $U_\Lambda(p') = EU_\Lambda^i(p)E^{-1}$, and thus $I_{g_i}(U_\Lambda^i)(p') = I_{g_i}(U_\Lambda^i)(p)$. (2). Theorem 2.10 would also hold in the group representation setting. Indeed, if one assumes that n is prime, k is a field of characteristic n , P is the homogeneous ideal of $k[X_{r,s}^i \mid 1 \leq r, s \leq d, 1 \leq i \leq c]$ generated by the entries of the matrices $(X_{r,s}^i)(X_{r,s}^j) - (X_{r,s}^j)(X_{r,s}^i)$, $i < j$, and $(X_{r,s}^i)^n$, $1 \leq i \leq c$, and V is the affine variety of P , then any point p in V corresponds to a module of dimension d over the group algebra of an elementary abelian p -group of rank c . Theorem 2.10 would then give a condition for modules over such algebras to have constant Jordan type.

3 AUSLANDER-REITEN THEORY

In this section, we use Corollary 2.7 to show that the property of having constant Jordan type is preserved under Auslander-Reiten translates. We then show that if one of the modules in a component of the stable Auslander-Reiten quiver of A_q^c has constant Jordan type, then so do all the other modules in that component.

Recall first that if A is any algebra and M an A -module, then from an algebra automorphism $\psi: A \rightarrow A$ we obtain a new A -module ${}_\psi M$, called the *twist* of M by ψ . The module structure is given by $a \cdot m = \psi(a)m$. The twist commutes with operations such as direct sum and syzygies. We shall be concerned with the homogeneous automorphisms of A_q^c , that is, automorphisms which map each generator x_j to a linear combination $\alpha_{1j}x_1 + \dots + \alpha_{cj}x_c$. Not all automorphisms are of this form. For example, by mapping x_1 to $x_1 + x_1^{n-1}x_2^{n-1} \dots x_c^{n-1}$ and x_j to itself for all $2 \leq j \leq c$, we have constructed a valid automorphism since all the relations in A_q^c are preserved. If we twist a module having constant Jordan type with such an automorphism, the result may be a module which does not have constant type. However, as the following lemma shows, constant Jordan type is preserved when we twist with homogeneous automorphisms.

LEMMA 3.1. *If M is an A_q^c -module of constant Jordan type, and $\psi: A_q^c \rightarrow A_q^c$ is a homogeneous automorphism, then the module ${}_\psi M$ also has constant Jordan type. Moreover, the Jordan types of M and ${}_\psi M$ are the same.*

Proof. For each $1 \leq j \leq c$ there are scalars $\alpha_{1j}, \dots, \alpha_{cj}$ with

$$x_j \mapsto \alpha_{1j}x_1 + \dots + \alpha_{cj}x_c$$

The $c \times c$ matrix $E = (\alpha_{ij})$ must have rank c ; otherwise, there would exist a nonzero c -tuple $\beta = (\beta_1, \dots, \beta_c) \in k^c$ with $E\beta^T = 0$. But then the automorphism ψ would map the nonzero element $\beta_1x_1 + \dots + \beta_cx_c$ to zero, which is impossible.

Now take a nonzero c -tuple $\lambda \in k^c$. Since $\psi(u_\lambda) = u_{(E\lambda^T)^T}$, the matrix for u_λ on ${}_\psi M$ is the same as the matrix for $u_{(E\lambda^T)^T}$ on M . As E has maximal rank, the c -tuple $(E\lambda^T)^T$ is nonzero, and so since M has constant Jordan type the matrix for u_λ on ${}_\psi M$ is independent of λ . Moreover, it is the same as the matrix for u_λ on M . □

REMARK 3.2. In most cases, namely when q is not ± 1 (thus $n \geq 3$), the homogeneous automorphisms on A_q^c are actually of a very simple form; they just map each generator x_j to a multiple of itself. To see this, take such an automorphism $\psi: A_q^c \rightarrow A_q^c$. Then for each $1 \leq j \leq c$ there are scalars $\alpha_{1j}, \dots, \alpha_{cj}$ with

$$x_j \mapsto \alpha_{1j}x_1 + \dots + \alpha_{cj}x_c$$

Suppose that α_{ij} is nonzero, and consider another generator x_s for $s \neq j$. If $j < s$, then the relation $x_jx_s - qx_sx_j = 0$ implies that

$$\begin{aligned} &(\alpha_{1j}x_1 + \dots + \alpha_{cj}x_c)(\alpha_{1s}x_1 + \dots + \alpha_{cs}x_c) - \\ & q(\alpha_{1s}x_1 + \dots + \alpha_{cs}x_c)(\alpha_{1j}x_1 + \dots + \alpha_{cj}x_c) \end{aligned}$$

must be zero. The term involving x_i^2 is $(1-q)\alpha_{ij}\alpha_{is}x_i^2$, and so since $(1-q)\alpha_{ij}\alpha_{is}$ is nonzero in A_q^c , we see that α_{is} must be zero. The same happens if $s < j$. This shows that if x_i occurs in the linear combination of $\psi(x_j)$, then x_i does *not* occur in the linear combinations of $\psi(x_s)$ for $s \neq j$. Consequently, since ψ is an automorphism, it must simply permute the generators x_1, \dots, x_c up to scalars; there are nonzero scalars $\alpha_1, \dots, \alpha_c$ and a permutation $\sigma \in S_c$ with $\psi(x_i) = \alpha_i x_{\sigma(i)}$ for every $1 \leq i \leq c$.

So far, we have only used that $q \neq 1$. Let us now use the fact that $q \neq \pm 1$ to show that the permutation σ must be the identity permutation. If not, there exist two integers $i, j \in \{1, \dots, c\}$ with $i < j$ and $\sigma(i) > \sigma(j)$. Then since $x_i x_j = q x_j x_i$ and $x_{\sigma(j)} x_{\sigma(i)} = q x_{\sigma(i)} x_{\sigma(j)}$, we obtain

$$\alpha_i \alpha_j x_{\sigma(i)} x_{\sigma(j)} = \psi(x_i x_j) = \psi(q x_j x_i) = q \alpha_i \alpha_j x_{\sigma(j)} x_{\sigma(i)} = q^2 \alpha_i \alpha_j x_{\sigma(i)} x_{\sigma(j)}$$

but this is impossible when $q \neq \pm 1$.

To sum up, when $q \neq \pm 1$, then a homogeneous automorphism $\psi: A_q^c \rightarrow A_q^c$ simply maps x_i to $\alpha_i x_i$ for some nonzero $\alpha_i \in k$. When $q^2 = 1$, however, then there are in general other kinds of homogeneous automorphisms. For example, over any ground field k , the quantum complete intersection (and exterior algebra)

$$k\langle x, y \rangle / (x^2, xy + yx, y^2)$$

admits the homogeneous morphisms $(x \mapsto y, y \mapsto x)$ and $(x \mapsto x+y, y \mapsto x-y)$.

We can now show that the property of having constant Jordan type is preserved under Auslander-Reiten translates. Recall that for a finite dimensional Frobenius algebra A , there is an automorphism $\nu: A \rightarrow A$, called the *Nakayama automorphism*, with the property that the bimodules $D(A)$ and ${}_\nu A_1$ are isomorphic. Here $D(A)$ denotes the vector space dual $\text{Hom}_k(A, k)$ of A , and the

action on the bimodule ${}_{\nu}A_1$ is defined by $a_1 \cdot a \cdot a_2 = \nu(a_1)aa_2$. The Nakayama automorphism is unique up to inner automorphisms. It is well known that for such an algebra A , the Auslander-Reiten translate τM of a module M is isomorphic to $\Omega_A^2({}_{\nu}M)$; see, for example, [SkY, Proposition 3.13 and Theorem 8.5]. We can now apply this to our quantum complete intersection A_q^c , which is Frobenius.

THEOREM 3.3. *An A_q^c -module M has constant Jordan type if and only if its Auslander-Reiten translate τM does. Moreover, if so, then their stable constant Jordan types are the same.*

Proof. By [Be2, Lemma 3.1], the Nakayama automorphism ν of A_q^c maps each generator x_i to $q^{m_i}x_i$ for some (possibly negative) integer $m_i \in \mathbb{Z}$. Therefore, by Lemma 3.1, the module M has constant Jordan type if and only if ${}_{\nu}M$ does, and with the same Jordan type. It now follows from Corollary 2.7 that M has constant Jordan type if and only if $\Omega_{A_q^c}^2({}_{\nu}M)$ does, and with the same stable Jordan type. Since $\tau M \simeq \Omega_{A_q^c}^2({}_{\nu}M)$, the result follows. \square

Next, we turn to the stable Auslander-Reiten quiver of A_q^c . Our aim is to show that when the ground field k is algebraically closed, then constant Jordan type is a property of the components of the quiver: if one of the modules has constant Jordan type, then so do all the others in that component. We also determine the stable Jordan types of the modules. The key to all this is the fact, proved in [BE2], that when either $n \geq 3$ or $c \geq 3$, then every component of the stable Auslander-Reiten quiver of A_q^c is of the form $\mathbb{Z}A_{\infty}$.

In order to prove this result, we need the following lemma and its corollary. They show that every Auslander-Reiten sequence over A_q^c ending in a module of constant Jordan type splits when we restrict to the subalgebras $k[u_{\lambda}]$.

LEMMA 3.4. *Let M be an indecomposable A_q^c -module of complexity at least 2, and*

$$0 \rightarrow \tau M \xrightarrow{f} E \xrightarrow{g} M \rightarrow 0$$

the Auslander-Reiten sequence ending in M . Then this sequence splits over $k[u_{\lambda}]$ for all nonzero $\lambda \in k^c$.

Proof. Fix a nonzero $\lambda \in k^c$, and denote the algebra A_q^c by just A . For every A -module L , the adjoint isomorphism

$$\text{Hom}_A(A \otimes_{k[u_{\lambda}]} M, L) \rightarrow \text{Hom}_{k[u_{\lambda}]}(M, \text{Hom}_A(A, L))$$

together with the natural isomorphism $\text{Hom}_A(A, L) \rightarrow L$ of $k[u_{\lambda}]$ -modules give an isomorphism

$$\text{Hom}_A(A \otimes_{k[u_{\lambda}]} M, L) \xrightarrow{\varphi_L} \text{Hom}_{k[u_{\lambda}]}(M, L)$$

which is natural in L . From the map g in the Auslander-Reiten sequence we therefore obtain a commutative diagram

$$\begin{CD}
 \mathrm{Hom}_A(A \otimes_{k[u_\lambda]} M, E) @>g_*>> \mathrm{Hom}_A(A \otimes_{k[u_\lambda]} M, M) \\
 @V\varphi_EVV @VV\varphi_MV \\
 \mathrm{Hom}_{k[u_\lambda]}(M, E) @>g_*>> \mathrm{Hom}_{k[u_\lambda]}(M, M)
 \end{CD}$$

Consider now the multiplication map $\mu: A \otimes_{k[u_\lambda]} M \rightarrow M$ given by $a \otimes m \mapsto am$. This is a surjective homomorphism of A -modules, and we claim that it cannot split. For if it did, then the A -module M would be a direct summand of the A -module $A \otimes_{k[u_\lambda]} M$. However, the complexity of M as a module over $k[u_\lambda]$ is at most one, since all the indecomposable non-projective $k[u_\lambda]$ -modules are periodic. Moreover, if we take any projective resolution of M over $k[u_\lambda]$, and apply $A \otimes_{k[u_\lambda]} -$ to it, then the result is a projective resolution of $A \otimes_{k[u_\lambda]} M$ over A , since A is free as a $k[u_\lambda]$ -module. Therefore the complexity of the A -module $A \otimes_{k[u_\lambda]} M$ is at most one. Then M cannot be a direct summand of this module, since the complexity of M is at least 2.

Since the multiplication map μ does not split, it factors through the map g , so that $\mu = g_*(\theta)$ for some map $\theta \in \mathrm{Hom}_A(A \otimes_{k[u_\lambda]} M, E)$. The image of μ under φ_M is the identity on M , and so the commutativity of the diagram implies that the map g splits as a homomorphism of $k[u_\lambda]$ -modules: $1_M = g \circ \varphi_E(\theta)$. This shows that the Auslander-Reiten sequence splits over $k[u_\lambda]$. □

It now follows from Theorem 2.6 that all the Auslander-Reiten sequences ending in modules of constant Jordan type must split over the subalgebras $k[u_\lambda]$.

COROLLARY 3.5. *Every Auslander-Reiten sequence over A_q^c ending in a module of constant Jordan type splits over $k[u_\lambda]$ for all nonzero $\lambda \in k^c$.*

Now we can prove the main result in this section: when the field k is algebraically closed, then the A_q^c -modules of constant Jordan type form complete components of the stable Auslander-Reiten quiver. As mentioned, the key ingredient is that in all cases except one, all the components are of the form $\mathbb{Z}A_\infty$. In such a component, a module is *quasi-simple* if it belongs to the τ -orbit at the end. For an arbitrary module in the component, there is a shortest sectional path to a quasi-simple module, and the number of modules in such a path is the *quasi-length* of the module. Thus the quasi-simple modules are precisely the ones having quasi-length one.

THEOREM 3.6. *Suppose that the ground field k is algebraically closed, and let Θ be a component of the stable Auslander-Reiten quiver of A_q^c containing a module of constant Jordan type. Then all the modules in Θ have constant Jordan type. In fact, the following hold:*

- (1) *If $n = c = 2$, then Θ is of the form $\mathbb{Z}\tilde{A}_{12}$, and its modules are precisely $\{\Omega_{A_q^c}^i(k) \mid i \in \mathbb{Z}\}$. For even i , the stable Jordan type of $\Omega_{A_q^c}^i(k)$ is $[1]$, and for odd i it is $[n - 1]$.*

(2) If either $n \geq 3$ or $c \geq 3$, then Θ is of the form $\mathbb{Z}A_\infty$. If the stable Jordan type of one of the quasi-simple modules in Θ is $[1]^{d_1} \cdots [n-1]^{d_{n-1}}$, then every module of quasi-length l has stable Jordan type $[1]^{ld_1} \cdots [n-1]^{ld_{n-1}}$.

Proof. (1) When $n = c = 2$, then $A_q^c/(x_1x_2)$ is just the commutative truncated polynomial algebra $k[x_1, x_2]/(x_1^2, x_2^2)$ modulo its socle, regardless of the characteristic of the field k . It is well known (see, for example, the proof of [Erd, Lemma II.7.3]) that the only indecomposable nonprojective and nonperiodic modules over this algebra are the syzygies of the simple module k , and that they form a component $\mathbb{Z}\tilde{A}_{12}$ in the stable Auslander-Reiten quiver. By Corollary 2.7 and the fact that k trivially has constant Jordan type $[1]$, every module in this component has constant stable Jordan type as given. By Theorem 2.6, there are no other indecomposable nonprojective modules of constant Jordan type.

(2) Suppose now that either $n \geq 3$ or $c \geq 3$. By [BE2, Theorem 3.6], every component of the stable Auslander-Reiten quiver of A_q^c is of the form $\mathbb{Z}A_\infty$. Let M be a module in Θ of constant Jordan type, and suppose that its quasi-length is l . Then by Theorem 3.3, all the modules in Θ of quasi-length l , that is, all the modules in the τ -orbit of M , also have constant Jordan type. Moreover, their stable Jordan types are all equal to that of M .

Consider the Auslander-Reiten sequence

$$0 \rightarrow \tau M \rightarrow E \rightarrow M \rightarrow 0$$

ending in M . If M is quasi-simple, that is, if $l = 1$, then E is indecomposable of quasi-length two. Otherwise, the module E is a direct sum $E \simeq E_1 \oplus E_2$ of two indecomposable modules, with E_1 of quasi-length $l - 1$ and E_2 of quasi-length $l + 1$. By Corollary 3.5, the sequence splits over $k[u_\lambda]$ for all nonzero $\lambda \in k^c$, hence since M and τM have constant Jordan type, so does E . It now follows from Corollary 2.4 and Theorem 3.3 that every module in Θ of quasi-length $l + 1$, or of quasi-length $l - 1$ if $l \geq 2$, must have constant Jordan type. Moreover, the stable Jordan types of the modules of quasi-length $l + 1$ are all the same, as are the stable Jordan types of the modules of quasi-length $l - 1$. By induction, we see that every module in Θ must have constant Jordan type. Moreover, two modules having the same quasi-length have the same stable Jordan type.

Suppose now that $[1]^{d_1} \cdots [n-1]^{d_{n-1}}$ is the stable Jordan type of one of (equivalently, all of) the quasi-simple modules. We show by induction on l that every module E of quasi-length l has stable Jordan type $[1]^{ld_1} \cdots [n-1]^{ld_{n-1}}$, the case $l = 1$ being the quasi-simple case. If $l = 2$, then there is an Auslander-Reiten sequence as above, in which M and τM are quasi-simple. Then since the sequence splits over $k[u_\lambda]$ for all nonzero $\lambda \in k^c$, and the stable Jordan type of both M and τM is $[1]^{d_1} \cdots [n-1]^{d_{n-1}}$, we see that the stable Jordan type of E must be $[1]^{2d_1} \cdots [n-1]^{2d_{n-1}}$. If $l \geq 3$, then there is an Auslander-Reiten sequence

$$0 \rightarrow \tau M \rightarrow E \oplus E' \rightarrow M \rightarrow 0$$

in which the quasi-length of E' is $l - 2$, and that of M and τM is $l - 1$. By induction, the stable Jordan type of E' is $[1]^{(l-2)d_1} \dots [n - 1]^{(l-2)d_{n-1}}$, and the stable Jordan type of both M and τM is $[1]^{(l-1)d_1} \dots [n - 1]^{(l-1)d_{n-1}}$. Using again that the sequence splits over $k[u_\lambda]$ for all nonzero $\lambda \in k^c$, we see that the direct sum $E \oplus E'$ must have stable Jordan type $[1]^{2(l-1)d_1} \dots [n - 1]^{2(l-1)d_{n-1}}$, hence the stable Jordan type of E must be $[1]^{ld_1} \dots [n - 1]^{ld_{n-1}}$. This completes the proof. \square

4 SYZYGIES OF THE SIMPLE MODULE

In this final section, we focus on the syzygies of the simple A_q^c -module k . Since this module trivially has constant stable Jordan type $[1]$, it follows from Corollary 2.7 that for every integer $i \in \mathbb{Z}$, the constant stable Jordan type of $\Omega_{A_q^c}^{2i}(k)$ is $[1]$, and for $\Omega_{A_q^c}^{2i+1}(k)$ it is $[n - 1]$. Our aim now is to show that these are the *only* modules having these constant stable Jordan types, in the case when $c = 2$, that is, when our algebra has two generators. The arguments we use are to a large extent adaptations of arguments from [Ca1].

Instead of writing x_1 and x_2 , we shall write x and y for the generators of A_q^2 , and just A for the algebra A_q^2 itself. Thus the algebra we are dealing with throughout this section is

$$A = k\langle x, y \rangle / (x^n, xy - qyx, y^n)$$

with k, q and n as before. Moreover, as in the previous sections we denote the radical of A by just \mathfrak{r} . For an A -module M , we denote by $\text{top } M$ its top $M/\mathfrak{r}M$, and by $\text{soc } M$ its socle, that is, the submodule $\{m \in M \mid \mathfrak{r}m = 0\}$. Note that the latter is the same as the set of all elements m in M with $xm = 0$ and $ym = 0$.

For an A -module M without any projective summands, consider its minimal complete resolution

$$\dots \rightarrow P_2 \xrightarrow{d_2} P_1 \xrightarrow{d_1} P_0 \xrightarrow{d_0} P_{-1} \xrightarrow{d_{-1}} P_{-2} \rightarrow \dots$$

of free A -modules with $\text{Im } d_i \subseteq \mathfrak{r}P_{i-1}$ and $M = \text{Im } d_0$. It is unique up to isomorphism, and so we define $\beta_i(M)$ to be the rank of the free module P_i . This is the same as the k -vector space dimension of $\text{top } \Omega_A^i(M)$, since $\Omega_A^i(M) = \text{Im } d_i$. In Section 2, we defined the complexity of a module over an arbitrary algebra in terms of the dimensions of the projective modules in its minimal projective resolution. Since the algebra we are working over is local, we may just as well use the integers $\beta_i(M)$:

$$\text{cx } M = \inf\{m \geq 0 \mid \text{there exists } b \in \mathbb{R} \text{ with } \beta_t(M) \leq bt^{m-1} \text{ for all } t \geq 0\}$$

The first result we prove compares $\beta_0(M)$ with $\beta_{-1}(M)$ when the module M has constant stable Jordan type $[1]$ or $[n - 1]$. By definition, for all $u_\lambda = \lambda_1x + \lambda_2y$ with $\lambda \neq 0$, such a module decomposes over $k[u_\lambda]$ into a direct sum $M_i \oplus F$,

where F is free and i is either 1 or $n - 1$ (with M_1 of dimension 1, and M_{n-1} of dimension $n - 1$). We call the module M_i the nonprojective component.

In the results and proofs to come, we write $k[x]$ and $k[y]$ for the subalgebras of A generated by x and y , respectively. These are not to be confused with polynomial rings: the are both isomorphic to $k[z]/(z^n)$.

PROPOSITION 4.1. *Suppose that M is an indecomposable A -module of constant stable Jordan type $[1]$ or $[n - 1]$. Then the following conditions are equivalent:*

- (1) $\beta_0(M) > \beta_{-1}(M)$;
- (2) *There exists a generator a for the nonprojective component of M as a $k[x]$ -module, with the following properties: $a \notin \mathfrak{r}M$, and if M is of stable Jordan type $[1]$ then $ya \neq 0$, whereas if M is of type $[n - 1]$ then $yx^{n-2}a \neq 0$;*
- (2') *There exists a generator b for the nonprojective component of M as a $k[y]$ -module, with the following properties: $b \notin \mathfrak{r}M$, and if M is of stable Jordan type $[1]$ then $xb \neq 0$, whereas if M is of type $[n - 1]$ then $y^{n-2}xb \neq 0$;*
- (3) *Any generator for the nonprojective component of M as a $k[x]$ -module satisfies (2);*
- (3') *Any generator for the nonprojective component of M as a $k[y]$ -module satisfies (2').*

Proof. Let a be a generator for the nonprojective component of M as a $k[x]$ -module, and consider the surjective map $M/xM \rightarrow x^{n-1}M$ induced by multiplication by x^{n-1} . Note that both the modules are A -modules, since y q -commutes with x . If we twist the module $x^{n-1}M$ with the automorphism $\psi: A \rightarrow A$ given by $x \mapsto x$ and $y \mapsto q^{n-1}y$, then the map above induces a homomorphism $M/xM \rightarrow \psi(x^{n-1}M)$ of A -modules. This gives an exact sequence

$$0 \rightarrow k \rightarrow M/xM \rightarrow \psi(x^{n-1}M) \rightarrow 0 \quad (\dagger)$$

over A , where the kernel of the map $M/xM \rightarrow \psi(x^{n-1}M)$ is of dimension one and generated by the coset of a . Now consider the socle of M as a $k[x]$ -module, i.e. $\text{soc}_{k[x]} M = \{m \in M \mid xm = 0\}$. Again, since y q -commutes with x , this is an A -module. There is an exact sequence

$$0 \rightarrow x^{n-1}M \rightarrow \text{soc}_{k[x]} M \rightarrow k \rightarrow 0 \quad (\dagger\dagger)$$

over A , where the one-dimensional module is generated by a if the Jordan type of M is $[1]$, and by $x^{n-2}a$ if the type is $[n - 1]$.

Note that in both of the exact sequences, the element x acts trivially on the modules. Consequently, we can view the modules involved as just $k[y]$ -modules. For such an A -module L , the top $L/\mathfrak{r}L$ is just L/yL , and so $\dim_k \text{top } L =$

$\dim_k \text{soc}_{k[y]} L$. From the sequence (†) we therefore obtain

$$\begin{aligned} \dim_k \text{top } M &= \dim_k \text{top } M/xM \\ &= \dim_k \text{soc}_{k[y]} M/xM \\ &\geq \dim_k \text{soc}_{k[y]} \psi(x^{n-1}M) \\ &= \dim_k \text{soc}_{k[y]} x^{n-1}M \\ &= \dim_k \text{top } x^{n-1}M \end{aligned}$$

with a strict inequality if and only if the sequence splits. This happens if and only if $a \notin \tau M$, and in this case $\dim_k \text{top } M = \dim_k \text{top } x^{n-1}M + 1$. On the other hand, if

$$0 \rightarrow M \hookrightarrow P \rightarrow \Omega_A^{-1}(M) \rightarrow 0$$

is exact with P projective, then from the sequence (††) we obtain

$$\begin{aligned} \dim_k \text{top } \Omega_A^{-1}(M) &= \dim_k \text{soc } Q \\ &= \dim_k \text{soc } M \\ &= \dim_k \text{soc}_{k[y]} (\text{soc}_{k[x]} M) \\ &\geq \dim_k \text{soc}_{k[y]} x^{n-1}M \\ &= \dim_k \text{top } x^{n-1}M \end{aligned}$$

Again, the inequality is strict if and only if the sequence (††) splits, which happens if and only if $a \in \text{soc } M$ when the stable Jordan type of M is $[1]$, and if and only if $x^{n-2}a \in \text{soc } M$ when the type is $[n-1]$. In other words, the sequence splits if and only if $ya = 0$ when the stable Jordan type is $[1]$, and if and only if $yx^{n-2}a = 0$ when the type is $[n-1]$. In this case, $\dim_k \text{top } \Omega_A^{-1}(M) = \dim_k \text{top } x^{n-1}M + 1$.

From what we have proved, we see that condition (1) holds if and only if $\dim_k \text{top } M = \dim_k \text{top } x^{n-1}M + 1$ and $\dim_k \text{top } \Omega_A^{-1}(M) = \dim_k \text{top } x^{n-1}M$. This happens if and only if (†) splits but (††) does not, and this is equivalent to condition (2) and (3), since a was an arbitrary generator for the nonprojective component of M as a $k[x]$ -module. A completely similar argument shows that (1), (2') and (3') are also equivalent. □

Before we proceed, we make the following definition.

DEFINITION. Let M be an indecomposable nonprojective A -module of stable constant Jordan type either $[1]$ or $[n-1]$.

(1) The module M satisfies the *rank property with respect to x* , abbreviated (RP x), if there exists a generator a for the nonprojective component of M over $k[x]$, with the following properties: if the stable Jordan type of M is $[1]$, then $a \notin \tau M$ and $y^{n-1}a \neq 0$, whereas if the stable Jordan type of M is $[n-1]$, then $y^{n-1}x^{n-2}a \neq 0$.

(2) Likewise, M satisfies the *rank property with respect to y* , abbreviated (RP y), if there exists a generator b for the nonprojective component of M over $k[y]$,

with the following properties: if the stable Jordan type of M is $[1]$, then $b \notin \mathfrak{r}M$ and $x^{n-1}b \neq 0$, whereas if the stable Jordan type of M is $[n-1]$, then $y^{n-2}x^{n-1}b \neq 0$.

(3) If M satisfies both (RP x) and (RP y), then we say that it satisfies the *rank property*, abbreviated (RP).

REMARK 4.2. If a module M satisfies (RP x), then *every* generator for the nonprojective component over $k[x]$ has the defining properties from part (1) of the definition. To see this, suppose that a is as in the definition, and take any other generator a' for the nonprojective component over $k[x]$. If the stable Jordan type of M is $[1]$, then $a' = a + x^{n-1}m$ for some $m \in M$. Then since a does not belong to $\mathfrak{r}M$, neither can a' . Moreover, since $y^{n-1}x^{n-1}M = 0$, we see that $y^{n-1}a' = y^{n-1}(a + x^{n-1}m) = y^{n-1}a \neq 0$. If the stable Jordan type of M is $[n-1]$, then $a' = a + xm$ for some $m \in M$. In this case $y^{n-1}x^{n-2}a' = y^{n-1}x^{n-2}(a + xm) = y^{n-1}x^{n-2}a \neq 0$. Thus in both cases the generator a' also has the defining properties from the rank property definition. Similarly, if M satisfies (RP y), then every generator for the nonprojective component over $k[y]$ has the defining properties from part (2) of the definition. In the proofs to come, we shall be using these facts without further mention.

As the terminology suggests, the rank property should of course have something to do with ranks. This is the following result.

PROPOSITION 4.3. *If an indecomposable nonprojective A -module satisfies either (RP x) or (RP y), then $\beta_0(M) > \beta_{-1}(M)$.*

Proof. Let M be a module that satisfies (RP x), and let a be any generator for the nonprojective component of M over $k[x]$. Note first that M is not projective, and so $y^{n-1}x^{n-1}M = 0$. If the stable constant Jordan type of M is $[1]$, then by definition of the rank property, the element a is not contained in $\mathfrak{r}M$, and $y^{n-1}a \neq 0$. Therefore ya , must be nonzero. If the stable constant Jordan type is $[n-1]$, then by definition $y^{n-1}x^{n-2}a \neq 0$, and in particular $yx^{n-2}a$ must be nonzero. In addition, also in this case the element a cannot belong to $\mathfrak{r}M$: if it did, then we could write $a = xa_1 + ya_2$ for some $a_1, a_2 \in M$, giving $y^{n-1}x^{n-2}a = 0$ since $y^n = 0$ and $y^{n-1}x^{n-1}M = 0$.

We have shown that both in the case the stable constant Jordan type of M is $[1]$, and in the case it is $[n-1]$, condition (3) of Proposition 4.1 holds. It follows that $\beta_0(M) > \beta_{-1}(M)$. A similar proof applies if M satisfies (RP y). \square

If the stable constant Jordan type of the module is $[n-1]$, then the converse also holds, under some extra conditions. To prove this, we need the first part of the following lemma.

LEMMA 4.4. *Let $\psi: A \rightarrow A$ be the automorphism defined by $x \mapsto x$ and $y \mapsto q^{-1}y$, and $\phi: A \rightarrow A$ the automorphism defined by $x \mapsto qx$ and $y \mapsto y$. Then for every A -module M with $\dim_k \underline{\mathrm{Hom}}_A(M, M) = 1$ and ${}_{\psi}M \simeq M \simeq {}_{\phi}M$, and every integer i , the following hold.*

- (1) For every surjective homomorphism $d: F \rightarrow \Omega_A^i(M)$ with F free, there exist homomorphisms $h_x: {}_\psi\Omega_A^i(M) \rightarrow F$ and $h_y: {}_\phi\Omega_A^i(M) \rightarrow F$, such that the composition $d \circ h_x: {}_\psi\Omega_A^i(M) \rightarrow \Omega_A^i(M)$ is multiplication by x , and the composition $d \circ h_y: {}_\phi\Omega_A^i(M) \rightarrow \Omega_A^i(M)$ is multiplication by y .
- (2) For every injective homomorphism $i: \Omega_A^i(M) \rightarrow F$ with F free, there exist homomorphisms $h_x: F \rightarrow {}_{\psi^{-1}}\Omega_A^i(M)$ and $h_y: F \rightarrow {}_{\phi^{-1}}\Omega_A^i(M)$, such that the composition $h_x \circ i: \Omega_A^i(M) \rightarrow {}_{\psi^{-1}}\Omega_A^i(M)$ is multiplication by x , and the composition $h_y \circ i: \Omega_A^i(M) \rightarrow {}_{\phi^{-1}}\Omega_A^i(M)$ is multiplication by y .

Proof. Consider the module $\Omega_A^i(M)$. When we twist a minimal projective resolution of a module by an automorphism θ , the result is a minimal projective resolution of the corresponding twisted module, hence ${}_\theta\Omega_A^i(M) \simeq \Omega_A^i({}_\theta M)$. Consequently, there are isomorphisms ${}_\psi\Omega_A^i(M) \simeq \Omega_A^i({}_\psi M) \simeq \Omega_A^i(M)$ and ${}_\phi\Omega_A^i(M) \simeq \Omega_A^i({}_\phi M) \simeq \Omega_A^i(M)$, due to the assumptions on M . Also, since $\underline{\text{Hom}}_A(\Omega_A^i(M), \Omega_A^i(M)) \simeq \underline{\text{Hom}}_A(M, M)$, the vector space $\underline{\text{Hom}}_A(\Omega_A^i(M), \Omega_A^i(M))$ is one-dimensional. This shows that the module $\Omega_A^i(M)$ satisfies the same assumptions as M , and so it suffices to prove the lemma for the latter.

For the first part, note that the automorphisms ψ and ϕ are precisely the ones we have to twist with in order for multiplication by x to induce an A -homomorphism $\mu_x: {}_\psi M \rightarrow M$, and multiplication by y to induce an A -homomorphism $\mu_y: {}_\phi M \rightarrow M$

$$\begin{aligned} \mu_x(y \cdot m) &= \mu_x(q^{-1}ym) = q^{-1}xym = yxm = y \cdot \mu_x(m) \\ \mu_y(x \cdot m) &= \mu_y(qxm) = qyxm = xym = x \cdot \mu_y(m) \end{aligned}$$

Since $M \simeq {}_\psi M$, there is an isomorphism $\underline{\text{Hom}}_A({}_\psi M, M) \simeq \underline{\text{Hom}}_A(M, M)$, and these are one-dimensional vector spaces by assumption. In the stable module category, an isomorphism ${}_\psi M \rightarrow M$ is nonzero, and so μ_x , which is not an isomorphism, must be zero in $\underline{\text{Hom}}_A({}_\psi M, M)$. Thus μ_x factors through some free A -module, and therefore also through F . The same argument works for the map μ_y .

The second part is proved similarly. Note that since ${}_\psi M \simeq M \simeq {}_\phi M$, there are also isomorphisms ${}_{\psi^{-1}}M \simeq M \simeq {}_{\phi^{-1}}M$. □

PROPOSITION 4.5. *Let M be an indecomposable A -module of stable constant Jordan type $[n - 1]$, and suppose that $\dim_k \underline{\text{Hom}}_A(M, M) = 1$ and ${}_\psi M \simeq M \simeq {}_\phi M$, where ψ and ϕ are the automorphisms from Lemma 4.4. Then M satisfies (RP) if $\beta_0(M) > \beta_{-1}(M)$.*

Proof. We show that M satisfies (RP x); the proof that M satisfies (RP y) is similar. Let a be any generator for the nonprojective component of M over $k[x]$. To show that M satisfies (RP x), we must show that $y^{n-1}x^{n-2}a$ is nonzero. To do this, we consider the projective cover

$$0 \rightarrow \Omega_A^1(M) \hookrightarrow P \xrightarrow{d} M \rightarrow 0$$

of M . Note that $\Omega_A^1(M)$ is also nonprojective and indecomposable, since M is. View the short exact sequence as a sequence of $k[x]$ -modules. As such, the module M is isomorphic to $\text{span}_k\{a, xa, \dots, x^{n-2}a\} \oplus F_1$ for some free module F_1 , where $\text{span}_k\{a, xa, \dots, x^{n-2}a\}$ is the nonprojective component of dimension $n - 1$. The module P is free, and $\Omega_A^1(M)$ is isomorphic to $k \oplus F_2$ for some free module F_2 . Consequently, there is an element p of P with $d(p) = a$, and such that $x^{n-1}p$ generates the one-dimensional nonprojective component of $\Omega_A^1(M)$. Note that since $a \in M \setminus \mathfrak{r}M$ by Proposition 4.1, the element p does not belong to $\mathfrak{r}P$, hence $y^{n-1}x^{n-1}p$ is nonzero. Moreover, since a generates the nonprojective component of M over $k[x]$, the element $x^{n-1}a$ is zero.

Now we use the first part of Lemma 4.4, with the automorphism ψ defined there. Let $h: {}_\psi M \rightarrow P$ be an A -homomorphism having the property that the composition $d \circ h: {}_\psi M \rightarrow M$ equals the multiplication map μ_x given by x . Then $d(h(a) - xp) = 0$, so that $h(a) = xp + u$ for some $u \in \Omega_A^1(M)$. Note that $x^{n-1}u = 0$ since $x^{n-1}u = x^{n-1}(h(a) - xp) = h(x^{n-1}a)$ and $x^{n-1}a = 0$, hence u belongs to xP . Now since $x^{n-1}p$ generates the nonprojective component of $\Omega_A^1(M)$ over $k[x]$, there is an equality

$$\Omega_A^1(M) \cap xP = \text{span}_k\{x^{n-1}p\} + x\Omega_A^1(M)$$

hence we can write u as $u = \alpha x^{n-1}p + xv$ for some scalar α and element $v \in \Omega_A^1(M)$. This gives

$$\begin{aligned} h(y^{n-1}x^{n-2}a) &= y^{n-1}x^{n-2}h(a) \\ &= y^{n-1}x^{n-2}(xp + u) \\ &= y^{n-1}x^{n-2}(xp + \alpha x^{n-1}p + xv) \\ &= y^{n-1}x^{n-1}p + y^{n-1}x^{n-1}v \end{aligned}$$

and so if $y^{n-1}x^{n-2}a$ were zero, then $y^{n-1}x^{n-1}p = -y^{n-1}x^{n-1}v$. As $y^{n-1}x^{n-1}p$ is nonzero, this would imply that $y^{n-1}x^{n-1}\Omega_A^1(M)$ was nonzero, but this is impossible since $\Omega_A^1(M)$ is a nonprojective indecomposable A -module. This shows that $y^{n-1}x^{n-2}a$ must be nonzero. \square

The idea behind the proof of the main result in this section is to start with a syzygy having the rank property (RP), and then show that its cosyzygies get smaller and smaller until we end up with the module k . The first step in this reduction process is the following result.

PROPOSITION 4.6. *Let M be an indecomposable A -module of stable constant Jordan type $[n - 1]$, and suppose that $\dim_k \underline{\text{Hom}}_A(M, M) = 1$ and ${}_\psi M \simeq M \simeq {}_\phi M$, where ψ and ϕ are the automorphisms from Lemma 4.4. Then if M satisfies (RP), the module $\Omega_A^{-1}(M)$ is either isomorphic to k , or satisfies (RP).*

Proof. We show that if M satisfies (RP x), then $\Omega_A^{-1}(M)$ is either isomorphic to k , or satisfies (RP x); the (RP y)-version is similar. As in the previous proof,

we decompose the module M over $k[x]$ as $\text{span}_k\{a, xa, \dots, x^{n-2}a\} \oplus F$ for some free module F , where $\text{span}_k\{a, xa, \dots, x^{n-2}a\}$ is the nonprojective component of dimension $n - 1$. By definition, as M satisfies (RP x), the element $y^{n-1}x^{n-2}a$ is nonzero. Now consider the injective envelope

$$0 \rightarrow M \hookrightarrow P \xrightarrow{d} \Omega_A^{-1}(M) \rightarrow 0$$

of M , viewed as a sequence of $k[x]$ -modules. Since $x^{n-1}a = 0$, we can write a as $a = xp$ for some $p \in P$, with $y^{n-1}x^{n-1}p$ nonzero since $y^{n-1}x^{n-2}a$ is. Thus p does not belong to $\mathfrak{r}P$, and so $d(p) \notin \mathfrak{r}\Omega_A^{-1}(M)$ since d is a projective cover. The element $d(p)$ generates the one-dimensional nonprojective component of $\Omega_A^{-1}(M)$ over $k[x]$, hence $xd(p) = 0$. If also $yd(p) = 0$, then this component is actually a direct summand of $\Omega_A^{-1}(M)$ as an A -module. Namely, since $d(p) \notin \mathfrak{r}\Omega_A^{-1}(M)$, there is a maximal submodule $W \subseteq \Omega_A^{-1}(M)$ with $d(p) \notin W$. The submodule generated by $d(p)$ is just $\text{span}_k\{d(p)\}$, and since this does not intersect W , the A -module $\Omega_A^{-1}(M)$ decomposes as a direct sum $W \oplus \text{span}_k\{d(p)\}$. But $\Omega_A^{-1}(M)$ is indecomposable over A , since M is, and so in this case $\Omega_A^{-1}(M) \simeq k$.

This shows that if $\Omega_A^{-1}(M)$ is not isomorphic to k , then $yd(p)$ must be nonzero, that is, the element yp cannot belong to M . We must now show that this forces $\Omega_A^{-1}(M)$ to satisfy (RP x). As the element $d(p)$ is a generator for the one-dimensional nonprojective component of $\Omega_A^{-1}(M)$ over $k[x]$, and we saw above that $d(p) \notin \mathfrak{r}\Omega_A^{-1}(M)$, it suffices to show that $y^{n-1}d(p)$ is nonzero.

Suppose, for a contradiction, that $y^{n-1}d(p) = 0$. Then $y^{n-1}p \in \text{Ker } d = M$. Now we use the second part of Lemma 4.4, with the automorphism ϕ defined there. There exists an A -homomorphism $h: P \rightarrow \phi^{-1}M$ such that the composition $h \circ i: M \rightarrow \phi^{-1}M$ is multiplication by y , where i is the inclusion map $M \hookrightarrow P$. Since xp belongs to M , it follows that $h(xp) = yxp$, and so when we view $\phi^{-1}M$ as a submodule of $\phi^{-1}P$ we obtain

$$x \cdot h(p) = h(xp) = yxp = q^{-1}xyp = x \cdot yp$$

Then $x \cdot (h(p) - yp) = 0$ in $\phi^{-1}P$, so with $u = h(p) - yp$ we obtain $h(p) = yp + u$ with $x \cdot u = 0$. From the above, the element $y^{n-1}p$ belongs to M , hence $y^{n-1}h(p) = h(y^{n-1}p) = y^n p = 0$. The expression $h(p) = yp + u$ then gives $y^{n-1} \cdot u = 0$ in $\phi^{-1}P$.

Since both $x \cdot u$ and $y^{n-1} \cdot u$ are zero in $\phi^{-1}P$, the element u can be written as $yx^{n-1}v$ for some element $v \in P$. Now consider the element $p' = p + x^{n-1}v$ in P . It does not belong to $\mathfrak{r}P$, since $p \notin \mathfrak{r}P$. Moreover, $yp' = yp + u = h(p)$, hence yp' belongs to M . Finally, $xp' = xp = a$. The latter means that we can start the argument all over again, but this time with p' instead of p . However, this time yp' belongs to M , and this is impossible when $\Omega_A^{-1}(M)$ is not isomorphic to k . □

In the proof of the second and final step in the reduction process, we need the following lemma. Recall that an indecomposable nonprojective A -module M

of stable constant Jordan type [1] satisfies (RP x) if the following hold: there exists a generator a for the nonprojective component of M over $k[x]$, such that $a \notin \mathfrak{r}M$ and $y^{n-1}a \neq 0$. By Remark 4.2, every such generator a has these properties. Similarly, the module M satisfies (RP y) if there exists a generator b for the nonprojective component of M over $k[y]$, with $b \notin \mathfrak{r}M$ and $x^{n-1}b \neq 0$. The lemma shows that when the assumptions from the last couple of results hold, then if we take any element $m \in M \setminus \mathfrak{r}M$, either $x^{n-1}m$ or $y^{n-1}m$ must be nonzero.

LEMMA 4.7. *Let M be an indecomposable A -module of stable constant Jordan type [1], satisfying either (RP x) or (RP y). Moreover, suppose that $\dim_k \underline{\text{Hom}}_A(M, M) = 1$ and ${}_{\psi}M \simeq M \simeq {}_{\phi}M$, where ψ and ϕ are the automorphisms from Lemma 4.4. Then for every element $m \in M \setminus \mathfrak{r}M$, either $x^{n-1}m$ or $y^{n-1}m$ is nonzero.*

Proof. We prove the (RP x)-version. Take such an element $m \in M \setminus \mathfrak{r}M$, and suppose, for a contradiction, that $x^{n-1}m = 0$ and $y^{n-1}m = 0$. Consider a projective cover

$$0 \rightarrow \Omega_A^1(M) \hookrightarrow P \xrightarrow{d} M \rightarrow 0$$

of M , and let $p \in P$ be an element with $m = d(p)$. Since $m \notin \mathfrak{r}M$, the element p does not belong to $\mathfrak{r}P$, and so $y^{n-1}x^{n-1}p \neq 0$. Note that since $0 = y^{n-1}d(p) = d(y^{n-1}p)$, the element $y^{n-1}p$ belongs to $\text{Ker } d = \Omega_A^1(M)$, and similarly so does $x^{n-1}p$.

We shall modify the element p and obtain a new element $p' \in P$, with the property that $d(p') \in M \setminus \mathfrak{r}M$, and with

$$y^{n-1}p' = y^{n-1}p \quad \text{and} \quad xd(p') = 0$$

It then follows that $y^{n-1}x^{n-1}M$ is nonzero, a contradiction since M is indecomposable and not projective. To see this, let $z \in P$ be an element with the property that $d(z)$ generates the one-dimensional nonprojective component of M over $k[x]$. Then $d(p') = \alpha d(z) + w$ for some scalar $\alpha \in k$ and some element w in the $k[x]$ -free summand. As $xd(p') = 0 = xd(z)$, we see that $xw = 0$, so $w = x^{n-1}w'$ for some w' in the free summand. This implies that $w \in x^{n-1}M$, and since $d(p') \notin \mathfrak{r}M$, the scalar α must be nonzero. Now

$$\begin{aligned} 0 &= y^{n-1}m = y^{n-1}d(p) = d(y^{n-1}p) = d(y^{n-1}p') = y^{n-1}d(p') \\ &= \alpha y^{n-1}d(z) + y^{n-1}w \end{aligned}$$

and so since $y^{n-1}d(z)$ is nonzero by (RP x), we see that $y^{n-1}x^{n-1}w' = y^{n-1}w \neq 0$. Thus we have produced a nonzero element in $y^{n-1}x^{n-1}M$.

To construct the element p' with the desired properties, we apply the second part of Lemma 4.4 to the module $\Omega_A^1(M)$: there exists an A -homomorphism $h: P \rightarrow {}_{\psi^{-1}}\Omega_A^1(M)$ such that the composition $h \circ i: \Omega_A^1(M) \rightarrow {}_{\psi^{-1}}\Omega_A^1(M)$ is multiplication by x , where i is the inclusion map $\Omega_A^1(M) \hookrightarrow P$. Therefore, in ${}_{\psi^{-1}}\Omega_A^1(M)$, there are equalities

$$q^{n-1}y^{n-1}h(p) = y^{n-1} \cdot h(p) = h(y^{n-1}p) = xy^{n-1}p = q^{n-1}y^{n-1}xp$$

and this is a nonzero element since $y^{n-1}x^{n-1}p \neq 0$ and x and y q -commute. Viewing ${}_{\psi^{-1}}\Omega_A^1(M)$ as a submodule of ${}_{\psi^{-1}}P$, we see that

$$q^{n-1}y^{n-1}(h(p) - xp) = 0$$

in the latter, and then also in P itself. Therefore, there is an element $u \in yP$ with $h(p) - xp = u$; say $u = yu'$ for some $u' \in P$.

Now we use the fact that $x^{n-1}p \in \Omega_A^1(M)$, and obtain

$$x^{n-1}u = x^{n-1}(h(p) - xp) = x^{n-1}h(p) = h(x^{n-1}p) = h \circ i(x^{n-1}p) = x^n p = 0$$

Thus $x^{n-1}yu' = 0$ in P , so we can write $u' = xu_1 + y^{n-1}u_2$ for some $u_1, u_2 \in P$. This, in turn, gives

$$u = yu' = yxu_1 = x(q^{-1}yu_1) = xv$$

where $v = q^{-1}yu_1 \in yP$. Now consider the element $p + v$ in P ; call it p' . It follows from the definition of u and v that

$$xp' = xp + xv = xp + u = h(p)$$

and

$$y^{n-1}p' = y^{n-1}p + y^{n-1}v = y^{n-1}p$$

since $v \in yP$. Moreover, since $y^{n-1}p$ belongs to $\Omega_A^1(M)$, so does $y^{n-1}p'$. Note also that p' cannot belong to $\mathfrak{r}P$, for $p \notin \mathfrak{r}P$ whereas $v = q^{-1}yu_1 \in \mathfrak{r}P$. Therefore, since d is a projective cover, the element $d(p')$ does not belong to $\mathfrak{r}M$. Moreover, as xp' equals $h(p)$, and the latter belongs to ${}_{\psi^{-1}}\Omega_A^1(M)$ and therefore also to $\Omega_A^1(M) = \text{Ker } d$, we see that $xd(p') = d(xp') = 0$. We have shown that the element p' has the desired properties. □

We can now prove the second and final step in the reduction process.

PROPOSITION 4.8. *Suppose that $n \geq 3$, and let M be an indecomposable A -module of stable constant Jordan type [1]. Furthermore, suppose that $\dim_k \underline{\text{Hom}}_A(M, M) = 1$ and ${}_{\psi}M \simeq M \simeq {}_{\phi}M$, where ψ and ϕ are the automorphisms from Lemma 4.4. Then if M satisfies (RP), so does $\Omega_A^{-1}(M)$.*

Proof. As before, we only show that $\Omega_A^{-1}(M)$ satisfies (RP x), since the (RP y)-version is completely similar. However, this time we need the fact that the module M satisfies both (RP x) and (RP y).

As a $k[x]$ -module, we can decompose M as $\text{span}_k\{a\} \oplus F$ for some free module F , where a generates the one-dimensional nonprojective component. The rank property (RP x) then gives $a \notin \mathfrak{r}M$ and $y^{n-1}a \neq 0$. As in the previous proof, consider the injective envelope

$$0 \rightarrow M \hookrightarrow P \xrightarrow{d} \Omega_A^{-1}(M) \rightarrow 0$$

of M , viewed as a sequence of $k[x]$ -modules. Since $xa = 0$, we can write a as $a = x^{n-1}p$ for some $p \in P$, and then since $y^{n-1}x^{n-1}p = y^{n-1}a \neq 0$, this element p does not belong to $\mathfrak{r}P$. Then $d(p)$ does not belong to $\mathfrak{r}\Omega_A^{-1}(M)$, and it generates the nonprojective component of dimension $n - 1$ over $k[x]$. By definition of the rank property (RP x), we must show that $y^{n-1}x^{n-2}d(p)$ is nonzero.

Suppose, for a contradiction, that $y^{n-1}x^{n-2}d(p) = 0$. Then $y^{n-1}x^{n-2}p \in \text{Ker } d = M$; call this element m . We shall show that $m = y^{n-1}m'$ for some other element $m' \in M$. To do this, note first that both ym and x^2m are zero. Now consider M as a module over $k[y]$; as such, it decomposes as $\text{span}_k\{b\} \oplus F'$ for some free module F' , where b generates the one-dimensional nonprojective component. Since $ym = 0$, we can write m as $m = \alpha b + y^{n-1}m'$ for some scalar α and some $m' \in F'$. Furthermore, since M satisfies (RP y), the element $x^{n-1}b$ is nonzero. Therefore, if α is nonzero, then so is $x^{n-1}m$, since $x^{n-1}y^{n-1}m' = 0$ as M is not projective. But this cannot be the case, for we saw above that $x^2m = 0$, and by assumption $n \geq 3$. The scalar α must therefore be zero, and this shows that $m = y^{n-1}m'$.

Finally, consider the element $x^{n-2}p - m'$ in P ; call it p' . Since $m' \in M$ and $x^{n-1}p = a \in M$, the element xp' belongs to M ; let us denote it by m'' . This element does not belong to $\mathfrak{r}M$, for $m'' = a + xm'$, and $a \notin \mathfrak{r}M$ since M satisfies (RP x). Therefore, by Lemma 4.7, either $y^{n-1}m''$ or $x^{n-1}m''$ must be nonzero. As $xa = 0$, it must be the case that $y^{n-1}m'' \neq 0$, but

$$\begin{aligned} y^{n-1}m'' &= y^{n-1}x^{n-1}p - y^{n-1}xm' = q^{1-n}x(y^{n-1}x^{n-2}p - y^{n-1}m') \\ &= q^{1-n}x(m - m) = 0 \end{aligned}$$

This is a contradiction, and so $y^{n-1}x^{n-2}d(p)$ must be nonzero. □

We now prove the main result in this section. It characterizes the syzygies of the simple A -module k . As mentioned, the idea behind the proof is to start with a syzygy having the rank property (RP), and then show that its cosyzygies get smaller and smaller, using Proposition 4.3, Proposition 4.6 and Proposition 4.8. We then end up with the module k itself.

THEOREM 4.9. *If the ground field k is infinite, then for an indecomposable nonprojective module M over $A = A_q^2$, the following are equivalent:*

- (1) $M \simeq \Omega_A^i(k)$ for some $i \in \mathbb{Z}$;
- (2) M has constant stable Jordan type either $[1]$ or $[n - 1]$. Moreover, $\underline{\text{Hom}}_A(M, M)$ is one-dimensional, and ${}_\psi M \simeq M$ for every homogeneous automorphism $\psi: A \rightarrow A$;
- (3) M has constant stable Jordan type either $[1]$ or $[n - 1]$. Moreover, $\underline{\text{Hom}}_A(M, M)$ is one-dimensional, and ${}_\psi M \simeq M \simeq {}_\phi M$, where ψ and ϕ are the automorphisms from Lemma 4.4.

Proof. Suppose that (1) holds. As we explained in the proof of Lemma 4.7, whenever we take an A -module X and automorphism $\psi: A \rightarrow A$, there is

an isomorphism ${}_{\psi}\Omega_A^1(X) \simeq \Omega_A^1({}_{\psi}X)$. Thus for every integer i , the modules ${}_{\psi}\Omega_A^i(X)$ and $\Omega_A^i({}_{\psi}X)$ are isomorphic. This implies that

$${}_{\psi}M \simeq {}_{\psi}\Omega_A^i(k) \simeq \Omega_A^i({}_{\psi}k) \simeq \Omega_A^i(k) \simeq M$$

for every homogeneous automorphism ψ , since ${}_{\psi}k \simeq k$. Moreover, from the isomorphisms

$$\underline{\text{Hom}}_A(M, M) \simeq \underline{\text{Hom}}_A(\Omega_A^i(k), \Omega_A^i(k)) \simeq \underline{\text{Hom}}_A(k, k)$$

we see that $\dim_k \underline{\text{Hom}}_A(M, M) = 1$, and from Corollary 2.7 we know that M has constant stable Jordan type either $[1]$ or $[n - 1]$. This shows that (2) holds, and since (2) trivially implies (3), we need to show that (3) implies (1).

Suppose therefore that (3) holds, and take any syzygy $\Omega_A^i(M)$ of M . As above, all the properties from (3) also hold for this module: the vector space $\underline{\text{Hom}}_A(\Omega_A^i(M), \Omega_A^i(M))$ is one-dimensional, there are isomorphisms ${}_{\psi}\Omega_A^i(M) \simeq \Omega_A^i(M) \simeq {}_{\phi}\Omega_A^i(M)$, and $\Omega_A^i(M)$ has constant stable Jordan type either $[1]$ or $[n - 1]$. We shall show that there exists a syzygy of Jordan type $[n - 1]$ satisfying the rank property (RP).

As mentioned before Theorem 2.6, the algebra A has finitely generated cohomology. This means that its Hochschild cohomology ring $\text{HH}^*(A)$ is Noetherian, and that $\text{Ext}_A^*(X, Y)$ is a finitely generated $\text{HH}^*(A)$ -module for all A -modules X and Y . As explained in [BE1, Section 2], it follows from [Op2, Corollary 3.5] that there exists a polynomial subalgebra $H = k[\eta_1, \eta_2]$ of $\text{HH}^*(A)$, with both η_1 and η_2 in degree 2, and such that $\text{Ext}_A^*(X, Y)$ is a finitely generated H -module for all A -modules X and Y . In particular, $\text{Ext}_A^*(M, A/\mathfrak{r})$ is finitely generated over H .

Since the ground field k is infinite, the first part of the proof of [Be1, Theorem 2.5] shows that there exists a homogeneous element $\eta \in H$, of degree 2, such that multiplication

$$\text{Ext}_A^i(M, A/\mathfrak{r}) \xrightarrow{\eta} \text{Ext}_A^{i+2}(M, A/\mathfrak{r})$$

is injective for $i \gg 0$. From [Be1, Proposition 2.2] and its proof, we see that this element η gives rise to an eventually surjective chain map $f_{\eta}: \mathbb{P} \rightarrow \mathbb{P}$ of degree -2 , where \mathbb{P} is the minimal projective resolution

$$\cdots \rightarrow P_2 \xrightarrow{d_2} P_1 \xrightarrow{d_1} P_0$$

of M . For each $i \geq 0$, this chain map gives a commutative diagram

$$\begin{array}{ccccccc} P_{i+3} & \xrightarrow{d_{i+3}} & P_{i+2} & \longrightarrow & \Omega_A^{i+2}(M) & \longrightarrow & 0 \\ \downarrow f_{i+3} & & \downarrow f_{i+2} & & & & \\ P_{i+1} & \xrightarrow{d_{i+1}} & P_i & \longrightarrow & \Omega_A^i(M) & \longrightarrow & 0 \end{array}$$

with exact rows, a diagram that induces a map $g_{i+2}: \Omega_A^{i+2}(M) \rightarrow \Omega_A^i(M)$. As f_i is surjective for all $i \gg 0$, so is g_i .

By Theorem 2.6, the complexity of the module M is 2. In particular, this means that M is not periodic, and so the maps g_i cannot be isomorphisms. Namely, if $\Omega_A^{i+2}(M)$ were isomorphic to $\Omega_A^i(M)$, then $\Omega_A^2(M)$ would be isomorphic to M . Thus for all $i \gg 0$, the map g_{i+2} is surjective but not injective, and therefore $\dim_k \Omega_A^{i+2}(M) > \dim_k \Omega_A^i(M)$. Taking the alternating sum of the dimensions of the modules in the exact sequence

$$0 \rightarrow \Omega_A^{i+2}(M) \rightarrow P_{i+1} \xrightarrow{d_{i+1}} P_i \rightarrow \Omega_A^i(M) \rightarrow 0$$

we then see that $\dim_k P_{i+1} > \dim_k P_i$. This shows that $\beta_{i+1}(M) > \beta_i(M)$ for all $i \gg 0$. Now since every second syzygy of M is of constant stable Jordan type $[n-1]$, we may now apply Proposition 4.5: there exists an integer i such that $\Omega_A^i(M)$ is of constant stable Jordan type $[n-1]$ and satisfies (RP).

Let i_0 be an integer with the property that $\Omega_A^{i_0}(M)$ is of constant stable Jordan type $[n-1]$ and satisfies (RP), and such that $\beta_{i_0}(M)$ is minimal; by the above, such an integer exists. By Proposition 4.6, the module $\Omega_A^{i_0-1}(M)$ is either isomorphic to k , or satisfies (RP). We claim that it must be isomorphic to k . If not, then by Proposition 4.8, the module $\Omega_A^{i_0-2}(M)$ also satisfies (RP), since the stable Jordan type of $\Omega_A^{i_0-1}(M)$ is $[1]$. However, when we apply Proposition 4.3 to both $\Omega_A^{i_0}(M)$ and $\Omega_A^{i_0-1}(M)$, we see that there are strict inequalities $\beta_{i_0}(M) > \beta_{i_0-1}(M) > \beta_{i_0-2}(M)$. Since the module $\Omega_A^{i_0-2}(M)$ is of constant stable Jordan type $[n-1]$, this contradicts the minimality of $\beta_{i_0}(M)$. Consequently, the module $\Omega_A^{i_0-1}(M)$ must be isomorphic to k . \square

We suspect that the condition $\dim \underline{\text{Hom}}_A(M, M) = 1$ on the stable homomorphism spaces in the statement of Theorem 4.9 is not needed. For a motivation for this condition, see [Ca2].

REFERENCES

- [Ben] D. Benson, *Representations of elementary abelian p -groups and vector bundles*, Cambridge Tracts in Mathematics, 208, Cambridge University Press, Cambridge, 2017, xvii+328 pp.
- [BEH] D. Benson, K. Erdmann, M. Holloway, *Rank varieties for a class of finite-dimensional local algebras*, J. Pure Appl. Algebra 211 (2007), no. 2, 497–510.
- [Bel] P.A. Bergh, *Complexity and periodicity*, Colloq. Math. 104 (2006), no. 2, 169–191.
- [Be2] P.A. Bergh, *Ext-symmetry over quantum complete intersections*, Arch. Math. (Basel) 92 (2009), no. 6, 566–573.

- [BE1] P.A. Bergh, K. Erdmann, *The Avrunin-Scott theorem for quantum complete intersections*, J. Algebra 322 (2009), no. 2, 479–488.
- [BE2] P.A. Bergh, K. Erdmann, *The stable Auslander-Reiten quiver of a quantum complete intersection*, Bull. Lond. Math. Soc. 43 (2011), no. 1, 79–90.
- [BeO] P.A. Bergh, S. Oppermann, *Cohomology of twisted tensor products*, J. Algebra 320 (2008), no. 8, 3327–3338.
- [Ca1] J.F. Carlson, *Endo-trivial modules over (p, p) -groups*, Illinois J. Math. 24 (1980), no. 2, 287–295.
- [Ca2] J.F. Carlson, *A characterization of endotrivial modules over p -groups*, Manuscripta Math. 97 (1998), no. 3, 303–307.
- [CFP] J.F. Carlson, E.M. Friedlander, J. Pevtsova, *Modules of constant Jordan type*, J. Reine Angew. Math. 614 (2008), 191–234.
- [Erd] K. Erdmann, *Blocks of tame representation type and related algebras*, Lecture Notes in Mathematics, 1428, Springer-Verlag, Berlin, 1990, xvi+312 pp.
- [EHSST] K. Erdmann, M. Holloway, N. Snashall, Ø. Solberg, R. Taillefer, *Support varieties for selfinjective algebras*, K-Theory 33 (2004), no. 1, 67–87.
- [Op1] S. Oppermann, *A lower bound for the representation dimension of kC_n^p* , Math. Z. 256 (2007), no. 3, 481–490.
- [Op2] S. Oppermann, *Hochschild cohomology and homology of quantum complete intersections*, Algebra Number Theory 4 (2010), no. 7, 821–838.
- [SkY] A. Skowroński, K. Yamagata, *Frobenius algebras. I. Basic representation theory*, EMS Textbooks in Mathematics, European Mathematical Society (EMS), Zürich, 2011, xii+650 pp.

Petter Andreas Bergh
Institutt for Matematiske Fag
NTNU
N-7491 Trondheim
Norway
petter.bergh@ntnu.no

Karin Erdmann
Mathematical Institute
University of Oxford
Andrew Wiles Building
Radcliffe Observatory Quarter
Oxford OX2 6GG
United Kingdom
erdmann@maths.ox.ac.uk

David A. Jorgensen
Department of Mathematics
University of Texas at Arlington
Arlington, TX 76019
USA
djorgens@uta.edu