



Optimal Double Stopping Problems for Maxima and Minima of Geometric Brownian Motions

Pavel V. Gapeev¹ · Peter M. Kort^{2,3} · Maria N. Lavrutich⁴ · Jacco J. J. Thijssen⁵

Received: 24 May 2021 / Revised: 23 February 2022 / Accepted: 8 April 2022
© The Author(s) 2022

Abstract

We present closed-form solutions to some double optimal stopping problems with payoffs representing linear functions of the running maxima and minima of a geometric Brownian motion. It is shown that the optimal stopping times are the first times at which the underlying process reaches some lower or upper stochastic boundaries depending on the current values of its running maximum or minimum. The proof is based on the reduction of the original double optimal stopping problems to sequences of single optimal stopping problems for the resulting three-dimensional continuous Markov process. The latter problems are solved as the equivalent free-boundary problems by means of the smooth-fit and normal-reflection conditions for the value functions at the optimal stopping boundaries and the edges of the three-dimensional state space. We show that the optimal stopping boundaries are determined as the extremal solutions of the associated first-order nonlinear ordinary differential equations. The obtained results are related to the valuation of perpetual real double lookback options with floating sunk costs in the Black-Merton-Scholes model.

Keywords Perpetual real double lookback options · the Black-Merton-Scholes model · Geometric Brownian motion · Double optimal stopping problem · First hitting time · Free-boundary problem · Instantaneous stopping and smooth fit · Normal reflection · A change-of-variable formula with local time on surfaces

Mathematics Subject Classification Primary 60G40 · 34B40 · 91G20 · Secondary 60J60 · 60J65 · 91B70

JEL Classification G13

✉ Pavel V. Gapeev
p.v.gapeev@lse.ac.uk

Extended author information available on the last page of the article

1 Formulation of the Problems

The main aim of this paper is to present closed-form solutions to the discounted optimal double stopping problems with the values:

$$\bar{V}_1 = \sup_{\tau \leq \zeta} E \left[e^{-r\tau} \left(L_1 X_\tau - \min_{0 \leq t \leq \tau} X_t \right) + e^{-r\zeta} \left(\max_{0 \leq t \leq \zeta} X_t - K_1 X_\zeta \right) \right] \tag{1}$$

and

$$\bar{V}_2 = \sup_{\tau \leq \zeta} E \left[e^{-r\tau} \left(\max_{0 \leq t \leq \tau} X_t - K_2 X_\tau \right) + e^{-r\zeta} \left(L_2 X_\zeta - \min_{0 \leq t \leq \zeta} X_t \right) \right] \tag{2}$$

for some given constants $L_i \geq 1 \geq K_i > 0$, for $i = 1, 2$. Here, for a precise formulation of the problem, we consider a probability space (Ω, \mathcal{F}, P) with a standard Brownian motion $B = (B_t)_{t \geq 0}$. We assume that the process $X = (X_t)_{t \geq 0}$ is defined by:

$$X_t = x \exp \left((\mu - \sigma^2/2) t + \sigma B_t \right) \tag{3}$$

so that it solves the stochastic differential equation:

$$dX_t = \mu X_t dt + \sigma X_t dB_t \quad (X_0 = x) \tag{4}$$

where $\mu < r$, $r > 0$ and $\sigma > 0$ are given constants, and $x > 0$ is fixed. In our application, the process X describes the current state of technological progress, which changes over time due to the active process of research and development in a branch of the industry, where r is the discount rate. The running maximum $S = (S_t)_{t \geq 0}$ and minimum $Q = (Q_t)_{t \geq 0}$ of the process X defined by:

$$S_t = s \vee \max_{0 \leq u \leq t} X_u \quad \text{and} \quad Q_t = q \wedge \min_{0 \leq u \leq t} X_u \tag{5}$$

for arbitrary $0 < q \leq x \leq s$, respectively, can be interpreted as the best and the worst market valuation of technology achieved so far. Suppose that the suprema in (1) and (2) are taken over all stopping times τ and ζ with respect to the natural filtration $(\mathcal{F}_t)_{t \geq 0}$ of the process X , and the expectations there are taken with respect to the risk-neutral probability measure P . In this case, the values of (1) and (2) can be interpreted as the rational (or no-arbitrage) values of (perpetual) real lookback compound options with present values, which are linear in the running maximum or minimum of X , as well as sunk cost investment amounts, which are constant or linear in X , in the Black-Merton-Scholes model, respectively (see, e.g. Dixit and Pindyck [Dixit and Pindyck (1994); Chapter X] for the examples of standard compound real options).

The problem of (1), which has its dual of (2), is a typical valuation problem for financial lookback options. It has, however, a broader interpretation related to capital budgeting of real investment decisions, that is, real lookback options. In particular, by utilising the approach presented in this paper, decision makers are able to quantify the financial value of investments in new promising technologies, as well as of the policy mechanisms that can be used to incentivise such investments. More specifically, we have in mind that the current state of technological progress is observable and is described by the process X , whereas r is the discount rate.

In order to adopt a new technology at time $t \geq 0$, a firm is required to pay sunk costs which are proportional to the value of the technology in the market denoted by $X = (X_t)_{t \geq 0}$. This implies that more valuable developments are also associated with higher investment cost (for example, due to competition for suppliers). Typically, however, there is a time lag between a technological breakthrough and the actual realisation of the full potential of a new technology. The latter often requires additional investment in complementary technologies, changes in business processes or proper infrastructure (see Brynjolfsson and Hitt (2003) and Brynjolfsson et al. (2017)). In our problem, this feature is encapsulated by a second option available to the firm upon adoption of the new technology. We call this feature the *commercialisation option*. More specifically, by paying the sunk costs $K_1 X$, the firm is able to realise the full potential of the technology, given by the running maximum S of the process X . These sunk costs are proportional to the current market value of the technology's potential due to, for example, the need to develop technology-specific infrastructure. Ideally, the firm would like to enter the market, adopt the technology, when the cost of acquisition is low, and undertake further investments, once the technology is valuable enough. In other words, a firm would like to identify future winners early on, as once the market identifies the winning technologies, the costs of their adoption will be high. Hence, if the firm realises the potential of a technology too late, it pays larger sunk costs and delays further investments in developing this technology. Consequently, some emerging technologies reach their productive potential later than it might be desirable from a social welfare point of view. In this paper, we propose a valuation framework that allows to quantify the value associated with implementing a specific incentive mechanism inspired by financial lookback options, which stimulates innovations by reducing the firm's regret of missing out on investment opportunities. This mechanism takes the form of an investment cost subsidy, which is equal to the difference between the current market value of the technology given by X , and the minimal value it has achieved until this moment Q . Then, upon the technology adoption, the firm receives the value $L_1 X$ which is proportional to its current value X , whereas the costs paid by the firm are equal to Q , which reduces its regret associated with failing to time the market.

From the derived closed-form solutions we conclude that under such a support mechanism, firms have an incentive to adopt a technology when X is moving away from its running minimum Q . In this case, the technology is more valuable. Upon adoption the following two cases can occur. If the technology is sufficiently promising, then the firm will wait with commercialisation. This happens, because the probability that a new, higher, maximum S will be reached soon is large and, thus, there is a larger potential for a higher payoff which induces the firm to wait with commercialisation. However, if the market value is small relative to its running maximum S , then the firm commercialises the technology immediately after adoption as it is now unlikely that the technology will improve in comparison to its best performance achieved so far to warrant waiting for a higher maximum. These results show that double lookback options allow a social planner to subsidise the most efficient technologies without having to pick winners *ex ante*. Rather, the benefit of such a subsidy is a direct support for the realised winners, that is, the most desirable technologies as evidenced by market value, which could result in a considerable welfare increase.

Discounted optimal stopping problems for certain reward functionals depending on the running maxima and minima of continuous Markov (diffusion-type) processes were initiated by Shepp and Shiryaev (1993) and further developed by Pedersen (2000); Guo and Shepp (2001); Gapeev (2007); Guo and Zervos (2010); Peskir (2012, 2014); Glover

et al. (2013); Rodosthenous and Zervos (2017); Gapeev (2019, 2020); Gapeev et al. (2021); Gapeev and Li (2021); Gapeev and Al Motairi (2021); Gapeev (2022) among others. The main feature in the analysis of such optimal stopping problems was that the normal-reflection conditions hold for the value functions at the diagonals of the state spaces of the multi-dimensional continuous Markov processes having the initial processes and the running extrema as their components. It was shown, by using the established by Peskir (1998) maximality principle for solutions of optimal stopping problems, which is equivalent to the superharmonic characterisation of the value functions, that the optimal stopping boundaries are characterised by the appropriate extremal solutions of certain (systems of) first-order nonlinear ordinary differential equations. Other optimal stopping problems in models with spectrally negative Lévy processes and their running maxima were studied by Asmussen et al. (2003); Avram et al. (2004); Ott (2013); Kyprianou and Ott (2014) among others.

We further consider the problems of (1) and (2) as the associated double (two-step) optimal stopping problems of (6) and (7) for the three-dimensional continuous Markov processes having the process X as well as its running maximum S and minimum Q as their state space components. The resulting problems turn out to be necessarily three-dimensional in the sense that they cannot be reduced to optimal stopping problems for Markov processes of lower dimensions. The original optimal double stopping problems are reduced to the appropriate sequences of single optimal stopping problems which are solved as the equivalent free-boundary problems for the value functions which satisfy the smooth-fit conditions at the optimal stopping boundaries and the normal-reflection conditions at the edges of the state space of the three-dimensional processes. The multiple (multi-step) optimal stopping problems for one-dimensional diffusion processes have recently drawn a considerable attention in the related literature. Duckworth and Zervos (2000) studied an investment model with entry and exit decisions alongside a choice of the production rate for a single commodity. The initial valuation problem was reduced to a double (two-step) optimal stopping problem which was solved through the associated dynamic programming differential equation. Carmona and Touzi (2008) derived a constructive solution to the problem of pricing of perpetual swing contracts, the recall components of which could be viewed as contingent claims with multiple exercises of American type, using the connection between optimal stopping problems and the associated with them Snell envelopes. Carmona and Dayanik (2008) then obtained a closed form solution of a multiple (multi-step) optimal stopping problem for a general linear regular diffusion process and a general payoff function among others. The problem of pricing of American compound standard put and call options in the classical Black-Merton-Scholes model was explicitly solved in Gapeev and Rodosthenous (2014a). The same problem in the more general stochastic volatility framework was studied by Chiarella and Kang (2009), where the associated two-step free-boundary problems for partial differential equations were solved numerically, by means of a modified sparse grid approach.

The rest of the paper is organised as follows. In Sect. 2, we embed the original problems with the values V_i^* , for $i = 1, 2$, in (1) and (2) into the optimal multiple stopping problems for the values functions $V_i^*(x, s, q)$, for $i = 1, 2$, in (6) and (7) for the three-dimensional continuous Markov process (X, S, Q) defined in (3) and (5), respectively. It is shown that the optimal exercise times $\tau_1^*(S, Q)$ and $\tau_2^*(S, Q)$ are the first times at which the process X reaches some upper or lower boundaries $b^*(S, Q)$ or $a^*(S, Q)$ depending on the current values of either the processes S and Q , respectively. In Sect. 3, we derive closed-form expressions for the candidate value functions for $V_i^*(x, s, q)$, for $i = 1, 2$, as solutions to the

equivalent free-boundary problems and apply the normal-reflection conditions at the edges of the three-dimensional state space for (X, S, Q) to characterise the candidate optimal stopping boundaries for $b^*(S, Q)$ and $a^*(S, Q)$ or as the minimal and maximal solutions of the appropriate first-order nonlinear ordinary differential equations, respectively. In Sect. 4, by applying the change-of-variable formula with local time on surfaces from Peskir (2007), it is verified that the resulting solutions to the free-boundary problem provide the expressions for the value functions and the optimal stopping boundaries for the underlying asset price process in the original problems. In Sect. 5, we recall the explicit solutions of the inner optimal stopping problems with the value functions $U_1^*(x, s)$ and $U_2^*(x, q)$ from (72). The main results of the paper are stated in Theorem 4.1. The resulting method is presented in Corollary 4.2 and described in Remark 4.3.

2 Preliminaries

In this section, we describe the structure of the three-dimensional optimal stopping problems of (1) and (2) which are related to the floating sunk costs real double lookback option pricing problems and formulate the equivalent free-boundary problems.

2.1 The Two-step Optimal Stopping Problems

It is seen that the problems of (1) and (2) can naturally be embedded into the optimal double stopping problems for the (time-homogeneous strong) Markov process $(X, S, Q) = (X_t, S_t, Q_t)_{t \geq 0}$ defined in (3) and (5) with the values:

$$\bar{V}_1 = \sup_{\tau \leq \zeta} E[e^{-r\tau} (L_1 X_\tau - Q_\tau) + e^{-r\zeta} (S_\zeta - K_1 X_\zeta)] \tag{6}$$

and

$$\bar{V}_2 = \sup_{\tau \leq \zeta} E[e^{-r\tau} (S_\tau - K_2 X_\tau) + e^{-r\zeta} (L_2 X_\zeta - Q_\zeta)] \tag{7}$$

for some $L_i \geq 1 \geq K_i > 0$, for $i = 1, 2$, fixed, where the suprema are taken over all stopping times τ and ζ with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$. In this case, by virtue of the strong Markov property of the process (X, S, Q) , the original problems of (6) and (7) can be reduced to the optimal stopping problems with the values:

$$\bar{V}_i = \sup_{\tau} E[e^{-r\tau} G_i(X_\tau, S_\tau, Q_\tau)] \tag{8}$$

where the suprema are taken over all stopping times τ of (X, S, Q) , and we set:

$$G_1(x, s, q) = L_1 x - q + U_1^*(x, s) \quad \text{and} \quad G_2(x, s, q) = s - K_2 x + U_2^*(x, q) \tag{9}$$

for some $L_1 \geq 1 \geq K_2 > 0$ fixed, respectively. Here, the functions $U_1^*(x, s)$ and $U_2^*(x, q)$ represent the values of the optimal stopping problems formulated in (72), where the optimal stopping times η_i^* , for $i = 1, 2$, have the form of (73), for some boundaries $0 < g^*(s) \equiv \lambda_* s < s$ and $h^*(q) \equiv v_* q > q > 0$ determined in Corollary 5.1 below.

2.2 The Outer Optimal Stopping Problems

Let us first transform the rewards in the expressions of (8) and (9) with the aim to formulate the associated optimal stopping problems. For this purpose, we use standard arguments based on an application of Itô’s formula (see, e.g. [Liptser and Shiryaev (2001); Theorem 4.4] or [Revuz and Yor (1999); Chapter II, Theorem 3.2]) to show that the infinitesimal operator \mathbb{L} of the process (X, S, Q) from (4) and (5) acts on an arbitrary function $V(x, s, q)$ from the class $C^{2,1,1}$ on E according to the rule:

$$(\mathbb{L}V)(x, s, q) = \mu x \partial_x V(x, s, q) + \frac{\sigma^2 x^2}{2} \partial_{xx} V(x, s, q) \quad \text{in } 0 < q < x < s \quad (10)$$

while we should also assume that:

$$\partial_q = 0 \quad \text{at } 0 < x = q < s \quad \text{and} \quad \partial_s = 0 \quad \text{at } 0 < q < x = s \quad (11)$$

in order to have the operator \mathbb{L} well-defined at d_1 and d_2 , respectively (see, e.g. [Peskir (1998); Subsection 3.1]). We first recall from the results of Beibel and Lerche (1997); Pedersen (2000) and Guo and Shepp (2001) (as well as Gapeev (2020)) on the expressions in (85)-(86) for the value functions $U_1^*(x, s)$ and $U_2^*(x, q)$ in (72) which solve the free-boundary problems in (74)-(80) that the processes $e^{-rt}U_1^*(X_t, S_t)$ and $e^{-rt}U_2^*(X_t, Q_t)$ admit the representations:

$$\begin{aligned} e^{-rt} U_1^*(X_t, S_t) &= U_1^*(x, s) + \int_0^t e^{-ru} (\mathbb{L}U_1^* - r U_1^*)(X_u, S_u) I(X_u < S_u) du \\ &\quad + \int_0^t e^{-ru} \partial_s U_1^*(X_u, S_u) I(X_u = S_u) dS_u \\ &\quad + \int_0^t e^{-ru} \partial_x U_1^*(X_u, S_u) I(X_u < S_u) dB_u \end{aligned} \quad (12)$$

and

$$\begin{aligned} e^{-rt} U_2^*(X_t, Q_t) &= U_2^*(x, q) + \int_0^t e^{-ru} (\mathbb{L}U_2^* - r U_2^*)(X_u, Q_u) I(X_u > Q_u) du \\ &\quad + \int_0^t e^{-ru} \partial_q U_2^*(X_u, Q_u) I(X_u = Q_u) dQ_u \\ &\quad + \int_0^t e^{-ru} \partial_x U_2^*(X_u, Q_u) I(X_u > Q_u) dB_u \end{aligned} \quad (13)$$

where the stochastic integrals with respect to the standard Brownian motion $B = (B_t)_{t \geq 0}$ are continuous square-integrable martingales. Let us now apply Itô’s formula to the processes $e^{-rt}G_i(X_t, S_t, Q_t)$, for $i = 1, 2$, to obtain:

$$\begin{aligned} e^{-rt} G_1(X_t, S_t, Q_t) &= G_1(x, s, q) \\ &\quad + \int_0^t e^{-ru} H_1(X_u, S_u, Q_u) I(Q_u < X_u < S_u) du \\ &\quad - \int_0^t e^{-ru} I(X_u = Q_u) dQ_u + N_t^1 \end{aligned} \quad (14)$$

with

$$H_1(x, s, q) = (\mathbb{L}G_1 - rG_1)(x, s, q) \equiv (rq - (r - \mu)L_1x) I(x > g^*(s)) - (r(s - q) + (r - \mu)(L_1 - K_1)x) I(x \leq g^*(s)) \tag{15}$$

and

$$e^{-rt} G_2(X_t, S_t, Q_t) = G_2(x, s, q) + \int_0^t e^{-ru} H_2(X_u, S_u, Q_u) I(Q_u < X_u < S_u) du + \int_0^t e^{-ru} I(X_u = S_u) dS_u + N_t^2 \tag{16}$$

with

$$H_2(x, s, q) = (\mathbb{L}G_2 - rG_2)(x, s, q) \equiv ((r - \mu)K_2x - rs) I(x < h^*(q)) - (r(s - q) + (r - \mu)(L_2 - K_2)x) I(x \geq h^*(q)) \tag{17}$$

for each $0 < q < x < s$, and all $t \geq 0$, where $I(\cdot)$ denotes the indicator function. Here, \mathbb{L} is the infinitesimal operator of the process (X, S, Q) having the form of (10)-(11) above, and the processes $N^i = (N_t^i)_{t \geq 0}$, for $i = 1, 2$, defined by:

$$N_t^i = \int_0^t e^{-ru} \partial_x G_i(X_u, S_u, Q_u) I(Q_u < X_u < S_u) \sigma X_u dB_u \tag{18}$$

for all $t \geq 0$, are continuous square-integrable martingales under the probability measure P . It also follows from the explicit expressions in (3)-(4) for the process X under the assumption $\mu < r$ as well as from the properties of the partial derivatives $\partial_x G_i(x, s, q)$, for $i = 1, 2$, and the structure of the other processes included into the expressions of (14) and (16) that the processes N^i , for $i = 1, 2$, defined in (18) are uniformly integrable. Note that the processes S and Q may change their values only at the times when $X_t = S_t$ and $X_t = Q_t$, for $t \geq 0$, respectively, and such times accumulated over the infinite horizon form the sets with zero Lebesgue measure, so that the indicators in the expressions of (14) and (16) as well as (18) can be ignored (see also Proof of Theorem 4.1 below for more explanations and references). Then, inserting τ in place of t and applying Doob's optional sampling theorem (see, e.g. [Liptser and Shiryaev (2001); Chapter III, Theorem 3.6] or [Revuz and Yor (1999); Chapter II, Theorem 3.2]) to the expressions in (14) and (16), we get that the equalities:

$$E[e^{-r\tau} G_1(X_\tau, S_\tau, Q_\tau)] = G_1(x, s, q) + E \left[\int_0^\tau e^{-ru} H_1(X_u, S_u, Q_u) du - \int_0^\tau e^{-ru} dQ_u \right] \tag{19}$$

and

$$E[e^{-r\tau} G_2(X_\tau, S_\tau, Q_\tau)] = G_2(x, s, q) + E \left[\int_0^\tau e^{-ru} H_2(X_u, S_u, Q_u) du + \int_0^\tau e^{-ru} dS_u \right] \tag{20}$$

hold, for any stopping time τ with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$. Hence, taking into account the expressions in (19) and (20), we conclude that the optimal stopping problems

with the values of (8) are equivalent to the optimal stopping problems with the value functions:

$$V_1^*(x, s, q) = \sup_{\tau} E_{x,s,q} \left[\int_0^{\tau} e^{-ru} H_1(X_u, S_u, Q_u) du - \int_0^{\tau} e^{-ru} dQ_u \right] \tag{21}$$

and

$$V_2^*(x, s, q) = \sup_{\tau} E_{x,s,q} \left[\int_0^{\tau} e^{-ru} H_2(X_u, S_u, Q_u) du + \int_0^{\tau} e^{-ru} dS_u \right] \tag{22}$$

where the functions $H_i(x, s, q)$, for $i = 1, 2$, are defined in (15) and (17), for $(x, s, q) \in E$, respectively. Here, we denote by $E_{x,s,q}$ the expectation with respect to the probability measures $P_{x,s,q}$ under which the three-dimensional (time-homogeneous strong Markov) processes (X, S, Q) starts at $(x, s, q) \in E$, and by $E = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q \leq x \leq s\}$ the state spaces of (X, S, Q) . We further obtain solutions to the optimal stopping problems in (21) and (22) and verify below that the value functions $V_i^*(x, s, q)$, for $i = 1, 2$, are the solutions of the problems in (8), and thus, give the solutions of the original multiple optimal stopping problems in (1) and (2), under $s = q = x$.

It follows from the results of general theory of optimal stopping problems for Markov processes (see, e.g. [Peskir and Shiryaev (2006); Chapter I, Subsection 2.2]) that the continuation and stopping regions of the optimal stopping problems in (8) have the form:

$$C_i^* = \{(x, s, q) \in E \mid V_i^*(x, s, q) > 0\} \quad \text{and} \quad D_i^* = \{(x, s, q) \in E \mid V_i^*(x, s, q) = 0\} \tag{23}$$

for every $i = 1, 2$, respectively. It is seen from the results of Theorem 4.1 below that the value function $V_i^*(x, s, q)$ is continuous, so that the set C_i^* is open and D_i^* is closed, for every $i = 1, 2$.

2.3 The Structure of Optimal Stopping Times

Let us now specify the structure of the optimal stopping times in the *outer* optimal stopping problems of (21) and (22).

- (i) It follows from the structure of the second integrals of (21) and (22) as well as the facts that the process S is increasing and the process Q is decreasing that it is not optimal to exercise the *outer* parts of the contracts (or exercise the double lookback options for the *first* time), whenever the appropriate integrands are positive. In other words, the diagonals $d_1 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < x = q < s\}$ and $d_2 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q < x = s\}$ belong to the continuation regions C_1^* and C_2^* in (23), respectively. Moreover, it follows from the structure of the first integrals of (21) and (22) that it is not optimal to exercise the *outer* parts of the contracts (or exercise the double lookback options for the *first* time) when the inequalities $H_i(X_t, S_t, Q_t) \geq 0$, for $i = 1, 2$ hold, which are equivalent to $0 < g^*(S_t) < X_t \leq rQ_t / ((r - \mu)L_1)$ and $0 < rS_t / ((r - \mu)K_2) \leq X_t < h^*(Q_t)$ with $Q_t < X_t < S_t$, for all $t \geq 0$, respectively. In other words, these facts mean that the set $\{(x, s, q) \in E \mid 0 < q \vee g^*(s) < x \leq r q / ((r - \mu)L_1) \wedge s\}$ belongs to the continuation region C_1^* , while the set $\{(x, s, q) \in E \mid 0 < q \vee r s / ((r - \mu)K_2) \leq x < h^*(q) \wedge s\}$ belongs to the continuation region C_2^* in (23).


(ii) We now observe that it follows from the definition of the process (X, S, Q) in (3) and (5) and the structure of the rewards in (21) and (22) that, for each $s > 0$ fixed, there exist $0 < q \leq x$ such that x is sufficiently close to s and the point (x, s, q) belongs to the stopping region D_i^* , for every $i = 1, 2$. Moreover, for each $q > 0$ fixed, there exist $0 < x \leq s$ such that x is sufficiently close to q and the point (x, s, q) belongs to D_i^* , for $i = 1, 2$. By virtue of arguments similar to the ones applied in [Dubins et al. (1993); Subsection 3.3] and [Peskir (1998); Subsection 3.3], these properties can be explained by the facts that the costs of waiting until the process X coming from either such a large $x > 0$ decreases to the current value of the running minimum process Q or such a small $x > 0$ increases to the current value of the running maximum process S may be too large due to the presence of the discounting factors in the reward functionals of (21) and (22). Furthermore, by virtue of the asymptotic distributional properties of the running maximum S and minimum Q from (5) of the geometric Brownian motion X from (3)-(4) on the infinitesimally small time intervals (see, e.g. [Dubins et al. (1993); Subsection 3.3] for similar arguments applied to the running maxima of the Bessel processes, [Peskir (1998); Proposition 2.1] for similar properties of the running maxima of general diffusion processes, and [Gapeev and Li (2021); Theorem 2.1, Part (i)] for similar properties of the running maxima and minima of geometric Brownian motions), it follows that the reward functionals in (21) and (22) infinitesimally increase when $X_t = Q_t$ or $X_t = S_t$, for each $t \geq 0$. This fact also shows that all points (x, s, q) from the diagonals $d_1 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < x = q < s\}$ and $d_2 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q < x = s\}$ belong to the continuation regions C_i^* , for $i = 1, 2$, in (23), respectively.

On the one hand, if we take some $(x, s, q) \in D_1^*$ from (23) such that $x > (1 \vee r / ((r - \mu)L_1))q$ and use the fact that the process (X, S, Q) started at some (x', s, q) such that $0 < (1 \vee r / ((r - \mu)L_1))q < x < x' < s$ passes through the point (x, s', q) , for some $s' \geq s$, before hitting the plane $d_1 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < x = q < s\}$, then the representation of (19) for the reward functional in (21) implies that $V_1^*(x', s, q) \leq V_1^*(x, s, q) = 0$ holds, so that $(x', s, q) \in D_1^*$. Moreover, if we take some $(x, s, q) \in D_2^*$ from (23) such that $0 < x < (r / ((r - \mu)K_2) \wedge 1)s$ and use the fact that the process (X, S, Q) started at some (x'', s, q) such that $0 < q < x'' < x < (r / ((r - \mu)K_2) \wedge 1)s$ passes through the point (x, s, q'') , for some $0 < q'' \leq q$, before hitting the diagonal $d_2 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q < x = s\}$, then the representation of (20) for the reward functional in (22) implies that $V_2^*(x'', s, q) \leq V_2^*(x, s, q) = 0$ holds, so that $(x'', s, q) \in D_2^*$. Thus, we may conclude that the stopping regions D_i^* , for $i = 1, 2$, from (23) have the right-hand and left-hand parts, respectively.

On the other hand, if we take some $(x, s, q) \in C_1^*$ from (23) and use the fact that the process (X, S, Q) started at (x, s, q) passes through some point (x'', s'', q) such that $0 < q < x'' < x < s \leq s''$ before hitting the plane d_1 , then the representation of (19) for the reward functional in (21) implies that $V_1^*(x'', s, q) > V_1^*(x, s, q) = 0$ holds, so that $(x'', s, q) \in C_1^*$. Moreover, if take some $(x, s, q) \in C_2^*$ from (23) and use the fact that the process (X, S, Q) started at (x, s, q) passes through some point (x', s, q') such that $0 < q' \leq q < x < x' < s$ before hitting the plane d_2 , then the representation of (20) for the reward functional in (22) implies that $V_2^*(x', s, q) > V_2^*(x, s, q) = 0$ holds, so that $(x', s, q) \in C_2^*$.

(iii) We may therefore conclude that there exist functions $b^*(s, q)$ and $a^*(s, q)$ such that the inequalities $H_i(x, s, q) < 0$, for $i = 1, 2$, hold, for $(x, s, q) \in E$ such that $x \geq b^*(s, q)$ or $x \leq a^*(s, q)$, respectively. In this respect, the continuation regions C_i^* , for $i = 1, 2$, in (23) have the form:

Fig. 1 A computer drawing of the optimal exercise boundary $b^*(s, q)$



$$C_1^* = \{(x, s, q) \in E \mid x < b^*(s, q)\} \quad \text{and} \quad C_2^* = \{(x, s, q) \in E \mid x > a^*(s, q)\} \quad (24)$$


while the stopping regions D_i^* , for $i = 1, 2$, in (23) are given by:

$$D_1^* = \{(x, s, q) \in E \mid x \geq b^*(s, q)\} \quad \text{and} \quad D_2^* = \{(x, s, q) \in E \mid x \leq a^*(s, q)\}. \quad (25)$$

(Figs. 1–2 on this page illustrate computer drawings of the optimal stopping boundaries $b^*(s, q)$ and $a^*(s, q)$.)

- (iv) Let us finally clarify the location of the boundaries $b^*(s, q)$ and $a^*(s, q)$ in relation to the optimal stopping boundaries $g^*(s)$ and $h^*(q)$ from (73) for the optimal stopping problems with the value functions $U_1^*(x, s)$ and $U_2^*(x, q)$ in (72) below. For this purpose, we use the notations of the functions $F_i(x, s, q)$, for $i = 1, 2$, from (36) and (37) below. Suppose that either the inequality $b^*(s, q) < h^*(q)$ or $a^*(s, q) > g^*(s)$ holds, for some $0 < q < g^*(s) < s$ and $0 < q < h^*(q) < s$. In this case, for each point

Fig. 2 A computer drawing of the optimal exercise boundary $a^*(s, q)$



(x, s, q) such that $x \in (b^*(s, q), h^*(q))$ and $0 < q < x \leq g^*(s) < s$, we would have $L_1x - q + U_1^*(x, s) > L_1x - q + s - K_1x \equiv s - q + (L_1 - K_1)x = V_1^*(x, s, q) + F_1(x, s, q)$ contradicting the fact that $L_1x - q + U_1^*(x, s) \leq V_1^*(x, s, q) + F_1(x, s, q)$, for all $(x, s, q) \in E$. Also, for each point (x, s, q) such that $x \in (g^*(s), a^*(s, q))$ and $0 < q < h^*(q) \leq x < s$, we would have $s - K_2x + U_2^*(x, q) > s - K_2x + L_2x - q \equiv s - q + (L_2 - K_2)x = V_2^*(x, s, q) + F_2(x, s, q)$ contradicting the fact that $s - K_2x + U_2^*(x, q) \leq V_2^*(x, s, q) + F_2(x, s, q)$, for all $(x, s, q) \in E$. Hence, we may conclude that the inequalities $b^*(s, q) \geq h^*(q)$ and $a^*(s, q) \leq g^*(s)$ should hold, for all $0 < q < g^*(s) < s$ and $0 < q < h^*(q) < s$, respectively.

- (v) Recall that the problem of (1) can be interpreted as the combined problem of finding the optimal time to adopt a new technology in the presence of an investment cost subsidy and then to determine when it is optimal to commercialise it. The subsidised technology adoption occurs when the process X hits the boundary $b^*(S, Q)$ from above, implying that the first investment occurs when X moves away from its running minimum Q and the technology becomes more valuable. Upon adoption the following two cases can occur. If the technology is sufficiently promising, that is, if X is sufficiently large, so that $X > g^*(S)$ holds, then the firm will wait with commercialisation. This is because the probability that a new, higher, maximum S will be reached soon is large, and thus, there is a larger potential for a higher payoff which induces the firm to wait with commercialisation. If however, X is relatively small, then the firm commercialise the technology immediately after adoption as it is now unlikely that the technology will improve in comparison to its best performance achieved so far to warrant waiting for a higher maximum.

2.4 The Free-boundary Problems

In order to find analytic expressions for the unknown value functions $V_i^*(x, s, q)$, for $i = 1, 2$, from (21) and (22) with the unknown boundaries $b^*(s, q)$ and $a^*(s, q)$ from (24)-(25), let us use the results of general theory of optimal stopping problems for Markov processes (see, e.g. [Peskir and Shiryaev (2006); Chapter IV, Section 8]) as well as optimal stopping problems for maximum processes (see, e.g. [Peskir and Shiryaev (2006); Chapter V, Sections 15–20] and references therein). We can therefore reduce the optimal stopping problems of (21) and (22) to the equivalent free-boundary problems:

$$(\mathbb{L}V_1 - rV_1)(x, s, q) = -H_1(x, s, q) \quad \text{for } q < x < b(s, q) \tag{26}$$

$$(\mathbb{L}V_2 - rV_2)(x, s, q) = -H_2(x, s, q) \quad \text{for } a(s, q) < x < s \tag{27}$$

$$V_1(x, s, q)|_{x=b(s,q)-} = 0, \quad V_2(x, s, q)|_{x=a(s,q)+} = 0 \tag{28}$$

$$\partial_x V_1(x, s, q)|_{x=b(s,q)-} = 0, \quad \partial_x V_2(x, s, q)|_{x=a(s,q)+} = 0 \tag{29}$$

$$\partial_q V_1(x, s, q)|_{x=q+} = 1, \quad \partial_s V_2(x, s, q)|_{x=s-} = -1 \tag{30}$$

$$V_1(x, s, q) = 0 \quad \text{for } x > b(s, q), \quad V_2(x, s, q) = 0 \quad \text{for } x < a(s, q) \quad (31)$$

$$V_1(x, s, q) > 0 \quad \text{for } x \leq b(s, q), \quad V_2(x, s, q) = 0 \quad \text{for } x \geq a(s, q) \quad (32)$$

$$(\mathbb{L}V_1 - rV_1)(x, s, q) < -H_1(x, s, q) \quad \text{for } x < b(s, q) \quad (33)$$

$$(\mathbb{L}V_2 - rV_2)(x, s, q) < -H_2(x, s, q) \quad \text{for } x > a(s, q) \quad (34)$$

where the instantaneous-stopping as well as the smooth-fit and normal-reflection conditions of (28)-(30) are satisfied, for each $0 < q < s$. Observe that the superharmonic characterisation of the value function (see, e.g. [Peskir and Shiryaev (2006); Chapter IV, Section 9]) implies that $V_i^*(x, s, q)$, for $i = 1, 2$, are the smallest functions satisfying (26)-(27) with (28) and (31)-(32) with the boundaries $b^*(s, q)$ and $a^*(s, q)$, respectively. Note that the inequalities in (33)-(34) follow directly from the arguments of parts (i) and (ii) of Subsection 2.3 above.

3 Solutions to the Free-boundary Problems

In this section, we obtain closed-form expressions for the value functions $V_i^*(x, s, q)$, for $i = 1, 2$, in (21) and (22) associated with the perpetual real double lookback options on maxima and minima and derive first-order nonlinear ordinary differential equations for the optimal stopping boundaries $b^*(s, q)$ and $a^*(s, q)$ from (24)-(25) as solutions to the free-boundary problems in (26)-(34).

3.1 The Candidate Value Functions

We first observe that the general solutions of the second-order ordinary differential equations in (10) + (26)-(27) have the form:

$$V_i(x, s, q) = C_{i,1}(s, q)x^{\gamma_1} + C_{i,2}(s, q)x^{\gamma_2} - F_i(x, s, q) \quad (35)$$

for every $i = 1, 2$, with the particular solutions:

$$F_1(x, s, q) = (L_1x - q)I(x > g^*(s)) + (s - q + (L_1 - K_1)x)I(x \leq g^*(s)) \quad (36)$$

and

$$F_2(x, s, q) = (s - K_2x)I(x < h^*(q)) + (s - q + (L_2 - K_2)x)I(x \geq h^*(q)) \quad (37)$$

for all $0 < q \leq x \leq s$, respectively. Here, $C_{i,j}(s, q)$, for $i, j = 1, 2$, are some continuously differentiable functions, and the numbers γ_j , $j = 1, 2$, are given by:

$$\gamma_j = \frac{1}{2} - \frac{\mu}{\sigma^2} - (-1)^j \sqrt{\left(\frac{1}{2} - \frac{\mu}{\sigma^2}\right)^2 + \frac{2r}{\sigma^2}} \quad (38)$$

so that $\gamma_2 < 0 < 1 < \gamma_1$, as well as the identity:

$$\frac{\gamma_1}{\gamma_1 - 1} \frac{\gamma_2}{\gamma_2 - 1} = \frac{r}{r - \mu} \tag{39}$$

holds. Then, by applying the conditions from (28)-(30) to the function in (35), we get that the equalities:

$$C_{1,1}(s, q) b^{\gamma_1}(s, q) + C_{1,2}(s, q) b^{\gamma_2}(s, q) = F_1(b(s, q), s, q) \tag{40}$$

$$C_{1,1}(s, q) \gamma_1 b^{\gamma_1}(s, q) + C_{1,2}(s, q) \gamma_2 b^{\gamma_2}(s, q) = \partial_x F_1(b(s, q), s, q) \tag{41}$$

$$C_{2,1}(s, q) a^{\gamma_1}(s, q) + C_{2,2}(s, q) a^{\gamma_2}(s, q) = F_2(a(s, q), s, q) \tag{42}$$

$$C_{2,1}(s, q) \gamma_1 a^{\gamma_1}(s, q) + C_{2,2}(s, q) \gamma_2 a^{\gamma_2}(s, q) = \partial_x F_2(a(s, q), s, q) \tag{43}$$

$$\partial_q C_{1,1}(s, q) q^{\gamma_1} + \partial_q C_{1,2}(s, q) q^{\gamma_2} = 0 \tag{44}$$

$$\partial_s C_{2,1}(s, q) s^{\gamma_1} + \partial_s C_{2,2}(s, q) s^{\gamma_2} = 0 \tag{45}$$

are satisfied, for $0 < q < s$, where the functions $F_i(x, s, q)$, for $i = 1, 2$, are defined in (36)-(37).

Now, by solving the system of equations in (40)-(41), we obtain that the candidate value function admits the representation:

$$V_1(x, s, q; b(s, q)) = C_{1,1}(s, q; b(s, q)) x^{\gamma_1} + C_{1,2}(s, q; b(s, q)) x^{\gamma_2} - F_1(x, s, q) \tag{46}$$

for $q \leq x < b(s, q)$, where

$$C_{1,j}(s, q; b(s, q)) = \frac{\gamma_{3-j}(L_1 b(s, q) - q) - L_1 b(s, q)}{(\gamma_{3-j} - \gamma_j) b^{\gamma_j}(s, q)} \tag{47}$$

for $0 < g^*(s) \leq q < s$, and

$$C_{1,j}(s, q; b(s, q)) = \frac{\gamma_{3-j}(s - q + (L_1 - K_1)b(s, q)) - (L_1 - K_1)b(s, q)}{(\gamma_{3-j} - \gamma_j) b^{\gamma_j}(s, q)} \tag{48}$$

for $0 < q < g^*(s) < s$, and every $j = 1, 2$. Also, by solving the system of equations in (42)-(43), we obtain that the candidate value function admits the representation:

$$V_2(x, s, q; a(s, q)) = C_{2,1}(s, q; a(s, q)) x^{\gamma_1} + C_{2,2}(s, q; a(s, q)) x^{\gamma_2} - F_2(x, s, q) \tag{49}$$

for $a(s, q) < x \leq s$, where

$$C_{2,j}(s, q; a(s, q)) = \frac{\gamma_{3-j}(s - K_2 a(s, q)) + K_2 a(s, q)}{(\gamma_{3-j} - \gamma_j) a^{\gamma_j}(s, q)} \tag{50}$$

for $0 < q < s \leq h^*(q)$, and

$$C_{2,j}(s, q; a(s, q)) = \frac{\gamma_{3-j}(s - q + (L_2 - K_2)a(s, q)) - (L_2 - K_2)a(s, q)}{(\gamma_{3-j} - \gamma_j) a^{\gamma_j}(s, q)} \tag{51}$$

for $0 < q < h^*(q) < s$, and every $j = 1, 2$.

Moreover, by means of straightforward computations, it can be deduced from the expression in (46) with (36) that the first-order and second-order partial derivatives $\partial_x V_1(x, s, q; b(s, q))$ and $\partial_{xx} V_1(x, s, q; b(s, q))$ of the function $V_1(x, s, q; b(s, q))$ take the form:

$$\begin{aligned} \partial_x V_1(x, s, q; b(s, q)) &= C_{1,1}(s, q; b(s, q)) \gamma_1 x^{\gamma_1 - 1} + C_{1,2}(s, q; b(s, q)) \gamma_2 x^{\gamma_2 - 1} \\ &\quad - L_1 I(x > g^*(s)) - (L_1 - K_1) I(x \leq g^*(s)) \end{aligned} \tag{52}$$

and

$$\begin{aligned} \partial_{xx} V_1(x, s, q; b(s, q)) &= C_{1,1}(s, q; b(s, q)) \gamma_1 (\gamma_1 - 1) x^{\gamma_1 - 2} \\ &\quad + C_{1,2}(s, q; b(s, q)) \gamma_2 (\gamma_2 - 1) x^{\gamma_2 - 2} \end{aligned} \tag{53}$$

on the interval $q < x < b(s, q)$, for each $0 < q < s$. Also, by means of straightforward computations, it can be deduced from the expression in (49) with (37) that the first-order and second-order partial derivatives $\partial_x V_2(x, s, q; a(s, q))$ and $\partial_{xx} V_2(x, s, q; a(s, q))$ of the function $V_2(x, s, q; a(s, q))$ take the form:

$$\begin{aligned} \partial_x V_2(x, s, q; a(s, q)) &= C_{i,1}(s, q; a(s, q)) \gamma_1 x^{\gamma_1 - 1} + C_{i,2}(s, q; a(s, q)) \gamma_2 x^{\gamma_2 - 1} \\ &\quad + K_2 I(x < h^*(q)) - (L_2 - K_2) I(x \geq h^*(q)) \end{aligned} \tag{54}$$

and

$$\begin{aligned} \partial_{xx} V_2(x, s, q; a(s, q)) &= C_{i,1}(s, q; a(s, q)) \gamma_1 (\gamma_1 - 1) x^{\gamma_1 - 2} \\ &\quad + C_{i,2}(s, q; a(s, q)) \gamma_2 (\gamma_2 - 1) x^{\gamma_2 - 2} \end{aligned} \tag{55}$$

on the interval $a(s, q) < x < s$, for each $0 < q < s$.

3.2 The Candidate Stopping Boundaries

In order to derive first-order nonlinear ordinary differential equations for the candidate boundary functions, we further assume that the functions $b(s, q)$ and $a(s, q)$ are continuously differentiable. Then, applying the condition of (44) to the functions $C_{1,j}(s, q; b(s, q))$, for $j = 1, 2$, in (47)-(48), we obtain the equalities:

$$\partial_q b(s, q) = \frac{\gamma_2 (q/b(s, q))^{\gamma_1} - \gamma_1 (q/b(s, q))^{\gamma_2}}{\gamma_1 \gamma_2 (q/b(s, q)) - (r - \mu) L_1 / r ((q/b(s, q))^{\gamma_1} - (q/b(s, q))^{\gamma_2})} \tag{56}$$

for $0 < g^*(s) \leq q < s$, and

$$\partial_q b(s, q) = \frac{\gamma_2 (q/b(s, q))^{\gamma_1} - \gamma_1 (q/b(s, q))^{\gamma_2}}{\gamma_1 \gamma_2 ((s - q)/b(s, q) + (r - \mu) (L_1 - K_1) / r) ((q/b(s, q))^{\gamma_1} - (q/b(s, q))^{\gamma_2})} \tag{57}$$

for $0 < q < g^*(s) < s$. Here, by virtue of the structure of the equation in (56), we have $b(s, q) = v_* q$ with $v_* > 1$ from (84), for all $q > g^*(s)$ and each $s > 0$. Note that the candidate value function $V_1(x, s, q; b(s, q))$ in (46) with (47)-(48) is (strictly) increasing in $b(s, q)$, so that we should take the candidate stopping boundary $b(s, q)$, for $i = 1, 2$, as the *minimal* solution of the first-order nonlinear ordinary differential equation in (57) located above the plane $d_1 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < x = q < s\}$.

Hence, applying the condition of (45) to the functions $C_{2,j}(s, q; a(s, q))$, for $j = 1, 2$, in (50)-(51), we obtain the equalities:

$$\partial_s a(s, q) = \frac{\gamma_2(s/a(s, q))^{\gamma_1} - \gamma_1(s/a(s, q))^{\gamma_2}}{\gamma_1 \gamma_2 (s/a(s, q) - (r - \mu)K_2/r)((s/a(s, q))^{\gamma_1} - (s/a(s, q))^{\gamma_2})} \tag{58}$$

for $0 < q < s \leq h^*(q)$, and

$$\partial_s a(s, q) = \frac{\gamma_2(s/a(s, q))^{\gamma_1} - \gamma_1(s/a(s, q))^{\gamma_2}}{\gamma_1 \gamma_2 ((s - q)/b(s, q) + (r - \mu)(L_2 - K_2)/r)((s/a(s, q))^{\gamma_1} - (s/a(s, q))^{\gamma_2})} \tag{59}$$

for $0 < q < h^*(q) < s$. Here, by virtue of the structure of the equation in (58), we have $a(s, q) = \lambda_* s$ with $0 < \lambda_* < 1$ from (82), for all $0 < s < h^*(q)$ and each $q > 0$. Note that the candidate value function $V_2(x, s, q; a(s, q))$ in (46) with (47)-(48) is (strictly) decreasing in $a(s, q)$, so that we should take the candidate stopping boundary $a(s, q)$, for $i = 1, 2$, as the maximal solution of the first-order nonlinear ordinary differential equation in (59) located below the plane $d_2 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q < x = s\}$.

3.3 The Minimal and Maximal Admissible Solutions $b^*(s, q)$ and $a^*(s, q)$.

We further consider the *minimal* and *maximal admissible* solutions of first-order nonlinear ordinary differential equations as the smallest and largest possible solutions $b^*(s, q)$ and $a^*(s, q)$ of the equations in (57) and (59), which satisfy the inequalities $0 < q < h^*(q) \leq b^*(s, q) \leq s$, and $0 < q \leq a^*(s, q) \leq g^*(s) < s$, for all $0 < q < g^*(s) < s$ and $0 < q < h^*(q) < s$. By virtue of the classical results on the existence and uniqueness of solutions for first-order nonlinear ordinary differential equations, we may conclude that these equations admit (locally) unique solutions, because of the facts that their right-hand sides represent (locally) continuous functions in $(s, q, b(s, q))$ and $(s, q, a(s, q))$ and (locally) Lipschitz functions in $b(s, q)$ and $a(s, q)$, for each $0 < q < g^*(s) < s$ and $0 < q < h^*(q) < s$ fixed (see also [Peskir (1998); Subsection 3.9] for similar arguments based on the analysis of other first-order nonlinear ordinary differential equations). Then, it is shown by means of technical arguments based on Picard’s method of successive approximations that there exist unique solutions $b(s, q)$ and $a(s, q)$ to the equations in (57) and (59) started at some points (q_0, s, q_0) and (s_0, s_0, q) , for each $0 < q_0 < g^*(s) < s$ and $0 < q < h^*(q) < s_0$ fixed (see also [Graversen and Peskir (1998); Subsection 3.2] and [Peskir (1998); Example 4.4] for similar arguments based on the analysis of other first-order nonlinear ordinary differential equations).

Hence, in order to construct the appropriate functions $b^*(s, q)$ and $a^*(s, q)$ which satisfy the equations in (56) and (58) and stays strictly above and below the appropriate diagonals $d_1 = \{(x, s, q) \in E \mid 0 < x = q < s\}$ or $d_2 = \{(x, s, q) \in E \mid 0 < q < x = s\}$, respectively, we construct the sequences of solutions satisfying such properties and intersecting d_1 and d_2 (see also [Peskir (2014); Subsection 3.5] (among others) for a similar procedure applied for solutions of other first-order nonlinear ordinary differential equations). For this purpose, for any decreasing and increasing sequences $(q_l)_{l \in \mathbb{N}}$ and $(s_l)_{l \in \mathbb{N}}$, such that $0 < q_l < g^*(s) < s$ and $0 < q < h^*(q) < s_l$, we can construct the sequences of solutions $b_l(s, q)$ and $a_l(s, q)$, for $l \in \mathbb{N}$, to the equations in (57) and (59), for all $0 < q < q_l$ and $s > s_l$ such that $b_l(s, q_l) = q_l$ and $a_l(s_l, q) = s_l$ holds, for each $0 < q_l < g^*(s) < s$ and $0 < q < h^*(q) < s_l$, and every $l \in \mathbb{N}$. It follows from the structure of the equations in (57) and (59) that the inequalities $\partial_q b_l(s, q_l) < 1$ and $\partial_s a_l(s_l, q) < 1$ should hold for the derivatives of the corresponding

functions, for each $0 < q_l < g^*(s) < s$ and $0 < q < h^*(q) < s_l$, and every $l \in \mathbb{N}$ (see also [Pedersen (2000); pages 979-982] for the analysis of solutions of another first-order nonlinear differential equation). Observe that, by virtue of the uniqueness of solutions mentioned above, we know that each two curves $q \mapsto b_l(s, q)$ and $q \mapsto b_m(s, q)$ as well as $s \mapsto a_l(s, q)$ and $s \mapsto a_m(s, q)$ cannot intersect, for each $0 < q < g^*(s) < s$ and $0 < q < h^*(q) < s$, and $l, m \in \mathbb{N}$, such that $l \neq m$, and thus, we see that the sequence $(b_l(s, q))_{l \in \mathbb{N}}$ is decreasing and the sequence $(a_l(s, q))_{l \in \mathbb{N}}$ is increasing, so that the limits $b^*(s, q) = \lim_{l \rightarrow \infty} b_l(s, q)$ and $a^*(s, q) = \lim_{l \rightarrow \infty} a_l(s, q)$ exist, for each $0 < q < s$, respectively. We may therefore conclude that $b^*(s, q)$ and $a^*(s, q)$ provides the minimal and maximal solutions to the equations in (57) and (59) such that $b^*(s, q) > q$ and $a^*(s, q) < s$ holds, for all $0 < q < g^*(s) < s$ and $0 < q < h^*(q) < s$.

Moreover, since the right-hand sides of the first-order nonlinear ordinary differential equations in (57) and (59) are (locally) Lipschitz in (s, q) , respectively, one can deduce by means of Gronwall's inequality that the functions $b_l(s, q)$ and $a_l(s, q)$, for each $l \in \mathbb{N}$, are continuous, so that the functions $b^*(s, q)$ and $a^*(s, q)$ are continuous too, for $0 < q < g^*(s) < s$ and $0 < q < h^*(q) < s$. The appropriate maximal admissible solutions of first-order nonlinear ordinary differential equations and the associated maximality principle for solutions of optimal stopping problems which is equivalent to the superharmonic characterisation of the payoff functions were established in Peskir (1998) and further developed in Graversen and Peskir (1998), Pedersen (2000), Guo and Shepp (2001), Gapeev (2007), Guo and Zervos (2010), Peskir (2012, 2014), Glover et al. (2013), Ott (2013), Kyprianou and Ott (2014), Gapeev and Rodosthenous (2014b, 2016a 2016b), Rodosthenous and Zervos (2017), and Gapeev et al. (2021) among other subsequent papers (see also [Peskir and Shiryaev (2006); Chapter I; Chapter V, Section 17] for other references).

4 Main Results and Proofs

In this section, based on the facts proved above, we formulate and prove the main result of the paper. Observe that, by means of the change-of-measure arguments from Shepp and Shiryaev (1994) and Gapeev (2019), the problems of (6)-(7) can be reduced the appropriate optimal stopping problems for the two-dimensional Markov process $(S/X, Q/X) = (S_t/X_t, Q_t/X_t)_{t \geq 0}$. However, we follow the classical approach initiated in Shepp and Shiryaev (1993) and to solve them as three-dimensional optimal stopping problems.

Theorem 4.1 *Let the process (X, S, Q) be given by (3)-(4) and (5) with $\sigma > 0$, $\mu < r$, and $r > 0$. Then, the value functions of the optimal stopping problems in (21) and (22), for some $L_i \geq 1 \geq K_i > 0$, for $i = 1, 2$, fixed, admit the representations:*

$$V_1^*(x, s, q) = \begin{cases} V_1(x, s, q; b^*(s, q)), & \text{if } q \leq x < b^*(s, q), \\ 0, & \text{if } x \geq b^*(s, q), \end{cases} \tag{60}$$

and

$$V_2^*(x, s, q) = \begin{cases} V_2(x, s, q; a^*(s, q)), & \text{if } a^*(s, q) < x \leq s, \\ 0, & \text{if } 0 < x \leq a^*(s, q), \end{cases} \tag{61}$$

while the optimal stopping times have the form:

$$\tau_1^* = \inf \{t \geq 0 \mid X_t \geq b^*(S_t, Q_t)\} \quad \text{and} \quad \tau_2^* = \inf \{t \geq 0 \mid X_t \leq a^*(S_t, Q_t)\} \quad (62)$$

where the candidate value functions and boundaries are specified as follows:

- (i) the function $V_1(x, s, q; b^*(s, q))$ is given by (46) with (47)-(48), where the boundary $b^*(s, q)$ satisfying the inequality $b^*(s, q) \geq h^*(q)$ represents the minimal solution of the first-order nonlinear ordinary differential equation in (57) such that $b^*(s, q) > q$, for $0 < q < g^*(s) < s$, while $b^*(s, q) = h^*(q) \equiv v_* q$ with $v_* > 1$ from (84), for $g^*(s) \leq q < s$;
- (ii) the function $V_2(x, s, q; a^*(s, q))$ is given by (49) with (50)-(51), where the boundary $a^*(s, q)$ satisfying the inequality $a^*(s, q) \leq g^*(s)$ represents the maximal solution of the first-order nonlinear ordinary differential equation in (59) such that $a^*(s, q) < s$, for $0 < q < h^*(q) < s$, while $a^*(s, q) = g^*(s) \equiv \lambda_* s$ with $0 < \lambda_* < 1$ from (82), for $q < s \leq h^*(q)$.

Recall that we can put $s = q = x$ to obtain the values of the original perpetual real floating-cost double lookback call-put and put-call option pricing problems of (1) and (2) from the values of the double optimal stopping problems of (6) and (7), which are equivalent to the sequence of single optimal stopping problems of (21)-(22) and (72). Note that, since the both parts of the assertion stated above are proved using similar arguments, we may only give a proof for the case of the three-dimensional single optimal stopping problem of (22), which is related to the outer perpetual real lookback put-call options.

Proof In order to verify the assertion of part (ii) stated above, it remains for us to show that the function defined in the right-hand side of (61) coincides with the value function in (22) and that the stopping time τ_2^* in (62) is optimal with the boundary $a^*(s, q)$ being the solution of the system in (42)-(43)+(45) specified in (49)-(51) with (58)-(59). For this purpose, let us denote by $V_2(x, s, q)$ the right-hand side of the expression in (61) associated with $a^*(s, q)$. Then, it is shown by means of straightforward calculations from the previous section that the function $V_2(x, s, q)$ solves the left-hand system of (26)-(34). Recall that the function $\overline{V}_2(x, s, q)$ is $C^{2,1,1}$ on the closure \overline{C}_2 of C_2 and is equal to 0 on D_2 , which are defined as \overline{C}_2^* , C_2^* and D_2^* in (24) and (25) with $a(s, q)$ instead of $a^*(s, q)$, respectively. Hence, taking into account the assumption that the boundary $a^*(s, q)$ is (at least piecewise) continuously differentiable, for all $0 < q < s$, by applying the change-of-variable formula from [Peskir (2007); Theorem 3.1] to the process $e^{-rt}V_2(X_t, S_t, Q_t)$ (see also [Peskir and Shiryaev (2006); Chapter II, Section 3.5] for a summary of the related results and further references), we obtain the expression:

$$\begin{aligned} e^{-rt} V_2(X_t, S_t, Q_t) &= V_2(x, s, q) + M_t^2 \\ &+ \int_0^t e^{-ru} (\mathbb{L}V_2 - rV_2)(X_u, S_u, Q_u) I(Q_u \vee a^*(S_u, Q_u) < X_u < S_u) du \\ &+ \int_0^t e^{-ru} \partial_s V_2(X_u, S_u, Q_u) I(X_u = S_u) dS_u \\ &+ \int_0^t e^{-ru} \partial_q V_2(X_u, S_u, Q_u) I(X_u = Q_u) dQ_u \end{aligned} \quad (63)$$

for all $t \geq 0$. Here, the process $M^2 = (M_t^2)_{t \geq 0}$ defined by:

$$M_t^2 = \int_0^t e^{-ru} \partial_x V_2(X_u, S_u, Q_u) I(Q_u < X_u < S_u) \sigma X_u dB_u \tag{64}$$

is a continuous local martingale with respect to the probability measure $P_{x,s,q}$. Note that, since the time spent by the process (X, S, Q) at the boundary surface $\{(x, s, q) \in E \mid x = a(s, q)\}$ as well as at the diagonals $d_1 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < x = q \leq s\}$ and $d_2 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q \leq x = s\}$ is of zero Lebesgue measure (see, e.g. [Borodin and Salminen (2002); Chapter II, Section 1]), the indicators in the second line of the formula in (63) as well as in the expression of (64) can be ignored. Moreover, since the component Q decreases only when the process (X, S, Q) is located on the diagonal $d_1 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < x = q < s\}$, while the component S increases only when the process (X, S, Q) is located on the diagonal $d_2 = \{(x, s, q) \in \mathbb{R}^3 \mid 0 < q < x = s\}$, the indicators appearing in the third line of (63) can be set equal to one. Finally, we observe from the expressions in (49) and (50)-(51) that the function $V_2(x, s, q)$ does not actually depend on the variable q , and thus, the partial derivative $\partial_q V_2(x, s, q)$ is equal to 0 in the region $\{(x, s, q) \in E \mid 0 < q < x \leq s \leq h^*(q)\}$. Therefore, since the diagonal d_1 lies outside to the region $\{(x, s, q) \in E \mid 0 < q < h^*(q) \leq x \leq s\}$, we may conclude that the second integral in the third line of (63) is actually equal to zero.

It follows from straightforward calculations and the arguments of the previous section that the function $V_2(x, s, q)$ satisfies the second-order ordinary differential equation in (27), which together with the left-hand conditions of (28)-(29) and (31) as well as the fact that the left-hand inequality in (34) holds imply that the inequality $(\mathbb{L}V_2 - rV_2)(x, s, q) \leq -H_2(x, s, q)$ is satisfied, for all $(x, s, q) \in E$ such that $0 < q < x < s$ and $x \neq a^*(s, q)$. Moreover, we observe directly from the expressions in (49) with (50)-(51) as well as (54)-(55) that the value function $V_2(x, s, q)$ is convex and increases from zero, because its first-order partial derivative $\partial_x V_2(x, s, q)$ is positive and increases from zero, while its second-order partial derivative $\partial_{xx} V_2(x, s, q)$ is positive, on the interval $q \vee a^*(s, q) < x \leq s$. Thus, we may conclude that the left-hand inequality in (32) holds, which together with the left-hand conditions of (28)-(29) and (31) imply that the inequality $V_2(x, s, q) \geq 0$ is satisfied, for all $(x, s, q) \in E$. Let $(\tau_n)_{n \in \mathbb{N}}$ be the localising sequence of stopping times for the process M^2 from (64) such that $\tau_n = \inf\{t \geq 0 \mid |M_t^2| \geq n\}$, for each $n \in \mathbb{N}$. It therefore follows from the expression in (63) that the inequalities:

$$\begin{aligned} \int_0^{\tau \wedge \tau_n} e^{-ru} H_2(X_u, S_u, Q_u) du + \int_0^{\tau \wedge \tau_n} e^{-ru} dS_u &\leq e^{-r(\tau \wedge \tau_n)} V_2(X_{\tau \wedge \tau_n}, S_{\tau \wedge \tau_n}, Q_{\tau \wedge \tau_n}) \\ &+ \int_0^{\tau \wedge \tau_n} e^{-ru} H_2(X_u, S_u, Q_u) du \\ &+ \int_0^{\tau \wedge \tau_n} e^{-ru} dS_u \leq V_2(x, s, q) + M_{\tau \wedge \tau_n}^2 \end{aligned} \tag{65}$$

hold, for any stopping time τ with respect to the natural filtration of (X, S, Q) and each $n \in \mathbb{N}$ fixed. Then, taking the expectation with respect to $P_{x,s,q}$ in (65), by means of Doob's optional sampling theorem, we get:

$$\begin{aligned}
 E_{x,s,q} \left[\int_0^{\tau \wedge X_n} e^{-ru} H_2(X_u, S_u, Q_u) du + \int_0^{\tau \wedge X_n} e^{-ru} dS_u \right] \\
 \leq E_{x,s,q} \left[e^{-r(\tau \wedge X_n)} V_2(X_{\tau \wedge X_n}, S_{\tau \wedge X_n}, Q_{\tau \wedge X_n}) \right. \\
 \left. + \int_0^{\tau \wedge X_n} e^{-ru} H_2(X_u, S_u, Q_u) du + \int_0^{\tau \wedge X_n} e^{-ru} dS_u \right] \\
 \leq V_2(x, s, q) + E_{x,s,q} [M_{\tau \wedge X_n}^2] = V_2(x, s, q)
 \end{aligned} \tag{66}$$

for all $(x, s, q) \in E$ and each $n \in \mathbb{N}$. Hence, letting n go to infinity and using Fatou’s lemma, we obtain from the expressions in (66) that the inequalities:

$$\begin{aligned}
 E_{x,s,q} \left[\int_0^{\tau} e^{-ru} H_2(X_u, S_u, Q_u) du + \int_0^{\tau} e^{-ru} dS_u \right] \leq E_{x,s,q} \left[e^{-r\tau} V_2(X_\tau, S_\tau, Q_\tau) \right. \\
 \left. + \int_0^{\tau} e^{-ru} H_2(X_u, S_u, Q_u) du \right] \\
 \left. + \int_0^{\tau} e^{-ru} dS_u \right] \leq V_2(x, s, q)
 \end{aligned} \tag{67}$$

hold, for any stopping time τ and all $(x, s, q) \in E$.

We now prove the fact that the boundary $a^*(s, q)$ specified above is optimal. By virtue of the fact that the function $V_2(x, s, q)$ from the right-hand side of the expression in (61) associated with the boundary $a^*(s, q)$ satisfies the equation of (27) and the right-hand condition of (28), and taking into account the structure of τ_2^* in (62), it follows from the expression in (63) that the equalities:

$$\begin{aligned}
 E_{x,s,q} \left[\int_0^{\tau_2^* \wedge X_n} e^{-ru} H_2(X_u, S_u, Q_u) du + \int_0^{\tau_2^* \wedge X_n} e^{-ru} dS_u \right] = E_{x,s,q} \left[e^{-r(\tau_2^* \wedge X_n)} V_2(X_{\tau_2^* \wedge X_n}, S_{\tau_2^* \wedge X_n}, Q_{\tau_2^* \wedge X_n}) \right. \\
 \left. + \int_0^{\tau_2^* \wedge X_n} e^{-ru} H_2(X_u, S_u, Q_u) du \right. \\
 \left. + \int_0^{\tau_2^* \wedge X_n} e^{-ru} dS_u \right] = V_2(x, s, q) + E_{x,s,q} [M_{\tau_2^* \wedge X_n}^2] = V_2(x, s, q)
 \end{aligned} \tag{68}$$

hold, for all $(x, s, q) \in E$ and each $n \in \mathbb{N}$. Observe that, by virtue of the arguments from [Shepp and Shiryaev (1993); pages 635–636], the property:

$$E_{x,s,q} \left[\sup_{t \geq 0} e^{-r(\tau_2^* \wedge t)} G_2(X_{\tau_2^* \wedge t}, S_{\tau_2^* \wedge t}, Q_{\tau_2^* \wedge t}) \right] \leq (1 + L_2) E_{x,s,q} \left[\sup_{t \geq 0} S_{\tau_2^* \wedge t} \right] < \infty \tag{69}$$

holds, where the function $G_2(x, s, q)$ is defined in (9), for all $(x, s, q) \in E$. We also note that the variable $e^{-r\tau_2^*} V_2(X_{\tau_2^*}, S_{\tau_2^*}, Q_{\tau_2^*})$ is finite on the event $\{\tau_2^* = \infty\}$ as well as recall from the arguments of Beibel and Lerche (1997) and Pedersen (2000) that the property $P_{x,s,q}(\tau_2^* < \infty) = 1$ holds, for all $(x, s, q) \in E$. Hence, letting n go to infinity and using the right-hand condition of (28), we can apply the Lebesgue dominated convergence theorem to the expression of (68) to obtain the equality:

$$E_{x,s,q} \left[\int_0^{\tau_2^*} e^{-ru} H_2(X_u, S_u, Q_u) du + \int_0^{\tau_2^*} e^{-ru} dS_u \right] = V_2(x, s, q) \tag{70}$$

for all $(x, s, q) \in E$, which together with the inequalities in (67) directly implies the desired assertion. We finally recall from the results of part (iii) of Subsection 2.3 above implied by standard comparison arguments applied to the value functions of the appropriate optimal stopping problems that the inequality $a^*(s, q) \leq g^*(s)$ should hold for the optimal stopping boundary, for $0 < q < h^*(q) < s$. Thus, taking into account the fact that $a^*(s, q) = g^*(s) \equiv \lambda_* s$ with $0 < \lambda_* < 1$ from (82), for $q < s \leq h^*(q)$, we may conclude that the inequality $a^*(s, q) \leq g^*(s)$ holds, for all $0 < q < s$, that completes the verification. □

Corollary 4.2 *The optimal method of exercising the perpetual real double lookback call-put and put-call options on the maxima and minima with the values in (1) and (2), which are equivalent to the ones of (6) and (7), acts as follows. After the outer options with the equivalent value functions from (21) and (22) are exercised at the first exit times τ_i^* , for $i = 1, 2$, from (62) with the boundaries $b^*(s, q)$ and $a^*(s, q)$ specified in Theorem 4.1 above, the inner options should be exercised at the first hitting times:*

$$\zeta_1^* = \inf \{t \geq \tau_1^* \mid X_t \leq g^*(S_t)\} \quad \text{and} \quad \zeta_2^* = \inf \{t \geq \tau_2^* \mid X_t \geq h^*(Q_t)\} \tag{71}$$

with the boundaries $g^*(s)$ and $h^*(q)$ specified in Corollary 5.1 below, respectively.

Remark 4.3 Note that in the cases in which one starts from the stretch, that is, when $x = s = q$ holds, the subsequent exercise of the *outer* and *inner* perpetual real lookback put and call options with the value functions in (21) and (22) may actually follow the subsequent exercise of the *standard* perpetual real lookback put and call options with the value functions in (72). More precisely, when the process X starts at some $x = s = q$, by virtue of the facts that the inequalities $0 < \lambda_* < 1$ and $\nu_* > 1$ hold for the unique solutions of the arithmetic equations in (82) and (84) below, the *outer* options *should be exercised* when the process X reaches an upper boundary $b^*(S, Q) \geq h^*(Q)$ or a lower boundary $a^*(S, Q) \leq g^*(S)$, respectively. However, in the cases in which the process X starts at some $x < g^*(s) < s$ or $x > h^*(q) > q$, the *outer* perpetual real lookback call-put and put-call options on the maxima and minima *should be exercised only* at the times at which the underlying asset price process reaches the upper boundary $[h^*(Q) <] b^*(S, Q) [< g^*(S)]$ or the lower boundary $[h^*(Q) <] a^*(S, Q) [< g^*(S)]$, respectively, and then, the appropriate *inner* options should be exercised at the same time.

5 Appendix

In this section, we derive explicit solutions to the optimal stopping problems of (72) below which are related to the perpetual American lookback options on the maximum and minimum. Although, by means of the change-of-measure arguments from Shepp and Shiryaev (1994) and Gapeev (2019), the problems of (72) can be reduced to the appropriate optimal stopping problems for one-dimensional Markov processes $S/X = (S_t/X_t)_{t \geq 0}$ and $Q/X = (Q_t/X_t)_{t \geq 0}$, we follow the arguments of Sects. 2–4 to solve them as two-dimensional optimal stopping problems, in order to spare some space of the paper. Note that the left-hand

optimal stopping problem in (72) was solved in Beibel and Lerche (1997), Pedersen (2000) and Guo and Shepp (2001), and we present its solution for completeness.

5.1 The Auxiliary Optimal Stopping and Free-boundary Problems

Let us finally consider the optimal stopping problems for the (time-homogeneous strong) Markov processes $(X, S) = (X_t, S_t)_{t \geq 0}$ and $(X, Q) = (X_t, Q_t)_{t \geq 0}$ with the value functions:

$$U_1^*(x, s) = \sup_{\eta} E_{x,s} [e^{-r\eta} (S_{\eta} - K_1 X_{\eta})] \quad \text{and} \quad U_2^*(x, q) = \sup_{\eta} E_{x,q} [e^{-r\eta} (L_2 X_{\eta} - Q_{\eta})] \tag{72}$$

for some given constants $L_2 \geq 1 \geq K_1 > 0$, where the suprema are taken over all stopping times η of (X, S) or (X, Q) . It can be shown by means of the same arguments as in Subsection 2.2 above that the optimal stopping times have the form:

$$\eta_1^* = \inf \{t \geq 0 \mid X_t \leq g^*(S_t)\} \quad \text{and} \quad \eta_2^* = \inf \{t \geq 0 \mid X_t \geq h^*(Q_t)\} \tag{73}$$

for some boundaries $0 < g^*(s) < s$ and $h^*(q) > q$ to be determined.

In order to find analytic expressions for the value functions $U_1^*(x, s)$ and $U_2^*(x, q)$ from (72) as well as the unknown boundaries boundaries $g^*(s)$ and $h^*(q)$ from (73), we formulate the equivalent free-boundary problems:

$$(\mathbb{L}U_1 - rU_1)(x, s) = 0 \quad \text{for } g(s) < x < s, \quad (\mathbb{L}U_2 - rU_2)(x, q) = 0 \quad \text{for } q < x < h(q) \tag{74}$$

$$U_1(x, s)|_{x=g(s)+} = s - K_1 g(s), \quad U_2(x, q)|_{x=h(q)-} = L_2 h(q) - q \tag{75}$$

$$\partial_x U_1(x, s)|_{x=g(s)+} = -K_1, \quad \partial_x U_2(x, q)|_{x=h(q)-} = L_2 \tag{76}$$

$$\partial_s U_1(x, s)|_{x=s-} = 0, \quad \partial_q U_2(x, q)|_{x=q+} = 0 \tag{77}$$

$$U_1(x, s) = s - K_1 x \quad \text{for } 0 < x \leq g(s), \quad U_2(x, q) = L_2 x - q \quad \text{for } x \geq h(q) \tag{78}$$

$$U_1(x, s) > s - K_1 x \quad \text{for } g(s) < x \leq s, \quad U_2(x, q) > L_2 x - q \quad \text{for } q \leq x < h(q) \tag{79}$$

$$(\mathbb{L}U_1 - rU_1)(x, s) < 0 \quad \text{for } 0 < x \leq g(s), \quad (\mathbb{L}U_2 - rU_2)(x, q) < 0 \quad \text{for } x \geq h(q) \tag{80}$$

where the left-hand and the right-hand conditions in (75)-(77) are satisfied for $s > 0$ and $q > 0$, respectively. The superharmonic characterisation of the value function implies that $U_1^*(x, s)$ and $U_2^*(x, q)$ are the smallest functions satisfying the left-hand and the right-hand sides of the equations in (74)-(75) with (78)-(79) with the boundaries $g^*(s)$ and $h^*(q)$, respectively.

5.2 Solutions to the Free-boundary Problems

In this case, by using straightforward calculations from Subsections 3.1-3.2 above, it can be shown that the candidate solution of the left-hand system in (74)-(80) takes the form:

$$U_1(x, s; g^*(s)) = \frac{\gamma_2(s - K_1 g^*(s)) + K_1 g^*(s)}{\gamma_2 - \gamma_1} \left(\frac{x}{g^*(s)}\right)^{\gamma_1} - \frac{\gamma_1(s - K_1 g^*(s)) + K_1 g^*(s)}{\gamma_2 - \gamma_1} \left(\frac{x}{g^*(s)}\right)^{\gamma_2} \tag{81}$$

for $0 < g^*(s) < x \leq s$, with $g^*(s) \equiv \lambda_* s$, for $s > 0$, where the value $0 < \lambda_* < 1$ is the unique root of the arithmetic equation:

$$\lambda^{\gamma_1 - \gamma_2} = \frac{(\gamma_1 - 1)(\gamma_2(1 - K_1 \lambda) + K_1 \lambda)}{(\gamma_2 - 1)(\gamma_1(1 - K_1 \lambda) + K_1 \lambda)} \tag{82}$$

on the interval $(0, 1)$ (see Beibel and Lerche (1997), Pedersen (2000) and Guo and Shepp (2001)).

By using straightforward calculations from Subsections 3.1–3.2 above, it can be shown that the candidate solution of the right-hand system in (74)–(80) takes the form:

$$U_2(x, q; h^*(q)) = \frac{\gamma_2(L_2 h^*(q) - q) - L_2 h^*(q)}{\gamma_2 - \gamma_1} \left(\frac{x}{h^*(q)}\right)^{\gamma_1} - \frac{\gamma_1(L_2 h^*(q) - q) - L_2 h^*(q)}{\gamma_2 - \gamma_1} \left(\frac{x}{h^*(q)}\right)^{\gamma_2} \tag{83}$$

for $q \leq x < h^*(q)$, with $h^*(q) \equiv v_* q$, for $q > 0$, where the value $v_* > 1$ is the unique root of the arithmetic equation:

$$v^{\gamma_1 - \gamma_2} = \frac{(\gamma_1 - 1)(\gamma_2(1 - L_2 v) + L_2 v)}{(\gamma_2 - 1)(\gamma_1(1 - L_2 v) + L_2 v)}. \tag{84}$$

on the interval $(1, \infty)$.

5.3 The Results

Summarising the facts shown above, we state the following result which can be proved by means of the same arguments as Theorem 4.1 above in combinations with the arguments from Gapeev (2020).

Corollary 5.1 *Let the processes (X, S) and (X, Q) be given by (3) and (5) with $\sigma > 0$, $\mu < r$, and $r > 0$. Then, the value functions of the inner optimal stopping problems in (72), for some $L_2 \geq 1 \geq K_1 > 0$ fixed, admit the representations:*

$$U_1^*(x, s) = \begin{cases} U_1(x, s; g^*(s)), & \text{if } g^*(s) < x \leq s, \\ s - K_1 x, & \text{if } 0 < x \leq g^*(s), \end{cases} \tag{85}$$

and

$$U_2^*(x, q) = \begin{cases} U_2(x, q; h^*(q)), & \text{if } 0 < q \leq x < h^*(q), \\ L_2 x - q, & \text{if } x \geq h^*(q), \end{cases} \tag{86}$$

while the optimal stopping times have the form of (73) above, where the candidate value functions and the candidate exercise boundaries are specified as follows:

- (i) *the function $U_1(x, s; g^*(s))$ is given by (81), while the boundary has the form $g^*(s) = \lambda_* s$, for each $s > 0$, with $0 < \lambda_* < 1$ being a unique solution of the arithmetic equation in (82) on $(0, 1)$;*
- (ii) *the function $U_2(x, q; h^*(q))$ is given by (83), while the boundary has the form $h^*(q) = \nu_* q$, for each $q > 0$, with $\nu_* > 1$ being a unique solution of the arithmetic equation in (84) on $(1, \infty)$.*

Acknowledgements The authors are grateful to the Editor and two anonymous Referees for their valuable suggestions which helped to improve the presentation of the paper.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Asmussen S, Avram F, Pistorius M (2003) Russian and American put options under exponential phase-type Lévy models. *Stochastic Processes and their Applications* 109:79–111
- Avram F, Kyprianou AE, Pistorius M (2004) Exit problems for spectrally negative Lévy processes and applications to (Canadized) Russian options. *Ann Appl Probab* 14(1):215–238
- Beibel M, Lerche HR (1997) A new look at warrant pricing and related optimal stopping problems. Empirical Bayes, sequential analysis and related topics in statistics and probability (New Brunswick, NJ, 1995). *Statistica Sinica* 7:93–108
- Borodin AN, Salminen P (2002) *Handbook of Brownian Motion*, 2nd edn. Birkhäuser, Basel
- Brynjolfsson E, Hitt L (2003) Computing productivity: firm-level evidence. *Rev Econ Stat* 85(4):793–808
- Brynjolfsson E, Rock D, Syverson C (2017) Artificial intelligence and the modern productivity paradox: a clash of expectations and statistics. The National Bureau of Economic Research. <https://doi.org/10.3386/w24001>
- Carmona R, Dayanik S (2008) Optimal multiple-stopping of linear diffusions. *Math Oper Res* 33(2):446–460
- Carmona R, Touzi N (2008) Optimal multiple stopping and valuation of swing options. *Math Finance* 18(2):239–268
- Chiarella C, Kang B (2009) The evaluation of American compound option prices under stochastic volatility using the sparse grid approach. Research Centre Research Paper No. 245, February, 2009, University of Technology, Sydney, Quantitative Finance
- Dixit AK, Pindyck RS (1994) *Investment under uncertainty*. Princeton University Press, New Jersey
- Dubins L, Shepp LA, Shiryaev AN (1993) Optimal stopping rules and maximal inequalities for Bessel processes. *Theory of Probability and its Applications* 38(2):226–261
- Duckworth JK, Zervos M (2000) An investment model with entry and exit decisions. *J Appl Probab* 37:547–559
- Gapeev PV (2007) Discounted optimal stopping for maxima of some jump-diffusion processes. *J Appl Probab* 44:713–731
- Gapeev PV (2019) Solving the dual Russian option problem by using change-of-measure arguments. *High Frequency (Special Issue in Memory of Larry Shepp)* 2(2):76–84
- Gapeev PV (2020) Optimal stopping problems for running minima with positive discounting rates. *Statist Probab Lett* 167:108899
- Gapeev PV (2022) Perpetual American double lookback options on drawdowns and drawups with floating strikes. To appear in *Methodology and Computing in Applied Probability*. pp 40
- Gapeev PV, Kort PM, Lavrutich MN (2021) Discounted optimal stopping problems for maxima of geometric Brownian motions with switching payoffs. *Adv Appl Probab* 53(1):189–219

- Gapeev PV, Li L (2021) Optimal stopping problems for maxima and minima in models with asymmetric information. Published online first in *Stochastics*. pp 28
- Gapeev PV, Al Motairi H (2021) Discounted optimal stopping problems in first-passage time models with random thresholds. To appear in *Journal of Applied Probability*. pp 20
- Gapeev PV, Rodosthenous N (2014a) On the pricing of perpetual American compound options. In: Kabanov, Yu M, Rutkowski M, Zariphopoulou TH (eds) *In the Volume Inspired by Finance - The Musiela Festschrift*. Springer, pp 283–304
- Gapeev PV, Rodosthenous N (2014b) Optimal stopping problems in diffusion-type models with running maxima and drawdowns. *J Appl Probab* 51(3):799–817
- Gapeev PV, Rodosthenous N (2016a) On the drawdowns and drawups in diffusion-type models with running maxima and minima. *J Math Anal Appl* 434(1):413–431
- Gapeev PV, Rodosthenous N (2016a) Perpetual American options in diffusion-type models with running maxima and drawdowns. *Stochastic Processes and their Applications* 126(7):2038–2061
- Graversen SE, Peskir G (1998) Optimal stopping and maximal inequalities for geometric Brownian motion. *J Appl Probab* 35(4):856–872
- Glover K, Hulley H, Peskir G (2013) Three-dimensional Brownian motion and the golden ratio rule. *Ann Appl Probab* 23:895–922
- Guo X, Shepp LA (2001) Some optimal stopping problems with nontrivial boundaries for pricing exotic options. *J Appl Probab* 38(3):647–658
- Guo X, Zervos M (2010) π options. *Stoch Process Appl* 120(7):1033–1059
- Kyprianou AE, Ott C (2014) A capped optimal stopping problem for the maximum process. *Acta Appl Math* 129:147–174
- Liptser RS, Shiryaev AN (2001) *Statistics of random processes I*. (Second Edition, First Edition 1977). Springer, Berlin
- Ott C (2013) Optimal stopping problems for the maximum process with upper and lower caps. *Ann Appl Probab* 23:2327–2356
- Pedersen JL (2000) Discounted optimal stopping problems for the maximum process. *J Appl Probab* 37(4):972–983
- Peskir G (1998) Optimal stopping of the maximum process: The maximality principle. *Ann Probab* 26(4):1614–1640
- Peskir G (2007) A change-of-variable formula with local time on surfaces. *Séminaire de Probabilité XL*. *Lecture Notes in Mathematics* 1899. Springer, pp 69–96
- Peskir G (2012) Optimal detection of a hidden target: The median rule. *Stochastic Processes and their Applications* 122:2249–2263
- Peskir G (2014) Quickest detection of a hidden target and extremal surfaces. *Ann Appl Probab* 24(6):2340–2370
- Peskir G, Shiryaev AN (2006) *Optimal stopping and free-boundary problems*. Birkhäuser, Basel
- Revuz D, Yor M (1999) *Continuous Martingales and Brownian motion*. Springer, Berlin
- Rodosthenous N, Zervos M (2017) Watermark options. *Finance Stochast* 21(1):157–186
- Shepp LA, Shiryaev AN (1993) The Russian option: reduced regret. *Ann Appl Probab* 3(3):631–640
- Shepp LA, Shiryaev AN (1994) A new look at the pricing of Russian options. *Theory Probability and Applications* 39(1):103–119

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Pavel V. Gapeev¹ · Peter M. Kort^{2,3} · Maria N. Lavrutich⁴ · Jacco J. J. Thijssen⁵

Peter M. Kort
kort@uvt.nl

Maria N. Lavrutich
maria.lavrutich@ntnu.no

Jacco J. J. Thijssen
jacco.thijssen@york.ac.uk

- ¹ Department of Mathematics, London School of Economics, Houghton Street, London WC2A 2AE, United Kingdom
- ² Department of Econometrics and Operations Research, Tilburg University CentER, PO Box 90153, 5000 LE Tilburg, The Netherlands
- ³ Department of Economics, University of Antwerp, Prinsstraat 13, 2000 Antwerp 1, Belgium
- ⁴ Department of Industrial Economics and Technology Management, Norwegian University of Science and Technology, 7491 Trondheim, Norway
- ⁵ Management School, University of York, Freboys Lane, York YO10 5GD, United Kingdom