



Prevalence of adverse childhood experiences and their co-occurrence in a large population of adolescents: a Young HUNT 3 study

Rosalie Broekhof^{1,2} · Hans M. Nordahl^{2,3} · Sigrid Bjørnelv² · Sara G. Selvik^{1,2}

Received: 1 July 2021 / Accepted: 31 March 2022 / Published online: 22 April 2022
© The Author(s) 2022

Abstract

Purpose Studies of adverse childhood experiences (ACEs) undertaken at the time of adolescence in the general population are not common. The aim of this study was to determine the prevalence and co-occurrence of the individual ACEs and sub-types of ACEs in a large population of adolescents.

Methods Data were used from the Young Nord-Trøndelag Health (Young HUNT 3) study, a population-based study of young adolescents. ACEs were operational defined as sexual, physical and/or emotional abuse; physical and/or emotional neglect; and/or household dysfunction. Co-occurrence was measured as the accumulation of ACEs and as an overlap analysis.

Results Of the 8199 evaluable adolescents, 65.8% had experienced at least one ACE and 28% of those had experienced more than one ACE. Household dysfunction was the most prevalent ACE subtype. The biggest overlaps among the three ACE sub-types were seen in those reporting neglect or abuse.

Conclusion There was a high degree of overlap between the three ACE sub-types and the individual ACEs, indicating that ACEs should be assessed together as a whole rather than separately. This study provides an opportunity to assess ACEs and their co-occurrences in relation to outcomes later in life.

Keywords ACE · Childhood maltreatment · Abuse · Neglect · Prevalence

Background

The impact and severity of adverse childhood experiences (ACEs) were largely underestimated until research from the 1990s showed that ACEs had severe short- and long-term consequences for mental and physical health [1–4]. In addition to the association of ACEs with health issues, such as cardiovascular diseases and premature death [1, 4], there is also an association with mental disorders, such as mood, anxiety, schizophrenia, impulse control and substance use disorders [5–9]. Patients with histories of ACEs seek care more often, make high use of healthcare services, and have

poor quality of life [10]. In a report from 2016, the World Health Organization (WHO) stated that it is a challenging task to study the scope of childhood maltreatment [11] because the prevalence of childhood maltreatment varies for different countries and numbers are often underestimated [11]. Variations in reported numbers might be explained by variations in the type of ACE studied, variations in the characteristics of the sample studied, or variations in the definition of ACEs. For example, the prevalence of reported child sexual abuse varied from 7 to 36% for women and 3 to 29% for men in one report [12]. However, the WHO concluded that at least 12% of children were sexually abused in 2015 [13]. The same report stated that 25% of all adults reported physical abuse in childhood [13]. In 2009, Gilbert et al. reported a prevalence of 4 to 16% for physical abuse and around 10% for neglect of children in high-income countries [14].

The WHO defines childhood maltreatment as “the abuse and neglect of people under 18 years of age” [11]. “It includes all forms of physical and/or emotional ill-treatment, sexual abuse, neglect or negligent treatment or commercial or other exploitation, resulting in actual or potential harm

✉ Rosalie Broekhof
Rosalie.Broekhof@hnt.no

¹ Department of Psychiatry, Namsos Hospital, Nord-Trøndelag Hospital Trust, Namsos, Norway

² Department of Mental Health, Faculty of Medicine and Health Sciences, NTNU-Norwegian University of Science and Technology, Trondheim, Norway

³ St. Olavs University Hospital, Avd. Østmarka, Trondheim, Norway

to the child's health, survival, development or dignity in the context of a relationship of responsibility, trust or power" [11]. The adverse childhood experiences (ACEs) study by Felitti and Anda were the first study that defined ACEs in more detail [1, 15]. ACE scores were assessed using (combined) questionnaires that followed the WHO's definition of childhood maltreatment [11], and household dysfunction was added as another form of maltreatment. Later studies on ACEs used assessment methods from the original ACE study [16], which assessed five types of interpersonal violence/childhood maltreatment: (a) sexual abuse; (b) physical abuse; (c) emotional abuse (also referred to as psychological abuse); (d) neglect and (e) household dysfunction [1]. Neglect can be divided into (d-1) physical neglect and (d-2) emotional neglect (see definitions in Table 1).

The ACE study found that 52% of those assessed had experienced at least one individual ACE and 27% had experienced two or more ACEs [1]. This shows the complexity and interactive character of ACE. More recent research has focused on the specific areas of the different individual ACEs. For example, sexually abused girls had more learning problems, more symptoms of depression and anxiety and lower self-esteem than those who had not been sexually abused [17]. In another study, women with a history of sexual abuse had a higher risk of a very broad spectrum of psychopathology, such as major depression, anxiety, and alcohol and drug dependence [18]. Physical abuse in childhood has been associated with increased frequency of depression, anxiety, anger, physical symptoms and a number of medical diagnoses, decades after the abuse [19]. Emotional neglect has been related to depression, dysthymia and social phobia, with the likelihood of developing more than one lifetime affective disorder [20]. Neglect is the least studied type of trauma [14].

Again, other studies focused only on the sum of the individual ACEs [21–23]. It was found that individuals with

a high sum of individual ACEs were at increased risk for sexual risk taking behavior, mental ill health, and problematic alcohol use [21–23]. This risk was even stronger for problematic drug use and interpersonal and self-directed violence [21–23].

Previous studies on ACEs have limitations. First, they tend to focus mainly on only one individual ACE [17–19]. Without measuring a broad range of ACEs, any long-term influence might be wrongly attributed to an individual type of abuse, neglect or household dysfunction. Second, the questionnaires were often based on retrospective reporting [1, 3, 15, 24]. Retrospective reports are vulnerable to recall bias [25]. While the ACE study did focus on the whole range of ACEs, the study population had presented for preventive health evaluations and were given questionnaires relating to retrospectively remembered ACE [1, 15, 26]. Thirdly, the early literature shows a dose–response relationship between an increase in the number of ACEs and poorer adult mental health [3, 15, 21, 24]. With only the number of ACEs, the ACEs interaction are not exposed.

Aims

This study is part of a more comprehensive research project. The overall aim of the project is to assess ACEs in association with adult mental health. The aims of this study are:

- to assess the prevalence of ACEs and their co-occurrence measured at the time of adolescence in a large population sample. ACEs will be operationally defined in the Young Nord-Trøndelag Health 3 questionnaire.
- to add to the collective understanding of the prevalence and co-occurrence of ACEs, measured as the accumulation of ACEs and an overlap analysis of the three subtypes of ACE at the time of adolescence.

Table 1 Definitions of adverse childhood experience (ACE) types

Type of ACE	Definition
Sexual abuse	Defined by the occurrence of sexual touching or fondling, attempted intercourse, or actual intercourse by any adult or other person when the child did not want the act to occur or was too young to understand what was happening
Physical abuse	Defined by the occurrence of a parent or other adult who pushed, grabbed, shoved, slapped, or hit the child; or hit the child so hard it left marks or bruises, or caused an injury
Emotional abuse	Defined by the occurrence of a parent or other adult who swore at, insulted, or said hurtful things to the child; or threatened to hit or throw something at the child (but did not do it); or acted in any other way that made the child afraid he/she would be physically hurt or injured
Physical neglect	Defined as the child being left unsupervised when too young to care for themselves or going without needed clothing, school supplies, food, or medical treatment
Emotional neglect	Defined as the child not feeling part of a close-knit family or there being no one in the child's family of origin who made the child feel special, wanted the child to succeed, believed in the child, or provided strength and support
Household dysfunction	Defined by the presence of life stressors, such as parental divorce, drug use by parents, psychiatric problems of parents, financial problems

Methods

Study population

The county of Nord-Trøndelag in Norway has 135,000 inhabitants. All the inhabitants were invited to join a large population study: the HUNT study. The purpose of the HUNT study was to gather information on a wide variety of conditions and lifestyle factors. Data and samples were obtained in three waves of data gathering: 1984–1986, 1995–1997 and 2006–2008. A new wave of data gathering has recently been started.

The epidemiological study of adolescents in the region was named Young HUNT. The third wave of this study (the Young HUNT 3 study) was conducted between 2006 and 2008; 10,464 adolescents were invited to participate [27].

In the Young HUNT 3 study, during a school hour, the students completed a self-administered questionnaire that contained over 100 health- and lifestyle-related questions, including items on exposure to traumatic events, loneliness, psychological distress, and family cohesion. In total, 8199 adolescents (78%) filled in the questionnaires: 4129 girls (50.4%) and 4070 (49.6%) boys. The mean age was 15.91 (SD 0.03) years for the girls and 15.85 (SD 0.03) years for the boys.

The study was approved by the Norwegian Research Committee for Medical and Health Research Ethics (REK 2016/1877). Inclusion was based on written consent from participants aged 16 years and older and from parents for those under 16 years, in accordance with Norwegian law.

Definition of adverse childhood experiences (ACEs)

The operationalization of ACEs was done by consensus of a group of experts in the HUNT material. These experts followed the outlines given in the ACE study by Felitti et al. [1], using the corresponding items in the questionnaire in the Young HUNT 3 study. To form an operational definition of household dysfunction, Young HUNT 3 participants were linked to their parents, who had participated in the HUNT 3 study, through a national family register code from Statistics Norway (SSB). See Fig. 1.

Co-occurrence of the ACEs

Co-occurrence was measured in two different ways. First, as the accumulation of individual ACEs and the accumulation of ACEs within a subtype. Second, as the overlap between the three sub-types of ACE, visualized using a Venn diagram.

Statistics

The ACEs were operationalized by combining the dichotomized variables, in a similar fashion to that of Felitti et al. [1]. The prevalence of the individual operational ACEs and the total scores for abuse, neglect and household dysfunction were calculated to account for potential cumulative effects. We used SPSS version 23.0 software for the descriptive analyses.

Results

In total, 88% of the adolescents lived together with their parents and 68% had siblings. The distribution of living in rural or urban situations was 72% versus 28%.

Prevalence of ACEs

Figure 2 gives an overview of the operational definitions of the ACEs.

Of the entire sample ($N=8199$), household dysfunction had the highest percentage of positive responses ($N=5048$, 61.6%). Of household dysfunction, parental divorce was most prevalent ($N=2421$, 29.5% of the entire sample), followed by having witnessed violence ($N=1810$, 22.1%), parental mental problems ($N=1299$, 18.8%), economic difficulties ($N=1271$, 15.5%), and parental substance abuse ($N=618$, 7.5%). More parental substance abuse and parental divorce were seen for girls and more witnessed violence for boys. Abuse in some form was reported by 1496 (18.2%) adolescents. Of abuse, physical abuse was most prevalent ($N=1177$, 14.7%), followed by sexual abuse ($N=430$, 5.2%) and emotional abuse ($N=230$, 2.8%). Girls reported significantly higher levels of emotional and sexual abuse and boys, significantly higher physical abuse. Neglect was reported by 581 (7.1%) adolescents: 126 reported emotional neglect and 471 reported physical neglect. Girls reported significantly more emotional neglect than boys.

Co-occurrence of the ACEs

The Venn diagram in Fig. 3 shows the overlap of reports of the three sub-types of ACE. Of the 8199 adolescents who completed the questionnaire, 65.8% had experienced at least one subtype of ACE and 28% of these had experienced two or more sub-types of ACE (Fig. 3). Of those who had experienced at least one subtype of ACE ($N=5398$), 4% had experienced all three sub-types. The biggest overlaps between the sub-types of ACE were seen for neglect and abuse. For example, of the adolescents who experienced neglect, 104 experienced only neglect, while 477 also experienced another of the subtype of ACEs.

Operational definitions of ACE			
Type of ACE	Young HUNT 3 study questions	Type of ACE is present	Type of ACE is not present
Sexual Abuse	Have any of the following things happened to you? ✓ been put in sexually uncomfortable/abusive situations by an adult OR ✓ been put in sexually uncomfortable/abusive situations by someone about your age.	Yes, last year/in my life	No
Physical Abuse	Have any of the following things happened to you? ✓ Have you been violently hurt? OR ✓ Have you been threatened or physically harassed by fellow students?	Yes, last year/in my life	No
Emotional Abuse	✓ Has there been much arguing in your family? AND ✓ I feel comfortable with my family AND ✓ I certainly feel useless at times	Yes, past month/previously (totally) disagree (totally) agree	No Average, (totally) agree Average, (totally) disagree
Emotional Neglect	✓ How good is your relationship with you immediate family? AND ✓ In my family we support each other AND ✓ Do you often feel lonely?	Not so good, bad (totally) disagree (very) often	(Very) good Average, (totally) agree Sometimes, seldom, never
Physical Neglect	✓ Do you use alcohol/have you been intoxicated? AND ✓ Outside school hours: How many hours a week do you play sports or exercise in your free time so much that you get out of breath or sweat? AND ✓ How often do you usually eat meals?	Yes...times <½ hour ≤ 1 meal a day	No never >½ hour > 2 meals a day
Parental substance abuse	✓ Alcohol consumption of parent(s), 2 CAGE questionnaires answered positively*	Yes	No
Parental mental illness	✓ Have you had or do you have any of the following: mental health problems you sought help for?*	Yes	No
Divorced parents	✓ Are your parents separated or divorced, or have they lived separately for more than one year?	Yes	No
Economic difficulties	✓ How well off do you think your family is compared to most others?	Worse financial situation	About the same as others/ better financial situation
Witnessed violence	✓ Have you seen others being violently hurt	Yes, last year/ in my life	No

Fig. 1 Overview of the questions from the Young Helse Nord-Trøndelag (HUNT) study questionnaire to obtain operational definitions of adverse childhood experiences (ACEs) * Questions for parents of the

Young HUNT 3 study adolescents who had participated in the HUNT 3 study

Table 2 shows the accumulation scores for the sub-types of ACEs as well as the accumulation scores for the individual ACEs. Of those who had experienced abuse ($N=1496$), 283 (19%) had experienced two sub-types and 29 (1.9%) had experienced all three. Sixteen (2.8%) adolescents experienced more than one subtype of neglect.

For household dysfunction, 3219 (63.8%) adolescents had experienced one subtype, 1350 (26.7%) had experienced two sub-types, and 479 (9.5%) had experienced three or more sub-types.

The average accumulation score for ACE, calculated from the total of all individual ACEs, was 1.8. Of those who had experienced ACE ($N=5398$), around 50% of the adolescents had experienced at least one type of individual ACE, almost 30% two, and 20% three or more individual ACEs.

Discussion

Our results show that in a non-clinical population study of adolescents aged 13–19 years, the prevalence of ACEs was high, and the ACEs often occurred together. In our sample, 65.8% of the participants ($N=5398$) had experienced at least one subtype of ACE, and 28% (1514) of those had experienced more than one subtype. Household dysfunction was the most prevalent subtype of ACE. Abuse and neglect had the biggest overlaps with the other sub-types of ACEs.

This high prevalence of ACEs is in line with earlier retrospective studies in adults, which found that over 50% had been exposed to at least one individual ACE





Fig.2 Operational definitions of adverse childhood experiences (ACEs). The prevalence is given as the number of adolescents responding positively (*N*) and also as a percentage of the total number of evaluable adolescents in the Young HUNT 3 study (*N*=8199; 10,464 were invited) The significant higher ACEs for girls are emo-

tional and sexual abuse, emotional neglect, parental substance use, and parental divorce. The significant higher ACEs for boys are physical abuse and witnessing violence. *m* mean, *SD* standard deviation, *y* years. * Questions for parents of the Young HUNT 3 study adolescents, who had participated in the HUNT 3 study

Fig.3 Venn diagram showing overlap between the three sub-types of adverse childhood experience (ACE). Results are given as the number of adolescents responding positively and as percentages of the number of adolescents in the Young HUNT 3 study who had experienced at least one of the three sub-types of ACE, *N*= 5398



and around 30% had been exposed to two or more individual ACEs [1, 3, 24]. A study by McLaughlin et al. also assessed 13- to 17-year-olds [7]. They defined childhood adversities as parental loss, maltreatment, parental maladjustment and economic adversity, and found that 58.3%

of the participants had experienced at least one individual childhood adversity and 59.7% of these had experienced multiple types.

The exposure rates for abuse and neglect in our study are in the lower range of those in earlier studies of adults

Table 2 Overview of the accumulation scores for the sub-types of ACEs and the accumulation scores for the individual ACEs

(Sub)types	1	2	3	4	5	≥ 6
Abuse <i>N</i> = 1496	1184 (79.1%)	283 (19.0%)	29 (1.9%)			
Neglect <i>N</i> = 581	565 (97.2%)	16 (2.8%)				
Household dysfunction <i>N</i> = 5048	3219 (63.8%)	1350 (26.7%)	420 (8.3%)	55 (1.1%)	4 (0.1%)	
Accumulation of individual ACEs SE = 1,8 <i>N</i> = 5398	2771 (51.3%)	1460 (27.0%)	705 (13.1%)	314 (5.8%)	103 (1.9%)	49 (0.9%)

Percentages were calculated from the total number of adolescents (*N*) who had experienced that type of ACE

[1, 14, 15, 28, 29]. As mentioned, most of those studies of ACEs were carried out retrospectively [1, 3, 15, 24, 26] and included other notions/reflections on the questions than those for the children aged 13–19 years in our study.

The questionnaire in our study sourced data that covered all but two of the household dysfunction events defined in the ACE study [1]. The adolescents in the Young HUNT study were not asked if their mothers were treated violently or about incarcerated household members. The Young HUNT questionnaire did enquire whether the adolescent had ‘witnessed violence’, including outside the family. We also expanded the household dysfunction individual ACEs by adding parental divorce and economic difficulties. Subsequently, we found a higher prevalence rate for household dysfunction in general than found in the ACE study [1] but in line with McLaughlin’s study [7] and the expanded ACE study [15]. In contrast, the prevalence of witnessed violence in Philadelphia, retrospectively assessed in the same way as in our study, was twice as high [29].

That ACEs interact with other ACEs is not a new concept [1, 21]. The “ACE score”, the sum of ACEs, has received comments on its simplicity and research is encouraged to focus also on the concept of timing, chronicity and duration, especially in relations to for example relative risks, where ACE scores were associated to health outcomes. [30–32]. Our aim was to investigate co-occurrence, not to make any claims about the severity of any given ACE itself. In this study, however, we not only assessed individual ACE accumulation scores, we also assessed the ACE accumulation scores for each of the sub-types of ACE separately and provided an overview of the overlap relationships. Our study showed overlap between both the individual ACEs and the three sub-types of ACE, with especially high overlap for neglect and abuse.

Strengths and limitations

An important strength of this study is that the Young HUNT 3 study was a population study, meaning that the sample was big enough to investigate and compare the individual ACEs as well as ACEs as a whole. Earlier research has mostly focused on individual ACEs [17–19]. In contrast to

earlier retrospective research, this study took into account the overlap and accumulation scores of ACEs at the age of 13–19 years [5, 6, 15, 20]. It is intended that prospective research on ACEs in association with other mental health problems will be pursued in this population. The Young HUNT 3 study will then be the largest prospective, population-based, cross-sectional Scandinavian study focusing on ACEs [33]. Because of the used population, the results of this study can be generalized in western world countries as Norway.

An important limitation of the Young HUNT 3 study is that the questionnaires were completed at school, with the inherent difficulties associated with attendance at school which can affect ACE scores, as adolescents who have experienced ACE’s are often more likely to be absent from school. However, 78% of all eligible school children participated in our study.

The Young HUNT 3 study is a population-based study with a broad approach to mental and physical health that has considered questions on a wide scale of topics. ACEs were therefore not defined with the childhood trauma questionnaire but were operationalized afterward. In this context, physical neglect should be mentioned. Neglect was the most challenging ACE to operationalize. Focus with this operationalization was to capture the definition of neglect, the adolescents not being met in their essential needs, as a whole.

However, it should be borne in mind that previous studies using HUNT data have successfully given operational definitions to depression, anxiety and interpersonal problems [34, 35].

Conclusion

This study shows that the prevalence of ACEs is high with serious interaction, already in adolescence. Especially neglect and abuse show substantial overlap. If ACEs are assessed separately, there is a possibility of obtaining the wrong impression, and feedback to clinical practice about the effects of that ACE could be misleading. This study provides an opportunity to assess those with ACEs, the

co-occurrence of ACEs and make correlations with their outcomes later in life.

Funding Open access funding provided by NTNU Norwegian University of Science and Technology (incl St. Olavs Hospital - Trondheim University Hospital). No funding was received to assist with the preparation of this manuscript.

Data availability The dataset reported in the current article is not publicly available. All data from HUNT are stored at the HUNT research center, with strict rules for access. All data that the researcher uses in their studies are anonymized. Researchers have no access to personal information. It is available on request by qualified scientists. Requests require a concept paper describing the purpose of data access, ethical approval at the applicant's institution, and provision for secure data access.

Declarations

Conflict of interest The authors have no relevant financial or non-financial interests to disclose.

Ethics approval This study was performed in line with the principles of the Declaration of Helsinki. The study was approved by the Norwegian Research Committee for Medical and Health Research Ethics (REK 2016/1877). Inclusion was based on written consent from participants aged 16 years and older and from parents for those under 16 years, in accordance with Norwegian law.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Felitti VJ, Anda RF, Nordenberg D, Williamson DF, Spitz AM, Edwards V et al (1998) Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *Am J Prev Med* 14(4):245–258
2. Collishaw S, Pickles A, Messer J, Rutter M, Shearer C, Maughan B (2007) Resilience to adult psychopathology following childhood maltreatment: evidence from a community sample. *Child Abuse Negl* 31(3):211–229
3. Kessler RC, McLaughlin KA, Green JG, Gruber MJ, Sampson NA, Zaslavsky AM et al (2010) Childhood adversities and adult psychopathology in the WHO World Mental Health Surveys. *Br J Psychiatry* 197(5):378–385
4. Hendrickson CM, Neylan TC, Na B, Regan M, Zhang Q, Cohen BE (2013) Lifetime trauma exposure and prospective cardiovascular events and all-cause mortality: findings from the Heart and Soul Study. *Psychosom Med* 75(9):849–855
5. Infurna MR, Reichl C, Parzer P, Schimmenti A, Bifulco A, Kaess M (2016) Associations between depression and specific childhood experiences of abuse and neglect: a meta-analysis. *J Affect Disord* 190:47–55
6. Hovens JG, Giltay EJ, Spinhoven P, van Hemert AM, Penninx BW (2015) Impact of childhood life events and childhood trauma on the onset and recurrence of depressive and anxiety disorders. *J Clin Psychiatry* 76(7):931–938
7. McLaughlin KA, Greif Green J, Gruber MJ, Sampson NA, Zaslavsky AM, Kessler RC (2012) Childhood adversities and first onset of psychiatric disorders in a national sample of US adolescents. *Arch Gen Psychiatry* 69(11):1151–1160
8. Pietrek C, Elbert T, Weierstall R, Muller O, Rockstroh B (2013) Childhood adversities in relation to psychiatric disorders. *Psychiatry Res* 206(1):103–110
9. Carr CP, Martins CM, Stingel AM, Lemgruber VB, Juruena MF (2013) The role of early life stress in adult psychiatric disorders: a systematic review according to childhood trauma subtypes. *J Nerv Ment Dis* 201(12):1007–1020
10. Chartier MJ, Walker JR, Naimark B (2010) Separate and cumulative effects of adverse childhood experiences in predicting adult health and health care utilization. *Child Abuse Negl* 34(6):454–464
11. World Health Organization. (2016) Child Maltreatment. <http://www.who.int/news-room/fact-sheets/detail/child-maltreatment>. Accessed 15 Dec 2021
12. Finkelhor D (1994) The international epidemiology of child sexual abuse. *Child Abuse Negl* 18(5):409–417
13. World Health Organization. (2016) Violence and injury prevention. https://www.who.int/violence_injury_prevention/violence/child/en/. Accessed 15 Dec 2021
14. Gilbert R, Widom CS, Browne K, Fergusson D, Webb E, Janson S (2009) Burden and consequences of child maltreatment in high-income countries. *Lancet (London, England)* 373(9657):68–81
15. Anda RF, Felitti VJ, Bremner JD, Walker JD, Whitfield C, Perry BD et al (2006) The enduring effects of abuse and related adverse experiences in childhood. A convergence of evidence from neurobiology and epidemiology. *Eur Arch Psychiatry Clin Neurosci* 256(3):174–186
16. Bethell CD, Carle A, Hudziak J, Gombojav N, Powers K, Wade R et al (2017) Methods to assess adverse childhood experiences of children and families: toward approaches to promote child well-being in policy and practice. *Acad Pediatr* 17(7s):S51–s69
17. Trickett PK, Noll JG, Putnam FW (2011) The impact of sexual abuse on female development: lessons from a multigenerational, longitudinal research study. *Dev Psychopathol* 23(2):453–476
18. Kendler KS, Bulik CM, Silberg J, Hettema JM, Myers J, Prescott CA (2000) Childhood sexual abuse and adult psychiatric and substance use disorders in women: an epidemiological and cotwin control analysis. *Arch Gen Psychiatry* 57(10):953–959
19. Springer KW, Sheridan J, Kuo D, Carnes M (2007) Long-term physical and mental health consequences of childhood physical abuse: results from a large population-based sample of men and women. *Child Abuse Negl* 31(5):517–530
20. Spinhoven P, Elzinga BM, Hovens JG, Roelofs K, Zitman FG, van Oppen P et al (2010) The specificity of childhood adversities and negative life events across the life span to anxiety and depressive disorders. *J Affect Disord* 126(1–2):103–112
21. Hughes K, Bellis MA, Hardcastle KA, Sethi D, Butchart A, Mikton C et al (2017) The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *Lancet Public Health* 2(8):e356–e366

22. Blum RW, Li M, Naranjo-Rivera G (2019) Measuring adverse child experiences among young adolescents globally: relationships with depressive symptoms and violence perpetration. *J Adolesc Health* 65(1):86–93
23. Ramaiya A, Choiriyah I, Heise L, Pulerwitz J, Blum RW, Levto R et al (2021) Understanding the relationship between adverse childhood experiences, peer-violence perpetration, and gender norms among very young adolescents in Indonesia: a cross-sectional study. *J Adolesc Health* 69(1s):S56–s63
24. Afifi TO, Mather A, Boman J, Fleisher W, Enns MW, Macmillan H et al (2011) Childhood adversity and personality disorders: results from a nationally representative population-based study. *J Psychiatr Res* 45(6):814–822
25. Frissa S, Hatch SL, Fear NT, Dorrington S, Goodwin L, Hotopf M (2016) Challenges in the retrospective assessment of trauma: comparing a checklist approach to a single item trauma experience screening question. *BMC Psychiatry* 16:20
26. Anda RF, Butchart A, Felitti VJ, Brown DW (2010) Building a framework for global surveillance of the public health implications of adverse childhood experiences. *Am J Prev Med* 39(1):93–98
27. Holmen TL, Bratberg G, Krokstad S, Langhammer A, Hveem K, Midthjell K et al (2014) Cohort profile of the Young-HUNT Study, Norway: a population-based study of adolescents. *Int J Epidemiol* 43(2):536–544
28. Thompson MP, Kingree JB, Lamis D (2019) Associations of adverse childhood experiences and suicidal behaviors in adulthood in a US nationally representative sample. *Child Care Health Dev* 45(1):121–128
29. Lê-Scherban F, Wang X, Boyle-Steed KH, Pachter LM (2018) Intergenerational associations of parent adverse childhood experiences and child health outcomes. *Pediatrics* 141(6):e20174274
30. Lacey RE, Minnis H (2020) Practitioner review: twenty years of research with adverse childhood experience scores—advantages, disadvantages and applications to practice. *J Child Psychol Psychiatry* 61(2):116–130
31. DeLisi M, Drury AJ, Elbert MJ (2021) Frequency, chronicity, and severity: New specification of adverse childhood experiences among federal sexual offenders. *Forensic Sci Int Mind Law* 2:100051
32. Briggs EC, Amaya-Jackson L, Putnam KT, Putnam FW (2021) All adverse childhood experiences are not equal: the contribution of synergy to adverse childhood experience scores. *Am Psychol* 76(2):243–252
33. Jahn A, Rysgaard TK, Andersen JH, Winding TN (2021) Adverse childhood experiences and future self-rated health: a prospective cohort study. *BMC Public Health* 21(1):895
34. Stensland SO, Thoresen S, Wentzel-Larsen T, Zwart J-A, Dyb G (2014) Recurrent headache and interpersonal violence in adolescence: the roles of psychological distress, loneliness and family cohesion: the HUNT study. *J Headache Pain* 15(1):35
35. Solhaug HI, Romuld EB, Romild U, Stordal E (2012) Increased prevalence of depression in cohorts of the elderly: an 11-year follow-up in the general population—the HUNT study. *Int Psychogeriatr* 24(1):151–158