

- Midshaft med. – lat. (16mm)

(probable) L ulna (pathology seems present, smaller than the other ulnas)

- Midshaft med. – lat. (13mm)

2 mandibles with different wear patterns

Cranium A (in bag)

- Part of occipital and part of temporal (L) (based on thickness, weight, colour and preservation these most likely belong to the mandible in the box)
- Mandible
 - o L: M3, M1, PM2, C, I2 (in occlusion) PM1, M2 (present but not in occlusion)
 - o R: M1, PM2, PM1, C (in occlusion) I2, M3, M2 (present but not in occlusion)
 - o M1(L) eroded all the way to the root, probable infection
 - o Extreme calculus on anterior teeth
- Maxilla (3 fragments)
 - o L: M2, M1, C (in occlusion) PM1, PM2 (present but not in occlusion)
 - o R: M2, PM2, PM1, C (in occlusion) M3, M1, I2 (present but not in occlusion)
 - o M2(L) cavity erosion between root and crown towards M3
- Sexing
 - o Mandible
 - Mandibular ramus = 2
 - Mental eminence = 3
 - Broad jawline
 - o Temporal
 - Mastoid process = 1
- Ageing
 - o Brothwell = 25-35
 - o Miles = 25-45

	Tooth		Presence	Wear		Notes
Maxillary Right	1	M ³	1	3	5	
				3	5	
	2	M ²	2	5	7	
				5	7	
	3	M ¹	1	9	9	
				9	9	
	4	p ²	2	5	5	
	5	p ¹	2	4	4	
	6	C	2	3	3	
	7	I ²	1	4	4	
	8	I ¹				